

Nitration Of Phenols Under Mild And Heterogeneous Conditions

Mohammad Ali Zolfigol*, Ezat Ghaemi, Elahe Madrakian

Department of Chemistry, College of Science, University of Bu-Ali Sina, Zip code 65174, P.O. Box No. 4135, Hamadan, I.R. Iran, Fax: (+98) 811 8272404.

*Author to whom correspondence should be addressed; e-mail: Zolfi@basu.ac.ir

Received: 14 November 2000; in revised form 24 May 2001 / Accepted: 1 June 2001 / Published: 30 June 2001

Abstract: Nitrophenols can be obtained in moderate to high yields using a combination of $\text{Mg}(\text{HSO}_4)_2$ or $\text{NaHSO}_4 \cdot \text{H}_2\text{O}$, NaNO_3 and wet SiO_2 in dichloromethane at room temperature.

Keywords: Nitration; Phenols; Heterogeneous conditions; Inorganic acidic salts

Introduction


The nitration of aromatic compounds may be achieved with many nitrating reagents and is a very useful method in organic synthesis [1]. Also, nitro compounds find use in many industrial applications [2-4]. Nitration of phenol as a special case has been studied using various nitrating agents under different conditions [4-22]. Recently, in this connection we have reported the use and reaction mechanisms of some hydrated metal nitrates (containing covalent nitrato groups) and their dinitrogen tetroxide complex analogues for the nitration of phenols under various conditions [23]. Our goal, in undertaking this line of work, is two-fold: (a) to overcome the limitations and drawbacks of the reported methods such as tedious work-up [9-11], strongly acidic media ($\text{H}_0 \sim -8$) [4b], oxidation ability of the reagents and safety problems (storage, handling, use and also the

presence of poisonous transition metal cations such as Cr^{+3} , Hg^{+2} , Cu^{+2} , etc. within the molecular structures of the reagents) [24, 25]; (b) moreover, constraining a reagent to the surface of a solid usually allows the use of milder conditions and increases its reactivity [26]. Very recently, we and others have demonstrated that heterogeneous reagent systems have many advantages such as simple experimental procedures, mild reaction conditions and minimization of chemical wastes as compared to their liquid phase counterparts [27]. Consequently, we decided to seek a heterogeneous system for the nitration of phenol, and we have investigated a number of different reaction conditions based upon the *in situ* generation of HNO_3 by *solid inorganic acidic salts* [$\text{NaHSO}_4 \cdot \text{H}_2\text{O}$, $\text{Mg}(\text{HSO}_4)_2$, $\text{p}K_a \sim 2$] with sodium nitrate. We report here a one-pot heterogeneous procedure for nitration of phenols.

Results and Discussion


Phenol (**1**) and different types of 4-substituted phenols (**4**) were subjected to a nitration reaction in the presence of inorganic acidic salts [e.g. $\text{Mg}(\text{HSO}_4)_2$ (**I**) or $\text{NaHSO}_4 \cdot \text{H}_2\text{O}$ (**II**)], NaNO_3 (**III**), and wet SiO_2 (50% w/w) in dichloromethane (Schemes 1 and 2). The nitration reactions were performed under mild and heterogeneous conditions at room temperature to give the products in moderate to excellent yields (Table 1).

Scheme 1



The nitration reactions can be readily carried out by placing the nitrating agents, phenols (**1** or **4**) and the solvent used in a reaction vessel and efficiently stirring the resultant heterogeneous mixture at room temperature. The resulting mononitrophenols can be obtained by simple filtration and evaporation of the solvent. This alternative method thus provides nitrated phenols directly, in short reaction times and good yields.

Scheme 2




4	X	4	X	4	X
a	F	e	Ph	i	COOH
b	Cl	f	CH ₃	j	CH ₂ Ph
c	Br	g	OCH ₃	k	NHOAc
d	CN	h	COCH ₃	l	4-HOC ₆ H ₄ -

In fact, a combination of sodium nitrate and inorganic acidic salts (**I** or **II**) can act as a *solid HNO₃ equivalent* which can be readily handled and used for different purposes in the presence of wet SiO₂ [28].

A competitive reaction between phenols and anisole was also performed. It was observed that exclusive phenol nitration proceeded, whereas anisole is remained intact in the reaction mixtures after 24 hours (Scheme 3).

Scheme 3



This method is also very mild as indicated by the fact that the hydrolysis product of 4-cyano-phenol was not observed (Scheme 4, Table 1, entries 9 and 10). Selective mononitration of 4,4'-dihydroxydiphenyl (**4I**) was also achieved by controlling the stoichiometry of the reagents (Table 1, entries 25 and 26).

Scheme 4




Table 1. Substrates and Products

Entry	Substrate	Products ^a	<u>Reag. /Subst. (mmol)^b</u>			Time (Min)	Yields ^c %
			I	II	III		
1	1	2[7]	1	--	1	30	26
		3[7]					36
2	1	2[7]	--	1	1	30	20
		3[7]					50
3	4a	5a[4] ^c	1	--	1	90	54
4	4a	5a[4c]	--	1	1	90	60
5	4b	5b[11, 18]	1	--	1	90	70
6	4b	5b[11, 18]	--	1	1	90	72
7	4c	5c[11, 18]	1	--	1	130	78
8	4c	5c[11, 18]	--	1	1	130	70
9	4d	5d[11, 18]	1	--	1	180	88
10	4d	5d[11, 18]	--	1	1	180	92
11	4e	5e[11, 18]	1	--	1	180	90
12	4e	5e[11, 18]	--	1	1	180	92
13	4f	5f[11, 18]	1	--	1	180	72
14	4f	5f[11, 18]	--	1	1	180	78
15	4g	5g[18]	1	--	1	45	84
16	4g	5g[18]	--	1	1	45	79
17	4h	5h[11, 18]	2	--	2	120	74
18	4h	5h[11, 18]	--	2	2	120	79
19	4i	5i[18]	2	--	2	90	80
20	4i	5i[18]	--	2	2	90	83
21	4j	5j[18]	2	--	2	180	91
22	4j	5j[18]	--	2	2	180	91
23	4k	5k[22]	--	1	1	180	55
24	4k	5k[22]	1	--	1	180	74
25	4l	5l[21]	--	1	1	90	56
26	4l	5l[21]	1	--	1	90	63
27	1	2, 3	--	--	1	No Reaction ^d	

^a All of the isolated products are known compounds and their spectra and physical data have been reported in the literature; ^b Wet SiO₂ : substrate (1) (0.2 g : 1 mmol); ^c Isolated pure Yields; ^d Nitration did not occur in the absence of inorganic acidic salts.

Although the nitration reaction also occurs in the absence of wet SiO₂, the reaction times are very long and the reactions only go to completion after several days. Therefore, we think that the

wet SiO₂ acts as a reaction medium providing a heterogeneous effective surface area for *in situ* generation of HNO₃ in low concentrations. It also makes work-up easy. On the other hand, nitration did not occur in the absence of the inorganic acidic salts (Table 1, entry 27).

Conclusions

In conclusion, the low cost and the availability of the reagents, easy and clean work-up, and high yields make this an attractive method for Organic Synthesis. This simple procedure is highly selective and contamination by oxidation side-products is avoided. Moreover, a key new feature is the heterogeneous nature of the reaction, which could be worthwhile in an industrial setting [26].

Acknowledgments

The authors gratefully acknowledge partial support of this work by the Research Affairs Office of Bu-Ali Sina University, Hamadan, I.R Iran.

Experimental section

General

Chemicals were purchased from the Fluka, Merck and Aldrich chemical companies. Melting points were taken on a Gallenkamp melting point apparatus and are uncorrected. Proton and carbon nuclear magnetic resonance spectra were recorded on a JEOL NMR-Spectrometer FX 90Q. IR spectra were recorded on a Shimadzu 435 IR spectrophotometer. Thin layer chromatography (TLC) on commercial aluminium-backed plates of silica gel 60 F₂₅₄ was used to monitor the progress of the reactions. Yields refer to isolated pure products. The nitration products were characterized by comparison of their spectral (IR, ¹H-NMR, ¹³C-NMR), TLC and physical data with authentic samples.

Typical Procedure for Mononitration of Phenol (1) with Mg(HSO₄)₂ (I), NaNO₃ (III) and wet SiO₂

A suspension of compound **1** (1.88 g, 0.02 mol), **I** (4.40 g, 0.02 mol), **III** (1.7 g, 0.02 mol) and wet SiO₂ (50% w/w, 4 g) in CH₂Cl₂ (20 mL) was stirred magnetically at room temperature. The reaction was complete after 30 min. and the reaction mixture was then filtered. The residue was washed with CH₂Cl₂ (2x10 mL). Anhydrous Na₂SO₄ (10 g) was added to the combined filtrate and washings. After 15 minutes the mixture was filtered. The solvent was removed by distillation using a water bath (35–40 °C). The residue is a mixture of 2- and 4-nitrophenol, which may be purified by addition of n-pentane as the latter is insoluble in this solvent. Yield of 4-nitrophenol (**2**): 1.44 g, 26%, mp 112–113 °C {lit [7] mp 114 °C}. The n-pentane was evaporated on a water bath (35–40 °C) to give the 2-nitrophenol (**3**): 2 g, 36%, mp 43–45 °C {lit. [7] mp 44 °C}.

Typical Procedure for Mononitration of 4-Cyanophenol (4d) with $Mg(HSO_4)_2$ (I), $NaNO_3$ (III) and wet SiO_2

A suspension of compound **4d** (0.238 g, 2 mmol), **I** (0.44 g 2 mmol), wet SiO_2 (50% w/w, 0.4 g) and **III** (0.17 g, 2 mmol) in dichloromethane (4 mL) was stirred at room temperature for 3 hours (the progress of the reaction was monitored by TLC) and then filtered. Anhydrous Na_2SO_4 (5 g) was added to the filtrate. After 15 minutes the resulting mixture was also filtered. Dichloromethane was removed by simple distillation using a water bath (35–40 °C). The yield was 0.288 g, (88%) of crystalline pale yellow solid (**5d**), mp 143–144 °C, {Lit. [4c] mp 145 °C}. 1H -NMR (FT-90 MHz, $CDCl_3$, TMS): δ 7.34 (d, 1H), 7.77 (dd, 1H), 8.48 (d, 1H), 10.87 (b, 1H). 1H -NMR spectra were identical with reference spectra [4c].

References and Notes

1. Gu, S. Jing, H. Wu, J. and Liang, y. *Synth. Commun.* **1997**, 27, 2793.
2. Smith, K. Musson, A. and DeBoos, G. A. *Chem. Commun.* **1996**, 469.
3. Waller, F. G. Barrett, A. G. M. Braddock, D. C. and Ramprasad, D. *Chem. Commun.*, **1997**, 613. and references cited therein.
4. (a) Delaude, L. Laszlo, P. and Smith, K. *Acc. Chem. Res.*, **1993**, 26, 607; (b) Laszlo, P. *Acc. Chem. Res.*, **1986**, 19, 121; (c) Cornelis, A.; Laszlo, P. and Pennetreau, P. *Bull. Soc. Chim. Belg.* **1984**, 93, 961.
5. Zeegers, P. J. *J. Chem. Ed.*, **1993**, 70, 1036.
6. Pervas, H. Onysiuka, S. O. Rees, L. Rooney, J. R. and Sukling, G. J. *Tetrahedron*, **1988**, 44, 4555.
7. Furniss, B. S., Hannaford, A. J., Smith, P. W. G. and Tatchell, A. R. " *Vogel's Textbook of Practical Organic Chemistry*" 4th Ed, Longman: London and New York; 1986.
8. Bruice, T. C. Gregor M. G. and Walters, S. L. *J. Am. Chem. Soc.* **1986**, 90, 1612.
9. Crivello, J. V. *J. Org. Chem.* **1981**, 46, 3056.
10. Oueartani, M. Girard, P. and Kagan, H. B. *Tetrahedron Lett.* **1982**, 23, 4315.
11. Poirier, J. M. and Vottero, C. *Tetrahedron* **1989**, 45, 1415.
12. Thompson, M. J. and Zeegers, P. J. *Tetrahedron* **1989**, 45, 191.
13. Tapia, R.; Torres, G. and Valderrama, J. A. *Synth. Commun.*, **1986**, 16, 681.
14. Gaude, D. Goallar, R. L. and Pierre, J. L. *Synth. Commun.*, **1986**, 16, 63.
15. Gigante, B. Prazeres, A. O. and Marcelo-Curto, M. J. *J. Org. Chem.* **1995**, 60, 3445.
16. Rodrigues, J. A. R. Filho, A. P. O. and Moran, P. J. S. *Tetrahedron* **1999**, 55, 6733.
17. Suboch, G. A. and Belyaev, E. Y., *Russ. Org. Chem.*, **1998**, 34, 288.
18. Nonoyama, N. Chiba, K. Hisatome, K. Suzuki, H. and Shintani, F. *Tetrahedron Lett.*, **1999**, 40, 6923.
19. Lehnig, M. *Tetrahedron Lett.*, **1999**, 40, 2299.

20. "Dictionary of Organic Compounds" 3th Ed, Eyre & Spottiswoode: London, **1965**, 2, 620.
21. Raiford, L. C. and Colbert, J. C., *J. Am. Chem. Soc.*, **1925**, 47, 1454.
22. Hancock, C. K. and Clagve, A. D., *J. Am. Chem. Soc.*, **1964**, 86, 4942.
23. (a) Zolfigol, M. A. Iranpoor, N. and Firouzabadi, H. *Orient. J. Chem.* **1998**, 14, 369; (b) Firouzabadi, H. Iranpoor, N. and Zolfigol, M. A. *Iran. J. Chem. & Chem. Eng.* **1997**, 16, 48; (c) Firouzabadi, H. Iranpoor, N. and Zolfigol, M. A. *Synth. Commun.* **1997**, 27, 3301. (d) Iranpoor, N. Firouzabadi, H. and Zolfigol, M. A. *Synth. Commun.* **1998**, 28, 2773.
24. (a) Firouzabadi, H. Iranpoor, N. and Zolfigol, M. A. *Synth. Commun.*, **1998**, 28, 377; (b) Firouzabadi, H. Iranpoor, N. and Zolfigol, M. A. *Synth. Commun.* **1998**, 28, 1179; (c) Iranpoor, N. Firouzabadi, H. and Zolfigol, M. A. *Synth. Commun.* **1998**, 28, 367; (d) Firouzabadi, H. Iranpoor, N. and Zolfigol, M. A. *Bull. Chem. Soc. Jpn.* **1998**, 71, 2169; (e) Iranpoor, N. Firouzabadi, H. and Zolfigol, M. A. *Bull. Chem. Soc. Jpn.* **1998**, 71, 905.
25. (a) Laszlo, P. and Cornelis, A. *Aldrichimica Acta*, **1988**, 21, 97; (b) Cornelis, A. and Laszlo, P. *Synthesis* **1985**, 909; (c) Laszlo, P. and Cornelis, A. *Synlett* **1994**, 155.
26. Riego, J. M. Sedin, Z. Zaldivar, J. M. Marziano, N. C. and Tortato, C. *Tetrahedron Lett.* **1996**, 37, 513.
27. (a) Zolfigol, M. A.; Kiany-Borazjani, M.; Sadeghi, M. M.; Mohammadpoor-Baltork, I.; Memarian, H. R. *Synth. Commun.*, **2000**, 30, 551; (b) Zolfigol, M. A.; Kiany-Borazjani, M.; Sadeghi, M. M.; Memarian, H. R.; Mohammadpoor-Baltork, I. *Synth. Commun.*, **2000**, 30, 2945; (c) Zolfigol, M. A. *Synth. Commun.*, **1999**, 29, 905; (d) Zolfigol, M. A.; Nematollahi, D.; Mallakpour, S. E. *Synth. Commun.*, **1999**, 29, 2277; (e) Zolfigol, M. A.; Mallakpour, S. E. *Synth. Commun.*, **1999**, 29, 4061; (f) Zolfigol, M. A. *Synth. Commun.*, **2000**, 30, 1593; (g) Zolfigol, M. A.; Ghaemi, E.; Madrakian, E.; Kiany-Borazjani, M. *Synth. Commun.*, **2000**, 30, 2057; (h) Zolfigol, M. A.; Kiany-Borazjani, M.; Mallakpour, S. E.; Nassr-Isfahani, H. *Synth. Commun.*, **2000**, 30, 2573; (i) Zolfigol, M. A.; Madrakian, E.; Ghaemi, E. *Indian J. Chem.*, **2000**, 39B, 308; (j) Zolfigol, M. A.; Ghaemi, E.; Madrakian, E. *Synth. Commun.*, **2000**, 30, 1689; (k) Zolfigol, M. A.; Kiany-Borazjani, M.; Sadeghi, M. M.; Mohammadpoor-Baltork, I.; Memarian, H. R. *Synth. Commun.*, **2000**, 30, 3919; (l) Zolfigol, M. A.; Shirini, F.; Ghorbani Choghamarani, A.; Taqian-nasab, A.; Keypour, H.; Salehzadeh, S. *J. Chem. Research (S)*, **2000**, 420.
28. For the application of this system to the oxidation of 1,4-dihydropyridines: Zolfigol, M. A. Kiany-Borazjani, M. Sadeghi, M. M. Memarian, H. R. and Mohammadpoor-Baltork, I. *J. Chem. Research (S)*, **2000**, 167.
29. Anderson, R. A. Dalgleish, D. T. Nonhebel, D. C. and Pauson, P. L. *J. Chem. Research (S)*, **1977**, 12.

Sample Availability: All products reported in this paper are available from the authors.