Copyrighted Material

WILEY-VCH

Christian Reichardt

Solvents and Solvent Effects in Organic Chemistry

Third, Updated and Enlarged Edition



Copyrighted Material

Christian Reichardt

Solvents and Solvent Effects in Organic Chemistry

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

Related Titles from WILEY-VCH

Bittner, C. et al.

Organic Synthesis Workbook II

2001. 3-527-30415-0

Jessop, P. G., Leitner, W. (Eds.)

Chemical Synthesis Using Supercritical Fluids

1999. ISBN 3-527-29605-0

Wasserscheid, P., Welton, T. (Eds.)

Ionic Liquids in Synthesis

2002. ISBN 3-527-30515-7

Drauz, K., Waldmann, H. (Eds)

Enzyme Catalysis in Organic Synthesis

2002. ISBN 3-527-29949-1

Christian Reichardt

Solvents and Solvent Effects in Organic Chemistry

Third, Updated and Enlarged Edition

WILEY-VCH

Prof. Dr. Christian Reichardt
Fachbereich Chemie
der Philipps-Universität Marburg
Hans-Meerwein-Straße
35032 Marburg
Germany
e-mail: Reichardt-Marburg@t-online.de

This book was carefully produced. Nevertheless, author and publisher do not warrant the information contained therein to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

First Reprint 2004

Library of Congress Card No.: applied for

A catalogue record for this book is available from the British Library.

Bibliographic information published by Die Deutsche Bibliothek
Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data is available in the Internet at http://dnb.ddb.de.

ISBN 3-527-30618-8

© 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
Printed on acid-free paper.
All rights reserved (including those of translation in other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Composition: Asco Typesetters, Hong Kong. Printing: Strauss Offsetdruck GmbH, Mörlenbach
Bookbinding: J. Schäffer GmbH & Co. KG, Grünstadt
Printed in the Federal Republic of Germany.

To Maria
and in memory of my parents

Preface to the Third Edition

Meeting the demand for the second edition of this book, which is – despite a reprint in 1990 – no longer available, and considering the progress that has been made during the last decade in the study of solvent effects in experimental and theoretical organic chemistry, this improved third edition is presented to the interested reader.

Following the same layout as in the second edition, all topics retained have been brought up to date, with smaller and larger changes and additions on nearly every page. Two Sections (4.4.7 and 5.5.13) are completely new, dealing with solvent effects on host/guest complexation equilibria and reactions in biphasic solvent systems and neoteric solvents, respectively. More than 900 new references have been added, giving preference to review articles, and many older ones have been deleted. New references either replace older ones or are added to the end of the respective reference list of each chapter. The references cover the literature up to the end of 2001.

From the vast number of published papers dealing with solvent effects in all areas of organic chemistry, only some illustrative examples from the didactic and systematic point of view could be selected. This book is not a monograph covering *all* relevant literature in this field of research. The author, responsible for this subjective selection, apologizes in advance to all chemists whose valuable work on solvent effects is not mentioned in this book. However, using the reviews cited, the reader will find easy access to the full range of papers published in a certain field of research on solvent effects.

Great progress has been made during the last decade in theoretical treatments of solvent effects by various quantum-chemical methods and computational strategies. When indicated, relevant references are given to the respective solution reactions or absorptions. However, a critical evaluation of all the theoretical models and methods used to calculate the differential solvation of educts, activated complexes, products, ground and excited states, is outside the expertise of the present author. Thus, a book on all kinds of theoretical calculations of solvent influences on chemical reactions and physical absorptions has still to be written by someone else.

Consistent use of the nomenclature,[a)] symbols,[b)] terms,[c)] and SI units[d)] recommended by the IUPAC commissions has also been made in this third edition.

For comments and valuable suggestions I have to thank many colleagues, in particular Prof. E. M. Kosower, Tel Aviv/Israel, Prof. R. G. Makitra, Lviv/Ukraine, Prof. N. O. Mchedlov-Petrossyan, Kharkiv/Ukraine, and Prof. K. Möckel, Mühlhausen/Germany. For their assistance in drawing formulae, preparing the indices, and providing me with difficult to obtain literature, I thank Mr. G. Schäfer (technician), Mrs. S. Schellenberg (secretary), and Mrs. B. Becht-Schröder (librarian), all at the Department

---

a) G. J. Leigh, H. A. Favre, and W. V. Metanomski: *Principles of Chemical Nomenclature – A Guide to IUPAC Recommendations*, Blackwell Science Publications, London, 1998.

b) I. Mills, T. Cvitas, K. Homann, N. Kallay, and K. Kuchitsu: *Quantities, Units and Symbols in Physical Chemistry*, 2nd ed., Blackwell Science Publications, London, 1993.

c) P. Müller: *Glossary of Terms used in Physical Organic Chemistry – IUPAC Recommendations 1994*, Pure Appl. Chem. *66*, 1077 (1994).

d) G. H. Aylward and T. J. V. Tristan: *SI Chemical Data*, 4th ed., Wiley, Chichester, 1999; *Datensammlung Chemie in SI-Einheiten*, 3rd ed., Wiley-VCH, Weinheim/Germany, 1999.

of Chemistry, Philipps University, Marburg/Germany. Special thanks are due to the staff of Wiley-VCH Verlag GmbH, Weinheim/Germany, particularly to Dr. Elke Westermann, for their fine work in turning the manuscript into the final book. Lastly, my biggest debt is to my wife Maria, not only for her assistance in the preparation of the manuscript, but also for her constant encouragement and support during the writing of this book.

Marburg (Lahn), Spring 2002 Christian Reichardt

Preface to the Second Edition

The response to the first English edition of this book, published in 1979, has been both gratifying and encouraging. Its mixed character, lying between that of a monograph and a textbook, has obviously made it attractive to both the industrial and academic chemist as well as the advanced student of chemistry.

During the last eight years the study of solvent effects on both chemical reactions and absorption spectra has made much progress, and numerous interesting and fascinating examples have been described in the literature. In particular, the study of ionic reactions in the gas phase – now possible due to new experimental techniques – has allowed direct comparisons between gas-phase and solution reactions. This has led to a greater understanding of solution reactions. Consequently, Chapters 4 and 5 have been enlarged to include a description of ionic gas-phase reactions compared to their solution counterparts.

The number of well-studied solvent-dependent processes, *i.e.* reactions and absorptions in solution, has increased greatly since 1979. Only a representative selection of the more instructive, recently studied examples could be included in this second edition.

The search for empirical parameters of solvent polarity and their applications in multiparameter equations has recently been intensified, thus making it necessary to rewrite large parts of Chapter 7.

Special attention has been given to the chemical and physical properties of organic solvents commonly used in daily laboratory work. Therefore, all Appendix Tables have been improved; some have been completely replaced by new ones. A new well-referenced table on solvent-drying has been added (Table A-3). Chapter 3 has been enlarged, in particular by the inclusion of solvent classifications using multivariate statistical methods (Section 3.5). All these amendments justify the change in the title of the book to *Solvents and Solvent Effects in Organic Chemistry*.

The references have been up-dated to cover literature appearing up to the first part of 1987. New references were added to the end of the respective reference list of each chapter from the first edition.

Consistent use of the nomenclature, symbols, terms, and SI units recommended by the IUPAC commissions has also been made in the second edition.*)

I am very indebted to many colleagues for corrections, comments, and valuable suggestions. Especially helpful suggestions came from Professors H.-D. Försterling, Marburg, J. Shorter, Hull/England, and R. I. Zalewski, Poznań/Poland, to whom I am very grateful. For critical reading of the whole manuscript and the improvement of my English I again thank Dr. Edeline Wentrup-Byrne, now living in Brisbane/Australia. Dr. P.-V. Rinze, Marburg, and his son Lars helped me with the author index. Finally, I would like to thank my wife Maria for her sympathetic assistance during the preparation of this edition and for her help with the indices.

Marburg (Lahn), Spring 1988 Christian Reichardt

---

* *Cf.* Pure Appl. Chem. *51*, 1 (1979); ibid. *53*, 753 (1981); ibid. *55*, 1281 (1983); ibid. *57*, 105 (1985).

Preface to the First Edition

The organic chemist usually works with compounds which possess labile covalent bonds and are relatively involatile, thereby often rendering the gas-phase unsuitable as a reaction medium. Of the thousands of reactions known to occur in solution only few have been studied in the gas-phase, even though a description of reaction mechanisms is much simpler for the gas-phase. The frequent necessity of carrying out reactions in the presence of a more or less inert solvent results in two main obstacles: The reaction depends on a larger number of parameters than in the gas-phase. Consequently, the experimental results can often be only qualitatively interpreted because the state of aggregation in the liquid phase has so far been insufficiently studied. On the other hand, the fact that the interaction forces in solution are much stronger and more varied than in the gas-phase, permits to affect the properties and reactivities of the solute in manifold modes.

Thus, whenever a chemist wishes to carry out a chemical reaction he not only has to take into consideration the right reaction partners, the proper reaction vessels, and the appropriate reaction temperature. One of the most important features for the success of the planned reaction is the selection of a suitable solvent. Since solvent effects on chemical reactivity have been known for more than a century, most chemists are now familiar with the fact that solvents may have a strong influence on reaction rates and equilibria. Today, there are about three hundred common solvents available, nothing to say of the infinite number of solvent mixtures. Hence the chemist needs, in addition to his intuition, some general rules and guiding-principles for this often difficult choice.

The present book is based on an earlier paperback "Lösungsmitteleffekte in der organischen Chemie" [1], which, though following the same layout, has been completely rewritten, greatly expanded, and brought up to date. The book is directed both toward the industrial and academic chemist and particularly the advanced student of chemistry, who on the one hand needs objective criteria for the proper choice of solvent but on the other hand wishes to draw conclusions about reaction mechanisms from the observed solvent effects.

A knowledge of the physico-chemical principles of solvent effects is required for proper bench-work. Therefore, a description of the intermolecular interactions between dissolved molecules and solvent is presented first, followed by a classification of solvents derived therefrom. Then follows a detailed description of the influence of solvents on chemical equilibria, reaction rates, and spectral properties of solutes. Finally, empirical parameters of solvent polarity are given, and in an appendix guidelines to the everyday choice of solvents are given in a series of Tables and Figures.

The number of solvent systems and their associated solvent effects examined is so enormous that a complete description of all aspects would fill several volumes. For example, in Chemical Abstracts, volume 85 (1976), approximately eleven articles per week were quoted in which the words "Solvent effects on ..." appeared in the title. In the present book only a few important and relatively well-defined areas of general importance have been selected. The book has been written from the point of view of practical use for the organic chemist rather than from a completely theoretical one.

In the selection of the literature more recent reviews were taken into account mainly. Original papers were cited in particular from the didactic point of view rather

than priority, importance or completeness. This book, therefore, does not only have the character of a monograph but also to some extent that of a textbook. In order to help the reader in his use of the literature cited, complete titles of the review articles quoted are given. The literature up until December 1977 has been considered together with a few papers from 1978. The use of symbols follows the recommendations of the Symbols Committee of the Royal Society, London, 1971 [2].

I am very grateful to Professor Karl Dimroth, Marburg, who first stimulated my interest in solvent effects in organic chemistry. I am indebted to Professors W. H. Pirkle, Urbana/Illinois, D. Seebach, Zürich/Switzerland, J. Shorter, Hull/England, and numerous other colleagues for helpful advice and information. Thanks are also due to the authors and publishers of copyrighted materials reproduced with their permission (cf. Figure and Table credits on page 495). For the careful translation and improvement of the English manuscript I thank Dr. Edeline Wentrup-Byrne, Marburg. Without the assistance and patience of my wife Maria, this book would not have been written.

Marburg (Lahn), Summer 1978 Christian Reichardt

References

[1] C. Reichardt: *Lösungsmitteleffekte in der organischen Chemie*. 2nd edition. Verlag Chemie, Weinheim 1973;
Effets de solvant en chimie organique (translation of the first-mentioned title into French, by I. Tkatchenko), Flammarion, Paris 1971;
Rastvoriteli v organicheskoi khimii (translation of the first-mentioned title into Russian, by E. R. Zakhsa), Izdatel'stvo Khimiya, Leningrad 1973.

[2] *Quantities, Units, and Symbols*, issued by The Symbols Committee of the Royal Society, London, in 1971.

Contents

1	**Introduction**	**1**
2	**Solute-Solvent Interactions**	**5**
2.1	Solutions	5
2.2	Intermolecular Forces	10
2.2.1	Ion-Dipole Forces	10
2.2.2	Dipole-Dipole Forces	11
2.2.3	Dipole-Induced Dipole Forces	13
2.2.4	Instantaneous Dipole-Induced Dipole Forces	13
2.2.5	Hydrogen Bonding	15
2.2.6	Electron-Pair Donor/Electron-Pair Acceptor Interactions (EPD/EPA Interactions)	19
2.2.7	Solvophobic Interactions	27
2.3	Solvation	30
2.4	Selective Solvation	38
2.5	Micellar Solvation (Solubilization)	42
2.6	Ionization and Dissociation	46
3	**Classification of Solvents**	**57**
3.1	Classification of Solvents according to Chemical Constitution	57
3.2	Classification of Solvents using Physical Constants	62
3.3	Classification of Solvents in Terms of Acid-Base Behaviour	73
3.3.1	Brønsted-Lowry Theory of Acids and Bases	73
3.3.2	Lewis Theory of Acids and Bases	79
3.4	Classification of Solvents in Terms of Specific Solute/Solvent Interactions	82
3.5	Classification of Solvents using Multivariate Statistical Methods	84
4	**Solvent Effects on the Position of Homogeneous Chemical Equilibria**	**93**
4.1	General Remarks	93
4.2	Solvent Effects on Acid/Base Equilibria	95
4.2.1	Brønsted Acids and Bases in Solution	95
4.2.2	Gas-Phase Acidities and Basicities	99
4.3	Solvent Effects on Tautomeric Equilibria	106
4.3.1	Solvent Effects on Keto/Enol Equilibria	106
4.3.2	Solvent Effects on other Tautomeric Equilibria	113
4.4	Solvent Effects on other Equilibria	121
4.4.1	Solvent Effects on Brønsted Acid/Base Equilibria	121
4.4.2	Solvent Effects on Lewis Acid/Base Equilibria	123
4.4.3	Solvent Effects on Conformational Equilibria	126
4.4.4	Solvent Effects on *cis*/*trans* or *E*/*Z* Isomerization Equilibria	132
4.4.5	Solvent Effects on Valence Isomerization Equilibria	135
4.4.6	Solvent Effects on Electron-Transfer Equilibria	137
4.4.7	Solvent Effects on Host/Guest Complexation Equilibria	139

5	**Solvent Effects on the Rates of Homogeneous Chemical Reactions**	**147**
5.1	General Remarks	147
5.2	Gas-Phase Reactivities	155
5.3	Qualitative Theory of Solvent Effects on Reaction Rates	162
5.3.1	The Hughes–Ingold Rules	163
5.3.2	Solvent Effects on Dipolar Transition State Reactions	173
5.3.3	Solvent Effects on Isopolar Transition State Reactions	187
5.3.4	Solvent Effects on Free-Radical Transition State Reactions	199
5.3.5	Limitations of the Hughes–Ingold Rules	215
5.4	Quantitative Theories of Solvent Effects on Reaction Rates	218
5.4.1	General Remarks	218
5.4.2	Reactions between Neutral, Apolar Molecules	219
5.4.3	Reactions between Neutral, Dipolar Molecules	225
5.4.4	Reactions between Neutral Molecules and Ions	233
5.4.5	Reactions between Ions	234
5.5	Specific Solvation Effects on Reaction Rates	237
5.5.1	Influence of Specific Anion Solvation on the Rates of S_N and other Reactions	238
5.5.2	Protic and Dipolar Aprotic Solvent Effects on the Rates of S_N Reactions	243
5.5.3	Quantitative Separation of Protic and Dipolar Aprotic Solvent Effects for Reaction Rates by Means of Solvent-Transfer Activity Coefficients	254
5.5.4	Acceleration of Base-Catalysed Reactions in Dipolar Aprotic Solvents	259
5.5.5	Influence of Specific Cation Solvation on Rates of S_N Reactions	262
5.5.6	Solvent Influence on the Reactivity of Ambident Anions	269
5.5.7	Solvent Effects on Mechanisms and Stereochemistry of Organic Reactions	273
5.5.8	Influence of Micellar and Solvophobic Interactions on Reaction Rates and Mechanisms	292
5.5.9	Liquid Crystals as Reaction Media	298
5.5.10	Solvent Cage Effects	303
5.5.11	External Pressure and Solvent Effects on Reaction Rates	308
5.5.12	Solvent Isotope Effects	315
5.5.13	Reactions in Biphasic Solvent Systems and in Neoteric Solvents	317
6	**Solvent Effects on the Absorption Spectra of Organic Compounds**	**329**
6.1	General Remarks	329
6.2	Solvent Effects on UV/Vis Spectra	330
6.2.1	Solvatochromic Compounds	330
6.2.2	Theory of Solvent Effects on UV/Vis Absorption Spectra	340
6.2.3	Specific Solvent Effects on UV/Vis Absorption Spectra	348
6.2.4	Solvent Effects on Fluorescence Spectra	352
6.2.5	Solvent Effects on ORD and CD Spectra	359
6.3	Solvent Effects on Infrared Spectra	363
6.4	Solvent Effects on Electron Spin Resonance Spectra	369

6.5 Solvent Effects on Nuclear Magnetic Resonance Spectra 375
6.5.1 Nonspecific Solvent Effects on NMR Chemical Shifts 375
6.5.2 Specific Solvent Effects on NMR Chemical Shifts 381
6.5.3 Solvent Effects on Spin-Spin Coupling Constants 387

7 Empirical Parameters of Solvent Polarity 389

7.1 Linear Gibbs Energy Relationships..... 389
7.2 Empirical Parameters of Solvent Polarity from Equilibrium Measurements..... 396
7.3 Empirical Parameters of Solvent Polarity from Kinetic Measurements . 402
7.4 Empirical Parameters of Solvent Polarity from Spectroscopic Measurements..... 411
7.5 Empirical Parameters of Solvent Polarity from other Measurements ... 443
7.6 Interrelation and Application of Solvent Polarity Parameters 445
7.7 Multiparameter Approaches 452

Appendix 471

A. Properties, Purification, and Use of Organic Solvents..... 471
A.1 Physical Properties 471
A.2 Purification of Organic Solvents..... 471
A.3 Spectroscopic Solvents..... 479
A.4 Solvents as Reaction Media..... 488
A.5 Solvents for Recrystallization 488
A.6 Solvents for Extraction and Partitioning (Distribution)..... 490
A.7 Solvents for Adsorption Chromatography 492
A.8 Solvents for Acid/Base Titrations in Non-Aqueous Media..... 496
A.9 Solvents for Electrochemistry 496
A.10 Toxicity of Organic Solvents..... 500

References..... 509

Figure and Table Credits 581

Subject Index 583

Author Index 599

List of Abbreviations

Abbreviations and Recommended Values of Some Fundamental Constants and Numbers[a,b)]

N_A	Avogadro constant	$6.0221 \cdot 10^{23}$ mol^{-1}
c_0	speed of light in vacuum	$2.9979 \cdot 10^{8}$ m $\cdot$ s^{-1}
ε_0	absolute permittivity of vacuum [$= 1/(\mu_0 \cdot {c_0}^2)$; μ_0 = permeability of vacuum]	$8.8542 \cdot 10^{-12}$ C^2 $\cdot$ J^{-1} $\cdot$ m^{-1}
e	elementary charge	$1.6022 \cdot 10^{-19}$ C
h	Planck constant	$6.6261 \cdot 10^{-34}$ J $\cdot$ s
R	gas constant	8.3145 J $\cdot$ K^{-1} $\cdot$ mol^{-1} (or 0.08206 L $\cdot$ atm $\cdot$ K^{-1} $\cdot$ mol^{-1})
k_B	Boltzmann constant ($= R/N_A$)	$1.3807 \cdot 10^{-23}$ J $\cdot$ K^{-1}
V_m	standard molar volume of an ideal gas (at $t = 0$ °C and $p = 100$ kPa)	22.711 L $\cdot$ mol^{-1}
T_0	zero of the Celsius scale	273.15 K
π	ratio of the circumference to the diameter of a circle	3.1416
e	exponential number and base of natural logarithms (ln)	2.7183
ln 10	natural logarithm of ten (ln x = ln 10 $\cdot$ lg x; lg = decadic logarithm)	2.303

Abbreviations and Symbols for Units[a,b)]

bar	bar ($= 10^5$ Pa $= 10^5$ N $\cdot$ m^{-2})	pressure
cg/g	centigram/gram	weight percent
cL/L, cl/l	centilitre/litre	volume percent
cmol/mol	centimol/mol	mole percent
cm	centimetre (10^{-2} m)	length
cm^3	cubic centimetre (millilitre mL; 10^{-6} m^3)	volume
C	coulomb	electric charge

[a)] I. Mills, T. Cvitaš, K. Homann, N. Kallay, and K. Kuchitsu: *Quantities, Units and Symbols in Physical Chemistry*. 2nd ed., Blackwell Scientific Publications, London, 1993.
[b)] G. H. Aylward and T. J. V. Tristan: *SI Chemical Data*. 4th ed., Wiley, Chichester, 1999; *Datensammlung Chemie in SI-Einheiten*. 3rd ed., Wiley-VCH, Weinheim/Germany, 1999.

°C	degrees centigrade (Celsius)	temperature
dm^3	cubic decimetre (litre L; 10^{-3} m^3)	volume
J	joule	energy
kJ	kilojoule (10^3 J)	energy
K	kelvin	temperature
L, l	litre (1 dm^3; 10^{-3} m^3)	volume
m	metre	length
min	minute	time
mol	mole	amount of substance
MPa	megapascal (10^6 Pa)	pressure
mT	millitesla (10^{-3} T)	magnetic flux density (magnetic field)
nm	nanometre (10^{-9} m)	length
Pa	pascal (1 $N \cdot m^{-2} = 10^{-5}$ bar)	pressure
percent (%)	part per hundred (10^{-2})	dimensionless fraction
ppm	part per million (10^{-6})	dimensionless fraction
s	second	time

Abbreviations and Symbols for Properties[c)]

a_i	activity of solute i	
$a(^1H)$	ESR hyperfine coupling constant (coupling with ^1H)	mT (= 10^{-3} T)
A_j	the solvent's anion-solvating tendency or 'acity' (Swain)	
AN	solvent acceptor number, based on ^{31}P NMR chemical shift of Et_3PO (Gutmann and Meyer)	
α	electric polarizability of a molecule, polarizability volume	$C^2 \cdot m^2 \cdot J^{-1}$ or $4\pi\varepsilon_0 \cdot cm^3$
α	empirical parameter of solvent hydrogen-bond donor acidity (Taft and Kamlet)	
B	empirical parameter of solvent Lewis basicity (Palm and Koppel)	
B_{MeOD}	IR based empirical parameter of solvent Lewis basicity (Palm and Koppel)	

[c)] P. Müller: *Glossary of Terms used in Physical Organic Chemistry – IUPAC Recommendations 1994*. Pure Appl. Chem. *66*, 1077 (1994).

B_{PhOH}	IR based empirical parameter of solvent Lewis basicity (Koppel and Paju; Makitra)	
B_j	the solvent's cation-solvating tendency or 'basity' (Swain)	
β	empirical parameter of solvent hydrogen-bond acceptor basicity (Taft and Kamlet)	
c	cohesive pressure (cohesive energy density) of a solvent	MPa ($= 10^6$ Pa)
$c_i, c(i)$	molar concentration of solute i	$mol \cdot L^{-1}$
C_A, C_B	Lewis acidity and Lewis basicity parameter (Drago)	
cmc	critical micelle concentration	$mol \cdot L^{-1}$
D_{HA}	molar bond-dissociation energy of the bond between H and A	$kJ \cdot mol^{-1}$
D_π	empirical parameter of solvent Lewis basicity, based on a 1,3-dipolar cycloaddition reaction (Nagai *et al.*)	
DN	solvent donor number (Gutmann) [$= -\Delta H(D—SbCl_5)$]	$kcal \cdot mol^{-1}$
DN^N	normalized solvent donor number (Marcus)	
δ, δ_H	Hildebrand's solubility parameter	$MPa^{1/2}$
δ	chemical shift of NMR signals	ppm
δ	solvent polarizability correction term (Taft and Kamlet)	
E	energy, molar energy	$kJ \cdot mol^{-1}$
E	electric field strength	$V \cdot m^{-1}$
E	enol constant (K. H. Meyer)	
E	empirical parameter of solvent Lewis acidity (Palm and Koppel)	
E_A, E_a	Arrhenius activation energy	$kJ \cdot mol^{-1}$
E_A, E_B	Lewis acidity and Lewis basicity parameter (Drago)	
EA	electron affinity	$kJ \cdot mol^{-1}$
E_B^N	empirical solvent Lewis basicity parameter, based on the $n \rightarrow \pi^*$ absorption of an aminyloxide radical (Mukerjee; Wrona)	
E_K	empirical solvent polarity parameter, based on the $d \rightarrow \pi^*$ absorption of a molybdenum complex (Walther)	$kcal \cdot mol^{-1}$

E^*_{MLCT}	empirical solvent polarity parameter, based on the $d \rightarrow \pi^*$ absorption of a tungsten complex (Lees)	
E_{T}	molar electronic *t*ransition *e*nergy, molar electronic excitation energy	$\mathrm{kJ \cdot mol^{-1}}$ or $\mathrm{kcal \cdot mol^{-1}}$
$E_{\mathrm{T}}(30)$	empirical solvent polarity parameter, based on the intramolecular CT absorption of a pyridinium-*N*-phenolate betaine dye (Dimroth and Reichardt)	$\mathrm{kcal \cdot mol^{-1}}$
$E^{\mathrm{N}}_{\mathrm{T}}$	normalized $E_{\mathrm{T}}(30)$ solvent polarity parameter (Reichardt)	
$E^{\mathrm{SO}}_{\mathrm{T}}$	empirical solvent polarity parameter, based on the $n \rightarrow \pi^*$ absorption of an *S*-oxide (Walter)	$\mathrm{kcal \cdot mol^{-1}}$
EPA	electron-pair acceptor	
EPD	electron-pair donor	
ε_{r}	relative permittivity ($=\varepsilon/\varepsilon_0$) ("dielectric constant")	
Φ	empirical solvent polarity parameter, based on the $n \rightarrow \pi^*$ absorption of ketones (Dubois)	
G	IR based empirical solvent polarity parameter (Schleyer and Allerhand)	
ΔG°	standard molar Gibbs energy change	$\mathrm{kJ \cdot mol^{-1}}$
$\Delta G^{\neq}$	standard molar Gibbs energy of activation	$\mathrm{kJ \cdot mol^{-1}}$
$\Delta G^\circ_{\mathrm{solv}}$	standard molar Gibbs energy of solvation	$\mathrm{kJ \cdot mol^{-1}}$
$\Delta G^\circ_{\mathrm{hydr}}$	standard molar Gibbs energy of hydration	$\mathrm{kJ \cdot mol^{-1}}$
$\Delta G^\circ_{\mathrm{t}}(\mathrm{X}, \mathrm{O} \rightarrow \mathrm{S})$, $\Delta G^\circ_{\mathrm{t}}(\mathrm{X}, \mathrm{W} \rightarrow \mathrm{S})$	standard molar Gibbs energy of transfer of solute X from a reference solvent (O) or water (W) to another solvent (S)	$\mathrm{kJ \cdot mol^{-1}}$
γ_{i}	activity coefficient of solute i	
ΔH°	standard molar enthalpy change	$\mathrm{kJ \cdot mol^{-1}}$
$\Delta H^{\neq}$	standard molar enthalpy of activation	$\mathrm{kJ \cdot mol^{-1}}$
ΔH_{v}	molar enthalpy (heat) of vapourization	$\mathrm{kJ \cdot mol^{-1}}$
H_0	acidity function (Hammett)	
HBA	hydrogen-bond acceptor	

HBD	hydrogen-bond donor	
HOMO	highest occupied molecular orbital	
E_i, I, IP	ionization energy	$kJ \cdot mol^{-1}$
I	gas-chromatographic retention index (Kováts)	
J	NMR spin-spin coupling constant	Hz
k	rate constant for monomolecular ($n = 1$) and bimolecular ($n = 2$) reactions	$(L \cdot mol^{-1})^{n-1} \cdot s^{-1}$
k_0	rate constant in a reference solvent or in the gas phase for monomolecular ($n = 1$) and bimolecular reactions ($n = 2$)	$(L \cdot mol^{-1})^{n-1} \cdot s^{-1}$
k_0	in Hammett equations the rate constant of unsubstituted substrates	$(L \cdot mol^{-1})^{n-1} \cdot s^{-1}$ with $n = 1$ or 2
K, K_c	equilibrium constant (concentration basis; v = stoichiometric number)	$(mol \cdot L^{-1})^{\Sigma v}$
K_a, K_b	acid and base ionization constants	$(mol \cdot L^{-1})^{\Sigma v}$
K_{auto}	autoionization ion product, autoprotolysis constant	$mol^2 \cdot L^{-2}$
$K_{Assoc}, K_{Dissoc}, K_{ion}, K_T$	equilibrium constants of association, dissociation, ionization, resp. tautomerization reactions	$(mol \cdot L^{-1})^{\Sigma v}$
$K_{o/w}$	1-octanol/water partition constant (Hansch and Leo)	
KB	kauri-butanol number	
L	desmotropic constant (K. H. Meyer)	
LUMO	lowest unoccupied molecular orbital	
λ	wavelength	nm ($= 10^{-9}$ m)
m	mass of a particle	g
M_r	relative molecular mass of a substance (“molecular weight”)	
M	miscibility number (Godfrey)	
MH	microscopic hydrophobicity parameter of substituents (Menger)	
μ	empirical solvent softness parameter (Marcus)	
μ	permanent electric dipole moment of a molecule	$C \cdot m$ (or D)
μ_{ind}	induced electric dipole moment of a molecule	$C \cdot m$ (or D)

μ_i°	standard chemical potential of solute i	$kJ \cdot mol^{-1}$
μ_i^∞	standard chemical potential of solute i at infinite dilution	$kJ \cdot mol^{-1}$
n, n_D	refractive index (at sodium D line) $(= c_0/c)$	
N	empirical parameter of solvent nucleophilicity (Winstein and Grunwald)	
N_+	nucleophilicity parameter for (nucleophile + solvent)-systems (Ritchie)	
ν	frequency	Hz, s^{-1}
ν°	frequency in the gas phase or in an inert reference solvent	Hz, s^{-1}
$\tilde{\nu}$	wavenumber $(= 1/\lambda)$	cm^{-1}
Ω	empirical solvent polarity parameter, based on a Diels-Alder reaction (Berson)	
p	pressure	Pa $(= 1N \cdot m^{-2})$, bar $(= 10^5$ Pa)
P	measure of solvent polarizability (Palm and Koppel)	
P	empirical solvent polarity parameter, based on ^{19}F NMR measurements (Taft)	
PA	proton affinity	$kJ \cdot mol^{-1}$
Py	empirical solvent polarity parameter, based on the $\pi^* \rightarrow \pi$ emission of pyrene (Winnik)	
$P_{o/w}$	1-octanol/water partition coefficient (Hansch and Leo)	
pH	$-\lg[H_3O^+]$, $-\lg\, c(H_3O^+)$ (abbreviation of *p*otentia *h*ydrogenii or *p*uissance d'*h*ydrogène (Sörensen 1909)	
pK	$-\lg\, K$	
π	internal pressure of a solvent	MPa $(= 10^6$ Pa)
π^*	empirical solvent dipolarity/ polarizability parameter, based on the $\pi \rightarrow \pi^*$ absorption of substituted aromatics (Taft and Kamlet)	

π^*_{azo}	empirical solvent dipolarity/ polarizability parameter, based on the $\pi \rightarrow \pi^*$ absorption of azo merocyanine dyes (Buncel)	
π_x	hydrophobicity parameter of substituent X in H_5C_6-X (Hansch)	
r	radius of sphere representing an ion or a cavity	cm ($= 10^{-2}$ m)
r	distance between centres of two ions or molecules	cm ($= 10^{-2}$ m)
ρ	density (mass divided by volume)	$g \cdot cm^{-3}$
ρ, ρ_A	Hammett reaction resp. absorption constants	
S	generalized for solvent	
S	empirical solvent polarity parameter, based on the Z-values (Brownstein)	
$\mathscr{S}$	lg k_2 for the Menschutkin reaction of tri-*n*-propylamine with iodomethane (Drougard and Decroocq)	
ΔS°	standard molar entropy change	$J \cdot K^{-1} \cdot mol^{-1}$
$\Delta S^{\neq}$	standard molar entropy of activation	$J \cdot K^{-1} \cdot mol^{-1}$
S_p	solvophobic power of a solvent (Abraham)	
SA	empirical parameter of solvent hydrogen-bond donor acidity (Catalán)	
SB	empirical parameter of solvent hydrogen-bond acceptor basicity (Catalán)	
SPP	empirical parameter of solvent dipolarity/polarizability, based on the $\pi \rightarrow \pi^*$ absorption of substituted 7-nitrofluorenes (Catalán)	
σ	Hammett substituent constant	
σ	NMR screening constant	
t	Celsius temperature	°C
T	thermodynamic temperature	K
t_{mp}	melting point	°C
t_{bp}	boiling point	°C
U	internal energy	kJ
ΔU_v	molar energy of vapourization	$kJ \cdot mol^{-1}$

V_m, $V_{m,i}$	molar volume (of i)	$cm^3 \cdot mol^{-1}$
$\Delta V^{\neq}$	molar volume of activation	$cm^3 \cdot mol^{-1}$
x_i, $x(i)$	mole fraction of i ($x_i = n_i / \sum n$)	
X	empirical solvent polarity parameter, based on an S_E2 reaction (Gielen and Nasielski)	
χ_R, χ_B	empirical solvent polarity parameters, based on the $\pi \rightarrow \pi^*$ absorption of merocyanine dyes (Brooker)	$kcal \cdot mol^{-1}$
${}^{O}y_X^S$, ${}^{W}y_X^S$	solvent-transfer activity coefficient of a solute X from a reference solvent (O) or water (W) to another solvent (S)	
Y	empirical parameter of solvent ionizing power, based on *t*-butyl chloride solvolysis (Winstein and Grunwald)	
Y_{OTs}	empirical parameter of solvent ionizing power, based on 2-adamantyl tosylate solvolysis (Schleyer and Bentley)	
Y	measure of solvent polarization (Palm and Koppel)	
z_i	charge number of an ion i	positive for cations, negative for anions
Z	empirical solvent polarity parameter, based on the intermolecular CT absorption of a substituted pyridinium iodide (Kosower)	$kcal \cdot mol^{-1}$

"*Agite, Auditores ornatissimi, transeamus alacres ad aliud negotii! quum enim sic satis excusserimus ea quatuor Instrumenta artis, et naturae, quae modo relinquimus, videamus quintum genus horum, quod ipsi Chemiae fere proprium censetur, cui certe Chemistae principem locum prae omnibus assignant, in quo se jactant, serioque triumphant, cui artis suae, prae aliis omnibus effectus mirificos adscribunt. Atque illud quidem Menstruum vocaverunt.*"*)

Hermannus Boerhaave (1668–1738)
De menstruis dictis in chemia, in:
Elementa Chemiae (1733) [1, 2].

1 Introduction

The development of our knowledge of solutions reflects to some extent the development of chemistry itself [3]. Of all known substances, water was the first to be considered as a solvent. As far back as the time of the Greek philosophers there was speculation about the nature of solution and dissolution. The Greek alchemists considered all chemically active liquids under the name "Divine water". In this context the word "water" was used to designate everything liquid or dissolved.

The alchemist's search for a universal solvent, the so-called "Alkahest" or "Menstruum universale", as it was called by Paracelsus (1493–1541), indicates the importance given to solvents and the process of dissolution. Although the eager search of the chemists of the 15th to 18th centuries did not in fact lead to the discovery of any "Alkahest", the numerous experiments performed led to the uncovering of new solvents, new reactions, and new compounds**). From these experiences arose the earliest chemical rule that "like dissolves like" *(similia similibus solvuntur)*. However, at that time, the words solution and dissolution comprised all operations leading to a liquid product and it was still a long way to the conceptual distinction between the physical dissolution of a salt or of sugar in water, and the chemical change of a substrate by dissolution, for example, of a metal in an acid. Thus, in the so-called chemiatry period (iatrochemistry period), it was believed that the nature of a substance was fundamentally lost upon dissolution. Van Helmont (1577–1644) was the first to strongly oppose this contention. He claimed that the dissolved substance had not disappeared, but was present in the solution, although in aqueous form, and could be recovered [4]. Nevertheless, the dissolution

* "Well then, my dear listeners, let us proceed with fervor to another problem! Having sufficiently analyzed in this manner the four resources of science and nature, which we are about to leave (*i.e.* fire, water, air, and earth) we must consider a fifth element which can almost be considered the most essential part of chemistry itself, which chemists boastfully, no doubt with reason, prefer above all others, and because of which they triumphantly celebrate, and to which they attribute above all others the marvellous effects of their science. And this they call the solvent (menstruum)."
** Even if the once famous scholar J. B. Van Helmont (1577–1644) claimed to have prepared this "Alkahest" in a phial, together with the adherents of the alkahest theory he was ridiculed by his contemporaries who asked in which vessel he has stored this universal solvent.

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

of a substance in a solvent remained a rather mysterious process. The famous Russian polymath Lomonosov (1711–1765) wrote in 1747: "Talking about the process of dissolution, it is generally said that all solvents penetrate into the pores of the body to be dissolved and gradually remove its particles. However, concerning the question of what forces cause this process of removal, there does not exist any somehow reasonable explanation, unless one arbitrarily attributes to the solvents sharp wedges, hooks or, who knows, any other kind of tools" [27].

The further development of modern solution theory is connected with three persons, namely the French researcher Raoult (1830–1901) [28], the Dutch physical chemist van't Hoff (1852–1911) [5], and the Swedish scientist Arrhenius (1859–1927) [6]. Raoult systematically studied the effects of dissolved nonionic substances on the freezing and boiling point of liquids and noticed in 1886 that changing the solute/solvent ratio produces precise proportional changes in the physical properties of solutions. The observation that the vapour pressure of solvent above a solution is proportional to the mole fraction of solvent in the solution is today known as Raoult's law [28].

The difficulty in explaining the effects of inorganic solutes on the physical properties of solutions led in 1884 to Arrhenius' theory of incomplete and complete dissociation of ionic solutes (electrolytes, ionophores) into cations and anions in solution, which was only very reluctantly accepted by his contemporaries. Arrhenius derived his dissociation theory from comparison of the results obtained by measurements of electroconductivity and osmotic pressure of dilute electrolyte solutions [6].

The application of laws holding for gases to solutions by replacing pressure by osmotic pressure was extensively studied by van't Hoff, who made osmotic pressure measurements another important physicochemical method in studies of solutions [5].

The integration of these three basic developments established the foundations of modern solution theory and the first Nobel prizes in chemistry were awarded to van't Hoff (in 1901) and Arrhenius (in 1903) for their work on osmotic pressure and electrolytic dissociation, respectively.

The further development of solution chemistry is connected with the pioneering work of Ostwald (1853–1932), Nernst (1864–1941), Lewis (1875–1946), Debye (1884–1966), E. Hückel (1896–1980), and Bjerrum (1879–1958). More detailed reviews on the development of modern solution chemistry can be found in references [29, 30].

The influence of solvents on the rates of chemical reactions [7, 8] was first noted by Berthelot and Péan de Saint-Gilles in 1862 in connection with their studies on the esterification of acetic acid with ethanol: "... l'éthérification est entravée et ralentie par l'emploi des dissolvants neutres étrangers à la réaction" [9]*). After thorough studies on the reaction of trialkylamines with haloalkanes, Menschutkin in 1890 concluded that a reaction cannot be separated from the medium in which it is performed [10]. In a letter to Prof. Louis Henry he wrote in 1890: "Or, l'expérience montre que ces dissolvants exercent sur la vitesse de combinaison une influence considérable. Si nous représentons par 1 la constante de vitesse de la réaction précitée dans l'hexane C_6H_{14}, cette constante pour la même combinaison dans $CH_3—CO—C_6H_5$, toutes choses égales d'ailleurs sera 847.7. La différence est énorme, mais, dans ce cas, elle n'atteint pas encore le maxi-

* "... the esterification is disturbed and decelerated on addition of neutral solvents not belonging to the reaction" [9].

mum.... Vous voyez que les dissolvants, soi-disant indifférents ne sont pas inertes; ils modifient profondément l'acte de la combinaison chimique. Cet énoncé est riche en conséquences pour la théorie chimique des dissolutions'' [26]*). Menschutkin also discovered that, in reactions between liquids, one of the reaction partners may constitute an unfavourable solvent. Thus, in the preparation of acetanilide, it is not without importance whether aniline is added to an excess of acetic acid, or *vice versa*, since aniline in this case is an unfavorable reaction medium. Menschutkin related the influence of solvents primarily to their chemical, not their physical properties.

The influence of solvents on chemical equilibria was discovered in 1896, simultaneously with the discovery of keto-enol tautomerism**) in 1,3-dicarbonyl compounds (Claisen [14]: acetyldibenzoylmethane and tribenzoylmethane; Wislicenus [15]: methyl and ethyl formylphenylacetate; Knorr [16]: ethyl dibenzoylsuccinate and ethyl diacetylsuccinate) and the nitro-isonitro tautomerism of primary and secondary nitro compounds (Hantzsch [17]: phenylnitromethane). Thus, Claisen wrote: ''Es gibt Verbindungen, welche sowohl in der Form —C(OH)=C(|)—CO— wie in der Form —CO—CH(|)—CO— zu bestehen vermögen; von der Natur der angelagerten Reste, von der Temperatur, bei den gelösten Substanzen auch von der Art des Lösungsmittels hängt es ab, welche von den beiden Formen die beständigere ist'' [14]***). The study of the keto-enol equilibrium of ethyl formylphenylacetate in eight solvents led Wislicenus to the conclusion that the keto form predominates in alcoholic solution, the enol form in chloroform or benzene. He stated that the final ratio in which the two tautomeric forms coexist must depend on the nature of the solvent and on its dissociating power, whereby he suggested that the dielectric constants were a possible measure of this ''power''. Stobbe was the first to review these results [18]. He divided the solvents into two groups according to their ability to isomerize tautomeric compounds. His classification reflects, to some extent, the modern division into protic and aprotic solvents. The effect of solvent on constitutional and tautomeric isomerization equilibria was later studied in detail

---

* ''Now, experience shows that solvents exert considerable influence on reaction rates. If we represent the rate constant of the reaction to be studied in hexane C_6H_{14} by 1, then, all else being equal, this constant for the same reaction in CH_3—CO—C_6H_5 will be 847.7. The increase is enormous, but in this case it has not even reached its maximum.... So you see that solvents, in spite of appearing at first to be indifferent, are by no means inert; they can greatly influence the course of chemical reactions. This statement is full of consequences for the chemical theory of dissolutions'' [26].

** The first observation of a tautomeric equilibrium was made in 1884 by Zincke at Marburg [11]. He observed that, surprisingly, the reaction of 1,4-naphthoquinone with phenylhydrazine gives the same product as that obtained from the coupling reaction of 1-naphthol with benzenediazonium salts. This phenomenon, that the substrate can react either as phenylhydrazone or as a hydroxyazo compound, depending on the reaction circumstances, was called *Ortsisomerie* by Zincke [11]. Later on, the name *tautomerism*, with a different meaning however from that accepted today, was introduced by Laar [12]. For a description of the development of the concept of tautomerism, see Ingold [13].

*** ''There are compounds capable of existence in the form —C(OH)=C(|)—CO— as well as in the form —CO—CH(|)—CO—; it depends on the nature of the substituents, the temperature, and for dissolved compounds, also on the nature of the solvent, which of the two forms will be the more stable'' [14].

by Dimroth [19] (using triazole derivatives, *e.g.* 5-amino-4-methoxycarbonyl-1-phenyl-1,2,3-triazole) and Meyer [20] (using ethyl acetoacetate).

It has long been known that UV/Vis absorption spectra may be influenced by the phase (gas or liquid) and that the solvent can bring about a change in the position, intensity, and shape of the absorption band*). Hantzsch later termed this phenomenon *solvatochromism***) [22]. The search for a relationship between solvent effect and solvent property led Kundt in 1878 to propose the rule, later named after him, that increasing dispersion (*i.e.* increasing index of refraction) is related to a shift of the absorption maximum towards longer wavelength [23]. This he established on the basis of UV/Vis absorption spectra of six dyestuffs, namely chlorophyll, fuchsin, aniline green, cyanine, quinizarin, and egg yolk in twelve different solvents. The – albeit limited – validity of Kundt's rule, *e.g.* found in the cases of 4-hydroxyazobenzene [24] and acetone [25], led to the realization that the effect of solvent on dissolved molecules is a result of electrical fields. These fields in turn originate from the dipolar properties of the molecules in question [25]. The similarities in the relationships between solvent effects on reaction rate, equilibrium position, and absorption spectra has been related to the general solvating ability of the solvent in a fundamental paper by Scheibe *et al.* [25].

More recently, research on solvents and solutions has again become a topic of interest because many of the solvents commonly used in laboratories and in the chemical industry are considered as unsafe for reasons of environmental protection. On the list of damaging chemicals, solvents rank highly because they are often used in huge amounts and because they are volatile liquids that are difficult to contain. Therefore, the introduction of cleaner technologies has become a major concern throughout both academia and industry [31–34]. This includes the development of environmentally benign new solvents, sometimes called *neoteric solvents* (neoteric = recent, new, modern), constituting a class of novel solvents with desirable, less hazardous, new properties [35, 36]. The term *neoteric solvents* covers supercritical fluids, ionic liquids, and also perfluorohydrocarbons (as used in fluorous biphasic systems). Table A-14 in Chapter A.10 (Appendix) collects some recommendations for the substitution of hazardous solvents, together with the relevant literature references.

For the development of a sustainable chemistry based on clean technologies, the best solvent would be no solvent at all. For this reason, considerable efforts have recently been made to design reactions that proceed under solvent-free conditions, using modern techniques such as reactions on solid mineral supports (alumina, silica, clays), solid-state reactions without any solvent, support, or catalyst between neat reactants, solid-liquid phase-transfer catalysed and microwave-activated reactions, as well as gas-phase reactions [37–42]. However, not all organic reactions can be carried out in the absence of a solvent; some organic reactions even proceed explosively in the solid state! Therefore, solvents will still be useful in mediating and moderating chemical reactions and this book on solvent effects will certainly not become superfluous in the foreseeable future.

---

* A survey of older works of solvent effects on UV/Vis absorption spectra has been given by Sheppard [21].

** It should be noted that the now generally accepted meaning of the term *solvatochromism* differs from that introduced by Hantzsch (*cf.* Section 6.2).

2 Solute-Solvent Interactions

2.1 Solutions

In a limited sense *solutions* are homogeneous liquid phases consisting of more than one substance in variable ratios, when for convenience one of the substances, which is called the *solvent* and may itself be a mixture, is treated differently from the other substances, which are called *solutes* [1]. Normally, the component which is in excess is called the solvent and the minor component(s) is the solute. When the sum of the mole fractions of the solutes is small compared to unity, the solution is called a *dilute solution**). A solution of solute substances in a solvent is treated as an *ideal dilute solution* when the solute activity coefficients γ are close to unity ($\gamma = 1$) [1, 171]. Solute/solvent mixtures $A + B$ that obey Raoult's law over the entire composition range from pure A to pure B are called *ideal solutions*. According to Raoult, the ratio of the partial pressure of component $A(p_A)$ to its vapour pressure as a pure liquid (p_A^*) is equal to the mole fraction of $A(x_A)$ in the liquid mixture, *i.e.* $x_A = p_A/p_A^*$. Many mixtures obey Raoult's law very well, particularly when the components have a similar molecular structure (*e.g.* benzene and toluene).

A solvent should not be considered a macroscopic continuum characterized only by physical constants such as density, dielectric constant, index of refraction *etc.*, but as a discontinuum which consists of individual, mutually interacting solvent molecules. According to the extent of these interactions, there are solvents with a pronounced internal structure (*e.g.* water) and others in which the interaction between the solvent molecules is small (*e.g.* hydrocarbons). The interactions between species in solvents (and in solutions) are at once too strong to be treated by the laws of the kinetic theory of gases, yet too weak to be treated by the laws of solid-state physics. Thus, the solvent is neither an indifferent medium in which the dissolved material diffuses in order to distribute itself evenly and randomly, nor does it possess an ordered structure resembling a crystal lattice. Nevertheless, the long-distance ordering in a crystal corresponds somewhat to the local ordering in a liquid. Thus, neither of the two possible models – the gas and crystal models – can be applied to solutions without limitation. There is such a wide gulf between the two models in terms of conceivable and experimentally established variants, that it is too difficult to develop a generally valid model for liquids. Due to the complexity of the interactions, the structure of liquids – in contrast to that of gases and solids – is the least-known of all aggregation states. Therefore, the experimental and theoretical examination of the structure of liquids is among the most difficult tasks of physical chemistry [2–7, 172–174].

Any theory of the liquid state has to explain – among others – the following facts: Except for water, the molar volume of a liquid is roughly 10% greater than that of the corresponding solid. According to X-ray diffraction studies, a short-range order of solvent molecules persists in the liquid state and the nearest neighbour distances are almost the same as those in the solid. The solvent molecules are not moving freely, as in the

* The superscript ∞ attached to the symbol for a property of a solution denotes the property of an *infinitely dilute solution*.

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

gaseous state, but instead move in the potential field of their neighbours. The potential energy of a liquid is higher than that of its solid by about 10%. Therefore, the heat of fusion is roughly 10% of the heat of sublimation. Each solvent molecule has an environment very much like that of a solid, but some of the nearest neighbours are replaced by holes. Roughly one neighbour molecule in ten is missing.

Even for the most important solvent – water – the investigation of its inner fine structure is still the subject of current research [8–15, 15a]*). Numerous different models, *e.g.* the "flickering cluster model" of Franck and Wen [16], were developed to describe the structure of water. However, all these models prove themselves untenable for a complete description of the physico-chemical properties of water and an interpretation of its anomalies [304]. Fig. 2-1 should make clear the complexity of the inner structure of water.

Liquid water consists both of bound ordered regions of a regular lattice, and regions in which the water molecules are hydrogen-bonded in a random array; it is permeated by monomeric water and interspersed with random holes, lattice vacancies, and cages. There are chains and small polymers as well as bound, free, and trapped water molecules [9, 176]. The currently accepted view of the structure of liquid water treats it as a dynamic three-dimensional hydrogen-bonded network, without a significant number of non-bonded water molecules, that retains several of the structural characteristics of ice (*i.e.* tetrahedral molecular packing with each water molecule hydrogen-bonded to four nearest neighbours), although the strict tetrahedrality is lost [176]. Its dynamic behaviour resembles that of most other liquids, with short rotational and translational correlation times of the order of 0.1 to 10 ps, indicating high hydrogen-bond exchange rates [176, 305].

In principle, other hydrogen-bonded solvents should possess similar complicated structures [306]. However, whereas water has been thoroughly studied [17, 176, 307], the inner structures of other solvents are still less well known [172, 177–179]. By way of example, the intermolecular structure of acetone is determined mainly by steric interactions between the methyl groups and, unexpectedly, only to a small extent by dipole/dipole forces [308], whereas the inner structure of dimethyl sulfoxide is dictated by strong dipole/dipole interactions [309].

From the idea that the solvent only provides an indifferent reaction medium, comes the *Ruggli-Ziegler dilution principle*, long known to the organic chemist. According to this principle, in the case of cyclization reactions, the desired intramolecular reaction will be favoured over the undesired intermolecular reaction by high dilution with an inert solvent [18, 310].

The assumption of forces of interaction between solvent and solute led, on the other hand, to the century-old principle that "like dissolves like" *(similia similibus solvuntur)*, where the word "like" should not be too narrowly interpreted. In many cases, the presence of similar functional groups in the molecules suffices. When a chemical

* The amusing story of "polywater," which excited the scientific community for a few years during the late 1960's and early 1970's, has been reviewed by Franks [175]. It turned out that polywater was not a new and more stable form of pure water, but merely dirty water. The strange properties of polywater were due to high concentrations of siliceous material dissolved from quartz capillaries in which it was produced.



Fig. 2-1. Two-dimensional schematic diagram of the three-dimensional structure of liquid water [9].

similarity is present, the solution of the two components will usually have a structure similar to that of the pure materials (*e.g.* alcohol-water mixtures [19]). This rule of thumb has only limited validity, however, since there are many examples of solutions of chemically dissimilar compounds. For example, methanol and benzene, water and *N*,*N*-dimethylformamide, aniline and diethyl ether, and polystyrene and chloroform, are all completely miscible at room temperature. On the other hand, insolubility can occur in spite of similarity of the two partners. Thus, polyvinyl alcohol does not dissolve in ethanol, acetyl cellulose is insoluble in ethyl acetate, and polyacrylonitrile is insoluble in acrylonitrile [20]. Between these two extremes there is a whole range of possibilities where the two materials dissolve each other to a limited extent. The system water/diethyl ether is such an example. Pure diethyl ether dissolves water to the extent of 15 mg/g at 25 °C, whereas water dissolves diethyl ether to the extent of 60 mg/g. When one of the two solvents is in large excess a homogeneous solution is obtained. Two phases occur when the ratio is beyond the limits of solubility. A more recent example of a reaffirmation of the old "like dissolves like" rule is the differential solubility of fullerene (C_{60}), consisting of a three-dimensional delocalized 60π-electron system, in solvents such as methanol ($s = 0.01$ mg/mL) and 1-chloronaphthalene ($s = 50$ mg/mL) [311].

However, rather than the "like dissolves like" rule, it is the intermolecular interaction between solvent and solute molecules that determines the mutual solubility. A compound *A* dissolves in a solvent *B* only when the intermolecular forces of attraction K_{AA} and K_{BB} for the pure compounds can be overcome by the forces K_{AB} in solution [21].

The sum of the interaction forces between the molecules of solvent and solute can be related to the so-called *polarity**) of *A* and *B*. Denoting compounds with large interactions $A \cdots A$ or $B \cdots B$, respectively, as polar, and those with small interactions as nonpolar, four cases allowing a qualitative prediction of solubility can be distinguished (Table 2-1).

An experimental verification of these simple considerations is given by the solubility data in Table 2-2.

Table 2-1. Solubility and polarity [22].

Solute A	Solvent B	Interaction			Solubility of A in B
		A···A	B···B	A···B	
Nonpolar	nonpolar	weak	weak	weak	can be high[a)]
Nonpolar	polar	weak	strong	weak	probably low[b)]
Polar	nonpolar	strong	weak	weak	probably low[c)]
Polar	polar	strong	strong	strong	can be high[a)]

[a)] Not much change for solute or solvent.
[b)] Difficult to break up B···B.
[c)] Difficult to break up A···A.

* For a more detailed definition of solvent polarity, see Sections 3.2 and 7.1.

Table 2-2. Solubilities of methane, ethane, chloromethane, and dimethyl ether in tetrachloro-methane (nonpolar solvent) and acetone (polar solvent) [22].

Solute	Solute polarity	Solubility/(mol · m^{-3}) at 25 °C	
		in CCl_4	in CH_3COCH_3
CH_4	nonpolar	29	25
CH_3CH_3	nonpolar	220	130
CH_3Cl	polar	1700	2800
CH_3OCH_3	polar	1900	2200

The solubilities of ethane and methane are higher in nonpolar tetrachloro-methane, whereas the opposite is true for chloromethane and dimethyl ether. A survey of the reciprocal miscibility of some representative examples of organic solvents is given in Fig. 2-2.

Solubility is commonly defined as the concentration of dissolved solute in a solvent in equilibrium with undissolved solute at a specified temperature and pressure. For a deeper and more detailed understanding of the diverse rules determining the solubility of organic compounds in various solvents, see references [312–316].

The *solubility parameter* δ of Hildebrand [4, 24], as defined in Eq. (2-1), can often be used in estimating the solubility of non-electrolytes in organic solvents.

$$\delta = \left(\frac{\Delta U_v}{V_m}\right)^{1/2} = \left(\frac{\Delta H_v - R \cdot T}{V_m}\right)^{1/2} \tag{2-1}$$

In this equation, V_m is the molar volume of the solvent, and ΔU_v and ΔH_v are the molar energy and the molar enthalpy (heat) of vapourization to a gas of zero pressure,



Fig. 2-2. Miscibility of organic solvents [23]. —— miscible in all proportions; – – – – limited miscibility; ..... little miscibility; without line: immiscible.

actions can be expressed as

$$U_{\text{dispersion}} = -\frac{1}{(4\pi \cdot \varepsilon_0)^2} \cdot \frac{3\alpha_1 \cdot \alpha_2}{2r^6} \cdot \left(\frac{I_1 \cdot I_2}{I_1 + I_2}\right) \tag{2-6a}$$

where α_1 and α_2 are the polarizabilities and I_1 and I_2 are the ionization potentials of the two different interacting species [33]. When applied to two molecules of the same substance, Eq. (2-6a) reduces to Eq. (2-6b).

$$U_{\text{dispersion}} = -\frac{1}{(4\pi \cdot \varepsilon_0)^2} \cdot \frac{3\alpha^2 \cdot I}{4r^6} \tag{2-6b}$$

Dispersion forces are extremely short-range in action (depending on $1/r^6$!).

Dispersion forces are universal for all atoms and molecules; they alone are responsible for the aggregation of molecules which possess neither free charges nor electric dipole moments. Due to the greater polarizability of π-electrons, especially strong dispersion forces exist between molecules with conjugated π-electron systems (*e.g.* aromatic hydrocarbons). For many other dipole molecules with high polarizability as well, the major part of the cohesion is due to dispersion forces. For example, the calculated cohesion energy of liquid 2-butanone at 40 °C consists of 8% orientational energy, 14% inductional energy, and 78% dispersion energy [35]. Two molecules with $\alpha = 3 \cdot 10^{-30}$ m^3, $I = 20 \cdot 10^{-19}$ J, and $r = 3 \cdot 10^{-10}$ m have an interaction potential of -11.3 kJ/mol (-2.7 kcal/mol) [35a]. These values of α, I, and the average intermolecular distance r correspond to those for liquid HCl. It is instructive to compare the magnitude of these dispersion forces with that of the dipole-dipole interactions. For two dipoles, both with dipole moments of $3.3 \cdot 10^{-30}$ Cm (1.0 D), separated by a distance of $r = 3 \cdot 10^{-10}$ m and oriented as in Fig. 2-3a, the interaction energy is only -5.3 kJ/mol (-1.1 kcal/mol) [35a]. Thus, for HCl and most other compounds, the dispersion forces are considerably stronger than the dipole-dipole forces of nearest neighbour distance in the liquid state. However, at larger distances the dispersion energy falls off rapidly.

As a result of the α^2 term in Eq. (2-6b), dispersion forces increase rapidly with the molecular volume and the number of polarizable electrons. The polarizability α is connected with the molar refraction and the index of refraction, according to the equation of Lorenz-Lorentz. Therefore, solvents with a large index of refraction, and hence large optical polarizability, should be capable of enjoying particularly strong dispersion forces. As indicated in Table A-1 (Appendix), all aromatic compounds possess relatively high indices of refraction, *e.g.* quinoline ($n = 1.6273$), iodobenzene ($n = 1.6200$), aniline ($n = 1.5863$), and diphenyl ether ($n = 1.5763$); of all organic solvents, carbon disulfide ($n = 1.6275$) and diiodomethane ($n = 1.738$) have the highest indices of refraction.

Solvents with high polarizability are often good solvators for anions which also possess high polarizability. This is due to the fact that the dispersional interactions between the solvents and the large, polarizable anions like $I_3^{\ominus}$, $I^{\ominus}$, $SCN^{\ominus}$ or the picrate anion are significantly larger than for the smaller anions like $F^{\ominus}$, $HO^{\ominus}$, or $R_2N^{\ominus}$ [36]. Perfluorohydrocarbons have unusually low boiling points because tightly held electrons in fluorine have only a small polarizability.

2.2.5 Hydrogen Bonding [37–46, 187–190, 306]

Liquids possessing hydroxy groups or other groups with a hydrogen atom bound to an electronegative atom X are strongly associated and have abnormal boiling points. This observation led to the contention that particular intermolecular forces apply here. These are designated as hydrogen bridges, or hydrogen bonds, characterized by a coordinative divalency of the hydrogen atom involved. A general definition of the hydrogen bond is: when a covalently bound hydrogen atom forms a second bond to another atom, the second bond is referred to as a *hydrogen bond* [38].

The concept of hydrogen bonding was introduced in 1919 by Huggins [37]. The first definitive paper on hydrogen bonding – applied to the association of water molecules – was published in 1920 by Latimer and Rodebush [191]. All three were working in the Laboratory of G. N. Lewis, University of California, Berkeley/USA.

A hydrogen bond is formed by the interaction between the partners R—X—H and :Y—R′ according to Eq. (2-7).



(2-7)

R—X—H is the proton donor and :Y—R′ makes available an electron pair for the bridging bond. Thus, hydrogen bonding can be regarded as a preliminary step in a Brønsted acid-base reaction which would lead to a dipolar reaction product $R—X^{\ominus} \cdots H—Y^{\oplus}—R'$. X and Y are atoms of higher electronegativity than hydrogen (*e.g.* C, N, P, O, S, F, Cl, Br, I). Both inter- and intramolecular hydrogen bonding are possible, the latter when X and Y belong to the same molecule.

The most important electron pair donors (*i.e.* hydrogen bond acceptors) are the oxygen atoms in alcohols, ethers, and carbonyl compounds, as well as nitrogen atoms in amines and *N*-heterocycles. Hydroxy-, amino-, carboxyl-, and amide groups are the most important proton donor groups. Strong hydrogen bonds are formed by the pairs $O—H \cdots O$, $O—H \cdots N$, and $N—H \cdots O$, weaker ones by $N—H \cdots N$, and the weakest by $Cl_2C—H \cdots O$ and $Cl_2C—H \cdots N$. The π-electron systems of aromatic compounds, alkenes, and alkynes can also act as weak hydrogen bond acceptors [189].

When two or more molecules of the same type associate, so-called *homo-intermolecular* hydrogen bonds are formed (Fig. 2-4). The association of different molecules (*e.g.* $R—O—H \cdots NR_3$) results in *hetero-intermolecular* hydrogen bonds. The designations *homo-* and *heteromolecular* [192] as well as *homo-* and *heteroconjugated* hydrogen bond are also in use. A remarkable example of a competitive solvent-dependent equilibrium between homo- and hetero-intermolecular hydrogen-bond associated species has been found in solutions of 4-hydroxyacetophenone and 2-(2-hexyloxyethoxy)ethanol [319].

Hydrogen bonds can be either *inter*molecular or *intra*molecular. Both types of hydrogen bonds are found in solutions of 2-nitrophenol, depending on the Lewis basicity of the solvent [298]. The intramolecularly hydrogen-bonded form exists in non-hydrogen-bonding solvents (*e.g.* cyclohexane, tetrachloromethane). 2-Nitrophenol breaks its intramolecular hydrogen bond to form an intermolecular one in electron-pair donor (EPD) solvents (*e.g.* anisole, HMPT).



Fig. 2-4. Homo-intermolecular hydrogen bonds in alcohols, carboxylic acids, and amides (the hydrogen bonds are denoted by dotted lines).



Circular hydrogen bonds have been found in the hexahydrate of α-cyclodextrin (cyclohexaamylose) [193]. Hydration water molecules and hydroxy groups of the macromolecule cooperate to form a network-like pattern with circular O—H · · · O hydrogen bonds. If the O—H · · · O hydrogen bonds run in the same direction, the circle is called *homodromic*. Circles with the two counter-running chains are called *antidromic*, and circles with more randomly oriented chains are designated *heterodromic* [193]; *cf.* Fig. 2-4a. Such circular hydrogen bonds can be of importance with respect to the inner molecular structure of water and alcohols (*cf.* also Fig. 2-1).

The question of the exact geometry of hydrogen bonds (distances, angles, lone-pair directionality) has been reviewed [194].

The bond dissociation enthalpy for normal hydrogen bonds is *ca.* 13 . . . 42 kJ/mol (3 . . . 10 kcal/mol)*). For comparison, covalent single bonds have dissociation enthalpies of 210 . . . 420 kJ/mol (50 . . . 100 kcal/mol). Thus, hydrogen bonds are approx. ten times weaker than covalent single bonds, but also approx. ten times stronger than the non-



Fig. 2-4a. Three types of circular hydrogen bonds: (a) homodromic, (b) antidromic, and (c) heterodromic hydrogen bonds [193].

* Bond dissociation enthalpies outside these limits are, however, known. Examples of weak, normal, and strong hydrogen bonds are found in the following pairs: phenol/benzene ($\Delta H = -5$ kJ/mol) [47], phenol/triethylamine ($\Delta H = -37$ kJ/mol) [47], and trichloroacetic acid/triphenylphosphane oxide ($\Delta H = -67$ kJ/mol) [48]. An extremely strong hydrogen bond is found in $Me_4N^+HF_2^-$ ($\Delta H = -155$ kJ/mol) [38]. The strength of a hydrogen bond correlates with the basicity of the proton-acceptor and the acidity of the proton-donor molecule. Compounds with very strong hydrogen bonds have been reviewed [320].

specific intermolecular interaction forces. The question as to whether or not a hydrogen bond is stronger than the equivalent deuterium bond is addressed in reference [321]: the D-bond seems to be somewhat stronger than the H-bond in the case of neutral hydrogen-bonded complexes, but the reverse is true for charged complexes.

Hydrogen bonds are characterized by the following structural and spectroscopic features [39]: (a) the distances between the neighbouring atoms involved in the hydrogen bond [X and Y in Eq. (2-7)] are considerably smaller than the sum of their van der Waals radii; (b) the X—H bond length is increased and hydrogen bond formation causes its IR stretching mode to be shifted towards lower frequencies (for exceptions see reference [190]); (c) the dipolarity of the X—H bond increases on hydrogen bond formation, leading to a larger dipole moment of the complex than expected from vectorial addition of its dipolar components R—X—H and Y—R′; (d) due to the reduced electron density at H-atoms involved in hydrogen bonds, they are deshielded, resulting in substantial downfield shifts of their ^{1}H NMR signals; (e) in hetero-molecular hydrogen bonds, a shift of the Brønsted acid/base equilibrium $R—X—H \cdots Y—R' \rightleftharpoons R—X^{\ominus} \cdots H—Y^{\oplus}—R'$ to the right-hand side with increasing solvent polarity is found (*cf.* Section 4.4.1 and references [195, 322] for impressive examples).

Up until now there has been no general agreement as to the best description of the nature of the forces in the hydrogen bond [42–46]. The hydrogen bond can be described as a dipole-dipole or resonance interaction. Since hydrogen bonding occurs only when the hydrogen is bound to an electronegative atom, the first assumption concerning the nature of the hydrogen bond was that it consists of a dipole-dipole interaction such as $R—X^{\delta\ominus}—H^{\delta\oplus} \cdots Y^{\delta\ominus}—R'$. This viewpoint is supported by the fact that the strongest hydrogen bonds are formed in pairs in which the hydrogen is bonded to the most electronegative elements (*e.g.* $F—H \cdots F^{\ominus}$, $\Delta H = -155$ kJ/mol). The greater strength of the hydrogen bond compared with non-specific dipole-dipole interactions is due to the much smaller size of the hydrogen atom relative to any other atom, which allows it to approach another dipole more closely. This simple dipole model accounts for the usual linear geometry of the hydrogen bond, because a linear arrangement maximizes the attractive forces and minimizes the repulsion.

However, there are reasons to believe that more is involved in hydrogen bonding than simply an exaggerated dipole-dipole interaction. The shortness of hydrogen bonds indicates considerable overlap of van der Waals radii and this should lead to repulsive forces unless otherwise compensated. Also, the existence of symmetrical hydrogen bonds of the type $F^{\delta\ominus} \cdots H \cdots F^{\delta\ominus}$ cannot be explained in terms of the electrostatic model. When the X—Y distance is sufficiently short, an overlap of the orbitals of the X—H bond and the electron pair of :Y can lead to a covalent interaction. According to Eq. (2-8), this situation can be described by two contributing "protomeric" structures, which differ only in the position of the proton*).

$$R\text{-}X\text{-}H\cdots Y\text{-}R' \quad \overset{p}{\longleftrightarrow} \quad R\text{-}X^{\ominus}\cdots H\text{-}Y^{\oplus}\text{-}R' \tag{2-8}$$

* The term "protomeric structure" was obviously introduced in analogy to the well-known "mesomeric structures", which are used to describe the electronic ground state of aromatic compounds such as benzene in terms of a resonance hybrid [323].

The approximate quantum mechanical description of proton states by linear combination of these protomeric structures has been called *protomerism* (symbol *p*) [323, 324]. It seems to be applicable to hydrogen bond systems in which a proton transfer may occur between two potential minima of equal depth [323, 324].

Solvents containing proton-donor groups are designated *protic* solvents [36] or *HBD* solvents [196]; solvents containing proton-acceptor groups are called *HBA* solvents [196]. The abbreviations *HBD* (hydrogen-bond donor) and *HBA* (hydrogen-bond acceptor) refer to donation and acceptance of the proton, and not to the electron pair involved in hydrogen bonding.

Solvents without proton-donor groups have been designated *aprotic* solvents [36]. However, this term is rather misleading, since, for example, solvents commonly referred to as *dipolar aprotic* (*e.g.* CH_3SOCH_3, CH_3CN, CH_3NO_2) are in fact *not* aprotic. In reactions where strong bases are employed, their protic character can be recognized. Therefore, the term *aprotic* solvents should be replaced by *nonhydroxylic* or better still by *non-HBD* solvents [197].

Typical protic or HBD solvents are water, ammonia, alcohols, carboxylic acids, and primary amides. Typical HBA solvents are amines, ethers, ketones, and sulfoxides. *Amphiprotic* solvents can act both as HBD and as HBA solvents simultaneously (*e.g.* water, alcohols, amides; *cf.* Fig. 2-4).

In *type-A* hydrogen bonding, the solute acts as a HBA-base and the solvent as a HBD-acid; in *type-B* hydrogen bonding, the roles are reversed [196].

Hydrogen bonding is responsible for the strong, temperature-dependent self- and hetero-association of amphiprotic solvents (*e.g.* water, alcohols, amides).

The molecular structure of binary HBD/HBA solvent mixtures is largely determined by intermolecular hydrogen bonding between the two components, which usually leads to pronounced deviations from ideal solution behaviour [306, 325–327]. Representative examples are trichloromethane/acetone [326] and trichloromethane/dimethyl sulfoxide mixtures [327], which readily form hydrogen-bonded 1:1 and 2:1 complexes, respectively, with distinct changes in their physical properties as a consequence.

Hydrogen bonding plays a particularly important role in the interactions between anions and HBD solvents. Hence, HBD solvents are good anion solvators. Due to the small size of the hydrogen atom, small anions like $F^{\ominus}$, $Cl^{\ominus}$, or $HO^{\ominus}$ are more effectively solvated by such solvents than the larger ones, *e.g.* $I_3^{\ominus}$, $I^{\ominus}$, $SCN^{\ominus}$, or the picrate ion [36]. This is also one of the reasons why the Gibbs energy of hydration, ΔG_{solv}, of the halide ions decreases in the series $F^{\ominus} > Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ [49].

Hydrogen bonding is of paramount importance for the stabilization and the shape of large biological molecules in living organisms (*e.g.* cellulose, proteins, nucleic acids). For instance, the anaesthetic properties of some halogen-containing solvents such as chloroform, halothane (CF_3—$CHClBr$), and methoxyflurane (CH_3O—CF_2—$CHCl_2$) have been connected with their ability to hinder the formation of biologically important hydrogen bonds. This is shown in the following equilibrium [300]:



Halohydrocarbon solvents containing an acidic C—H bond shift this equilibrium in favour of free or less associated species, thus perturbing the ion channels which determine the permeability of neuron membranes to $K^{\oplus}/Na^{\oplus}$ ions in the nervous system. Hydrogen bonds play a decisive role in determining the structure and dimension of these ion channels, on which this permeability depends [300].

Hydrogen-bonding also seems to be the molecular basis of sweetness. All sweet compounds seemingly have a H-bond donor and a H-bond acceptor ca. 250 . . . 400 pm apart, which can form hydrogen bonds with a complementary pair on the sweet receptor in the tastebuds of the tongue [328].

The effectiveness of solvents (and solutes) as hydrogen-bond donors and/or acceptors has been studied experimentally using suitable reference compounds, comprising representative HBDs or HBAs, in order to construct quantitative scales of solvent (and solute) hydrogen-bond acidity and hydrogen-bond basicity, respectively. For reviews on their construction and application to physicochemical and biochemical processes, see references [329–334] as well as Chapter 7. Scales of hydrogen-bond acidity and basicity have mostly been set up using complex formation constants, as determined in inert solvents [329–332]. For example, the strength of HBAs has been measured from the Gibbs energy change ΔG_{HB} for the formation of 1:1 hydrogen-bonded complexes between all kinds of HBAs (bases) and the reference HBD 4-fluorophenol in tetrachloromethane at 25 °C [331, 332]. Other attempts to construct scales of HBD and HBA strengths, *e.g.* the α and β scale of Taft and Kamlet [333, 334], are described in Chapter 7. Not unexpectedly, the pK_{HB} scales derived in this way do not correspond to the common pK_a and pK_b scales, *i.e.* to the normal acidity or basicity constants.

2.2.6 Electron-Pair Donor/Electron-Pair Acceptor Interactions (EPD/EPA Interactions) [50–59, 59a, 59b]

When tetrachloromethane solutions of yellow chloranil and colourless hexamethylbenzene are mixed, an intensely red solution is formed ($\lambda_{max} = 517$ nm [50]). This is due to the formation of a complex between the two components, and is only one example of a large number of so-called *electron-pair donor/electron-pair acceptor complexes (EPD/EPA complexes)**). It is generally accepted that the characteristic long-wavelength absorptions of these EPD/EPA complexes are associated with an electron transfer from the donor to the acceptor molecule. Mulliken termed these absorptions "charge-transfer (CT) absorptions" [51].

A necessary condition for the formation of an additional bonding interaction between two valency-saturated molecules is the presence of an occupied molecular

* Synonyms for *EPD/EPA complex* are *electron donor acceptor (EDA) complex* [50], *molecular complex* [57, 58], and *charge-transfer (CT) complex* [51]. Since normally the term *molecular complex* is only used for weak complexes between neutral molecules, and the appearance of a charge-transfer absorption band does not necessarily prove the existence of a stable complex, the more general expression *EPD/EPA complex*, proposed by Gutmann [53], will be used here. This will comprise all complexes whose formation is due to an interaction between electron-pair donors (Lewis bases) and electron-pair acceptors (Lewis acids), irrespective of the stabilities of the complexes or the charges of the components.

orbital of sufficiently high energy in the EPD molecule, and the presence of a sufficiently low unoccupied orbital in the EPA molecule*). Based on the type of orbitals involved in bonding interactions, all EPD molecules can be divided into three groups [51, 53]: *n*-, σ-, and π-EPD. In the first group, the energetically highest orbital is that of the lone pair of *n*-electrons on the heteroatoms (R_2O, R_3N, R_2SO), in the second it is that of the electron pair of a σ-bond (R—Hal, cyclopropane), and in the third it is that of the pair of π electrons of unsaturated and aromatic compounds (alkenes, alkylbenzenes, polycyclic aromatics). Similarly, EPA molecules can also be divided into three groups [51, 53]: *v*-, σ-, and π-EPA. The lowest orbital in the first group is a vacant valency-orbital of a metal atom ($Ag^{\oplus}$, certain organometallic compounds), in the second it is a non-bonding σ-orbital (I_2, Br_2, ICl), and in the third it is a system of π-bonds (aromatic and unsaturated compounds with electron-withdrawing substituents such as aromatic polynitro compounds, halobenzoquinones, tetracyanoethene). Because, in principle, any donor is able to form a complex with any acceptor, there exist nine different types of EPD/EPA complexes. The largest number of investigations have been concerned with complexes of the type π-EPD/π-EPA (*cf.* the above-mentioned hexamethylbenzene/chloranil complex) and π-EPD/σ-EPA (*cf.* complexes of aromatic hydrocarbons and alkenes with halogens and interhalogens).

More recent interesting examples of π-EPD/π-EPA complexes can be found in references [335, 336] and of π-EPD/*v*-EPA complexes (*i.e.* π/cation interactions) in references [337, 338]. For the synthesis of the first free, non-coordinated silyl cation in solution [*i.e.* trimesitylsilylium tetrakis(pentafluorophenyl)borate], the careful selection of a *non-coordinating* solvent, which nevertheless dissolves educts and product, was of crucial importance. Only with arenes as weak EPD solvents, bulky substituents around the silicon atom, and a weak nucleophilic anion, was the synthesis of $(Mes)_3Si^+$ $(F_5C_6)_4B^-$ in solution possible [338].

The reaction enthalpies, ΔH, for the formation of strong EPD/EPA complexes, often used as a measure of the bond energies, lie between -42 and -188 kJ/mol (-10 to -45 kcal/mol) [59]. *n*-EPD/*v*-EPA complexes are particular members of this group (*e.g.* Et_2O—BF_3, $\Delta H = -50$ kJ/mol or -11.9 kcal/mol [60]). For weak complexes, ΔH is usually larger than the dispersion energies but smaller than about 42 kJ/mol (10 kcal/mol) [59]. π-EPD/π-EPA complexes between neutral molecules are examples ($-\Delta H = 0 \ldots 21$ kJ/mol or $0 \ldots 5$ kcal/mol), *e.g.* benzene/1,3,5-trinitrobenzene ($\Delta H = -8$ kJ/mol or -1.9 kcal/mol [57]).

No general agreement exists as to the relative importance of the different intermolecular forces in making up the EPD/EPA complexes. According to Mulliken's VB description of weak EPD/EPA complexes, the electronic ground state can be considered as a hybrid of two limiting structures (*a*) and (*b*) in Fig. 2-5.

The non-ionic structure (*a*) represents a state without any donor-acceptor interactions, in which only non-specific intermolecular forces hold D and A together. The mesomeric structure (*b*) characterizes a state in which an ionic bond has been formed by

* The fundamental difference between this EPD/EPA bonding interaction and a normal chemical bond is that in an ordinary chemical bond each atom supplies one electron to the bond, whereas in EPD/EPA bonding one molecule (the donor) supplies the pair of electrons, while the second molecule (the acceptor) provides the vacant molecular orbital.



Fig. 2-5. Formation and optical excitation of an EPD/EPA complex between donor D and acceptor A (the predominating mesomeric structure in the ground and excited states is underlined).

transfer of an electron from D to A. This electron transfer will be easier the lower the ionization potential of the donor [61, 63], and the higher the electron affinity of the acceptor [62, 63]. The ionic limiting structure (*b*) is relatively energy-rich and contributes only slightly to the ground state. Nevertheless, this small contribution is sufficient in establishing an extra bonding interaction in addition to the non-specific van der Waals forces. However, subsequent investigations showed that these charge-transfer forces are weaker than was previously believed, and that the classical van der Waals forces (including electrostatic forces) suffice in explaining the stabilities of EPD/EPA complexes [59, 64, 198]. The relative importance of contributions from the electrostatic and charge-transfer forces in the ground state of EPD/EPA complexes has been studied by many authors. For a review, see reference [183; Vol. 1, p. 6ff.]. It seems that both electrostatic and charge-transfer interactions are important in the ground state of EPD/EPA complexes. Their relative contribution, however, varies widely in different EPD/EPA complexes [183].

Another description of EPD/EPA interactions, particularly useful for strong complexes, is based on the coordinative interaction between Lewis bases or nucleophiles (as EPD) and Lewis acids or electrophiles (as EPA) [53, 58]. The intermolecular bonding is seen not as a hybrid of electrostatic and charge-transfer forces, but as one of electrostatic and covalent ones. The interaction of the acceptor A with the electron pair of the donor D is a result of an overlap of the orbitals of the two molecules; consequently, a finite electron density is created between the two partners according to Eq. (2-9).



(2-9)

Hence, the structure $D^{\oplus}—A^{\ominus}$ is a covalent one and the EPD/EPA interaction between D and A can be described as a Lewis acid/base interaction [65].

Of the solvents, aromatic and olefinic hydrocarbons are π-donors (π-EPD); alcohols, ethers, amines, carboxamides, nitriles, ketones, sulfoxides and *N*- and *P*-oxides are *n*-donors (*n*-EPD), and haloalkanes are σ-donors (σ-EPD). Boron and antimony trihalides are acceptor solvents (*v*-EPA), as are halogens and mixed halogens (σ-EPA), and liquid sulfur dioxide (π-EPA). In principle, all solvents are amphoteric in this respect, *i.e.* they may act as a donor (nucleophile) and an acceptor (electrophile) simultaneously. For example, water can act as a donor (by means of the oxygen atom) as well as as an acceptor (by forming hydrogen bonds). This is one of the reasons for the exceptional importance of water as a solvent.

n-Donor solvents are particularly important for the solvation of cations. Examples are hexamethylphosphoric triamide, pyridine, dimethyl sulfoxide, *N*,*N*-dimethylformamide, acetone, methanol, and water. Their specific EPD properties make them excellent cation solvators, and they are, therefore, good solvents for salts. They are

also known as *coordinating solvents* [66]. The majority of inorganic reactions are carried out in coordinating solvents.

An empirical semiquantitative measure of the nucleophilic properties of EPD solvents is provided by the so-called *donor number DN* (or *donicity*) of Gutmann [53, 67] (*cf.* also Section 7.2). This donor number has been defined as the negative ΔH values for 1:1 adduct formation between antimony pentachloride and electron-pair donor solvents (D) in dilute solution in the non-coordinating solvent 1,2-dichloroethane, according to Eq. (2-10)*).

$$D: + SbCl_5 \underset{\text{in } Cl\text{-}CH_2CH_2\text{-}Cl}{\overset{\text{room temp.}}{\rightleftharpoons}} \overset{\oplus}{D}—\overset{\ominus}{SbCl_5} \qquad (2\text{-}10)$$

Solvent Donor Number $DN = -\Delta H_{D\text{–}SbCl_5}/(\text{kcal} \cdot \text{mol}^{-1})$

The linear relationship between $-\Delta H_{D—SbCl_5}$ and the logarithm of the corresponding equilibrium constant ($\lg K_{D—SbCl_5}$) shows that the entropy contributions are equal for all the studied acceptor/donor solvent reactions. Therefore, one is justified in considering the donor numbers as semiquantitative expressions for the degree of coordination interaction between EPD solvents and antimony pentachloride. Antimony pentachloride is regarded as an acceptor on the borderline between hard and soft Lewis acids. A list of organic solvents ordered according to increasing donicity is given in Table 2-3. From this it is seen that, for example, nitromethane and acetonitrile are weak donor solvents, whereas dimethyl sulfoxide and triethylamine are very strong donors. The higher the donor number, the stronger the interaction between solvent and acceptor.

Unfortunately, donor numbers have been defined in the non-SI unit kcal $\cdot$ mol^{-1}. Marcus has presented a scale of dimensionless, normalized donor numbers DN^N, which are defined according to $DN^N = DN/(38.8 \text{ kcal} \cdot \text{mol}^{-1})$ [200]. The non-donor solvent 1,2-dichloroethane ($DN = DN^N = 0.0$) and the strong donor solvent hexamethylphosphoric triamide (HMPT: $DN = 38.8$ kcal $\cdot$ mol^{-1}; $DN^N = 1.0$) have been used to define the scale. Although solvents with higher donicity than HMPT are known (*cf.* Table 2-3), it is expedient to choose the solvent with the highest directly (*i.e.* calorimetrically) determined *DN* value so far as the second reference solvent [200]**). The DN^N values are included in Table 2-3.

A visual estimate of the different donicities of EPD solvents can easily be made using the colour reaction with copper(II), nickel(II), or vanadyl(IV) complexes as acceptor solutes [204].

The donor number has proven very useful in coordination chemistry, since it can be correlated with other physical observables for such reactions, *e.g.* thermodynamic

* An analogous approach was first used by Lindqvist and Zackrisson [67a]. The authors established a series of EPD solvents calorimetrically, based on their increasing donor capacities relative to a standard acceptor ($SbCl_5$ or $SnCl_4$) with which the given donor was combined in 1,2-dichloroethane.

** The donor number of 38.8 kcal $\cdot$ mol^{-1} for HMPT was given by Gutmann [67]. It should be mentioned, however, that a much higher *DN* value of 50.3 kcal $\cdot$ mol^{-1} was subsequently measured for this solvent by Bollinger *et al.* [214]. This shows that serious problems arise in measuring the Lewis basicity of this EPD solvent towards $SbCl_5$.

Table 2-3. Donor numbers (donicities) *DN* [199, 200, 212, 241, 339] and normalized DN^N values [200] of a selection of thirty-six organic EPD solvents[a], determined calorimetrically in dilute 1,2-dichloroethane solutions at room temperature and valid for isolated EPD solvent molecules[b].

Solvents	$DN/(\text{kcal} \cdot \text{mol}^{-1})$[c]	DN^N[d]
1,2-Dichloroethane *(reference solvent)*	*0.0*[e]	*0.00*[e]
Nitromethane	2.7	0.07
Nitrobenzene	4.4	0.11
Acetic anhydride	10.5	0.27
Cyanobenzene, Benzonitrile	11.9	0.31
Ethanenitrile, Acetonitrile	14.1	0.36
Tetrahydrothiophene-1,1-dioxide, Sulfolane	14.8	0.38
1,4-Dioxane	14.8	0.38
4-Methyl-1,3-dioxol-2-one, Propylene carbonate	15.1	0.39
(Cyanomethyl)benzene, Benzylcyanide	15.1	0.39
2-Methylpropanenitrile, *i*-Butanenitrile	15.4	0.40
Diethyl carbonate	16.0	0.41
Propanenitrile	16.1	0.41
Methyl acetate	16.3	0.42
1,3-Dioxol-2-one, Ethylene carbonate	16.4	0.42
Butanenitrile	16.6	0.43
3,3-Dimethyl-2-butanone, *t*-Butyl methyl ketone	17.0	0.44
Acetone	17.0	0.44
Ethyl acetate	17.1	0.44
3-Methyl-2-butanone, Methyl *i*-propyl ketone	17.1	0.44
2-Butanone	17.4	0.45
Diethyl ether	19.2	0.49
Tetrahydrofuran	20.0	0.52
Trimethyl phosphate	23.0	0.59
Tri-*n*-butyl phosphate	23.7	0.61
N,*N*-Dimethylformamide	26.6	0.69
1-Methylpyrrolidin-2-one	27.3	0.70
N,*N*-Dimethylacetamide	27.8	0.72
Tetramethylurea	29.6	0.76
Dimethyl sulfoxide	29.8	0.77
N,*N*-Diethylformamide	30.9	0.80
N,*N*-Diethylacetamide	32.2	0.83
Pyridine	33.1	0.85
Hexamethylphosphoric triamide	38.8	*1.00*[f]
Triethylamine	61.0	1.57

[a] A compilation of 170 resp. 134 *DN*-values taken from different sources can be found in references [200, 339]. Further 14 *DN* values, determined indirectly from the ^{1}H NMR shift of chloroform, are given in reference [293].

[b] As the basic donor numbers were measured in an inert diluent, they reflect the donicity of the isolated EPD solvent molecules. In neat, associated EPD solvents an increase in the donicity should occur [199]. For such highly-structured solvents (*e.g.* water, alcohols, amines) the term *bulk donicity* has been introduced [201] in order to rationalize the deviations of these solvents in plots of $^{23}Na^{\oplus}$ NMR shifts [202] and ESR parameters [203] *vs.* the donor numbers. Because of the great discrepancies which exist between the DN_{bulk} values given in the literature, they are not included in this table. For a collection of bulk donicities, DN_{bulk}, see reference [200], Table II.

[c] For the definition of *DN cf.* Eq. (2-10). For conversion into SI units: $1\ \text{kcal} \cdot \text{mol}^{-1} = 4.184\ \text{kJ} \cdot \text{mol}^{-1}$.

[d] $DN^N = DN/(38.8\ \text{kcal} \cdot \text{mol}^{-1})$ [200]; $DN = 38.8\ \text{kcal} \cdot \text{mol}^{-}$ for hexamethylphosphoric triamide as reference solvent.

[e] Zero by definition.

[f] Unity by definition [200].

(ΔG or K), kinetic (rates), electrochemical (polarographic half-wave and redox potentials), and spectroscopic data (chemical shifts of NMR signals) [53, 67–69, 205–207].

The donor number approach has been criticized for conceptual [208] and experimental reasons [200, 209–212]. For this and other reasons, other Lewis basicity parameters have been sought.

Another remarkable Lewis basicity scale for 75 non-HBD solvents has been established by Gal and Maria [211, 212]. This involved very precise calorimetric measurements of the standard molar enthalpies of 1:1 adduct formation of EPD solvents with gaseous boron trifluoride, $\Delta H^{\circ}_{D—BF_3}$, in dilute dichloromethane solution at 25 °C, according to Eq. (2-10a).



(2-10a)

A selection of $\Delta H^{\circ}_{D—BF_3}$ values is given in Table 2-4. This new Lewis basicity scale is more comprehensive and seems to be more reliable than the donor number scale. A comparison of various Lewis basicity scales has been given by Persson [301].

Persson, Sandström, and Goggin have proposed an empirical solvent scale, called the D_S scale, ranking the donor strength of 64 EPD solvents towards a soft acceptor

Table 2-4. Molar enthalpies of complex formation between boron trifluoride and several non-HBD solvents, determined in dichloromethane at 25 °C, according to Eq. (2-10a) [211, 212].

Solvents	$-\Delta H^{\circ}_{D-BF_3}/(kJ \cdot mol^{-1})$[a]
Dichloromethane	10.0
Nitrobenzene	35.79
Nitromethane	37.63
Tetrahydrothiophene-1,1-dioxide	51.32
Acetonitrile	60.39
Propylene carbonate	64.19
3-Pentanone	72.28
1,4-Dioxane	74.09
Ethyl acetate	75.55
Acetone	76.03
Di-*i*-propyl ether	76.61
Diethyl ether	78.77
Tetrahydrofuran	90.40
1,3-Dimethylimidazolidin-2-one, DMEU	98.93
Dimethyl sulfoxide	105.34
N,*N*,*N*′,*N*′-Tetramethylurea	108.62
N,*N*-Dimethylformamide	110.49
3,4,5,6-Tetrahydro-1,3-dimethylpyrimidin-2(1 *H*)-one, DMPU	112.13
1-Methylpyrrolidin-2-one	112.56
Hexamethylphosphoric triamide	117.53
Tris(pyrrolidino)phosphane oxide	122.52
Pyridine	128.08
Triethylamine	135.87
1-Methylpyrrolidine	139.51

[a] See reference [212] for a set of 75 $\Delta H^{\circ}_{D—BF_3}$ values. At present, $\Delta H^{\circ}_{D—BF_3}$ values for *ca.* 350 organic EPD compounds are known (J.-F. Gal and P.-C. Maria, private communication).

such as mercury(II) bromide [303]. The D_S values correspond to the Raman wavenumber shift of the symmetric IR stretching vibration on going from the gas phase to a solution of $HgBr_2$. Further measurements of $\Delta\nu$(Hg—Br) of $HgBr_2$ and the relationships between the corresponding D_S values and other soft EPD solvent parameters can be found in reference [340]. An additional D_H scale of donor strength towards hard acceptors (*e.g.* $Na^{\oplus}$) has been derived for 24 EPD solvents [303].

An analogous empirical quantity for characterizing the electrophilic properties of EPA solvents has been derived by Gutmann and coworkers from the ^{31}P NMR chemical shifts produced by the electrophilic actions of acceptor solvents A in triethylphosphane oxide, according to Eq. (2-11) (*cf.* also Section 7.4) [70, 199, 207, 213].

$$\{Et_3P{=}O \longleftrightarrow Et_3\overset{\oplus}{P}-\overset{\ominus}{O}\} + A \rightleftharpoons Et_3\overset{\delta\oplus}{P}\cdots\overset{\delta\ominus}{O}-A \tag{2-11}$$

$$AN = \frac{\delta_{corr}(A) - \delta_{corr}(n\text{-}C_6H_{14})}{\delta_{corr}(Et_3PO{—}SbCl_5) - \delta_{corr}(n\text{-}C_6H_{14})} \cdot 100 = \Delta\delta_{corr} \cdot 2.348/\text{ppm}$$

These quantities have been termed *acceptor number AN* (or *acceptivity*) and they were obtained from the relative ^{31}P NMR chemical shift values δ_{corr} (*n*-hexane as reference solvent) with respect to that of the 1:1 adduct $Et_3PO{—}SbCl_5$ dissolved in 1,2-dichloroethane, which has been arbitrarily taken to have the value of 100. The acceptor numbers are dimensionless numbers expressing the acceptor property of a given solvent relative to those of $SbCl_5$, which is also the reference compound for assessing the donor numbers. A compilation of organic solvents in order of increasing acceptor number is given in Table 2-5.

Acceptor numbers are less than 10 for nonpolar non-HBD solvents, they vary between about 10 . . . 20 for dipolar non-HBD solvents, and they cover a wide range of about 25 . . . 105 for protic solvents (*cf.* Table 2-5). Surprisingly, benzene and tetrachloromethane have stronger electrophilic properties than diethyl ether and tetrahydrofuran. Acceptor numbers are also known for binary solvent mixtures [70, 213].

Using the neutral Fe(II) complex $[Fe(phen)_2(CN)_2]$, the different Lewis acidities of EPA solvents can easily be visualized by its colour change: solutions of this Fe(II) complex are blue in HMPT, violet in dichloromethane, red in ethanol, and yellow in trifluoroacetic acid [204].

Another approach to the estimation of EPD/EPA interactions between a Lewis acid A and a Lewis base B was given by Drago [71]. Drago proposed the four-parameter Eq. (2-12) to correlate the standard enthalpy of the reaction of an acceptor A with a donor B to give a neutral 1:1 adduct in an inert solvent (tetrachloromethane or *n*-hexane).

$$-\Delta H^{\circ}_{AB}/(\text{kJ} \cdot \text{mol}^{-1}) = E_A \cdot E_B + C_A \cdot C_B \tag{2-12}$$

E_A and C_A are empirical acceptor parameters and E_B and C_B are empirical donor parameters. The E parameters are measures of the tendency of an acid or a base to participate in *e*lectrostatic interactions, while the C parameters are measures of their tendency to form *c*ovalent bonds.

Table 2-5. Acceptor numbers (acceptivities) *AN* [70, 213, 339] of forty-eight organic EPA solvents, determined ^{31}P-NMR spectroscopically at 25 °C.

Solvents	*AN*[a)]
n-Hexane *(reference solvent)*	*0.0*
Triethylamine	1.4
Diethyl ether	3.9
Tetrahydrofuran	8.0
Benzene	8.2
Tetrachloromethane	8.6
Ethyl acetate	9.3
Diethylamine	9.4
Hexamethylphosphoric acid triamide	9.8
Tri-*n*-butyl phosphate	9.9
Diethylene glycol dimethyl ether	9.9
1,2-Dimethoxyethane	10.2
Methyl acetate	10.7
1,4-Dioxane	10.8
Acetone	12.5
1-Methylpyrrolidin-2-one	13.3
N,*N*-Dimethylacetamide	13.6
Pyridine	14.2
Nitrobenzene	14.8
Cyanobenzene	15.5
N,*N*-Dimethylformamide	16.0
Trimethyl phosphate	16.3
1,2-Dichloroethane	16.7
4-Butyrolactone	17.3
Morpholine	17.5
4-Methyl-1,3-dioxol-2-one, Propylene carbonate	18.3
N,*N*-Dimethylthioformamide	18.8
Ethanenitrile, Acetonitrile	18.9
Tetrahydrothiophene-1,1-dioxide, Sulfolane	19.2
Dimethyl sulfoxide	19.3
Dichloromethane	20.4
Nitromethane	20.5
1,2-Diaminoethane	20.9
Chloroform	23.1
2-Methyl-2-propanol, *t*-Butanol	27.1
N-Methylformamide	32.1
1-Butanol	32.2
2-Propanol	33.5
1-Propanol	33.7
2-Aminoethanol	33.7
Ethanol	37.1
Formamide	39.8
Methanol	41.5
Acetic acid	52.9
2,2,2-Trifluoroethanol	53.8
Water	54.8
Formic acid	83.6
$Et_3PO \cdot SbCl_5$ *in 1,2-dichloroethane as reference compound*	*100.0*
Trifluoroacetic acid	105.3

[a)] For the definition of *AN*, see Eq. (2-11). All δ values have been extrapolated to zero concentration and corrected for differences in volume susceptibilities.

Table 2-6. Some E and C parameters expressing Lewis acid/base strength according to Drago [217][a]; *cf.* Eq. (2-12).

Lewis acids	E_A	C_A	Lewis bases	E_B	C_B
$SbCl_5$	14.4[b]	1.17[b]	$[(CH_3)_2N]_3PO$	1.52	3.55
BF_3 (g)	9.88	1.62	CH_3SOCH_3	1.34	2.85
$(CF_3)_2CHOH$	5.93	0.62	$CH_3CON(CH_3)_2$	*1.32*[c]	2.58
C_6H_5OH	4.33	0.42	C_5H_5N	1.17	6.40
CF_3CH_2OH	3.88	0.45	$CH_3CO_2C_2H_5$	0.975	1.74
$CHCl_3$	3.02	0.16	CH_3COCH_3	0.94	2.33
$(CH_3)_3C—OH$	2.04	0.30	$(C_2H_5)_2O$	0.94	3.25
H_2O	1.64	0.57	CH_3CN	0.89	1.34
I_2	*1.00*[c]	*1.00*[c]	$(C_2H_5)_2S$	0.34	*7.40*[c]
SO_2	0.92	0.81	C_6H_6	0.28	0.59

[a] For a more complete list see references [71, 215, 217].
[b] Corrected values; see reference [217].
[c] Used to define the E/C scale.

The original set of E and C parameters was determined mainly with the help of enthalpies of adduct formation of iodine and phenol as acceptors with alkylamines as donors. Subsequently, the best set of E and C parameters has been obtained by computer optimization of a large data base of enthalpies and four arbitrarily fixed reference values [71, 215]: $E_A = C_A = 1$ for iodine, $E_B = 1.32$ for *N*,*N*-dimethylacetamide, and $C_B = 7.40$ for diethyl sulfane. Table 2-6 gives a selection of E and C parameters for Lewis acids and bases commonly used as solvents.

On the basis of these parameters, it is possible to predict the enthalpies of Lewis acid/base reactions, even those reactions which might be inaccessible experimentally, with remarkable accuracy (within ± 0.8 kJ $\cdot$ mol^{-1}) [216].

Drago's E/C analysis and Gutmann's donor/acceptor approach [53, 67] have been compared [200, 217, 218]. Eq. (2-12) has been extended for specific and nonspecific interactions between solutes and polar solvents [219]. Various Lewis acidity and basicity scales for polar solvents have been examined and compared by Fawcett, who concluded that the donor/acceptor scales of Gutmann seem to be the most appropriate [341].

Finally, an attempt was made to establish a measure of the electron-donating and electron-accepting power of organic solvents by means of infrared [72, 73] and ^{1}H NMR measurements [73]. Further empirical Lewis acid and base parameters will be discussed in Chapters 7.2 ... 7.5. A thorough and critical compilation of empirical solvent scales, including Lewis acidity and basicity scales, has recently been made for non-HBD solvents [342].

2.2.7 Solvophobic Interactions [74–77, 176, 220–225]

Hydrocarbons have extremely low solubilities in water. Accordingly, the dissolution of a hydrocarbon in water is usually associated with an increase in the Gibbs energy G of the system ($\Delta G > 0$). Since it is known experimentally that the dissolution of a hydrocarbon



Fig. 2-6. The formation of a hydrophobic interaction between two hydrocarbon molecules A and B (the circles represent water molecules) [78].

in water is exothermic ($\Delta H < 0$) it follows from $\Delta G = \Delta H - T \cdot \Delta S$ that the entropy of the system must decrease. This can be interpreted as a consequence of the highly ordered structure of the water molecules around the dissolved hydrocarbon molecules. In other words, the water molecules are more tightly packed around the dissolved hydrocarbon molecules than in pure water. This is called a structure increase. If aqueous solutions of two hydrocarbons are mixed, the two hydrocarbons may form an aggregate with simultaneous partial reconstruction of the original undisturbed water structure. This is shown schematically in Fig. 2-6.

Due to the contact between A and B, fewer water molecules are now in direct contact with the hydrocarbon molecules. Thus, the ordering influence of the hydrophobic molecules will be diminished and the entropy increases ($\Delta S > 0$). Although thermal energy is required for the destructuring of the hydration shells around A and B ($\Delta H > 0$), the free energy diminishes upon aggregation ($\Delta G < 0$). Therefore, it is energetically advantageous for apolar molecules, or apolar groups in otherwise polar molecules, when dissolved in water, to aggregate with expulsion of water molecules from their hydration shells. In order to minimise the unfavourable solute/water interactions, the apolar solute molecules (or apolar groups) will interact preferentially, thus reducing the number of their water contacts [176]. This effect has been called *hydrophobic interaction.**) The water molecules around an inert apolar solute have a higher coordination and are thus more ordered than in the bulk liquid, which is entropically unfavourable. The aggregation of apolar solutes as shown in Fig. 2-6 releases water molecules into the bulk water, which is entropically very favourable.

This hydrophobic interaction can be illustrated by considering the thermodynamic parameters for the dissolution of the archetypal apolar hydrocarbon methane in cyclohexane (an apolar, non-associated solvent) and in water (a polar, strongly self-associated solvent); Table 2-7 [225].

The unfavorable Gibbs energy ($\Delta G_s^\circ \gg 0$) for the dissolution of methane in water is the result of a strongly negative entropy of solution ($\Delta S_s^\circ \ll 0$), which prevails over

* Glass beads can be used as an illustration of hydrophobia interactions. Thus, glass beads covered with dichloro-dimethylsilane can be regarded as solid hydrocarbon particles. Only hydrophobic interactions are possible. In a structured solvent such as water or formamide, the beads cluster together. When the polarity of the solvent is decreased by addition of alcohols the clusters disintegrate [79].

Table 2-7. Thermodynamic parameters for dissolution of gaseous methane in cyclohexane and water at 25 °C [225].

Solvents	$\Delta G_s^\circ/(kJ \cdot mol^{-1})$	$\Delta H_s^\circ/(kJ \cdot mol^{-1})$	$\Delta S_s^\circ/(J \cdot mol^{-1} \cdot K^{-1})$
Water	25.5	−13.8	−132
Cyclohexane	14.2	−3.0	−58

the favorable enthalpic contribution ($\Delta H_s^\circ < 0$). The negative enthalpy and entropy of transfer of methane from cyclohexane to water can be interpreted in terms of an increased degree of water-water hydrogen bonding in the solvation shell surrounding the apolar solute molecule.

Generally, the introduction of apolar molecules (such as hydrocarbons or noble gases), or apolar residues in otherwise polar molecules (such as alkyl side chains in biopolymers) into water leads to a reduction of the degrees of freedom (spatial, orientational, dynamic) of the neighbouring water molecules. This effect is called the *hydrophobic effect* or *hydrophobic hydration* [176]. *Hydrophobic* means 'water-fearing'. It should be noted that the interaction between hydrophobic molecules and water molecules is actually attractive because of the dispersion interactions. However, the water/water interaction is much more attractive. Water molecules simply love themselves too much to let some other compounds get in the way [26b]! Therefore, from the point of view of the water molecules, the term "hydrophobic" is rather a misnomer; it would be better to refer to water as being "lipophobic".

This hydrophobic hydration was first postulated by Frank and Evans in 1945. They wrote: "The nature of deviation found for non-polar solutes in water leads to the idea that the water forms frozen patches or microscopic icebergs around such solute molecules. The word 'iceberg' represents a microscopic region, surrounding the solute molecule, in which water molecules are tied together in some sort of quasi-solid structure" [226].

The model of "icebergs" around nonpolar solute molecules in aqueous solution is clearly not a very realistic one. However, if solutions of hydrocarbons (or noble gases) are cooled, then the solid phase that sometimes separates out consists of a so-called gas hydrate (clathrate), in which water provides a particular kind of hydrogen-bonded framework containing cages that are occupied by the nonpolar solute molecules. Obviously, such gas hydrates (clathrates) represent more realistic models for the phenomenon of hydrophobic hydration [176].

In principle, such interactions should also apply to other solvents resembling water, and therefore the more general term *solvophobic interactions* has been proposed [80, 343]. In fact, analogous water-like behaviour has been observed with self-associated solvents other than water, *e.g.* ethanol [81], glycerol [82], ethylammonium nitrate [227], and some dipolar non-HBD solvents [228].

Although there is overwhelming experimental evidence that the hydrophobic interaction is "entropy-driven", this classical view is still a matter of debate [79a, 167, 227, 229–231, 343–347]. For example, it has been claimed that the major contribution to the hydrophobic interaction between the methylene groups of *n*-alkanes is an enthalpic and not an entropic effect [230]. In other words, the poor solubility of non-

polar solutes in water should be due to unfavourable enthalpy and not to unfavourable entropy [227, 231].

Furthermore, what is the real origin of hydrophobicity (or solvophobicity), that is, which molecular property of water is primarily responsible for the positive Gibbs energy of hydration of nonpolar solutes and their tendency to associate? The two main physical explanations are:

(a) the high cohesive pressure of water (see Table 3-2 in Section 3.2), caused by the strong hydrogen-bonding interactions compared to the weak interactions between water and nonpolar solutes; and/or
(b) the small size of the water molecules, which increases the entropic cost of opening up a cavity to accommodate the solute. Opening up a cavity for solute molecules is entropically unfavourable in any solvent. However, the small size of water molecules exacerbates this situation and gives rise to entropies more negative that in other solvents with larger molecule size.

For a more detailed discussion of these questions, see references [76, 77, 176, 343–347] and references cited therein. More recent results [346, 347] have shown that the classical view (a) seems to be basically correct. The essential condition for solvophobicity is that solvent/solvent interactions are much stronger than solute/solvent interactions. However, the solvophobic effect is not necessarily always an entropic phenomenon; it can be enthalpic or entropic depending on the temperature and the geometrical size of the solute molecules [346].

Hydrophobic interactions are important in the aggregation of polymethine dyes [81] and in the stabilization of particular conformations of polypeptides and proteins in aqueous solution [222, 232]. They also play an important role in the biochemical complexation between an enzyme and a substrate [77, 78, 83, 84, 348].

Hydrophobicity parameters for organic substituents have been developed by Hansch *et al.* using partitioning phenomena [296], and by Menger *et al.* using kinetic measurements (hydrolysis of long-chain esters) [297]; see Section 7.2. Further results connected with the presence of hydrophobic interactions in solutions are discussed in Sections 2.5 and 5.4.8.

2.3 Solvation [49, 85–98, 98a, 233–241]

The term *solvation* refers to the surrounding of each dissolved molecule or ion by a shell of more or less tightly bound solvent molecules. This solvent shell is the result of intermolecular forces between solute and solvent. For aqueous solutions the term used is *hydration*. Intermolecular interactions between solvent molecules and ions are particularly important in solutions of electrolytes, since ions exert specially strong forces on solvent molecules. Crude electrostatic calculations show that the field experienced by nearest neighbours of dissolved ions is $10^6 \ldots 10^7$ V/cm. Fig. 2-7 shows a highly simplified picture of such an interaction between ions and dipolar solvent molecules.

The *solvation energy* is considered as the change in Gibbs energy when an ion or molecule is transferred from a vacuum (or the gas phase) into a solvent. The Gibbs



Fig. 2-7. Solvation of ions in a solvent consisting of dipolar molecules [99]. The charges of the dipolar molecules are in fact partial charges $\delta\oplus$ and $\delta\ominus$.

energy of solvation, $\Delta G^{\circ}_{\text{solv}}$, a measure of the solvation ability of a particular solvent, is the result of a superimposition of four principal components of a different nature [100]:

(a) the cavitation energy associated with the hole that the dissolved molecule or ion produces in the solvent;
(b) the orientation energy corresponding to the phenomenon of partial orientation of the dipolar solvent molecules caused by the presence of the solvated molecule or ion (*cf.* Fig. 2-7);
(c) the isotropic interaction energy corresponding to the unspecific intermolecular forces with a long radius of activity (*i.e.* electrostatic, polarisation, and dispersion energy);
(d) the anisotropic interaction energy resulting from the specific formation of hydrogen bonds or electron-pair donor/electron-pair acceptor bonds at well localized points in the dissolved molecules.

The dissolution of a substance requires that not only the interaction energy of the solute molecules (for crystals the lattice energy*)) be overcome but also the interaction energy between the solvent molecules themselves. This is compensated by the gain in Gibbs energy of solvation, $\Delta G^{\circ}_{\text{solv}}$. The standard molar Gibbs energy of solvation, $\Delta G^{\circ}_{\text{solv}}$, can be formulated as the difference between the Gibbs energy of solution, $\Delta G^{\circ}_{\text{soln}}$,

* The lattice energy is the work required to separate to infinity the elements of the lattice from their equilibrium position at 0 K. For ionic lattices of the alkali halides it is of the order 628 ... 837 kJ/mol (150 ... 200 kcal/mol) [49]. For molecular lattices of organic compounds such as benzene, naphthalene, and anthracene it is of the order 42 ... 105 kJ/mol (10 ... 25 kcal/mol) [101]. The experimental heat of sublimation of benzene is 44.6 kJ/mol (10.7 kcal/mol) [102].



Fig. 2-8. The relationship between standard molar Gibbs energies of solvation and solution and the crystal lattice energy of an ionophore $A^{\oplus}B^{\ominus}$: $\Delta G^{\circ}_{solv} = \Delta G^{\circ}_{soln} - \Delta G^{\circ}_{latt}$.

and the crystal lattice energy, ΔG°_{latt}, as shown by means of the customary Born-Haber cycle in Fig. 2-8.

If the liberated solvation energy is higher than the lattice energy, then the overall process of dissolution is exothermic. In the opposite case the system uses energy and the dissolution is endothermic. The values for NaCl are typical: lattice energy +766 kJ/mol, hydration energy −761 kJ/mol, and energy of solution +3.8 kJ/mol. The energies of solution are generally small because interaction within the crystal lattice is energetically similar to interaction with the solvent.

The Gibbs energies of solvation of individual ions cannot be directly measured but they can be calculated [49]. The Gibbs energies of hydration of some representative ions are collected in Table 2-8. It can be seen that these values can be as high as bond energies or even higher (209 . . . 628 kJ/mol; 50 . . . 150 kcal/mol). Consequently, the solvent is often considered a direct reaction partner and should really be included in the reaction equation. The isolation of numerous solvates such as hydrates, alcoholates, etherates, and ammoniates, especially of inorganic or organometallic compounds, are examples. Between the two extremes, *viz*. the simple solvation resulting from weak

Table 2-8. Standard molar Gibbs energies of hydration, ΔG°_{hydr}, of some representative single ions at 25 °C [241, 242][a].

Cations	$\Delta G^{\circ}_{hydr}/(kJ \cdot mol^{-1})$	Anions	$\Delta G^{\circ}_{hydr}/(kJ \cdot mol^{-1})$
$H^{\oplus}$	−1056	$F^{\ominus}$	−472
$Li^{\oplus}$	−481	$Cl^{\ominus}$	−347
$Na^{\oplus}$	−375	$Br^{\ominus}$	−321
$K^{\oplus}$	−304	$I^{\ominus}$	−283
$Mg^{\oplus\oplus}$	−1838	$HO^{\ominus}$	−439
$Al^{\oplus\oplus\oplus}$	−4531	$SO_4^{\ominus\ominus}$	−1090

[a] For a comprehensive compilation of Gibbs energies of solvation, see C. M. Criss and M. Salomon: *Thermodynamic Measurements – Interpretation of Thermodynamic Data.* In A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London & New York 1973, p. 253 ff. – *Cf.* also D. W. Smith: *Ionic Hydration Enthalpies*, J. Chem. Educ. *54*, 540 (1977). – A critical selection of standard molar heat capacities of hydration, $\Delta_{hyd}C^{\circ}_p/(J \cdot K^{-1} \cdot mol^{-1})$, of single ions has been given by M. H. Abraham and Y. Marcus, J. Chem. Soc., Faraday Trans. I *82*, 3255 (1986).

intermolecular interactions, and the *bona fide* chemical modification of the substrate by the solvent, all other possibilities exist.

The most direct measure of the energetics of ion solvation is, without doubt, their *standard molar Gibbs energy of solvation*, *i.e.* transfer from the gas phase to the solvent (*cf.* Fig. 2-8). However, this quantity is generally unknown, particularly for ions in nonaqueous solvents. Therefore, $\Delta G^\circ_{\text{solv}}$ is advantageously replaced by the *standard molar Gibbs energy of transfer* of the ion X from water, W, as reference solvent, to another solvent, S, $\Delta G^\circ_t(\text{X}, \text{W} \rightarrow \text{S})$, as defined by Eq. (2-12a):

$$\Delta G^\circ_t(\text{X}, \text{W} \rightarrow \text{S}) = \mu^\infty_{\text{X}}(\text{in S}) - \mu^\infty_{\text{X}}(\text{in W}) = R \cdot T \cdot \ln {}^{\text{W}} y^{\text{S}}_{\text{X}} \tag{2-12a}$$

μ^∞_{X} is the standard (*i.e.* infinite dilution) chemical potential of X and ${}^{\text{W}} y^{\text{S}}_{\text{X}}$ the so-called *solvent-transfer activity coefficient* of X.

In order to obtain the $\Delta G^\circ_t(\text{X}, \text{W} \rightarrow \text{S})$ of individual ions from experimental data on complete electrolytes, the extrathermodynamic assumption that $\Delta G^\circ_t(\text{Ph}_4\text{As}^\oplus, \text{W} \rightarrow \text{S}) = \Delta G^\circ_t(\text{Ph}_4\text{B}^\ominus, \text{W} \rightarrow \text{S})$ for all solvents has been made, using $\text{Ph}_4\text{As}^\oplus\text{Ph}_4\text{B}^\ominus$ as reference electrolyte ($\text{Ph} = \text{C}_6\text{H}_5$). This seems reasonable because the large symmetrical ions of tetraphenylarsonium tetraphenylborate are of comparable size, structure, and charge, and are, therefore, similarly solvated on transfer from one solvent to another. Arguments in favour of and against this extrathermodynamic assumption have been reviewed [235, 241, 243, 244]*).

Experimentally, the molar Gibbs energy of transfer of an anion $\text{X}^\ominus$ is obtained from the combined results of four solubility measurements, namely of the salts $\text{Ph}_4\text{As}^\oplus\text{Ph}_4\text{B}^\ominus$ and $\text{Ph}_4\text{As}^\oplus\text{X}^\ominus$ in water, W, and of the same salts in the solvent S. The Gibbs energy of transfer is then:

$$\begin{aligned}\Delta G^\circ_t(\text{X}^\ominus, \text{W} \rightarrow \text{S}) = R \cdot T[&2 \cdot \ln s(\text{Ph}_4\text{AsX}, \text{W}) - 2 \cdot \ln s(\text{Ph}_4\text{AsX}, \text{S}) \\ &+ \ln s(\text{Ph}_4\text{AsPh}_4\text{B}, \text{S}) - \ln s(\text{Ph}_4\text{AsPh}_4\text{B}, \text{W})]\end{aligned} \tag{2-12b}$$

where s is the solubility, expressed on the molar scale ($\text{mol} \cdot \text{l}^{-1}$).

Table 2-9 collects selected values of $\Delta G^\circ_t(\text{X}, \text{W} \rightarrow \text{S})$ obtained on this basis, taken from the extensive and critically evaluated compilations of Marcus [244, 349] and Gritzner [350]. A nice graphical representation of the changes in ΔG°_t, ΔH°_t, and ΔS°_t for

---

* Analogously, the following extrathermodynamic "reference electrolyte" assumptions are widely used:

$$\Delta H^\circ_t(\text{Ph}_4\text{As}^\oplus, \text{W} \rightarrow \text{S}) = \Delta H^\circ_t(\text{Ph}_4\text{B}^\ominus, \text{W} \rightarrow \text{S}),$$

and similarly

$$\Delta S^\circ_t(\text{Ph}_4\text{As}^\oplus, \text{W} \rightarrow \text{S}) = \Delta S^\circ_t(\text{Ph}_4\text{B}^\ominus, \text{W} \rightarrow \text{S}),$$

for the transfer from water to all solvents at any temperature [244]. This is equivalent to assuming that the molar Gibbs energy of transfer, $\Delta G^\circ_t(\text{X}, \text{W} \rightarrow \text{S})$, at a given reference temperature (usually 298.15 K) is valid for all temperatures [244].

Table 2-9. Selected standard molar Gibbs energies of transfer of single ions X from water (W) to seven nonaqueous solvents (S), $\Delta G_t^\circ(X, W \rightarrow S)$/(kJ · mol^{-1})[a], at 25 °C (molar scale), taken from the compilation of Marcus [244]. Values for $F^\ominus$ were taken from G. T. Hefter, Pure Appl. Chem. *63*, 1749 (1991).

X	S						
	CH_3OH	C_2H_5OH	CH_3COCH_3	$HCON(CH_3)_2$	CH_3CN	CH_3SOCH_3	$[(CH_3)_2N]_3PO$
$H^\oplus$	10.4	11.1		−18	46.4	−19.4	
$Li^\oplus$	4.4	11		−10	25	−15	
$Na^\oplus$	8.2	14		−9.6	15.1	−13.4	
$K^\oplus$	9.6	16.4	4	−10.3	8.1	−13.0	−16
$Ag^\oplus$	6.6	4.9	9	−20.8	−23.2	−34.8	−44
$(CH_3)_4N^\oplus$	6	10.9	3	−5.3	3	−2	
$(C_6H_5)_4As^\oplus$ [b]	−24.1	−21.2	−32	−38.5	−32.8	−37.4	−39
$F^\ominus$	16	26	79	85	71	73	
$Cl^\ominus$	13.2	20.2	57	48.3	42.1	40.3	58
$Br^\ominus$	11.1	18.2	42	36.2	31.3	27.4	46
$I^\ominus$	7.3	12.9	25	20.4	16.8	10.4	30
$CN^\ominus$	8.6	7	48	40	35	35	
$ClO_4^\ominus$	6.1	10		4	2		−7
$(C_6H_5)_4B^\ominus$ [b]	−24.1	−21.2	−32	−38.5	−32.8	−37.4	−39

[a] A positive value of $\Delta G_t^\circ(X, W \rightarrow S)$ means that the ion is better solvated by water than by solvent S; a negative value means that the ion is more strongly solvated after transfer from water to solvent S.

[b] See text for the so-called tetraphenylarsonium tetraphenylborate assumption.

the transfer of univalent single ions from water to other solvents has been given by Persson [301]. See Section 5.5.3 for further discussions.

The following three aspects are also of importance in solvation: the stoichiometry of the solvate complexes (normally described by the coordination or solvation number), the lability of the solvate complexes (usually described by the rate of exchange of the molecules of the solvent shell with those of the bulk solvent), as well as the fine structure of the solvation shell (for water often described by the simple model of ion solvation of Frank and Wen [16]).

Coordination and solvation numbers reflect the simple idea that the solvation of ions or molecules consists of a coordination of solute and solvent molecules. The *coordination number* is defined as the number of solvent molecules in the first coordination sphere of an ion in solution [103]. This first coordination sphere is composed only of solvent molecules in contact with or in bonding distance of the ion such that no other solvent molecules are interposed between them and the ion. This kind of solvation is sometimes termed *primary* or *chemical solvation*. Coordination numbers, determined by different experimental techniques [103], range in water from approx. 4 for $Be^{2\oplus}$ to approx. 9 for $Th^{4\oplus}$, although the majority of the values are close to 6 (*e.g.* for $Al^{3\oplus}$).

The *solvation number* is defined as the number of solvent molecules per ion which remain attached to a given ion long enough to experience its translational movements [94, 97, 104]. The solvation number depends upon the reference ion and its assumed solvation number as well as upon the method of measurement. Depending on the method of measurement, solvent molecules loosely bound in the second or in a higher sphere may be included. The partial ordering of more distant solvent molecules beyond the primary solvation shell is termed *secondary* or *physical solvation*. For example, mobility measurements indicate the number of solvent molecules moving with the ion, while dielectric measurements indicate only the number of solvent molecules in the first sphere. The solvation number of $Li^{\oplus}$ in water, determined using different electrolytic transference methods, varies therefore between 5 and 23. An inspection of the solvation numbers measured by electrolytic transport methods shows that the order of hydration numbers of the alkali metal cations is: $Li^{\oplus} > Na^{\oplus} > K^{\oplus} > Rb^{\oplus} > Cs^{\oplus}$. The alkaline earth metal cations are more highly solvated than the alkali cations ($Mg^{\oplus\oplus} > Ca^{\oplus\oplus} > Sr^{\oplus\oplus} > Ba^{\oplus\oplus}$). The more dilute the solution the greater the solvation of a given ion. The halogen anions are hydrated in the order $F^{\ominus} > Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$. Therefore, as a rule it can be stated that the smaller the ion and the greater its charge, the more highly it is solvated [94, 97, 104]. Conductance data show, that the solvation number for a given ion varies strongly with the solvent. Thus, the solvation number of $Li^{\oplus}$ varies from 1.4 in sulfolane, 7 in methanol, 9 in acetonitrile to 21 in water. The conductance data also indicate that in all organic solvents used, the solvation of the alkali metal cations is in the order: $Li^{\oplus} > Na^{\oplus} > K^{\oplus} > Rb^{\oplus} > Cs^{\oplus}$. The order of solvation of the halogen anions in the organic solvents studied is, in general, $Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ [94, 97, 104].

Even in the case of strong interactions between solvent and solute, the life time of each solvate is brief since there is continuous rotation or exchange of the solvent shell molecules. The time required for reorientation of hydrates in water is of the order $10^{-10} \ldots 10^{-11}$ s at 25 °C [91]. If the exchange between bulk solvent molecules and those in the inner solvation shell of an ion is slower than the NMR time scale, then it is possible to observe two different resonance signals for the free and bound solvent. In this

way, it has been shown, using ^{17}O NMR spectroscopy, that the hexaaquo hydration spheres of $Al^{3\oplus}$ and $Cr^{3\oplus}$, and the four water molecules bound by $Be^{2\oplus}$ exchange at a rate of less than 10^4/s, while those of the alkali metal cations exchange at a rate faster than 10^4/s [96, 105].

In general, since solvent molecules directly bound to an ion have different chemical shifts from those of the bulk solvent, NMR spectroscopy is a very useful method for studying solvation shells [106–111]. If the exchange rates are too high, however, the NMR signals coalesce to a single time-averaged resonance signal. It is usually assumed that solvent molecules in environments other than the first coordination sphere are exchanging at diffusion-controlled rates and therefore appear in the environmentally averaged bulk solvent resonance. A variety of different solvent nuclei have been used for this purpose: ^1H, ^{13}C, ^{17}O, and ^{31}P. As an example, the ^1H NMR spectrum of 2.1 M aqueous solution of $Al(NO_3)_3$ at -40 °C shows two signals [112]. The low-field signal arises from the coordinated solvent, and the high-field resonance from the bulk solvent. Two ^{13}C NMR signals are also observed for aqueous dimethyl sulfoxide containing $AlCl_3$ at 30 °C, one for the bulk and one for the bound solvent (1.94 ppm upfield) [113].

The ^1H NMR spectrum of an aqueous $Al(ClO_4)_3$ solution in $[D_6]$acetone shows nicely the two different signals of bulk water and hydration water in the $Al^{3\oplus}$ inner shell, even at room temperature [245]. The addition of acetone slows down the proton exchange rate. A primary hydration number of six for $Al^{3\oplus}$ has been obtained in this way [245].

Another approach to the study of ion-solvent interactions involves the determination of the solvent effect on the resonance frequency of the solute ion, using nuclei of spin $I \neq 0$ such as ^7Li, ^{23}Na, ^{27}Al, ^{35}Cl, ^{59}Co, ^{69}Ga, ^{133}Cs, ^{195}Pt, and ^{205}Tl [106–111, 111a, 246, 247, 294]. $^{205}Tl^{\oplus}$ is an exceptionally sensitive ion [294]. In going from water to pyridine the change in resonance frequency is approximately 782 ppm (!) [114]. In comparison, the change in chemical shift for $^{23}Na^{\oplus}$ in these two solvents is only about 1.3 ppm [115]. Therefore, $^{205}Tl^{\oplus}$ and other ions are very useful probes for the study of solvation and solvent structure. The greater the Lewis basicity of the solvent, the higher the resonance frequency of the $^{205}Tl^{\oplus}$ ion. The increase in resonance frequency with increasing solvent Lewis basicity can be considered as a measure of the strength of interaction between the solute ion and solvent molecules [294].

Other spectroscopic methods have also been used to study the statics and dynamics of solvation shells of ions and molecules [351–354]. In this respect, *solvation dynamics* refers to the solvent reorganization (*e.g.* rotation, reorientation, and residence time of solvent molecules in the first solvation shell) in response to an abrupt change in the solute properties, *e.g.* by photoexcitation of the solute with ultra-short laser-light pulses. Provided that this excitation is accompanied by an electron transfer or a change in the dipole moment, then the dynamics of this process correspond to how quickly the solvent molecules rearrange around the instantaneously created charge or the new dipole.

A number of models have been developed to describe the fine structure of the solvent shells of ions and molecules. While the agreement with experimental findings is more or less satisfactory, it is for the most part only qualitative (for reviews, see references [85, 91, 94, 95, 98]). According to the influence of the solute on the solvent structure, two different types of solvent can be distinguished (Fig. 2-9) [98]. In the former case, the pure solvent does not show a high degree of order. The directional properties of



Fig. 2-9. Schematic multizone models for ion solvation in solvents: (a) with low degree of order such as hydrocarbons, consisting of solvation shell A and disordered bulk solvent B [98]; (b) in highly ordered solvents such as water, consisting of solvation shell A with immobilized solvent molecules, followed by a structure-broken region B, and the ordered bulk solvent C (Frank and Wen [16]).

the dissolved ion dominate in a rather large region around the center and decrease gradually on proceeding into the unperturbed bulk solvent. The solution consists of an ordered sphere – the primary solvation shell A – and the disordered bulk solvent B (Fig. 2-9a) [98].

In the latter case, the solvent possesses a highly ordered structure such as that found in water. Frank and Wen [16] distinguish between three different regions in the solvent surrounding a solute. In the first coordination sphere A, the solvent molecules are strongly bound to the ion and therefore appear less mobile than the molecules in the bulk solvent. At some distance from the ion there exists the normal structure of the pure ordered solvent C. Between A and C, according to Frank and Wen [16], lies an intermediate region of disorder B, with highly mobile solvent molecules. This has been introduced in order to explain the "structure making" and "structure breaking" properties of ions of different charge and size in aqueous solutions. The concept of different regions around the dissolved ion was developed by Gurney [116], who introduced the term *cosphere* for the zone surrounding a spherical ion in which significant differences in structure and properties of solvent molecules are to be expected*). In contrast to the ordinary strong *positive hydration* of small spherical ions possessing a structure-making effect on the solvent molecules (*cf.* Fig. 2-9a), water molecules around a dissolved ion are in some cases more mobile than in pure water. In other words, the exchange frequency of water molecules around the ions is greater than in regions of pure water (*cf.* region B in Fig. 2-9b). This explains the experimental observation that aqueous solutions of certain salts such as potassium iodide show a greater fluidity than pure water at the same temperature. This effect has been called *negative hydration* [85] and it is con-

* In a liquid, the formation of temporary solvent molecule groups which have some crystalline character has been called *cybotaxis* (Greek, κυβεύω, dice-play, τάξις, an arrangement) by Stewart [116a]; see also [116b]. A *cybotactic region* may then be defined as the volume around a solute molecule in which the ordering of the solvent molecules has been influenced by the solute, including both the first solvation shell and the transition region; *cf.* [129].

nected with the structure-breaking effect of large singly-charged spherical ions on solvent molecules [91, 117]. The structure-breaking effect of large ions is not restricted to water as a solvent. Ethylene glycol and glycerol are liquids which also show this effect for a number of salts that cause structure-breaking in water [117]. To date, however, the validity of the multizone models for ion solvation proposed by Frank and Wen [16] and others has lacked direct experimental proof [117]. Consequently, owing to the lack of detailed knowledge of the solvents' structure and of satisfactory molecular theories for associated liquids, all attempts at a detailed description of solvation shells are still imperfect.

The solubility of a dissolved non-electrolyte solute can be reduced by the addition of a salt. This phenomenon, known as the *salting-out effect*, is of practical importance for the isolation of organic compounds from their solutions. In the presence of a dissolved dissociated salt, a fraction of the solvent molecules becomes involved in solvational interaction with the ions of the electrolyte, whereby their activity is diminished, leading to salting-out of the dissolved non-electrolyte solute. In other words, the salting-out can be considered as the difference in solubility in two kinds of solvents, the ion-free and the ion-containing one [248].

Theoretical chemists have developed a variety of methods and computational strategies for describing and understanding the complex phenomenon of solvation [27d, 355–358]. Altogether, three general approaches have been used for the theoretical description of solute/solvent interactions:

(a) quantum-chemical continuum models, where the solvent is treated as a structureless, *i.e.* continuous and homogeneous, medium that surrounds the solute ions or molecules like a bath, characterized solely by its relative permittivity ε_r. The solvated species (ions, polar molecules) induce polarization charges in the surrounding solvent continuum that in turn give rise to an extra electric field in the vicinity of the solute, called (Onsager) reaction field [357, 359; and references cited therein];
(b) supramolecular models, which treat the solvent molecules around the solute on the same footing as the solute, *i.e.* as discrete particles in an ensemble of solute and solvent species, using Monte Carlo statistical mechanics or molecular dynamics techniques [360, 361; and references cited therein];
(c) semicontinuum quantum-chemical models, which retain the reaction field contribution, but the direct electrostatic solute/solvent interactions in the first solvation shell are modeled differently. That is, the supermolecule (solute + first solvation shell) is surrounded by a continuum solvent [362–364; and references cited therein].

Particularly during the last decade, much progress has been made in the theoretical description of solvation. However, when applied to actual solutes, all models still have their limitations and flaws. For comprehensive reviews on theoretical treatments of solvation phenomena, see references [27d, 355–358].

2.4 Selective Solvation [89, 94, 96, 118–120, 241, 249, 250]

The description of solvation of ions and molecules in solvent mixtures is even more complicated. Besides the interaction between solvent and solute, the interaction between unlike solvent molecules plays an important supplementary role. This leads to large



Fig. 2-10. Schematic model for the selective solvation of ions by one component of a binary 1:1 mixture of the solvents A and B [119].
(a) Homoselective solvation: both ions are preferentially solvated by the same solvent A.
(b) Heteroselective solvation: the cation is preferentially solvated by A and the anion by B.

deviations from the ideal behaviour expected from Raoult's law of vapour pressure depression of binary mixtures; see references [365, 366] for reviews on the physicochemical properties of solvent mixtures. Typical examples of the non-ideal behaviour of binary solvent mixtures are water/alcohol [19], dimethyl sulfoxide/methanol [367], as well as water/methanol and water/acetonitrile mixtures [368]. When water is mixed with methanol an exothermic mixing enthalpy is observed, whereas in the case of acetonitrile the enthalpy of mixing is endothermic [368].

From investigations of the solvation of ions and dipolar molecules in binary solvent mixtures it has been found that the ratio of the solvent components in the solvent shell can be different from that in the bulk solution. As expected, the solute is surrounded preferably by the component of the mixture which leads to the more negative Gibbs energy of solvation, ΔG°_{solv}. The observation that the solvent shell has a composition other than the macroscopic ratio is termed *selective* or *preferential solvation* (*cf.* Fig. 2-10). These terms are generally used to describe the molecular-microscopic local solute-induced inhomogeneity in a multicomponent solvent mixture. They include both (i) nonspecific solute/solvent association caused by dielectric enrichment in the solvent shell of solute ions or dipolar solute molecules, and (ii) specific solute/solvent association such as hydrogen-bonding or EPD/EPA interactions.

When in a mixture of two solvents, both ions of a binary salt are solvated preferably by the same solvent, the term applied is *homoselective solvation* (Fig. 2-10a). Similarly, the preferred solvation of the cation by one, and the anion by the other solvent, is termed *heteroselective solvation* (Fig. 2-10b) [119]. Thus, in a solution of silver nitrate in the binary solvent mixture acetonitrile/water, a preferential solvation of $Ag^{\oplus}$ by acetonitrile and of $NO_3^{\ominus}$ by water was observed (heteroselective solvation) [121, 369]*). In contrast, in solutions of calcium chloride in water/methanol mixtures, both $Ca^{2\oplus}$ and $Cl^{\ominus}$ are solvated largely by water (homoselective solvation) [122]. $Zn^{2\oplus}$ (from $ZnCl_2$) in

* The reasons for preferential solvation of $Ag^{\oplus}$ ions by acetonitrile in acetonitrile/water mixtures and the solvation shell structure of silver ions have been discussed [251].

the solvent mixture water/hydrazine is preferentially solvated by hydrazine; in an acetonitrile/water mixture solvation is largely by water [123]. Ag_2SO_4 is heteroselectively solvated in methanol/dimethyl sulfoxide mixtures: the silver ion is preferentially solvated by dimethyl sulfoxide, whereas the sulfate ion is preferably solvated by methanol. The Ag_2SO_4 salt is only sparingly soluble in methanol and in dimethyl sulfoxide. Its solubility is higher in mixtures of the two solvents than in the neat liquids, since both the cation and the anion can be solvated with the solvent component for which it has a greater affinity [123a]. A study of silver(I) salts in the *isodielectric* mixture of methanol ($\varepsilon_r = 32.7$) and *N*-methylpyrrolidin-2-one (NMP; $\varepsilon_r = 32.2$) showed heteroselective solvation of Ag^+ by NMP and the anions (SO_4^{2-}, BrO_3^-, IO_3^-) by methanol [370]. The $Cu^{\oplus}$ ion (from $CuClO_4$) shows strong preferential solvation by acetonitrile in acetonitrile/acetone mixtures, which may be of interest in the hydrometallurgical purification of copper [252]. Even protons exhibit preferential solvation by amines in mixed water/amine ion clusters studied in the gas phase [253].*)

In a binary mixture of solvents S_1 and S_2, a cation $M^{z\oplus}$ with a coordination number k and charge $z^{\oplus}$ forms $(k+1)$ cations of the type $[M(S_1)_i(S_2)_{k-i}]^{z\oplus}$ with $i = 0 \ldots k$, differently solvated in the first solvation shell. These differently solvated species have been called *solvatomers* [254]. For example, with octahedrally coordinated cations ($k = 6$), $k + 1 + 3 = 10$ solvatomers are to be expected (including three *cis*/*trans* isomeric solvatomers with $i = 2$, 3, or 4). In favourable cases, the concentrations of all solvatomers have been obtained as a function of the solvent mole fraction by NMR measurements [254].

Preferential solvation is not restricted to ions of electrolytes dissolved in multicomponent solvent systems. Even for dipolar nonelectrolyte solutes the composition of the solvation shell can deviate from that of the bulk solvent mixture, as shown for *β*-disulfones [255] and *N*-methylthiourea [256].

Different methods for the study of selective solvation have been developed [118, 120]: conductance and Hittorf transference measurements [119], NMR measurements (especially the effect of solvent composition on the chemical shift of a nucleus in the solute) [106–109], and optical spectra measurements like IR absorption shifts [111] or UV/Vis absorption shifts of solvatochromic dyes in binary solvent mixtures [124, 249, 371]. Recently, the preferential solvation of ionic (tetralkylammonium salts) and neutral solutes (phenol, nitroanilines) has been studied particularly successfully by ^1H NMR spectroscopy through the analysis of the relative intensities of intermolecular ^1H NOESY cross-peaks [372].

A convenient measure of the degree of selective solvation is the bulk solvent composition at which both solvents of a binary mixture participate equally in the contact solvation shell. This is the solvent composition at which the NMR chemical shifts lie midway between the values for the two pure solvents. This composition has been called the *equisolvation* or *iso-solvation point* (usually expressed in mole fractions of one solvent) [125]. According to Fig. 2-10, this point describes the bulk solvent composition at which both solvents A and B participate equally in the solvation shell of the cation or the anion, respectively.

---

* A comprehensive tabulation of selective solvation of ions in a number of binary solvent systems is given by Gordon [96] (p. 256).



Fig. 2-11. NMR chemical shift of $^{23}Na^{\oplus}$ as a function of the mole fraction of dimethyl sulfoxide (DMSO) in a binary mixture of DMSO and acetone (according to [295]). *Straight line:* ideal case without preferential solvation, primary solvation shell of the same composition as the bulk solvent mixture. *Curved line:* real case with preferential solvation of $^{23}Na^{\oplus}$ by DMSO and iso-solvation point at $x_{DMSO}/(cmol \cdot mol^{-1}) \approx 0.21$, that is, the mole fraction of the bulk solvent for which the solvated ion chemical shift is the average of the shifts obtained in the pure solvents ($\Delta\delta = \delta_{DMSO} - \delta_{Acetone}$).

A useful probe of the immediate chemical environment of solute ions is the NMR chemical shift of alkali metal ions obtained in binary solvent mixtures [111, 126, 295]. These measurements are based on the assumption that the chemical shift of the solute cation is determined in an additive fashion by the solvent molecules comprising the first solvation shell. For example (*cf.* Fig. 2-11), the iso-solvation point of $^{23}Na^{\oplus}$ in dimethyl sulfoxide/acetone mixtures occurs at $x \approx 0.21$ cmol/mol dimethyl sulfoxide, indicating the higher solvating ability of this solvent relative to acetone. As shown schematically in Fig. 2-11, the preferential solvation of $^{23}Na^{\oplus}$ by dimethyl sulfoxide displaces its chemical shift towards δ_{DMSO} and a deviation from the straight line is observed.

The iso-solvation points obtained from $^{23}Na^{\oplus}$ chemical shifts of sodium tetraphenylborate in different binary solvent mixtures indicate the following relationships between the solvating abilities of pairs of organic solvents: $CH_3SOCH_3 \gg CH_3NO_2$; pyridine $> CH_3NO_2$; $CH_3SOCH_3 > CH_3CN$; pyridine $> CH_3CN$; $C_6H_5CN > CH_3NO_2$; $CH_3SOCH_3 >$ pyridine [126].

The term selective solvation also applies when one and the same dipolar molecule is preferentially solvated at two different loci by two different solvents. An example is the



(1)

chloro-oxalato-tripyridine-rhodium(III) complex (*1*), which dissolves in a 1:1 mixture of pyridine and water, but not in either pure water or pyridine [127]. Presumably, a Gibbs energy of solvation large enough to overcome the lattice forces is attained only by selective solvation of the three pyridine ligands by pyridine, and of the oxalato ligand by water.

Many macromolecular compounds dissolve in mixtures better than in pure solvents [20]. Thus, poly(vinyl chloride) is insoluble in acetone as well as in carbon disulfide, but soluble in a mixture of the two. The opposite situation is also known. Malononitrile and *N*,*N*-dimethylformamide both dissolve polyacrylonitrile but a mixture of the two does not [20]. Soaps dissolve neither in ethylene glycol nor in hydrocarbons at room temperature but are quite soluble in a mixture of the two. Here, ethylene glycol solvates the ionic end, and the hydrocarbon the apolar end of the fatty acid chain [128].



A great variety of models for a quantitative description of the composition dependence of the physicochemical properties of solutes dissolved in binary solvent mixtures have been developed [257–261, 373–378]. For example, using a rather simple two-step solvent-exchange model [374, 377], the behaviour of seventy binary solvent mixtures towards a solvatochromic betaine dye (structure see Fig. 6-2 in Section 6.2.1) can be quite precisely described, even for so-called *synergetic* solvent mixtures [377].

A binary solvent mixture exhibits *synergistic effects* on a physicochemical solute property P if for some mixtures this property P has a value higher or lower than either properties P_1 and P_2 corresponding to the neat solvents S_1 and S_2 [379]. For example, the empirically determined solvent polarity of binary mixtures of HBD and HBA solvents is often larger than the polarities of the two neat components. Clearly, the formation of hydrogen-bonded 1:1 complexes between HBD ad HBA solvent molecules leads to a new, more polar medium [124, 249, 377] (see Chapter 7 for a definition of the term *solvent polarity*).

The non-ideal behaviour of a wide selection of binary solvent mixtures has been studied experimentally mainly by means of suitable solvatochromic dyes, the UV/Vis absorptions of which are solvent-dependent (*cf.* Section 6.2.1); see references [380–385] for some more recent examples. Conversely, the largely non-ideal solute behaviour in binary solvent mixtures has been used for the quantitative determination of the compositions of such solvent mixtures, *e.g.* for the determination of small water contents in organic solvents [386–388].

2.5 Micellar Solvation (Solubilization) [96, 128, 130–132, 220, 262–267]

Special conditions are found in solutions of large cations and anions possessing a long unbranched hydrocarbon chain, *e.g.* $CH_3—(CH_2)_n—CO_2^{\ominus}M^{\oplus}$, $CH_3—(CH_2)_n—SO_3^{\ominus}M^{\oplus}$, or $CH_3—(CH_2)_n—N(CH_3)_3^{\oplus}X^{\ominus}$ (with $n > 7$). Such compounds are known as *amphiphiles*, reflecting the presence of distinct polar and nonpolar regions in the



Fig. 2-12. Schematic two-dimensional representation of spherical micelle formation by an anionic amphiphile such as $CH_3—(CH_2)_{11}—CO_2^{\ominus}M^{\oplus}$ in water. The head group ($\ominus$), the counterions ($\oplus$), and the hydrocarbon chains are only schematically indicated to denote their relative position. The highly charged interface (ionic head groups plus bound counterions) between the micelle's hydrophobic core and the bulk solution is called the *Stern layer*. For a more realistic three-dimensional picture of a micelle, see references [264, 389].

molecule. Salts of such large organic ions are often highly aggregated in dilute aqueous solution. The resulting structured aggregates, together with counterions localized near their periphery by coulomb forces, are termed *micelles**). Fig. 2-12 gives a schematic representation of the formation of a spherical micelle by an anionic amphiphile.

The hydrophobic part of the aggregate molecules forms the core of the micelle while the polar head groups are located at the micelle-water interface in contact with the water molecules. Such micelles usually have average radii of $2 \ldots 4$ nm and contain $50 \ldots 100$ monomers in water. Their geometric structure is usually roughly spherical or ellipsoidal. In non-aqueous nonpolar solvents, the micellar structures are generally the inverse of those formed in water. In these solvents, the polar head groups form the interior of the micelle while the hydrocarbon chains of the ions are in contact with the nonpolar solvent.

At very low concentrations, ionic amphiphiles behave as normal strong electrolytes, but if the concentration is raised above the so-called *c*ritical *m*icelle *c*oncentration (*cmc*; usually $10^{-4} \ldots 10^{-2}$ mol $\cdot$ l^{-1}), spherical aggregates are formed. Increasing the amphiphile (surfactant) concentration results in two different effects [264, 389]: (a) the increasing amphiphile concentration leads to an increased ionic strength of the aqueous bulk solution, thus decreasing the electrostatic repulsion between the head groups due to screening of their negative or positive charges; (b) for the hydrophobic hydrocarbon tails, an increase in the amphiphile concentration is unfavourable because of the increasing hydrophobic amphiphile/water interactions (see Section 2.2.7). Eventually,

* The term *micelle* was introduced in 1877 by Nägeli (from the Latin *mica*, a crumb) for a molecular organic aggregate of limited size without exact stoichiometry [270]. The existence of surfactant aggregates in aqueous soap solutions was established in 1896 by Krafft [271], and the first description of a surfactant micelle was given in 1913 by Reychler [272].

Table 2-10. Some typical surfactants (*surf*ace *act*ive *a*gents) with their critical micelle concentrations (*cmc*) and aggregation numbers in aqueous solutions at 25 °C [268].

Surfactants	*cmc*/(mol · l^{-1})	Aggregation number
Anionic Sodium 1-dodecyl sulfate (SDS) $CH_3—(CH_2)_{11}—OSO_3^{\ominus}Na^{\oplus}$	0.0081	62
Cationic 1-Hexadecyl (=Cetyl)-trimethylammonium bromide (CTAB) $CH_3—(CH_2)_{15}—\overset{\oplus}{N}(CH_3)_3Br^{\ominus}$	0.0013	78
Nonionic Hexa(oxyethylene)dodecanol $CH_3—(CH_2)_{11}—(OCH_2CH_2)_6—OH$	0.00009	400
Zwitterionic 3-(*N*-1-Dodecyl-*N*,*N*-dimethylammonio)propane-1-sulfonate (SB 12) $CH_3—(CH_2)_{11}—\overset{\oplus}{N}(CH_3)_2—(CH_2)_3—SO_3^{\ominus}$	0.003	55

the driving force for dissolution of the amphiphile will be completely balanced by the forces working against the dissolution of the hydrophobic tails. Now, two different scenarios are possible: either a macroscopic phase separation will occur (*i.e.* formation of aggregates of infinite size), or micelles will be formed (*i.e.* formation of aggregates of finite size). Such micelles are thermodynamically stable, microheterogeneous, supramolecular species, dissolved in the aqueous bulk. They are characterized by the aforementioned *cmc* and the *micellar aggregation number*, both of which are dependent on the hydrocarbon tail length, the nature of the counter ion, and the ionic strength of the bulk solution.

Typical surfactants are listed in Table 2-10 along with their respective *cmc* values and aggregation numbers [268].

In reality, micellar systems are more complex than is implied by the simple static picture given in Fig. 2-12 (also known as the Hartley model [390]). A more realistic picture of micellar structures has been given by Menger [269]. According to his ''porous cluster'' or ''reef'' model, micelles possess rugged, dynamic surfaces, water-filled pockets, nonradial distribution of chains, and random distribution of terminal methyl groups. In fact, a micelle is a highly disorganized structure with multiple bent hydrocarbon chains, cavities, and even hydrocarbon/water contacts, and shows deviations from a precise spherical shape. A micelle is a dynamic molecular assembly which exists in equilibrium with its monomer, where monomer units are both leaving and entering the micelle. A monomer remains in a micelle for only $10^{-8} \ldots 10^{-3}$ s depending on the chain length of the surfactant molecule. Another, so-called surfactant-block model of micelles, has been given by Fromherz [273].



Fig. 2-13. Schematic two-dimensional representation of the solubilization of (b) *n*-nonane as a nonpolar substrate, and (c) 1-pentanol as another amphiphile, by a spherical ionic micelle (a) of an *n*-decanoic acid salt in water.

The existence of micelles in solutions of large ions with hydrocarbon chains is responsible for the observation that certain substances, normally insoluble or only slightly soluble in a given solvent, dissolve very well on addition of a surfactant (detergent or tenside). This phenomenon is called *solubilization* and implies the formation of a thermodynamically stable isotropic solution of a normally slightly soluble substrate (the *solubilizate*) on the addition of a surfactant (the *solubilizer*) [128, 133]. Non-ionic, nonpolar solubilizates such as hydrocarbons can be trapped in the hydrocarbon core of the micelle. Other amphiphilic solutes are incorporated alongside the principal amphiphile and oriented radially, and small ionic species can be adsorbed on the surface of the micelle. Two modes of solubilizate incorporation are illustrated in Fig. 2-13.

Because the micellar interior is far from being rigid, a solubilized substrate is relatively mobile. Like micelle formation, solubilization is a dynamic equilibrium process. Representative recent examples are the solubilization of benzene, naphthalene, anthracene, and pyrene in aqueous solution by the addition of 1-dodecanesulfonic acid [391], the solubilization of fullerene C_{60} in aqueous solutions of the non-ionic surfactant Triton X-100 [392], and the solubilization of a cholesteryl-group bearing pullulane (a hydrophobized polysaccharide) [393].

Essentially spherical micelles are not the only aggregates that can be formed in aqueous solution above the critical micelle concentration. Depending on the molecular structure of the amphiphile and the solution conditions (*e.g.* electrolyte concentration,

pH, temperature), inverted micelles, bilayers, vesicles, and biological membranes can readily be formed by spontaneous self-association [394] of certain amphiphilic molecules [130–132, 264]. In contrast to infinite *planar* bilayers, vesicles or liposomes (vesicles formed from lipids) are closed *spherical* bilayer aggregates that are much larger than micelles. Eventually, the mutual interaction of these aggregates at high concentrations (low water content) can lead to a transition to larger and more ordered *mesophases* or *lyotropic liquid crystalline* structures [264].

Not only solubilities, but also the rates and selectivities of organic reactions can be affected by the addition of surfactants to the reaction medium. The modification of chemical reactions by incorporating the reactant molecules into organized assemblies as a kind of microreactor has been the subject of considerable attention [274–277, 395, 396] (*cf.* also Section 5.4.8).

2.6 Ionization and Dissociation [49, 96, 134–139, 241, 278, 279]

Solutions of non-electrolytes contain neutral molecules or atoms*) and are non-conductors. Solutions of electrolytes are good conductors due to the presence of anions and cations. The study of electrolytic solutions has shown that electrolytes may be divided into two classes: *ionophores* and *ionogens* [134]. Ionophores (like alkali halides) are ionic in the crystalline state and they exist only as ions in the fused state as well as in dilute solutions. Ionogens (like hydrogen halides) are substances with molecular crystal lattices which form ions in solution only if a suitable reaction occurs with the solvent. Therefore, according to Eq. (2-13), a clear distinction must be made between the *ionization* step, which produces ion pairs by heterolysis of a covalent bond in ionogens, and the *dissociation* process, which produces free ions from associated ions [137, 397, 398].

$$\underset{\text{Ionogen}}{(A\text{-}B)_{solv}} \overset{K_{Ion}}{\rightleftharpoons} \underset{\text{Ion Pair}}{(A^{\oplus}B^{\ominus})_{solv}} \overset{K_{Dissoc}}{\underset{K_{Assoc}}{\rightleftharpoons}} \underset{\text{Free Ions}}{(A^{\oplus})_{solv} + (B^{\ominus})_{solv}} \tag{2-13}$$

$$K_{Ion} = [A^{\oplus}B^{\ominus}]/[A - B] \tag{2-14}$$

$$K_{Dissoc} = [A^{\oplus}] \cdot [B^{\ominus}]/[A^{\oplus}B^{\ominus}] \tag{2-15}$$

The index "solv" indicates that the species in parentheses are within one solvent cage.

Ionophores may exist in solution as an equilibrium mixture containing ion pairs and free ions. *Ion pairs* are defined as pairs of oppositely charged ions with a common solvation shell, whose life times are sufficiently long to render them recognizable kinetic entities in solution and for which only electrostatic binding forces are assumed [135]. Experimentally, ion pairs behave as one unit in determining electric conductivity, kinetic behaviour, and some thermodynamic properties (*e.g.* activity coefficient; osmotic pres-

* An example of a monoatomic un-ionized substrate solution is that of mercury in air-free water, which contains zero-valent mercury atoms [140].

sure) of electrolyte solutions. In an external electric field such paired ions do not move individually but reorient themselves as an electric dipole. The ion-pair concept was introduced in 1926 by Bjerrum [280] to account for the behaviour of ionophores in solvents of low relative permittivity.

It is possible to distinguish between free ions from associated and covalently bonded species by conductivity measurements, because only free ions are responsible for electrical conductivity in solution [136, 399]. Spectrophotometric measurements distinguish between free ions and ion pairs on the one hand, and covalent molecules on the other, because in a first approximation the spectroscopic properties of ions are independent of the degree of association with the counterion [141]. The experimental equilibrium constant K_{exp}, obtained from conductance data, may then be related to the ionization and dissociation constants by Eq. (2-16).

$$K_{exp} = \frac{[A^{\oplus}] \cdot [B^{\ominus}]}{[A - B] + [A^{\oplus}B^{\ominus}]} = \frac{K_{Ion} \cdot K_{Dissoc}}{1 + K_{Ion}} \tag{2-16}$$

When the extent of ionization is small, then $K_{exp} = K_{Ion} \cdot K_{Dissoc}$ ($K_{Ion} \ll 1$ or $[A^{\oplus}B^{\ominus}] \approx 0$). For strong electrolytes, where $K_{Ion} \gg 1$, Eq. (2-16) reduces to $K_{exp} = K_{Dissoc}$.

Another equation, more comprehensive than Eq. (2-16), has been developed by Izmailov according to $K_{exp} = K_{dissoc}/(1 + K_{ion} + K^*)$, where K^* describes the equilibrium $A—B + solv \rightleftharpoons (A—B)_{solv}$ [412].

The two steps of Eq. (2-13), ionization and dissociation, are influenced in different ways by solvents. The coulombic force of attraction between two oppositely charged ions is inversely proportional to the relative permittivity of the solvent, according to Eq. (2-17). Therefore, only solvents with sufficiently high relative permittivities will be capable

$$U_{ion\text{-}ion} = -\frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{z^{\oplus} \cdot z^{\ominus} \cdot e^2}{\varepsilon_r \cdot r} \tag{2-17}$$

(U = potential energy of an ion-ion interaction; $z \cdot e$ = charge on the ion; r = distance between the ions; ε_0, ε_r = permittivity of the vacuum and of the medium, resp.)

of reducing the strong electrostatic attraction between oppositely charged ions to such an extent that ion pairs can dissociate into free solvated ions. These solvents are usually called *dissociating solvents**).

* Nernst [141a] and Thomson [141b] first showed independently that solvents of high relative permittivity promote the dissociation of ionic solutes. The term "dissociating solvent" was first used by Beckmann [141c] in connection with his ebullioscopic determination of the molecular mass of dissolved substances. Later on the term "smenogenic solvent" was proposed by Fuoss for solvents of low relative permittivity which favor the formation of ion pairs. Conversely, "smenolytic solvents" are those whose relative permittivities are high enough to prevent ion association [134]. The latter two terms have, however, found little application.

According to Eq. (2-17), for two isolated ions such as Na^+ and Cl^- in contact in vacuum ($\varepsilon_r = 1$), with $r = 276$ pm as the sum of the two ionic radii, the electrostatic binding energy is -8.4×10^{-19} J [26b]. This binding energy is of the order $200 \cdot kT$ per ion pair in vacuum, as compared to the thermal energy $kT = 4.1 \times 10^{-21}$ J at 300 K. Only at ion separation $r > 56000$ pm will the Coulomb energy fall below kT, which means that electrostatic Coulomb interactions are very strong and of long range [26b]. At ca. 500 kJ/mol ($=8.4 \times 10^{-19} \times 6.0 \times 10^{23}$), this interionic binding energy is similar to the energies of covalent bonds (*i.e.* 200–600 kJ/mol). The electrostatic interaction between oppositely charged ions can only be overcome by liberation of the molar Gibbs energy of solvation, ΔG°_{solv}, in transferring the ion pair from the gas phase (vacuum) into a medium with $\varepsilon_r \gg 1$.

Ion association is only noticeable in aqueous solutions at very high concentrations because of the exceptionally high relative permittivity of water ($\varepsilon_r = 78.4$), but are found at much lower concentrations in alcohols, ketones, carboxylic acids, and ethers. In solvents of relative permittivities less than $10 \ldots 15$, practically no free ions are found (*e.g.* in hydrocarbons, chloroform, 1,4-dioxane, acetic acid); on the other hand, when the relative permittivity exceeds 40, ion associates barely exist (*e.g.* water, formic acid, formamide). In solvents of intermediate relative permittivity ($\varepsilon_r = 15 \ldots 20$, *e.g.* ethanol, nitrobenzene, acetonitrile, acetone, *N*,*N*-dimethylformamide), the ratio between free and associated ions depends on the structure of the solvent as well as on the electrolyte (*e.g.* ion size, charge distribution, hydrogen-bonded ion pairs, specific ion solvation, *etc.*) [96]. Thus, lithium halides in acetone ($\varepsilon_r = 20.6$) are very weak electrolytes, whereas tetraalkylammonium halides are strongly dissociated in the same solvent [142–144]. In solvents of very low relative permittivity like benzene ($\varepsilon_r = 2.3$), very large association constants are usually found. This indicates that most ion pairs in such solutions exist in the form of higher aggregates [96].

The ability of a solvent to transform the covalent bond of an ionogen into an ionic bond, *i.e.* its *ionizing power*, is not determined in the first instance by its relative permittivity. Rather, the ionizing power of a solvent depends on its ability to function as an electron-pair acceptor or donor [53, 137]. A dissociating solvent is not necessarily an ionizing one – and *vice versa*. In most cases, ionization of bonds of the type $H^{\delta\oplus}—X^{\delta\ominus}$ (*e.g.* ionization of hydrogen halides), $R^{\delta\oplus}—X^{\delta\ominus}$ (*e.g.* ionization of haloalkanes in S_N1 reactions), or $M^{\delta\oplus}—R^{\delta\ominus}$ (*e.g.* ionization of organometallic compounds) is strongly assisted by electron-pair donor (EPD) and electron-pair acceptor (EPA) solvents (*cf.* Section 2.2.6), according to (R = H, alkyl):





The ionization of an ionogen can therefore be regarded as a coordinative interaction between substrate and solvent [281]. The polarization of the covalent bond to be ionized can occur *via* a nucleophilic attack of the EPD solvent on the electropositive end of the bond, or by an electrophilic attack of an EPA solvent on the electronegative end. Both attacks can, of course, also occur simultaneously. The following examples are illustrative.







In EPD solvents, ionization depends on the stabilization of the cation through coordination and, in some solvents, on solvation of the anions as well. In EPA solvents, the anion is stabilized through coordination and, to a lesser extent, additional solvation of the cation may occur.

An evaluation of the ionizing power of a solvent requires knowledge, not only of its coordinating abilities, but also of its relative permittivity. According to Eq. (2-13), solvents of high relative permittivity promote the dissociation of ion pairs. The consequential decrease in ion pair concentration displaces the ionization equilibrium in such a way that new ion pairs are formed from the substrate. Thus, a good ionizing solvent must not only be a good EPD or EPA solvent but also possess a high relative permittivity. The donor and acceptor properties of ionizing solvents can be described empirically in a quantitative way by donor numbers [67] or acceptor numbers [70] (*cf.* Section 2.2.6).

The extraordinary ionizing ability of water is above all due to the fact that it may act as an EPD as well as an EPA solvent. Thus, water is both an ionizing and dissociating medium whereas nitromethane, nitrobenzene, acetonitrile, and sulfolane are mainly dissociating. *N*,*N*-Dimethylformamide, dimethyl sulfoxide, and pyridine are mildly dissociating but good ionizing solvents. Hexamethylphosphoric triamide is an excellent ionizing medium due to its exceptional donor properties, particularly in the case of metal-carbon bonds [145, 146]. Alcohols and carboxylic acids, as hydrogen-bond donors are good EPA solvents and, therefore, good ionizing solvents for suitable substrates.

Chloro-triphenylmethane constitutes a classical example for distinguishing the ionizing and dissociating ability of a solvent. In 1902, Walden used it in liquid sulfur dioxide in the first demonstration of the existence of carbenium ions [147]. The colourless chloro-triphenylmethane dissolves in liquid sulfur dioxide ($\varepsilon_r = 15.6$ at 0 °C), giving an intense yellow colour ($\lambda_{max} = 430$ nm). This is caused by a partial formation of ion pairs, which do not conduct electricity. At low concentrations, the ion pairs partially dissociate into free ions, which do conduct electricity [148, 149].



(2-18)

colorless yellow yellow

$K_{Ion} = 1.46 \cdot 10^{-2}$ (0 °C); $K_{Dissoc} = 2.88 \cdot 10^{-3}$ mol/L (0 °C);

$K_{exp} = 4.1 \cdot 10^{-5}$ mol/L (0 °C) [148].

Table 2-11. Ionization equilibrium constants K_{Ion} of chloro-triphenylmethane in various solvents at 0 . . . 25 °C [150]. *Cf.* also [282].

Solvents	ε_r [a] (at 0 . . . 25 °C)	$K_{Ion} \cdot 10^4$	References
Nitrobenzene	34.8 (25 °C)	Too low to measure (25 °C)[c]	[151]
Acetonitrile	35.9 (25 °C)	Too low to measure (25 °C)	[152]
Dichloromethane	8.9 (25 °C)	0.07	[153]
1,1,2,2-Tetrachloroethane	8.2 (20 °C)	0.48 (18.5 °C)	[154]
1,2-Dichloroethane	10.4 (25 °C)	0.56 (20 °C)	[154]
Nitromethane	35.9 (25 °C)	2.7 (25 °C)	[155]
Sulfur dioxide	15.6 (0 °C)[b]	146 (0 °C)	[148]
Formic acid	58.5 (16 °C)	3100 (20.5 °C)	[156]
m-Cresol	11.8 (25 °C)	5600[d] (18 °C)	[157]

[a] J. A. Riddick, W. B. Bunger, and T. K. Sakano: *Organic Solvents*, 4th edition, in A. Weissberger (ed.), *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986.
[b] A. A. Maryott and E. R. Smith: *Table of Dielectric Constants of Pure Liquids*, NBS Circular 514, Washington, 1951.
[c] Because nitrobenzene absorbs strongly at the wavelength of the carbenium ion maximum from chlorotriphenylmethane, this result was obtained with chloro-diphenyl-4-tolylmethane.
[d] This K_{Ion} value corresponds to $36 \pm 4\%$ ionization of chlorotriphenylmethane in *m*-cresol [157].

Sulfur dioxide is a π-electron-pair acceptor. The standard explanation for the strong ionizing power of SO_2 is the formation of an EPD—EPA complex between the halide anion and the sulfur dioxide molecules [148]. Table 2-11 summarizes some of the available data concerning the comparative efficiencies of various solvents in promoting the ionization of chloro-triphenylmethane [150].

The K_{Ion} of chloro-triphenylmethane varies in different solvents by at least a factor of 10^5. In the protic solvents *m*-cresol and formic acid, which have relative permittivities of 11.8 and 58.5, respectively, chloro-triphenylmethane is strongly ionized but is only slightly dissociated in the former. The remarkable ionizing power of phenols and carboxylic acids has been attributed to their EPA properties, *i.e.* their ability to form a hydrogen bond between the hydroxyl group and the halide ion. Solvents with high relative permittivities but lacking pronounced EPA properties, such as acetonitrile and nitrobenzene, are barely capable of ionizing chloro-triphenylmethane. In the case of tri(4-anisyl)-chloromethane, the K_{Ion} value in the EPA solvent sulfur dioxide at 0 °C is about $5 \cdot 10^{10}$ times greater than that in nitrobenzene at 25 °C [151].

On the other hand, the ionization of chloro-triphenylmethane is also favored by EPD solvents. Since the developing carbenium ion is an electrophilic species, it readily interacts with nucleophilic solvents. Thus, the extent of ionization of chloro-triphenylmethane in nitrobenzene increases on the addition of aprotic EPD solvents in direct relation to the donor number [158]. See reference [299] for a study of ionization and dissociation equilibria of other halo-triphenylmethanes in solution ($Ph_3C—X$ with X = F, Cl, Br).

Another remarkable example of the solvent effect on the ionization of ionogens is the Friedel-Crafts intermediate antimony pentachloride/4-toluoyl chloride. It can exist as two distinct well-defined adducts depending on the solvent from which it is recrystallized, the donor-acceptor complex *(2)* or the ionic salt *(3)* [159].



(2) (3)

The donor-acceptor complex *(2)* is isolated from tetrachloromethane solution ($\varepsilon_r = 2.2$), the ionic salt *(3)* from chloroform solution ($\varepsilon_r = 4.9$). When dissolved in chloroform, the donor-acceptor complex recrystallizes as the ionic salt. Similarly, the ionic salt is converted to the donor-acceptor complex when dissolved in tetrachloromethane. This result shows that in solution an equilibrium exists between the two forms. The isolation depends on the solvent used for recrystallization. Similar results have been obtained in the case of the adduct between acetyl chloride and aluminium trichloride, which is un-ionized in chloroform, but completely ionized in nitrobenzene [160].

Other nice examples of well-studied solvent-dependent ionization equilibria of ionogens are azidocycloheptatriene ⇌ tropylium azide [282, 283] and (triphenylcyclopropen-1-yl) (4-nitrophenyl)malononitrile *(2a)* ⇌ triphenylcyclopropenium dicyano(4-nitrophenyl)methide *(3a)*, the latter being one of the first examples of direct heterolysis of a weak carbon-carbon bond to a carbocation and carbanion in solution [284].



(2a) (3a)

When dissolved in nonpolar solvents such as benzene or diethyl ether, the colourless *(2a)* forms an equally colourless solution. However, in more polar solvents (*e.g.* acetone, acetonitrile), the deep-red colour of the resonance-stabilized carbanion of *(3a)* appears ($\lambda = 475 \ldots 490$ nm), and its intensity increases with increasing solvent polarity. The carbon-carbon bond in *(2a)* can be broken merely by changing from a less polar to a more polar solvent. Cation and anion solvation provides the driving force for this heterolysis reaction, whereas solvent displacement is required for the reverse coordination reaction. The Gibbs energy for the heterolysis of *(2a)* correlates well with the reciprocal solvent relative permittivity in accordance with the Born electrostatic equation [285], except for EPD solvents such as dimethyl sulfoxide, which give larger ΔG°_{het} values than would be expected for a purely electrostatic solvation [284].

The first purely organic salt, $C_{48}H^{\oplus}_{51}C_{67}H^{\ominus}_{39}$, consisting solely of carbon and hydrogen atoms and being fully ionised in the crystalline state and in solution, was prepared by mixing tris[1-(5-isopropyl-3,8-dimethylazulenyl)]cyclopropenylium perchlorate with potassium tris(7*H*-dibenzo[*c*,*g*]fluorenylidenemethyl)methide (Kuhn's anion) in tetrahydrofuran solution [292]. An analogous green hydrocarbon salt, $C_{12}H^{\oplus}_{15}$ $C_{67}H^{\ominus}_{39}$, which consists of the tri(cyclopropyl)cyclopropenylium cation and Kuhn's anion, is completely ionized in the solid and in DMSO solution. However, in chloroform, tetrachloromethane, and benzene solutions at room temperature, a covalent hydrocarbon is formed from the two ions. Surprisingly, cooling the chloroform solution to −78 °C or evaporation of the solvent regenerates the original green hydrocarbon salt.

In acetonitrile, the ionic and covalent forms coexist in a clean equilibrium. This compound is the first hydrocarbon that only exists covalently in solution [292]. In acetone, dichloromethane, and tetrahydrofuran, a radical, derived from Kuhn's anion by single-electron transfer (SET), was detected in addition to the two ionic species. Thus, all three types of elementary organic species (ion, radical, and a covalent compound) are shown to be able to coexist in a solution equilibrium, depending on the solvent used [292]. For reviews on solvent-dependent equilibria, including radical pairs (produced by bond heterolysis) and radical ion pairs (produced by electron transfer), see references [291, 400, 401].

Another type of ion pairs, called *penetrated ion pairs* [402], has been found by studying the conductivity of tetraalkylammonium tetrafluoroborates (with variable alkyl-chain lengths) [399, 403], and the UV/Vis spectroscopic behaviour of salts with a trimethinium cyanine cation and the tetrakis(phenylethynyl)borate anion [404], in non-dissociating solvents of low relative permittivity. Clearly, in solutions of such low relative permittivity any ionic species will be highly associated. However, it has been found that the ion pairs formed can be smaller than the sum of the van der Waals radii of the components. Clearly, the ions of the ion pair interpenetrate each other depending on their molecular structure: in the first case, the BF_4^- ion penetrates into the voids between the alkyl chains of the tetraalkylammonium ion, and in the second case the cyanine cation penetrates into the crevices of the borate ion.

It should be mentioned that the ionization step in Eq. (2-13) is analogous to that involved in S_N1 and S_N2 reactions of aliphatic substrates. For example, in solvolytic reactions of haloalkanes, the process of going from a covalently bonded initial state to a dipolar or ionic activated complex (transition state) is similar to the ionization step in Eq. (2-13). Therefore, those solvent properties that promote ionization are also important in the estimation of solvent effects on nucleophilic displacement reactions [161] (*cf.* Section 5.4.1).

The ionization of an ionogen and its subsequent dissociation according to Eq. (2-13) can be further elaborated. Between the ion pair immediately formed on heterolysis of the covalent bond and the independently solvated free ions, there are several steps of progressive loosening of the ion pair by penetration of solvent molecules between the ions. At least four varieties of ion interactions representing different stages of dissociation have been postulated [96, 134, 138, 141]; *cf.* Eq. (2-19) and Fig. 2-14.



(2-19)

Based on the mutual geometric arrangement of the two ions and the solvent molecules, the following definitions of ion pairs have been given (*cf.* Fig. 2-14).



Fig. 2-14. Schematic representation of the equilibrium between (a) a solvated contact ion pair, (b) a solvent-shared ion pair, (c) a solvent-separated ion pair, and (d) unpaired solvated ions of a 1:1 ionophore in solution, according to reference [241]. Hatched circles represent solvent molecules of the primary solvation shell.

First, immediately after ionization, *contact ion pairs**) are formed, in which no solvent molecules intervene between the two ions that are in close contact. The contact ion pair constitutes an electric dipole having only one common primary solvation shell. The ion pair separated by the thickness of only one solvent molecule is called a *solvent-shared ion pair**). In solvent-shared ion pairs, the two ions already have their own primary solvation shells. These, however, interpenetrate each other. Contact and solvent-shared ion pairs are separated by an energy barrier which corresponds to the necessity of creating a void between the ions that grows to molecular size before a solvent molecule can occupy it. Further dissociation leads to *solvent-separated ion pairs**). Here, the primary solvation shells of the two ions are in contact, so that some overlap of secondary and further solvation shells takes place. Increase in ion-solvating power and relative permittivity of the solvent favours solvent-shared and solvent-separated ion pairs. However, a clear experimental distinction between solvent-shared and solvent-separated ion pairs is not easily obtainable. Therefore, the designations solvent-shared and solvent-separated ion pairs are sometimes interchangeable. Eventually, further dissociation of the two ions leads to *free*, *i.e.* unpaired solvated *ions* with independent primary and secondary solvation shells. The circumstances under which contact, solvent-shared, and solvent-separated ion pairs can exist as thermodynamically distinct species in solution have been reviewed by Swarcz [138] and by Marcus [241].

Interestingly, theoretical calculations of Gibbs energy profiles for the separation of *tert*-butyl cation and chloride ion during the hydrolysis of 2-chloro-2-methylpropane have given support for the existence of a contact ion pair, while solvent-separated ion pairs and free, unpaired ions do not appear as energetically distinct species [302]. Monte Carlo simulations predict the occurrence of a contact ion pair at a C—Cl distance of 290 pm and the onset of the solvent-separated ion pair regime near 550 pm (*cf.* the normal C—Cl bond length of ca. 180 pm). A significant barrier of ca. 8 kJ/mol (2 kcal/mol) between the contact and solvent-separated ion pairs has been calculated [302]. For tetramethylammonium chloride in dilute aqueous solution at 25 °C, the contact and

* Some authors use the designations *intimate ion pair*, *internal ion pair* (Winstein [162]), *cage ion pair* (Kosower [129]), or *inner-sphere ion pair* (Marcus [241]) instead of *contact ion pair*, and *external ion pair* (Winstein [162]) or *outer-sphere ion pair* (Marcus [241]) for *solvent-shared* and *solvent-separated ion pairs*. The more general designation *tight* and *loose ion pair* (Swarcz [138]) implies that, in principle, more than two different kinds of ion pairs may exist in solution. An IUPAC glossary recommends the designations *tight ion pair* (or *intimate* or *contact ion pair*) and *loose ion pair* [286].

solvent-separated ion pairs are separated by a calculated activation barrier of only 2.9 kJ/mol (0.7 kcal/mol) [302]. Analogous Monte Carlo simulations for sodium iodide ion pairs in water clusters substantiate the existence of distinct contact and solvent-separated ion pairs, showing that the $Na^+ I^-$ contact ion pair is quite stable with respect to dissociation into free ions [405].

The suggestion that ion pairs may exist in more than one distinct form was made by Winstein [162] and by Fuoss [163] in 1954, but direct evidence for the existence of contact and solvent-separated ion pairs came from UV/Vis spectroscopic investigations of sodium fluorenide in tetrahydrofuran solution [141, 164]. Further evidence for the existence of a dynamic equilibrium between contact and solvent-separated ion pairs (*e.g.* hyperfine splitting of radical-anion ESR lines by cationic nuclei; electronic spectra of mesomeric anions; *etc.*) has been summarized by Gordon [96], Szwarc [138], and Marcus [241]. Increasing association of ions in solution greatly affects their chemical behaviour. A large variety of possible ion-pair effects on rate constants, mechanism and stereochemistry is known, especially in reactions of ion pairs containing carbenium ions [161, 165] or carbanions [166, 168, 168a].

The observation that the rate of loss of optical activity during the solvolysis of certain chiral substrates $R^{\delta\oplus} - X^{\delta\ominus}$ exceeded the rate of acid production and the occurrence of a special salt effect led to the postulation of two distinct ion-pair intermediates [161, 162]. The basic Winstein solvolysis scheme is given by Eq. (2-20).

$$\begin{array}{ccccccc} (R^{\delta\oplus}-X^{\delta\ominus})_{solv} & \rightleftharpoons & (R^{\oplus}\,X^{\ominus})_{solv} & \rightleftharpoons & (R^{\oplus}//\,X^{\ominus})_{solv} & \rightleftharpoons & (R^{\oplus})_{solv} + (X^{\ominus})_{solv} \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \text{products} & & \text{products} & & \text{products} \end{array} \qquad (2\text{-}20)$$

According to this scheme, the solvolysis products are not only obtained from free unpaired ions, but also from the two different ion pairs, depending on the solvent-dependent degree of dissociation.

An analogous scheme holds for the reactions of certain dipolar organometallics $R^{\delta\ominus} - M^{\delta\oplus}$, according to Eq. (2-21) [138, 167, 168, 168a].

$$\begin{array}{ccccc} (R^{\delta\ominus}-M^{\delta\oplus})_{n,solv} & \rightleftharpoons & (R^{\delta\ominus}-M^{\delta\oplus})_{solv} & \rightleftharpoons & (R^{\ominus}\,M^{\oplus})_{solv} \\ \text{Higher Aggregates} & & \text{Ionogen} & & \text{Contact Ion Pair} \end{array}$$

$$\rightleftharpoons (R^{\ominus}//\,M^{\oplus})_{solv} \rightleftharpoons (R^{\ominus})_{solv} + (M^{\oplus})_{solv} \qquad (2\text{-}21)$$

Solvent-separated Ion Pair Free Ions

Whereas the spectral behavior of solvent-separated ion pairs and free ions is very similar, the UV/Vis spectra of contact and solvent-separated ion pairs are usually different from each other, as has been shown with sodium fluorenide [141, 164]. Due to the penetration of solvent molecules between the ion-pair couples, the direct influence of the metal cation on the π-electron system of the carbanion is lost. With increasing dissociation, the absorption maximum of sodium fluorenide in tetrahydrofuran solution is shifted bathochromically in the direction of the absorption maximum of the free

fluorenide ion: $\lambda_{max} = 356$ nm $\rightarrow$ 373 nm $\rightarrow$ 374 nm, for the contact ion pairs, solvent-separated ion pairs, and free fluorenide ions, respectively [164]. The equilibrium between contact and solvent-separated ion pairs is shifted in the direction of increased dissociation by the addition of cation solvators such as EPD solvents. Thus, the proportion of solvent-separated ion pairs for sodium fluorenide at 25 °C in tetrahydrofuran is 5 cmol/mol, whereas in 1,2-dimethoxyethane, a better cation solvator, it is 95 cmol/mol. In strong EPD solvents such as dimethyl sulfoxide, hexamethylphosphoric triamide, or polyethyleneglycol dimethyl ethers, most of the fluorenide salt exists as solvent-separated ion pairs only. Small quantities of dimethyl sulfoxide, when added to the sodium fluorenide solution in 1,4-dioxane, convert the contact ion pairs to dimethyl sulfoxide-separated ion pairs [141, 164].

Sodium naphthalenide behaves similarly when the solvent is changed from tetrahydrofuran to 1,2-dimethoxyethane. The formation of solvent-separated from contact ion pairs is shown by a dramatic simplification of the ESR spectrum: the 100-line spectrum of the contact ion pair, due to the spin-spin coupling of the unpaired electron with the four equal hydrogen nuclei in the α- and β-positions, together with the sodium nucleus ($I = 3/2$), collapses to a 25-line spectrum as the interaction with the sodium ion is disrupted [169, 170].

Other illustrative examples of carbanionic ion-pair dissociation/aggregation are: lithium triphenylmethide, which exists as a tight ion pair in diethyl ether and as a solvent-separated ion pair in tetrahydrofuran, as shown by UV/Vis spectrophotometric measurements [287], and lithium 10-phenylnonafulvene-10-oxide, which exists as a tight ion pair *(2b)* in tetrahydrofuran solution and as a solvent-separated ion pair *(3b)* when hexamethylphosphoric triamide or dimethyl sulfoxide are added (^{1}H and ^{13}C NMR measurements) [288].

This second case is particularly interesting since the addition of an EPD solvent is connected with a shift from the olefinic nonafulvenoxide anion in *(2b)* to the aromatic benzoyl [9] annulene anion in *(3b)*. Without association of the lithium cation with the enolate oxygen atom, the negative charge is preferably delocalized in the [9] annulene ring. Therefore, the aromatic character of this ionophore depends on its ion-pair character [288].



(2b) *(3b)*

The degree of aggregation of organolithium compounds (alkyl-, aryl-, and alkynyl-lithium compounds as well as lithium enolates) in dilute tetrahydrofuran solution at −108 °C has been determined by means of cryoscopic [289] and NMR spectroscopic measurements [290]; for a review on the solution structure of lithium enolates and phenolates, see reference [406].

The enolate and iminate ions of tetra-*n*-butylammonium salts of carbonyl compounds (*e.g.* malonates) and nitriles (*e.g.* 2-phenylpropionitrile) exhibit special dimeric molecular structures in the solid state and in solution (benzene), held together by multi-

ple C—H···O and C—H···N hydrogen bonds, resp., with the α-methylene units of the $(n\text{-Bu})_4N^+$ cation. Thus, these anions are not truly 'naked' carbanions; they interact with one another through hydrogen bonds in a highly ordered manner, leading to another type of ion pairs, called *supramolecular ion pairs* [407]. For a review on genuine non-coordinating anions, see reference [408].

Of particular interest for regio- and stereoselective C—C bond-forming synthetic reactions are lithium organocuprates, the detailed molecular structure of which was unknown for a long time. Application of sophisticated NMR techniques has shown that a representative salt-containing lithium dimethylcuprate, $Me_2CuLi \cdot LiCN$, exists in solution (S = THF, Et_2O) in an equilibrium between homo-dimeric contact ion pairs and monomeric solvent-separated ion pairs such as $[LiS_4]^+$ [Me—Cu—Me]$^-$ [409]. A systematic X-ray study of solid-state structures of lithium organocuprates has substantiated the formation of monomeric solvent-separated ion pairs in good-solvating solvents for Li^+ (*e.g.* THF, crown ethers, amines), while in poor-solvating solvents for Li^+ (*e.g.* Et_2O, Me_2S) a dimeric contact ion pair is found. This is of practical relevance because it seems to be only the lithium organocuprate dimer of the contact-ion type that undergoes C—C bond-forming reactions such as addition to enones [409].

The ^{7}Li NMR spectra of solutions of the dilithium salts of the (*R*)- and (*S*)-configured (*sec*-butoxy)cyclooctatetraene dianion, $Li_2^+[C_8H_7—OC_4H_9]^{2-}$, in the chiral solvent 1,4-bis(dimethylamino)-2,3-dimethoxybutane (DDB; see Table A-2 in the Appendix) were found to be remarkably different. The chiral dilithium salt exists in DDB solution as a mixture of contact and solvent-separated ion pairs. Interestingly, the relative concentration of the contact ion pair is much greater for the (*R*)-enantiomer of $[C_8H_7—OC_4H_9]^{2-}$ than for the (*S*)-enantiomer, indicating a solvent/ion-pair chiral recognition. Thus, the interaction between the chiral solvent and the (*R*)- and (*S*)-*sec*-butoxy groups results in the DDB solvent being more capable of partially separating Li^+ from the (*S*)-enantiomer of $[C_8H_7—OC_4H_9]^{2-}$ than from the (*R*)-enantiomer [410].

3 Classification of Solvents

Due to the physical and chemical differences between the numerous organic and inorganic solvents it is difficult to organise them in a useful scheme. Here, five attempts at a classification of solvents are presented, which should prove useful to the chemist. Due to broad definitions, some overlapping of these is unavoidable. As has been customary in previous reviews non-aqueous organic solvents will receive particular attention [1–15, 103–108, 172, 174–177]. Extensive compilations of chemical and physical properties of non-aqueous solvents can be found in references [11–14, 104, 106, 175, 177].

3.1 Classification of Solvents according to Chemical Constitution

Solvents can be classified according to their chemical bonds: (a) molecular liquids (molecule melts; covalent bonds only), (b) ionic liquids (molten salts; only ionic bonds), and (c) atomic liquids (low-melting metals like liquid mercury or liquid sodium; metallic bonds) [16]. Numerous transitions are possible by mixing solvents of these three classes (Fig. 3-1). However, research into this area is still far from exhausted.

The customary non-aqueous organic solvents belong to the group of molecular melts and, according to their chemical constitutions, to the following classes of compounds (*cf.* Table A-1, Appendix): aliphatic and aromatic hydrocarbons and their halogen and nitro derivatives, alcohols, carboxylic acids, carboxylic esters, ethers, ketones, aldehydes, amines, nitriles, unsubstituted and substituted amides, sulfoxides, and sulfones. The classification of solvents according to chemical constitution allows certain qualitative predictions, summarized in the old rule "similia similibus solvuntur". In general, a compound dissolves far more easily in a solvent possessing related functional groups than in one of a completely different nature. A proper choice of solvent, based on the knowledge of its chemical reactivity, helps to avoid undesired reactions between solute and solvent. For example, condensations should not be carried out in solvents possessing carbonyl groups (*e.g.* ketones) or hydrolyses in carboxylic esters, amides, or nitriles.

Further well-known molecular liquids that have recently found renewed interest as solvents for many applications are water and perfluorohydrocarbons. These occupy the two extreme positions on empirical solvent polarity scales, with water being the most polar solvent and perfluorohydrocarbons being among the least polar of solvents (*cf.* Chapter 7).

Plain water has undergone a magnificent renaissance as a solvent for organic reactions, not only because it is an environmentally safe solvent, but also due to its pronounced capacity for hydrophobic hydration of apolar solutes [178–181]; *cf.* Sections 2.2.7 and 5.5.8 as well as Table A-14 in the Appendix.

Perfluorohydrocarbons are nonpolar, hydrophobic, chemically inert, and nontoxic solvents with a higher density than the corresponding hydrocarbons. They show a high capacity to dissolve gases (*e.g.* oxygen) and a temperature-dependent mutual miscibility with common organic solvents, mostly forming biphasic systems at ambient temperature [182]. These properties make them useful solvents for spectroscopic measurements [183] and for various kinds of organic reactions [184]. With different solubil-

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8



Fig. 3-1. Classification of solvents according to their characteristic chemical bonds [16].

ities for educts, reagents, products, and catalysts, biphasic solvent systems with a 'fluorous' phase can facilitate the separation of products from reaction mixtures [182, 185, 186]; *cf.* Section 5.5.13 and Table A-14 in the Appendix.

Liquid crystals or mesomorphic compounds occupy a special position [17–22, 22a, 109, 110]. Compounds capable of forming liquid crystals are long, flat, and fairly rigid along the axis of the molecule. Most known mesomorphic solvents are systems of the following general structure with polarizable aromatic nuclei held in a planar skeleton:



Common central bridges:
—CH_2—CH_2, —CH=CH—,
—C≡C—, —CH=N—,
—CH=N(O)—, —CO—O—,
—N=N—, —N=N(O)—,



Common terminal substituents:
R—, RO—, HO—, R—CO—O—,
R—O—CO—O—,
R—O—$(CH_2)_n$—O—, H_2N—,
O_2N—,
N≡C—, Cl—, Br—, I—

Unlike normal isotropic liquids, which possess a completely random arrangement of molecules, liquid crystals are considerably ordered. The degree of order in the latter lies somewhere between that of isotropic liquids and crystals. Liquid crystals are classified into lyotropic and thermotropic crystals depending on the way in which the mesomorphic phase is generated. *Lyotropic* liquid-crystalline solvents are formed by addition of controlled amounts of polar solvents to certain amphiphilic compounds. *Thermotropic* liquid-crystalline solvents, simply obtained by temperature variations, can be further classified into nematic, smectic, and cholesteric solvents depending on the type of molecular order present. In *nematic* mesophases, the long molecular axes of the component molecules are arranged, for the most part, parallel to one another. There is no further ordering present and these are the most fluid of liquid-crystalline solvents. *Cholesteric* mesophases are optically active nematic phases, additionally characterized by a gradual twist in orientational alignment as one proceeds through the bulk solvent, and forming a twisted helical macrostructure. In *smectic* mesophases, the component mole-

cules are further arranged in layers, with their long molecular axes parallel to one another and perpendicular to the plane of the layers. As a result of this additional molecular ordering, smectic solvents are considered to be the most ordered and the less fluid liquid-crystalline solvents. A typical example of a liquid-crystalline nematic solvent, at room temperature, is *N*-(4-methoxybenzylidene)-4-*n*-butylaniline ("MBBA") [23]. At 21 °C, the crystalline structure is lost and an ordered fluid, thermally stable up to about 48 °C, is formed (nematic range 21 to 48 °C).



$$\text{Solid phase} \overset{21\ ^\circ\text{C}}{\rightleftharpoons} \text{Nematic phase} \overset{48\ ^\circ\text{C}}{\rightleftharpoons} \text{Isotropic phase}$$

Liquid crystals are usually excellent solvents for other organic compounds. Non-mesomorphic solute molecules may be incorporated into liquid-crystalline solvents without destruction of the order prevailing in the liquid-crystalline matrix. The anisotropic solute-solvent interaction leads to an appreciable orientation of the guest molecules with respect to the axis of preferred solvent alignment. The consequences may be useful as shown by the use of liquid crystals as anisotropic solvents for spectroscopic investigations of anisotropic molecular properties [166]. Ordered solvent phases such as liquid crystals have also been used as reaction media, particularly for photochemical reactions; *cf.* for example [111, 155, 163] and Section 5.5.9.

The second corner of the triangle in Fig. 3-1 is occupied by ionic liquids. Of these, molten salts are becoming increasingly important as solvents for inorganic as well as organic reactions [3, 24–30, 112–114]. High thermal stability, good electrical conductivity*), low viscosity, wide liquid range, low vapour pressure and the resulting possibility of high working temperatures, together with their excellent ability to dissolve salts and metals, make them extremely useful reaction media. Therefore, such systems are becoming increasingly important technologically. A further advantage of molten salts is their high heat conductivity, which permits a very rapid dispersal of the heat of reaction.

Often, these are the only solvents capable of dissolving salt-like hydrides such as NaH and CaH_2, carbides, nitrides, various oxides, sulfides, and cyanamides. Many metals form atomic dispersions in the melts of their halogenides, yielding extremely strongly reducing solutions. The working temperature for oxide and silicate melts lies above 1500 °C, for normal salts it is between 100 and 1000 °C, and for eutectic mixtures it is often at even lower temperatures. Salts with large organic cations such as tetraalkylammonium, tetraalkylphosphonium, 1,3- and 1,4-dialkylpyridinium, or 1,3-dialkylimidazolium ions, and with suitable inorganic anions, have melting points

* Ionic melts possess electrical conductivities roughly a factor ten larger than those of concentrated aqueous solutions of strong electrolytes (ionophores).

Table 3-1. Melting points (t_{mp}), boiling points (t_{bp}), and liquid working ranges, $\Delta t = t_{bp} - t_{mp}$, of some fused inorganic and organic salt systems [27, 28], including a few room temperature ionic liquids (entries 17, 18, 21–28) [187–190]. For binary or ternary eutectics, the figures in parentheses give the portions in cmol/mol.

Fused salts	t_{mp}/°C	t_{bp}/°C	Δt/°C
(1) NaCl[a]	801	1465	664
(2) KCl[a]	771	1413	642
(3) LiCl[a]	610	1383	773
(4) KOH[a]	406	1327	921
(5) NaOH[a]	323	1388	1065
(6) $NaNO_3$[a]	308	380	73
(7) $K^+H_3C—CO_2^-$[a]	292	–	–
(8) $ZnCl_2$[a]	290	732	442
(9) $NaNH_2$[a]	210	500 (dec.)	290
(10) $ZnCl_2$(60)—NaCl(20)—KCl(20)	203	–	–
(11) KCl(33)—$AlCl_3$(67)	128	–	–
(12) *N*-Butyl-3-methylpyridinium chloride	98	–	–
(13) $AlBr_3$[a]	97	255	158
(14) $LiNO_3$(25.8)—NH_4NO_3(66.7)—NH_4Cl(7.5)	86	–	–
(15) $AgNO_3$(52)—$TlNO_3$(48)	82.5	–	–
(16) $(n\text{-}C_6H_{13})_4N^+NO_3^-$	69	–	–
(17) $(n\text{-}C_4H_9)_3NH^+NO_3^-$[b]	21.5	119 (dec.)	–
(18) $CH_3—CH_2—NH_3^+NO_3^-$[b,c]	12	170 (dec.)	–
(19) $[EMIM]^+Cl^-$[d]	87	–	–
(20) $[EMIM]^+PF_6^-$[d]	62	–	–
(21) $[EMIM]^+AlCl_4^-$[d]	7	–	–
(22) $[EMIM]^+F_3C—SO_3^-$[d]	−9	–	–
(23) $[EMIM]^+F_3C—CO_2^-$[d]	−14	–	–
(24) $[EMIM]^+(F_3C—SO_2)_2N^-$[d]	−15	–	–
(25) $Li^+(F_3C—SO_2)_2N^-$/Urea (1:3)[e]	−37.6	–	–
(26) $(n\text{-}C_6H_{13})_4N^+H_5C_6—CO_2^-$[f]	−50[a]	–	–
(27) $(n\text{-}C_6H_{13})(C_2H_5)_3N^+(n\text{-}C_6H_{13})(C_2H_5)_3B^-$[g]	<-75[a]	87 (dec.)	–
(28) $(CH_3)_2NH_2^+(CH_3)_2N—CO_2^-$ ("Dimcarb")[h]	–	60 (dec.)	–

[a] D. R. Lide (ed.): *CRC Handbook of Chemistry and Physics*, 77th ed., CRC Press, Boca Raton/FL, USA, 1996.
[b] C. F. Poole, B. R. Kersten, S. S. J. Ho, M. E. Coddens, K. G. Furton, J. Chromatogr. *352*, 407 (1986).
[c] H. Weingärtner, A. Knocks, W. Schrader, U. Kaatze, J. Phys. Chem. A *105*, 8646 (2001).
[d] $[EMIM]^+$ = 1-*E*thyl-3-*m*ethyl*im*idazolium ion.
[e] H. Liang, H. Li, Z. Wang, F. Wu, L. Chen, X. Huang, J. Phys. Chem. B *105*, 9966 (2001).
[f] C. G. Swain, A. Ohno, D. K. Roe, R. Brown, T. Maugh, J. Am. Chem. Soc. *89*, 2648 (1967); T. G. Coker, J. Ambrose, G. J. Janz, ibid. *92*, 5293 (1970).
[g] For other liquid tetraalkylammonium tetraalkylborides, see W. T. Ford, R. J. Hauri, D. J. Hart, J. Org. Chem. *38*, 3916 (1973).
[h] W. Schroth, J. Andersch, H.-D. Schädler, R. Spitzner, Chemiker-Ztg. *113*, 261 (1989); Z. Chem. *29*, 56, 129 (1989).

under 100 °C and are often liquid even at room temperature. Table 3-1 gives a somewhat arbitrary selection of inorganic and organic salts, eutectic salt mixtures [27, 28], and room temperature liquid salts [187–189], together with their physical constants.

Although the relative permittivities of molten salts are generally quite small ($\varepsilon_r = 2 \ldots 3$), they behave as strongly dissociating solvents! This is due to the ability of the solvent ions to exchange places with solute ions of the same charge.

Three examples of the many reactions that can be advantageously carried out in high-temperature molten salts have been selected to illustrate this molten-salt technique. When 1,1-dichloroethane is passed through a $ZnCl_2/KCl$ melt at 330 °C, chloroethene is formed in 97 cmol/mol yield by dehydrohalogenation. Likewise, the addition of hydrogen chloride to acetylene proceeds with 89 cmol/mol yield in the same melt. A combination of these two steps allows a ready synthesis of the technically important chloroethene from acetylene and 1,1-dichloroethane according to Eq. (3-1) [25, 31].

$$HC{\equiv}CH + H_3C\text{-}CHCl_2 \xrightarrow[\text{yield 68.5 cmol/mol}]{\text{in } ZnCl_2/KCl/HgCl_2} 2\ H_2C{=}CH\text{-}Cl \qquad (3\text{-}1)$$

Ionic liquids such as the eutectic $AgNO_3/KNO_3/AgCl$ (mp 113 °C) are the best solvents for effecting silver ion-catalyzed isomerizations of cage-like organic molecules, *e.g.* basketane → snoutane [31a], *cf.* Eq. (3-2).



(3-2)

Molten sodium tetrachloroaluminate (a 1:1 mixture of NaCl and $AlCl_3$) is a good reaction medium for the Friedel-Crafts acylation reaction given in Eq. (3-3) [115].



(3-3)

Whereas the classical procedure for the synthesis of 1-indanone from 3-phenylpropanoic acid consists of three reaction steps with a total reaction time of *ca.* six hours [116], the molten salt reaction is finished in five minutes and gives an even better yield [115].

Ethylammonium nitrate (entry 18 in Table 3-1) was shown in 1914 to have m.p. 12 °C and was hence the first room temperature ionic liquid [156]; this was followed in 1967 by tetra-*n*-hexylammonium benzoate with m.p. −50 °C (entry 26) [169]. Ambient-temperature ionic liquids based on 1-alkyl-3-methylimidazolium salts (entries 19–24) were first reported by Wilkes *et al.* in 1982 as tetrachloroaluminates [162a]. Replacement of this moisture-sensitive anion by the tetrafluoroborate ion and other anions led, in 1992, to air- and water-stable, room temperature ionic liquids [162b], which have since found increasing application as reaction media for various kinds of organic reactions, mainly owing to the work of Seddon [167, 190] and Hussey [187]. Suitably selected

combinations of cation and anion (*cf.* entries 19–24 in Table 3-1) allow the design of ionic liquids that meet all requirements for the chemical reaction under study. Therefore, they have also been called *designer solvents.* The polarities of room temperature ionic liquids have been determined empirically using solvatochromic dyes, which revealed polarities varying between that of acetonitrile and those of lower alcohols [191, 192]. For the use of room temperature ionic liquids in synthesis and catalysis, see reviews [187–190] and Section 5.5.13. Section 5.5.13 describes applications of some *neoteric solvents* such as ionic liquids, perfluorohydrocarbons, etc., in organic synthesis. Neoteric (=recent, new, modern) indicates a class of novel reaction media that have remarkable new properties [167].

A remarkable solvent on the borderline between ionic and molecular liquids is dimethylammonium *N*,*N*-dimethylcarbamate (''Dimcarb''; see entry 28 in Table 3-1), a colourless distillable liquid that readily decomposes at its b.p. into dimethylamine and carbon dioxide, from which it is prepared at lower temperatures. Being formally an ionic liquid, owing to the weak bond between the two components, it has a variable hydrogen-bonded molecular structure with one dimethylamine moiety rapidly exchanging between the ammonium and carbamate ions. Its physical and chemical properties have been reviewed [193]. Reaction of dimethylamine with sulfur dioxide also yields an ionic liquid, presumably dimethylammonium *N*,*N*-dimethylamidosulfinate, (''Dimsulf''; b.p. 112 °C), a useful electrolyte for organic electrochemistry [194].

Another type of solvents on the borderline between ionic and molecular liquids are highly concentrated solutions of ionophores in nonpolar organic solvents, such as solutions of lithium perchlorate in diethyl ether (LPDE) [195–197]. Anhydrous lithium perchlorate is a solid with m.p. 247 °C; a 5 M solution of $LiClO_4$ in dry diethyl ether contains only about one to two ether molecules per lithium cation and can be considered as a diluted ionic liquid with the structure of a dietherate, $[Li(OEt_2)_2]^+ ClO_4^-$, below 4.25 M, and a mixture of both the dietherate and monoetherate, $[Li(OEt_2)]^+ ClO_4^-$, above 4.25 M. The lithium cation is not fully coordinated by solvent molecules and such ions are sometimes mistakenly called 'naked' ions; naked ions only exist in the gas phase. Nevertheless, the partially coordinated lithium cation is highly electrophilic and ethereal lithium perchlorate solutions can be used for many organic reactions requiring electrophilic catalysis. As early as 1959, Winstein recognized that addition of lithium perchlorate to less well coordinating organic solvents such as diethyl ether increases not only their polarity but also their Lewis acid/base properties [198]. The manifold applications of LPDE in organic synthesis have been reviewed [195–197].

The third corner of the triangle in Fig. 3-1 represents liquid metals, *e.g.* mercury or liquid sodium, which until now have received little attention as reaction media. Chemical reactions in liquid alkali metals have been reviewed [164, 199].

3.2 Classification of Solvents using Physical Constants

The following physical constants can be used to characterize the properties of a solvent: melting and boiling point, vapour pressure, heat of vapourization, index of refraction, density, viscosity, surface tension, dipole moment, relative permittivity, polarizability, specific conductivity, *etc.* A compilation of data of usual organic solvents is given in

Table A-1 (Appendix). For more comprehensive recent compilations of physical data for common organic solvents, see references [175] (260 solvents) and [177] (1141 solvents).

Solvents can be broadly classified as low, middle, or high boiling, *viz.* $t_{bp} < 100$ °C, 100 ... 150 °C, or >150 °C at 1 bar. Similarly, liquids can be classified according to their *evaporation number* using diethyl ether as reference (evaporation number = 1 at 20 °C and 65 cL/L relative air humidity). Thus, high volatility is signified by evaporation numbers <10, moderate volatility by numbers 10 ... 35, low volatility by numbers 35 ... 50, and very low volatility by numbers >50 [32, 177]. Using viscosity as a criterion, solvents are of low viscosity when their dynamic viscosity is <2 mPa · s at 20 °C, of medium viscosity between 2 and 10 mPa · s, and of high viscosity above 10 mPa · s [32].

The degree of association of molecules in a liquid can be estimated by means of its *Trouton constant* [117]. At the normal boiling temperature, T_{bp}, vapourization proceeds with standard molar changes of enthalpy, ΔH°_{bp}, and entropy, ΔS°_{bp}, from which Trouton's rule is derived as given in Eq. (3-4).

$$\Delta S^{\circ}_{bp} = \frac{\Delta H^{\circ}_{bp}}{T_{bp}} \approx 21 \text{ cal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1} \text{ or } 88 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1} \quad (3\text{-}4)^{*)}$$

This rule works best for apolar, quasi-spherical molecules. Large deviations occur when chemical association is involved (*e.g.* carboxylic acids), from molecular dipolarity (*e.g.* dimethyl sulfoxide), and from molecular asphericity (*e.g.* neopentane/*n*-pentane). Strongly associating solvents (*e.g.* HF, H_2O, NH_3, alcohols, carboxylic acids) have Trouton constants which are higher than the average value of 88 J · mol^{-1} · K^{-1} found for non-associating solvents such as diethyl ether and benzene.

In addition to Trouton's rule, some other parameters for measuring the *structuredness* of solvents have been recommended, for example a solvent dipole orientation correlation parameter [175, 200], the solvent's heat capacity density [175, 200], and a so-called S_p parameter derived from the solvent's enthalpy of vapourization minus EPD/EPA and van der Waals interactions [201]. According to these parameters, solvents can be classified as highly structured (*e.g.* water, formamide), weakly structured (*e.g.* DMSO, DMF), and practically non-structured (*e.g. n*-hexane and other hydrocarbons) [200, 201].

In this connection, two other physical solvent properties are important: the *cohesive pressure c* (also called *cohesive energy density*) and the *internal pressure* π of a solvent [98–100, 175].

The *cohesive pressure c* is a measure of the total molecular cohesion per unit volume, given by Eq. (3-5),

$$c = \frac{\Delta U_v}{V_m} = \frac{\Delta H_v - R \cdot T}{V_m} \quad (3\text{-}5)$$

* Trouton's rule can also be written independent of units: $\Delta S^{\circ}_{bp}/R = \Delta H^{\circ}_{bp}/(R \cdot T_{bp}) \approx 11$.

where ΔU_v and ΔH_v are respectively the energy and enthalpy (heat) of vapourization of the solvent to a gas of zero pressure, and V_m is the molar volume of the solvent. On vapourization of a solvent to a non-interacting vapour, *all* intermolecular solvent-solvent interactions will be broken. Therefore, c represents the total strength of the intermolecular solvent structure. Cohesive pressure has very high values for solvents of high polarity and low values for nonpolar solvents such as perfluorohydrocarbons with weak interaction forces. Intermolecular hydrogen bonding in a solvent increases the cohesive pressure (*cf.* Table 3-2). Cohesive pressure is related to the energy required to create cavities in a liquid in order to accommodate solute molecules during the process of dissolution.

On the other hand, the *internal pressure* π is defined as the change in internal energy of a solvent as it undergoes a very small isothermal expansion, as seen in Eq. (3-6).

Table 3-2. Cohesive pressures, c, internal pressures, π, and their ratio $n = \pi/c$ for thirty organic solvents, arranged in order of decreasing n, that is, in order of increasing "structuredness", at 20 °C [99, 154, 175].

Solvents	c/MPa[a)]	π/MPa[a)]	$n = \pi/c$
Perfluoro-*n*-heptane	151	215	1.42
Perfluoro(methylcyclohexane)	161	228	1.41
1,4-Dioxane	388	494	1.27
2,2,4-Trimethylpentane	200	236	1.18
Methylcyclohexane	260	297	1.14
Cyclohexane	285	326	1.14
Hexamethylphosphoric acid triamide (HMPT)	365	403	1.10
Tetrachloromethane	310	339	1.09
Ethyl acetate	331	356	1.08
n-Hexane	225	239	1.06
Benzene	353	369	1.05
Toluene	337	355	1.05
Diethyl ether	251	264	1.05
1,2-Dichloroethane	400	419	1.05
Trichloromethane (Chloroform)	362	370	1.02
Acetylbenzene (Acetophenone)	456	457	1.00
Dichloromethane	414	408	0.99
Acetic acid	357	348	0.97
Carbon disulfide	412	372	0.90
N,*N*-Dimethylformamide (DMF)	581	480	0.83
Dimethyl sulfoxide (DMSO)	708	521	0.74
2-Methyl-2-propanol, *t*-Butanol	467	339	0.73
Acetone	488	331	0.68
Acetonitrile	581	395	0.68
1-Butanol	485	300	0.62
Ethane-1,2-diol	1050	502	0.48
Ethanol	676	293	0.43
Formamide	1568	554	0.35
Methanol	858	288	0.34
Water	2294	151	0.07

[a)] 1 MPa $= 10^6$ Pa $= 1$ J $\cdot$ cm$^{-3} = 9.87$ atm.

$$\pi = \left(\frac{\partial U}{\partial V_m}\right)_T \tag{3-6}$$

(U = molar internal energy; V_m = molar volume; T = absolute temperature). This small expansion does not necessarily disrupt all the intermolecular solvent-solvent interactions. The internal pressure results from the forces of attraction between solvent molecules exceeding the forces of repulsion, *i.e.* mainly dispersion and dipole-dipole interactions (*cf.* Table 3-2).

Although there is obviously a close connection between cohesive pressure and internal pressure, they are not equivalent, as shown by the compilation of c and π values for some selected organic solvents shown in Table 3-2 [99, 100, 154]. It has been assumed that π is mainly a reflection of dispersion and dipole-dipole interactions within the solvent, whereas c additionally includes specific solvent-solvent interactions such as hydrogen bonding. Hydrogen bonding in a solvent increases the cohesive pressure, while the internal pressure is comparable to that of solvents without hydrogen bonding. Therefore, the hydrogen-bonding pressure or energy density contribution can be measured by the difference $(c - \pi)$ [99, 100]. Values of π approach those of c only for weakly polar solvents with dipole moments less than *ca.* $7 \cdot 10^{-30}$ Cm (*ca.* 2 D) and without specific solvent/solvent interactions. As shown in Table 3-2, the ratio $n = \pi/c$ approaches a value of unity for nonpolar solvents (*e.g.* hydrocarbons), but it can be less than or greater than unity for other solvents. High values of n are obtained for the noninteracting fluorohydrocarbons, while at the other end of the solvent spectrum HBD solvents with very low values of n are found.

The square root of the cohesive pressure c as defined in Eq. (3-5) has been termed the *solubility parameter* δ by Hildebrand and Scott [98] because of its value in correlating and predicting the solvency of solvents for non-electrolyte solutes [*cf.* Eqs. (2-1) and (5-77) in Sections 2.1 and 5.4.2, respectively]. Solvency is defined as the ability of solvents to dissolve a compound [118]. A selection of δ values is given in Table 3-3; see also references [99, 177].

The solvency increases as the δ value of the solvent approaches that of the solute. Two liquids are miscible if their solubility parameters differ by no more than *ca.* 3 units.

Because the term *solubility parameter* is too restrictive for this quantity, which can be used to correlate a wide range of physical an chemical properties (for instance, *cf.* Section 5.4.2), the term *cohesion parameter* has been proposed by Barton [99]. The term solubility parameter suggests a close relationship between the phenomenon "solubility" or "miscibility" and that of "cohesion" or "vapourization". This seems to be reasonable, considering what happens in a mixing process: the like molecules of each component in a mixture are separated from one another to an infinite distance, comparable to what happens in the vapourization process. A comprehensive review on the determination and application of solubility parameters has been given by Barton [99].

An alternative approach for the prediction of mutual miscibility of solvents has been given by Godfrey [119] (*cf.* also Appendix, Chapter A-1). As a measure of lipophilicity (*i.e.* affinity for oil-like substances), the so-called *miscibility numbers* (M numbers, with values between 1 and 31) have been developed. These are serial numbers of 31 classes of organic solvents, ordered empirically by means of simple test tube miscibility

Table 3-3. Hildebrand solubility parameters, δ, of thirty organic solvents at 25 °C, taken from reference [99] (Table 2 in Chapter 8).

Solvents	δ/MPa$^{1/2}$[a)]	Solvents	δ/MPa$^{1/2}$[a)]
Water	47.9	Acetic acid	20.7
Formamide	39.3	1,4-Dioxane	20.5
N-Methylformamide	32.9	Carbon disulfide	20.4
Ethane-1,2-diol	29.9	Cyclohexanone	20.3
Methanol	29.6	Acetone	20.2
Tetrahydrothiophene-1,1-dioxide	27.4	1,2-Dichloroethane	20.0
Ethanol	26.0	Chlorobenzene	19.4
N,*N*-Dimethylformamide	24.8	Trichloromethane	19.0
Dimethyl sulfoxide	24.5	Benzene	18.8
Acetonitrile	24.3	Ethyl acetate	18.6
1-Butanol	23.3	Tetrahydrofuran	18.6
Cyclohexanol	23.3	Tetrachloromethane	17.6
Pyridine	21.9	Cyclohexane	16.8
t-Butanol	21.7	*n*-Hexane	14.9
Aniline	21.1	Perfluoro-*n*-heptane	11.9

[a)] $1\ \mathrm{MPa}^{1/2} = 1\ \mathrm{J}^{1/2} \cdot \mathrm{cm}^{-3/2}$.

experiments and critical solution temperature measurements. There is a close correlation between M numbers and Hildebrand's δ values [99].

The solvency of hydrocarbon solvents used in paint and lacquer formulations is empirically described by their *kauri butanol numbers*, *i.e.* the volume in milliliters at 25 °C of the solvent required to produce a defined degree of turbidity when added to 20 g of a standard solution of kauri resin in 1-butanol [120]. Standard values are $KB = 105$ for toluene and $KB = 40$ for *n*-heptane/toluene (75:25 cL/L). A high KB number corresponds to high solvent power. An approximately linear relationship exists between Hildebrand's δ values and KB numbers for hydrocarbons with $KB > 35$: $\delta = 0.06 \cdot KB + 12.9$ [99, 177].

Solvents whose molecules possess a permanent *dipole moment* are designated *dipolar* as opposed to *apolar* or *nonpolar* for those lacking a dipole moment. Unfortunately, in the literature the terms "polar" and "apolar" or "nonpolar" are used indiscriminately to characterize a solvent by its relative permittivity as well as its permanent dipole moment, even though dipole moment and relative permittivity are not directly related. Molecules possessing a centre of symmetry in all possible conformations, more than one *n*-fold axis of symmetry, or a plane of symmetry perpendicular to an *n*-fold axis of symmetry, cannot exhibit a permanent dipole moment for symmetry reasons. Therefore, only those molecules which belong to the point groups C_1, C_s, C_n, or C_{nv} can have a permanent dipole moment. The permanent dipole moments of organic solvents vary from 0 to $18.5 \cdot 10^{-30}$ Cm (0 to 5.5 D); *cf.* Appendix, Table A-1. Values of dipole moments increase steadily on going from hydrocarbon solvents to solvents containing dipolar groups such as $C^{\delta+}\cdots O^{\delta-}$, $C^{\delta+}\cdots N^{\delta-}$, $N^{\delta+}\cdots O^{\delta-}$, $S^{\delta+}\cdots O^{\delta-}$, or $P^{\delta+}\cdots O^{\delta-}$. The solvents with the highest dipole moments found to date are zwitterionic 3-alkyl-1,2,3-oxadiazolium-5-olates (sydnones). For example, 4-ethyl-3-(1-propyl)sydnone, a

high-boiling liquid ($t_{bp} = 155$ °C/3 Torr) with a high relative permittivity ($\varepsilon_r = 64.6$ at 25 °C), exhibits a dipole moment of $\mu = 35.7 \cdot 10^{-30}$ Cm (10.7 D) [202]. The orientation of dipolar solvent molecules around the solute molecule in the absence of specific solute/solvent interactions is largely determined by the dipole moment.

It should not be forgotten that the solution value of a solute dipole moment (μ_s) differs from its gas-phase value (μ_g) and depends on the nature of the solvent. For theoretical approaches to relating the difference ($\mu_s - \mu_g$) to various physical solvent parameters (*e.g.* ε_r, n_D, *etc.*), see reference [122].

The importance of electric moments of orders higher than two (dipoles) such as quadrupoles and even octupoles in solute/solvent interactions between multipolar molecules has been stressed [121]. Depending on the charge distribution, there exist multipoles (2^n-poles) such as monopoles (n = 0; *e.g.* $Na^{\oplus}$, $Cl^{\ominus}$), dipoles (n = 1; *e.g.* HF, H_2O), quadrupoles (n = 2; *e.g.* CO_2, C_6H_6), octupoles (n = 3; *e.g.* CH_4, CCl_4), and hexadecapoles (n = 4; *e.g.* SF_6). According to Reisse [121], only neutral species with a spherical charge distribution (*e.g.* rare gases) should be designated as *apolar*. All others, with non-spherical charge distribution, should be called *polar*, *i.e.* dipolar, quadrupolar, octupolar, *etc.*, depending on the first non-zero electric moment. In this respect, methane and tetrachloromethane are polar molecules, as are *cis*- and *trans*-1,2-dichloroethene (dipolar and quadrupolar, respectively), 1,3-dioxane (dipolar) and 1,4-dioxane (quadrupolar). It has been shown that the 2^n-polar contributions ($n > 1$) to solute/solvent interactions are in many cases non-negligible [121].

The *relative permittivities* play a particular role in the characterization of solvents. Their importance over other criteria is due to the simplicity of electrostatic models of solvation and they have become a useful meaure of solvent polarity. In this connection, it is important to realize what exactly is represented by the macroscopic relative permittivity of a solvent (called relative permittivity because of $\varepsilon_r = \varepsilon/\varepsilon_0$ where ε_0 is a constant, the permittivity of vacuum). Relative permittivities are determined by inserting the solvent between the two charged plates of a condenser. The strength of the electric field E between the plates is lower than the value E_0 measured when the plates are in a vacuum, and the ratio E_0/E gives the numerical value of the relative permittivity. If the solvent molecules do not have permanent dipole moments of their own, then the external field will separate the charge within the molecules thereby inducing dipoles. Molecules with induced or permanent dipoles are forced into an ordered arrangement by the charged plates, causing what is known as polarization. The larger the polarization, the larger the drop in the electric field strength. Therefore, the relative permittivity represents the ability of a solvent to separate charge and to orient its dipoles. The relative permittivities of organic solvents vary from about 2 (*e.g.* hydrocarbons) to about 180 (*e.g.* secondary amides); *cf.* Appendix, Table A-1. Solvents with large relative permittivities may act as dissociating solvents (*cf.* Section 2.6) and are therefore called *polar* solvents, in contrast to the *apolar* or *nonpolar* solvents with low relative permittivities. Relative permittivity values often run parallel to the dissolving power of the solvent, because in the case of ionic solutes (*i.e.* ionophores) solvents of high relative permittivity facilitate dissolution by separating the ions.

Since both the relative permittivity ε_r and the dipole moment μ are important complementary solvent properties, it has been recommended that organic solvents should be classified according to their *electrostatic factor EF* (defined as the product of ε_r

and μ), which takes into account the influence of both properties [101]. Considering the *EF* values and the structures of solvents, a four-part classification of organic solvents has been established: hydrocarbon solvents (*EF* $0 \ldots 7 \cdot 10^{-30}$ Cm), electron-donor solvents (*EF* $7 \ldots 70 \cdot 10^{-30}$ Cm), hydroxylic solvents (*EF* $50 \ldots 170 \cdot 10^{-30}$ Cm), and dipolar non-HBD solvents (*EF* $\geq 170 \cdot 10^{-30}$ Cm) [99, 101]. Another approach towards defining the polarity of solvents by means of their physical constants uses the ratio $\beta = A \cdot \mu^2 / V_m$, derived from electrostatic solvation theory (A = constant, μ = dipole moment, V_m = molar volume of the solvent) [203].

The *refractive index n* of a solvent is the ratio of the speed of light of a specified wavelength (usually sodium D-line; $\lambda = 589$ nm) in vacuum, c_0, to its speed in the liquid, c, according to $n_D = c_0/c$. The refractive index at a specified frequency typical of visible light is related to the relative permittivity by $n_D = {\varepsilon_r}^{1/2}$. The molar polarization P_m of a solvent is then given by the Lorenz-Lorentz equation: $P_m = (n^2 - 1)/(n^2 + 2) \cdot M_r/\rho$ (where M_r = relative molar mass, and ρ = mass density of the solvent). There is a general tendency for the refractive index to increase as the molar mass and the polarizability of the solvent increase [177, 204]. For example, diiodomethane, with $n_D = 1.748$, has one of the highest refractive indices, as compared with $n_D = 1.620$ for iodobenzene and 1.628 for carbon disulfide (see Table A-1 in the Appendix). This high refractive index for diiodomethane reflects a particularly high electron mobility around the two iodine atoms [205]. Solvents with high refractive indices and hence large polarizabilities are particularly suitable for solute/solvent dispersion interactions; see Section 2.2.4. The refractive index is closely related to the property of optical activity of chiral solvents. Optically active, chiral solvents (see later) rotate the plane of plane-polarized light, which arises from the differences in the refractive indices for right- and left-circularly polarized light, n_R and n_L, respectively. Therefore, it is not unexpected that specific rotations $[\alpha]_D$ of chiral solutes are often solvent-dependent as well [206]; see also Section 6.2.5.

As mentioned before, relative permittivities as well as dipole moments are often used in the quantitative characterization of *solvent polarity*. However, the characterization of a solvent by means of its "polarity" is an unsolved problem since the term "polarity" itself has, until now, not been precisely defined. Polarity might be interpreted as: a) the permanent dipole moment of a compound, b) its relative permittivity, or c) the sum of all those molecular properties responsible for all the interaction forces between solvent and solute molecules (*e.g.* Coulombic, directional, inductive, dispersion, hydrogen-bonding, and EPD/EPA interaction forces) [33]. The important thing concerning the so-called polarity of a solvent is its *overall solvation ability*. This, in turn, depends on the sum of all specific as well as non-specific interactions between solvent and solute. Therefore, in the following, the term "solvent polarity" will be applied according to the definition c) above. It should be noted, however, that all interactions which lead to a chemical change of the solute are excluded under this definition (*e.g.* protonation, oxidation, reduction, and complexation).

Evidently, "solvent polarity", as so-defined, is badly described in a quantitative manner by means of individual physical constants such as relative permittivity, dipole moment, *etc.* It is no surprise therefore, that the macroscopic relative permittivities are an unsuitable measure of molecular-microscopic interactions. This has often been demonstrated experimentally. One reason is that the molecular-microscopic relative permit-

tivity of the solvent in the vicinity of the solute is lower than that for the bulk solvent, because solvent dipoles in a solvation shell are less free to orientate themselves in a direction imposed by charged condenser plates. In the extreme case, complete dielectric saturation can occur for solvent molecules around an ionic solute.

The failure of the solvent relative permittivity to represent solute/solvent interactions has led to the definition of polarity in terms of empirical parameters. Such attempts at obtaining better parameters of solvent polarity by choosing a solvent-dependent standard system and looking at the changes in parameters of that system when the solvent is changed (*e.g.* rate constants of solvent-dependent reactions or spectral shifts of solvatochromic dyes) are treated in Chapter 7.

The aforementioned macroscopic physical constants of solvents have usually been determined experimentally. However, various attempts have been made to calculate bulk properties of liquids from pure theory. By means of quantum chemical methods, it is possible to calculate some thermodynamic properties (*e.g.* molar heat capacities and viscosities) of simple molecular liquids without specific solvent/solvent interactions [207]. A quantitative structure-property relationship treatment of normal boiling points, using the so-called CODESSA technique (*i.e. co*mprehensive *de*scriptors for *s*tructural and *s*tatistical *a*nalysis), leads to a four-parameter equation with physically significant molecular descriptors, allowing rather accurate predictions of the normal boiling points of structurally diverse organic liquids [208]. Based solely on the molecular structure of solvent molecules, a non-empirical solvent polarity index, called the *first-order valence molecular connectivity index*, has been proposed [137]. These purely calculated solvent polarity parameters correlate fairly well with some corresponding physical properties of the solvents [137].

Optically active organic solvents, consisting of chiral molecules*) (chiral solvents), which rotate the plane of linearly polarized light, have become increasingly important lately [35]. In principle, diastereomeric solvates are formed when a mixture of enantiomers is dissolved in an optically active solvent. Consequently, these solvates should possess slightly different physical and chemical properties. In fact, optically active solvents have already found use in stereoselective syntheses [36–38, 123], as NMR shift reagents [39, 124], in the evaluation of the optical purity of enantiomers [40], and in the gaschromatographic separation of enantiomers on chiral phases [41, 124, 125]. A selection of such optically active solvents is given in Table A-2 (Appendix).

Some recent examples of enantioselective syntheses in chiral solvents are: the high-pressure (10^4 bar) Wagner-Meerwein rearrangement of a racemic oxirane derivative in (−)-(2*S*,3*S*)-diethyl tartrate (optical yield 6.7%) [42]; the chlorination of substituted aziridines with *t*-butyl hypochlorite to give *N*-chloroaziridines in the presence of chiral trifluoromethylcarbinols (highest enantiomeric excess, $ee = 28.7\%$) [126]; the Grignard reaction of 2,2-dimethylpropanal with phenylmagnesium bromide in (−)-menthyl methyl ether ($ee = 19.4\%$) [127] and the addition of *n*-butyllithium to benzaldehyde in the presence of (+)-(*S*,*S*)-1,4-bis(dimethylamino)-2,3-dimethoxybutane (DDB) (optical yield 30%) [128], according to the following equation:

* A molecule is chiral, and hence optically active, when it does not possess planes, centres, or alternating axes of symmetry (order $n > 2$) [34].



Aldol additions between *achiral* reactants in *chiral* solvents have also been examined [157–159]. Only low asymmetric inductions ($ee = 2 \ldots 22\%$) have been found [158]. A twofold stereodifferentiation is observed in aldol additions between *chiral* reactants carried out in *chiral* solvents [159]. However, asymmetric inductions caused by chiral solvents or cosolvents are usually rather small [157].

In addition to the aforementioned conventional liquid organic solvents, two other types of molecular liquids have attracted much interest as supplementary or alternative media for synthesis and processing, namely *supercritical fluids* (sc-fluids) [209] and *metastable liquids* [210].

The importance of *supercritical fluids* as media for chemical reactions and in separation technology (extractions and materials processing) originates from their unique solvent properties [209, 211–217]. A fluid is termed supercritical when its temperature exceeds the so-called critical temperature, T_C. Under these experimental conditions, the two fluid phases, liquid and vapour, become indistinguishable. They exist as a single phase without any phase boundaries, possessing the favourable properties of both a liquid and a gas. The observation that above a certain temperature pure substances do not condense or evaporate, but exist only as fluids, was made as long ago as 1822 by Baron Charles Cagniard de la Tour [218]. The ability of sc-fluids to dissolve other compounds was first noted in 1879, when Hannay and Hogarth investigated the solubility of iron and cobalt chlorides in sc-ethanol [219].

Most notably, sc-fluids exhibit properties intermediate between those of gases and liquids. Their densities can be finely tuned by varying the pressure and temperature; see Fig. 3-2 [220]. The viscosities and diffusivities of sc-fluids are comparable with those of gases, yet their densities are closer to those of liquids. The low viscosity facilitates the high mass-transfer rates in sc-fluids. The density of sc-fluids is sufficiently high to allow the adequate solvation of a variety of solutes, but their solvating power is much less than that of conventional solvents. The possibility of varying the density of sc-fluids, and therefore the solubility of the solute, by adjusting the temperature and pressure, makes them ideal candidates as solvents for extraction and partitioning processes (*cf.* Section A.6 in the Appendix). The addition of normal co-solvents (modifiers, entrainers) can enhance the solubility of involatile substances in sc-fluids.

Although the unique properties of sc-fluids were known for a long time, their use as reaction media has only become more popular in recent years [211–217, 220]. Reasons for carrying out reactions in sc-fluids are [216]: (a) enhanced reaction rates for reactions with negative activation volumes; (b) homogenization of reactions that would otherwise be carried out in more than one phase (heterogeneous reactions); (c) enhanced mass transfer, important for diffusion-limited reactions; (d) increased catalyst activity in heterogeneous reactions through in situ regeneration of surfaces with tuning of the operation conditions; (e) tunable reaction rates through changing the polarities of sc-fluids by manipulating their densities; and (f) low toxicity of most sc-fluids, which makes



Fig. 3-2. P/V/T phase diagram of a pure substance (pure solvent) showing domains in which it exists as solid, liquid, gas (vapour), and/or sc-fluid (CP = critical point; TP = triple point; ρ = mass density). The inserted isotherms T_2 ($T_2 > T_C$) and T_3 ($T_3 \gg T_C$) illustrate the pressure-dependent density ρ of sc-fluids, which can be adjusted from that of a gas to that of a liquid. The influence of pressure on density is greatest near the critical point, as shown by the greater slope of isotherm T_2 compared to that of T_3, which is further away from T_C. Isotherm T_1 demonstrates the discontinuity in the density at subcritical conditions due to the phase change. This figure is taken from reference [220].

them ideal candidates for the replacement of environmentally problematic solvents (see Table A-14 in the Appendix). A selection of common sc-fluids is given in Table 3-4.

The most common sc-fluid for industrial processing and benchtop research is supercritical carbon dioxide, chosen because of its moderate and easily attained critical temperature and pressure and its non-toxicity. Reactions in sc-CO_2 produce similar results as reactions in nonpolar organic solvents. Its solvent polarity, empirically determined using solvatochromic dyes as polarity indicators (see Section 7.4), corresponds to that of hydrocarbons such as cyclohexane [221, 222]. Carbon dioxide has no dipole moment and only a small quadrupole moment, a small polarizability volume, and a low relative permittivity (ε_r = 1.4–1.6 at 40 °C and 108–300 bar) [221, 223]. Thus, sc-CO_2 is only suitable as a solvent for nonpolar substances, which unfortunately imposes considerable limitations on its practical applications. To overcome this limitation, more polar co-solvents (modifiers) such as methanol can be added to sc-CO_2.

Another environmentally benign solvent is supercritical water, which has a great potential to replace conventional solvents, although it requires rather harsh experimental conditions (t_C = 374 °C and p_C = 218 atm; see Table 3-4) [224, 225]. At the critical point of sc-H_2O, its volume is three times larger than that at ambient temperature, its relative permittivity is only 5.3 (compared to ε_r = 78.4 at 25 °C), and its intermolecular hydrogen-bond network is partially broken. Under supercritical conditions, water consists of small clusters, oligomers, and even monomeric gas-like water molecules [226,

Table 3-4. Critical temperature, t_C, critical pressure, p_C, critical density, ρ_C, and critical molar volume, V_C, for twenty supercritical fluids, arranged in order of increasing t_C.[a)]

Solvents	t_C/°C	p_C/MPa[b)]	ρ_C/g · cm^{-3}	V_C/cm^3 · mol^{-1}
Methane	−82.6	4.60	0.16	99
Ethene	9.2	5.04	0.22	131
Xenon	16.6	5.84	1.15	118
Trifluoromethane	26.2	4.86	0.62	133
Carbon dioxide	31.0	7.38	0.47	94
Ethane	32.2	4.87	0.20	146
Tetrafluoroethene	33.4	3.94	–	172
Dinitrogen oxide	36.4	7.26	0.45	97
Fluoromethane	44.7	5.88	–	113
Sulfur hexafluoride	45.5	3.77	0.74	199
Difluoromethane	78.5	5.83	–	121
Propane	96.7	4.25	0.22	200
1,1,1,2-Tetrafluoroethane	101.0	4.07	–	198
Dichloro-difluoromethane	111.8	4.14	0.56	217
Perfluorobutane	113.3	2.32	–	378
Ammonia	132.4	11.4	0.24	72
n-Butane	152.0	3.80	0.23	255
Methanol	239.4	8.08	0.27	117
Ethanol	240.9	6.14	0.28	168
Water	374.0	22.1	0.32	56

[a)] Values taken from (a) D. Ambrose: *Critical Constants*, in D. R. Lide (Ed.): *CRC Handbook of Chemistry and Physics*, 77th ed., CRC Press, Boca Raton/FL, USA, 1996, p. **6**-54ff.; (b) Y. Marcus: *The Properties of Solvents*. Wiley, Chichester, 1998, p. 78.
[b)] 1 MPa = 10^6 Pa = 1 J · cm^{-3} = 9.87 atm.

227]. Because oxygen, carbon dioxide, methane and other alkanes are completely miscible with dense sc-H_2O, combustion can occur in this fluid phase; even flames can burn in sc-H_2O!

In addition to supercritical water, *superheated water* can be used as a reaction medium. The reactivity of various organics in high-temperature water under autogenic pressure has been studied between 200 and 460 °C [228–230]. As the critical temperature of water is 374 °C, these experiments are both sub- and supercritical. In this particular chemistry, water participates as solvent, reactant, and catalyst. For example, ethers and esters undergo facile hydrolysis in neutral superheated water at 250–350 °C. The physical and chemical properties of water change dramatically as the temperature increases to such a large extent. For example, as the temperature rises from 25 to 300 °C, the density of water decreases from $\rho = 0.997$ to 0.713 g/cm^3, its relative permittivity decreases from $\varepsilon_r = 78.4$ to 19.7, and its solubility parameter decreases from $\delta = 47.9$ to 29.7 MPa$^{1/2}$ (see Table 3-3). Over the same temperature range, the ionic product of water increases by three orders of magnitude from 10^{-14} to $10^{-11.3}$. This means that water becomes both a stronger acid and a stronger base as the temperature increases. The change in the relative permittivity makes the solvent polarity properties of water at 300 °C roughly equivalent to those of acetone at 25 °C ($\varepsilon_r = 20.6$). Therefore, under superheated conditions, water can act as a powerful acid/base bicatalyst and ionic reactions

are preferred over free-radical reactions. The ability of superheated water to effect condensation, cleavage, and hydrolysis reactions, as well as selective ionic reactions, is mainly due to the changes in its physical and chemical properties, which approach those of polar organic solvents at room temperature [228–230].

For the use of sc-fluids as reaction media in synthesis, see reviews [211–217, 224, 225, 228–230] and Section 5.5.13, which describes applications of some neoteric solvents such as ionic liquids, perfluorohydrocarbons, and sc-fluids in organic synthesis and catalysis.

Metastable liquids constitute another remarkable manifestation of the liquid state. Liquids can also exist in conditions under which the *stable* state is either a solid, a vapour, or a liquid of different composition. When this occurs, a liquid is said to be *metastable* with respect to the stable phase. For example, metastability with respect to crystalline or vapour phases defines *supercooled* and *superheated* liquids, respectively [210]. Water can exist as a liquid below 0 °C (=supercooled water). It is also possible to heat liquid water above 100 °C without boiling (=superheated water). However, care must be taken to prevent supercooled water from freezing and superheated water from boiling. Supercooled and superheated water exist in a kind of precarious equilibrium. The temperature range over which supercooled and superheated water can exist at atmospheric pressure extends from ca. −41 °C to ca. 280 °C [210, 231]. This possibility of superheating a liquid above its boiling point, or of supercooling it below its freezing point, is not limited to water. Supercooled and superheated liquids are said to be metastable. They are of no importance as reaction media, but supercooled water is found abundantly in Nature [210, 231]. Organisms and plants exposed to prolonged sub-0 °C conditions have developed strategies to prevent the potential freezing of their water content, as formation of ice crystals inside their cells would be lethal. To avoid its freezing in animals, there exist two possibilities: colligative depression of freezing point by so-called cryoprotectants (*e.g.* glycerol) or through the action of antifreeze proteins. Antarctic fish are protected down to −2 °C by antifreeze proteins; for a more detailed discussion and other examples of metastable liquids, see references [210, 231].

3.3 Classification of Solvents in Terms of Acid-Base Behaviour

3.3.1 Brønsted-Lowry Theory of Acids and Bases [43–50, 107, 168]

According to the Brønsted-Lowry definition, acids and bases are proton donators and -acceptors, respectively, as expressed in the following equilibrium

$$\underset{\text{Acid}}{HA^{z+1}} \rightleftharpoons \underset{\substack{\text{Conjugate}\\ \text{Base}}}{A^{z}} + H^{+} \tag{3-7}$$

where $z = 0, \pm 1, \ldots$ [51, 52]. Since, in solution, the isolated proton cannot exist [129], an acid-base reaction will take place only in the presence of a base possessing a higher proton affinity than the conjugate base A^z. As most solvents possess acid or base properties, the strengths of acids and bases depend on the medium in which they are dissolved.

The equilibrium shown in Eq. (3-8) will be established when an acid HA is dissolved in a basic solvent SH.

$$\underset{\text{Acid}}{HA^{z+1}} + \underset{\text{Solvent}}{SH} \rightleftharpoons \underset{\substack{\text{Lyonium}\\ \text{Ion}}}{SH_2^+} + A^z \tag{3-8}$$

The strength of the acid HA in the solvent SH is given by the acidity contant K_a according to Eq. (3-9)*) [53].

$$K_a = \frac{[SH_2^+] \cdot [A^z]}{[HA^{z+1}]} \tag{3-9}$$

In an acidic solvent SH, the acid-base equilibrium shown in Eq. (3-10) will be established.

$$\underset{\text{Base}}{A^z} + \underset{\text{Solvent}}{SH} \rightleftharpoons HA^{z+1} + \underset{\substack{\text{Lyate}\\ \text{Ion}}}{S^-} \tag{3-10}$$

Increasing basicity or acidity of the solvent displaces the equilibria (3-8) and (3-10) to the right. The addition of these two equations gives a new equilibrium describing the self-ionization (autoprotolysis) of the solvent.

$$\underset{\text{Solvent}}{2\,SH} \rightleftharpoons \underset{\substack{\text{Lyonium}\\ \text{Ion}}}{SH_2^+} + \underset{\substack{\text{Lyate}\\ \text{Ion}}}{S^-} \tag{3-11}$$

Eq. (3-11) reflects both the acidic and basic properties of a solvent, which are described quantitatively by the *autoprotolysis constant* K_{auto}.

$$K_{auto} = [SH_2^+] \cdot [S^-] \tag{3-12}**)$$

Autoprotolysis constants for some representative solvents are collected in Table A-12 (Appendix). The useful pH range of a solvent increases as the autoprotolysis constant decreases. The smaller the autoprotolysis constant, the greater the range of acid or base strengths which can exist in a solvent***).

Self-ionizing solvents possessing both acid and base characteristics (*e.g.* water) are designated *amphiprotic solvents*, in contrast to *aprotic solvents*, which do not self-

---

* Since, in general, the solvent SH is in large excess, its concentration [SH] remains almost constant and is therefore included in the constant K_a.

** Since the solvent concentration [SH] remains practically constant due to the large excess of unionized solvent, this term is usually included in K_{auto}.

*** Thus, the magnitude of K_{auto} is an important criterion for the proper choice of solvent for titrations in non-aqueous solvents; *cf.* Section A-8 (Appendix).

Table 3-5. Classification of organic solvents according to their Brønsted acid-base behaviour [56].

Solvent designation		Relative acidity[a)]	Relative basicity[a)]	Examples
	Neutral	+	+	H_2O, CH_3OH, $(CH_3)_3COH$, $HOCH_2CH_2OH$, C_6H_5OH
Amphiprotic	Protogenic	+	−	H_2SO_4, HCOOH, CH_3COOH
	Protophilic	−	+	NH_3, $HCONH_2$, $CH_3CONHCH_3$, $H_2N—CH_2CH_2—NH_2$
	Dipolar Protophilic	−	+	$HCON(CH_3)_2$, CH_3SOCH_3, pyridine, 1,4-dioxane $(C_2H_5)_2O$, tetrahydrofuran
Aprotic	Dipolar Protophobic	−	−	CH_3CN, CH_3COCH_3, CH_3NO_2, $C_6H_5NO_2$, sulfolane
	Inert	−	−	Aliphatic hydrocarbons, C_6H_6, $Cl—CH_2CH_2—Cl$, CCl_4

[a)] − indicates weaker and + indicates stronger acid or base than water.

ionize to a measurable extent (*e.g.* aliphatic hydrocarbons, tetrachloromethane) [47, 53–55].

It is not possible to draw a sharp line between amphiprotic and aprotic solvents since, in practice, amphiprotic solvents with extremely small K_{auto} values behave like aprotic solvents. It has been suggested that solvents with K_{auto} values greater than 20 should be called aprotic rather than amphiprotic [44].

This classification of solvents was first proposed by Brønsted, who distinguished between four types of solvents on the basis of their acid and base properties [54]. Davies extended the Brønsted classification and distinguished between solvents with relative permittivities greater or smaller than 20, thus arriving at eight classes of solvents [47]. Kolthoff's classification in a slightly simplified fashion is given in Table 3-5 [56].

Water is the prototype of an amphiprotic solvent and all other solvents with similar acid-base properties are called *neutral solvents*. Solvents that are much stronger acids and much weaker bases than water are called *protogenic solvents*, while those that are much stronger bases and much weaker acids than water are designated *protophilic solvents*. This division is somewhat arbitrary since, by agreement, water is the reference taken to define neutral.

From Eq. (3-8), it is seen that the ionization of an acid depends on the basicity of the solvent. In other words, the effective strength of an acid is greater, the higher the proton affinity of the medium. However, the ionization of the acid depends not only on the basicity of the solvent, but also on its relative permittivity and its ion-solvating ability. The dependence of the acidity and basicity constants of a compound on the basicity and acidity, respectively, of the solvents, leads to a distinction between *levelling* and *differentiating solvents* [49, 57, 58].

All mineral acids ionize to the same extent in aqueous solution: they are essentially completely ionized due to almost quantitative reaction with the base water. It would be a strange coincidence, however, if all these acids had exactly the same acid



Fig. 3-3. Schematic description of the levelling effect of water on (a) acids and (b) bases in aqueous solution [2]. Relative orders of acidity and basicity are not invariable to changes of solvent and of the conjugated acid or base, respectively.

strength in spite of their different constitutions. The explanation is that water exerts a levelling effect on the acid strengths. Fig. 3-3 shows that in water the strengths of all acids stronger than H_3O^+ are adjusted to that of H_3O^+ itself. Solvents exhibiting such behaviour are called *levelling solvents**). In order to establish the relative strengths of the mineral acids, it is necessary to perform the measurements in solvents of very low basicity and ionizing ability. Logically enough, such solvents are designated *differentiating solvents*. Fig. 3-3 makes it clear that the strongest acid that can exist in a solvent is the lyonium ion $SH_2^{\oplus}$ (in case of (a) $H_3O^{\oplus}$), while the lyate ion, $S^{\ominus}$, is the strongest base that can exist in a solvent (in case of (b) $HO^{\ominus}$). Thus, in methanol, hydrochloric acid is completely ionized, whereas nitric acid is only partly so. In the less basic formic acid, hydrochloric acid is also only partially ionized, whereas the first H-atom of sulfuric acid is still completely ionized. Acetonitrile is a very weak base and an exceptionally weak acid, thus, only slight levelling of acids and bases occurs in this solvent, making it a good differentiating solvent. Perchloric acid appears to be strong in acetonitrile, whereas

* The expression "levelling solvents" ("chemisch nivellierende Lösungsmittel") was introduced by A. Hantzsch, Z. Elektrochem. *29*, 221 (1923).

other acids are differentiated [59]: $HBr > H_2SO_4 > HNO_3 > HCl$ and picric acid. When the solvent is a stronger base than water, its levelling effect will apply also to weaker acids. Thus, in liquid ammonia, even the carboxylic acids are practically fully ionized.

Similarly, strong bases will also have equal basicities in sufficiently acidic solvents. As shown in Fig. (3-3b), all bases stronger than the $HO^{\ominus}$ ion are adjusted to the basicity of this ion in water. Consequently, the effective basicities of guanidines and carbanions cannot be measured in water, but only in less acidic differentiating solvents such as liquid ammonia or diethyl ether.

Evidently, in a given solvent the more highly ionized acid or base is also the strongest one. However, if the same acid is examined in different solvents, one finds, surprisingly enough, that the most acidic solution is the one in which the acid is the least ionized. For example, a solution of hydrochloric acid in benzene is a stronger acid than in aqueous solution; in the former, ionization is slight, in the latter, complete. In the aqueous solution, an indicator would have to compete with the base H_2O for the protons; however, in the benzene solution, competition involves only the much weaker base $Cl^{\ominus}$. Therefore, a larger proportion of the indicator would be transformed into its conjugate acid in benzene than in water, thus making the benzene solution of HCl the better proton donator.

Numerous acidity and basicity scales have been elaborated for water and other solvents. However, there is no one single scale of acidity and basicity, equally valid and useful for all types of solvents and applicable to both equilibrium and kinetic situations. Excellent reviews on different acidity functions are given by Boyd [60] and Bates [50].

In dilute aqueous solution, the acidity is measured using pH values. For concentrated acid solutions and non-aqueous acid solutions, pH values are no longer available. Hence, the *Hammett acidity function* H_0 is used as a measure of the acidity of such media [130]. The proton donor ability of an acid in such media is measured by studying the equilibria of a series of indicator bases B (mostly nitroanilines), the UV/Vis absorption spectra of which are markedly different from those of their conjugate acids, so that the indicator concentration ratio $I = [B]/[BH^+]$ can be measured spectrophotometrically. The acidity function H_0 is then given by $H_0 = pK_{BH+} + \lg I$, with the subscript zero indicating that the H_0 function applies only to neutral bases B [130, 170]. For dilute solutions, H_0 corresponds exactly to pH; in concentrated solutions, the two functions differ appreciably.

Brønsted acids stronger than pure (100%) sulfuric acid ($H_0 = -11.9$) are classified as *super acids* [131, 132]. Thus, perchloric acid ($HClO_4$), fluorosulfonic acid ($F—SO_3H$), and trifluoromethanesulfonic acid ($CF_3—SO_3H$) are considered as super acids. Even exceedingly weak basic solvents (*e.g.* carbonyl compounds; aromatic, olefinic, and saturated hydrocarbons) are protonated by these super acids to give the corresponding carbocations [131].

Streitwieser *et al.* [160] and Bordwell *et al.* [161] used the lyate ions of organic solvents such as cyclohexylamine and dimethyl sulfoxide in the determination of the C—H acidity of weak organic carbon acids. Using *super base* systems such as alkali metal salts of cyclohexylamine (*i.e.* lithium and cesium cyclohexylamides) [160] and dimethyl sulfoxide (sodium dimsyl) [161] in an excess of these non-HBD solvents, relative acidity scales for weak carbon acids have been established. In this way, pK_a^{DMSO} values for the ionization of over a thousand Brønsted acids in dimethyl sulfoxide have

become available, covering a range of *ca.* 35 pK_a units [161]. The upper limits of measurements in cyclohexylamine and dimethyl sulfoxide are imposed by the acidities of the two solvents, corresponding to maximum determinable pK_a^{CHA} and pK_a^{DMSO} values of *ca.* 39 and 32, respectively.

The acidity and basicity of solvents can be measured in different ways [49]. Besides the usual experimental methods of measuring acid-base equilibrium constants, another possible approach is the determination of solvent basicities and acidities by measuring the change in some physical property (like an IR or UV/Vis absorption or NMR chemical shift) of the molecules of a standard substrate when transferred from a reference solvent to another solvent. For example, the shift in wavenumber of the ≡C—H valence vibration band of phenylacetylene when transferred from tetrachloromethane to nineteen other solvents has been measured, giving a relative order of basicity ranging from tetrachloromethane (low) to hexamethylphosphoric triamide (high) [61]. The basicity of solvents has also been measured using the ^{1}H NMR chemical shift of the chloroform H-atom $\Delta\delta_\infty(CHCl_3)$ obtained by extrapolation to infinite $CHCl_3$ dilution in the solvent in question and in an inert reference solvent (cyclohexane), respectively. The results, summarized in Table 3-6, establish an order of solvent basicity using chloroform as the standard proton donor [62].

For other, more recent quantitative scales of the hydrogen-bond donor acidities and hydrogen-bond acceptor basicities of solvents, see the end of Section 2.2.5 (and references [329–334] cited therein).

The protonation equilibria of a variety of solvents in strong acids such as aqueous sulfuric acid at 25 °C have been measured ^{1}H NMR spectroscopically [232]. For exam-

Table 3-6. Solvent order of increasing basicity relative to chloroform, measured using the relative ^{1}H NMR chemical shift and extrapolated to infinite dilution with cyclohexane as reference solvent [62].

Solvents	$\Delta\delta_\infty(CHCl_3)$[a)] ppm
Cyclohexane	0.00
Tetrachloromethane	0.18
Chloroform	0.20
Nitromethane	0.47
Acetonitrile	0.56
1,4-Dioxane	0.63
Diethyl ether	0.74
Tetrahydrofuran	0.79
Acetone	0.92
Cyclohexanone	0.97
Triethylamine	1.22
N,*N*-Dimethylformamide	1.30
Dimethyl sulfoxide	1.32
Pyridine	1.56
Hexamethylphosphoric acid triamide	2.06

[a)] $\Delta\delta_\infty(CHCl_3) = \delta_\infty(CHCl_3, \text{solvent}) - \delta_\infty(CHCl_3, \text{cyclohexane})$

ple, the relative order of increasing basicity for some dipolar non-hydroxylic solvents determined this way is: nitromethane < sulfones < acetonitrile ≪ acetone < dimethyl sulfoxide < amides [232].

Recent gas-phase studies of proton-transfer reactions with stepwise solvation of the reactants (*i.e.* incremental addition of solvent molecules to form supermolecular clusters) have demonstrated that the acid/base behaviour of isolated solvent molecules can be dramatically different from their performance as bulk liquids. Water, the classical amphiprotic solvent, shall serve as an example.

In contrast to the acid/base behaviour of "polymeric" bulk water, "monomeric" water is a relatively weak acid and base in the gas phase compared to its substituted derivatives (R—OH, R—O—R, *etc.*), whose conjugated base or acid ions are stabilized by polarization of the alkyl groups. The gas-phase basicity of water is 138 kJ/mol (33 kcal/mol) below that of ammonia. Its gas-phase acidity is comparable to that of propene and it is less acidic than phenol by about 167 kJ/mol (40 kcal/mol). With respect to the well-known acid/base properties of water, ammonia, and phenol in aqueous solution, one has to conclude that enormous solvation energies must contribute to the difference from the behaviour of isolated water molecules. See Section 4.2.2 for further discussions and references.

3.3.2 Lewis Theory of Acids and Bases [44–65, 65a]

According to Lewis, acids are electron-pair acceptors (EPA) and bases electron-pair donors (EPD), connected through the following equilibrium [63, 65a]:

A	+	:D	⇌	A—D	(3-13)
Acid		Base		Acid-Base	
EPA		EPD		Complex	
Electrophile		Nucleophile			
Hard	———	Hard			
	╲ ╱		[——— favorable combination		
	╳		----- unfavorable combination]		
Soft	———	Soft			

The Lewis acid/base complex is formed through an overlap between a doubly occupied orbital of the donor D and a vacant orbital of the acceptor A (*cf.* also Section 2.2.6). This acid/base approach was extended by Pearson, who divided Lewis acids and bases into two groups, hard and soft, according to their electronegativity and polarizability (principle of hard and soft acids and bases; HSAB concept) [66, 67]. Hard acids (*e.g.* $H^{\oplus}$, $Li^{\oplus}$, $Na^{\oplus}$, BF_3, $AlCl_3$, hydrogen-bond donors HX) and hard bases (*e.g.* $F^{\ominus}$, $Cl^{\ominus}$, $HO^{\ominus}$, $RO^{\ominus}$, H_2O, ROH, R_2O, NH_3) are those derived from small atoms with high electronegativities and are generally of low polarizability. Soft acids (*e.g.* $Ag^{\oplus}$, $Hg^{\oplus}$, I_2, 1,3,5-trinitrobenzene, tetracyanoethene) and soft bases (*e.g.* $H^{\ominus}$, $I^{\ominus}$, $R^{\ominus}$, $RS^{\ominus}$, RSH, R_2S, alkenes, C_6H_6) are usually derived from large atoms with low electronegativities and are usually polarizable. The usefulness of this division arises from a simple rule concerning the stability of Lewis acid/base complexes: *hard acids prefer to coordi-*

nate to hard bases and soft acids to soft bases [66, 67]. This HSAB concept describes a wide range of chemical phenomena in a qualitative way and has found many applications in organic chemistry [66–70] (for a criticism of the HSAB concept, see refs. [71, 72]). Since the hard species are generally of small size and high charge density, and the soft species are large in size with a low charge density, *ionic bonding* predominates in hard/hard reactions, whereas *covalent bonding* between the reactants, involving their frontier orbitals (*i.e.* HOMO/LUMO interactions according to Fukui), predominates in soft/soft reactions [233, 234]. Based on ionization potentials and electron affinities, even numerical values of absolute hardness, *i.e.* resistance to deformation or change in the electronic charge cloud, can now be assigned to various Lewis acids and bases.

Solvents can be classified as EPD or EPA according to their chemical constitution and reaction partners [65]. However, not all solvents come under this classification since *e.g.* aliphatic hydrocarbons possess neither EPD nor EPA properties. An EPD solvent preferably solvates electron-pair acceptor molecules or ions. The reverse is true for EPA solvents. In this respect, most solute/solvent interactions can be classified as generalized Lewis acid/base reactions. A dipolar solvent molecule will always have an electron-rich or basic site, and an electron-poor or acidic site. Gutmann introduced so-called donor numbers, *DN*, and acceptor numbers, *AN*, as quantitative measures of the donor and acceptor strengths [65]; *cf.* Section 2.2.6 and Tables 2-3 and 2-4. Due to their coordinating ability, electron-pair donor and acceptor solvents are, in general, good ionizers; *cf.* Section 2.6.

The HSAB behaviour is dependent on the medium in which EPD/EPA reactions are carried out. For example, the order of stability of complexes of metal ions with halide ions in the gas phase is $F^- > Cl^- > Br^- > I^-$, which makes all metal ions appear hard in the gas phase. However, in aqueous solution, the stability order is reversed to $F^- < Cl^- < Br^- < I^-$ for those metal ions classified as soft [69].

The application of the HSAB concept to solutions leads to the rule that hard solutes dissolve in hard solvents and soft solutes dissolve in soft solvents [66]. This rule can be considered as a modern version of "similia similibus solvuntur". For example, benzene is considered a very soft solvent since it contains only a basic function. Contrary to benzene, water is a very hard solvent, with respect to both its basic and acidic properties. It is the ideal solvent for hard bases and hard acids. The hardness of water is reduced by the introduction of alkyl substituents in proportion to the size of the alkyl group. In alcohols, therefore, softer solutes become soluble. Whereas oxalate salts are quite insoluble in methanol, the corresponding softer bisthiooxalate salts are quite soluble.

Since hydrogen-bonding is a hard acid-hard base interaction, small basic anions prefer specific solvation by protic solvents. Hence, the reactivity of $F^{\ominus}$, $HO^{\ominus}$, or $CH_3O^{\ominus}$ is reduced most on going from a dipolar non-HBD solvent such as dimethyl sulfoxide to a protic solvent like methanol. Dipolar non-HBD solvents are considered as fairly soft compared to water and alcohols [66].

Similar considerations can be made in the case of cation solvation. Small, hard cations of high oxidation state will be preferably solvated by hard EPD solvents like H_2O or ROH. In principle, relative to the gas state, all ions become softer in the solute state as a result of solvation [66].

Replacement of the oxygen in hard oxygen-donor solvents (such as THF, DMF,

and 1-methylpyrrolidine-2-one) by sulfur to give the corresponding sulfur-donor solvents (*i.e.* thiolane [235], *N*,*N*-dimethylthioformamide [236], and 1-methylpyrrolidine-2-thione [237]) leads to much softer EPD solvents, more suitable for the solvation of soft cations (*e.g.* Ag^+, Tl^+, Hg^{2+}, *etc.*).

Based on single-ion transfer properties, various quantitative empirical scales describing the softness of EPD solvents have been proposed; for reviews, see references [173, 238, 239].

Marcus proposed a μ scale of solvent softness (from the Greek *malakos* = soft), defined as the difference between the mean of the standard molar Gibbs energies of transfer of Na^+ and K^+ ions from water (W) to a given solvent (S), $\Delta G^\circ_t(Me^+, W \rightarrow S)/(kJ \cdot mol^{-1})$, and the corresponding Gibbs transfer energy for Ag^+ ions, divided by 100 [171, 239]. Since water is a hard solvent, the Gibbs energy of transfer of ions from water as a reference solvent to other solvents, should depend on the softness of these EPD solvents to a different extent for hard and soft ions. Provided that the charge and size of the solvated ions are equal, hard ions prefer water and soft ions the softer solvents. The definition of μ was selected because the size of the soft Ag^+ ion ($r = 115$ pm) is intermediate between those of the hard Na^+ ($r = 102$ pm) and K^+ ions ($r = 138$ pm). The degree of softness among solvents with oxygen, nitrogen, and sulfur donor atoms increases in the sequence *O*-donor (alcohols, ketones, amides) < *N*-donor (nitriles, pyridines, amines) < *S*-donor solvents (thioethers, thioamides) [171].

The soft donor properties of EPD solvents have also been quantified by the softness parameter *SP* of Gritzner [173, 240]. This parameter is solely based on the standard molar Gibbs energies of transfer of Ag^+ ions from benzonitrile as reference solvent to other soft solvents and should be used for soft/soft interactions only.

Finally, the D_S and D_H solvent scale of Persson *et al.* should be mentioned [241]. The empirical D_S scale, ranking the EPD solvent's donor strength towards a soft acceptor, is based on the shift of the Raman IR absorption for the symmetrical stretching vibration of the Hg—Br bond of soft mercury(II) bromide in the solvent of interest relative to that in the gas phase: $D_S = \Delta\tilde{\nu}(Hg—Br)/cm^{-1} = [\tilde{\nu}(Hg—Br)_{gas} - \tilde{\nu}(Hg—Br)_{solvent}]/cm^{-1}$. D_S values are known for more than sixty solvents with widely varying solvating properties [241]. An additional solvent donor strength scale for hard acceptors, D_H, has been derived for twenty-four solvents from published molar Gibbs' energies for transfer of Na^+ from a given solvent to 1,2-dichloroethane as a reference solvent [241].

A detailed discussion and comparison of all these and further solvent softness scales can be found in references [173, 238, 239]. For other Lewis acid/base parameters of EPD and EPA solvents, derived from calorimetric measurements (*e.g.* Gutmann's donor and acceptor numbers), see reference [65] and Section 2.2.6.

The concept of the superacidity of Brønsted acids has been extended to Lewis acids [131, 132]. It is suggested that those Lewis acids stronger than anhydrous aluminium trichloride (the most commonly used Friedel-Crafts catalyst) should be designated as *super acids*. These superacidic Lewis acids include such higher-valence halides as antimony, arsenic, tantalum, and niobium pentafluorides. Frequently used conjugated Brønsted-Lewis super acids are $F—SO_3H/SbF_5$ "magic acid" and HF/SbF_5 (fluoroantimonic acid) [131]. These superacidic systems are considered to be *ca.* 10^{16} times stronger than 100 percent sulfuric acid [131].

3.4 Classification of Solvents in Terms of Specific Solute/Solvent Interactions

Parker divided solvents into two groups according to their specific interactions with anions and cations, namely *dipolar aprotic solvents* and *protic solvents* [73]. The distinction lies principally in the dipolarity of the solvent molecules and their ability to form hydrogen bonds. The origin of this solvent classification was the experimental finding that certain S_N2 reactions at saturated carbon atoms involving anions as nucleophiles are much faster in the so-called dipolar aprotic solvents than in protic solvents. This is because in dipolar aprotic solvents most anions are much less solvated and hence more reactive than in protic solvents [74]. It appears appropriate to add to these two groups a third one, namely, the *apolar aprotic solvents*, according to Fig. 3-4.

An apolar aprotic solvent is characterized by a low relative permittivity ($\varepsilon_r < 15$), a low dipole moment ($\mu < 8.3 \cdot 10^{-30}$ Cm = 2.5 D), a low E_T^N value (E_T^N *ca.* 0.0 . . . 0.3); *cf.* Table A-1, Appendix), and the inability to act as a hydrogen-bond donor. Such solvents interact only slightly with the solute since only the non-specific directional, induction, and dispersion forces can operate. To this group belong aliphatic and aromatic hydrocarbons, their halogen derivatives, tertiary amines, and carbon disulfide.

In contrast, dipolar aprotic solvents*) possess large relative permittivities ($\varepsilon_r > 15$), sizeable dipole moments ($\mu > 8.3 \cdot 10^{-30}$ Cm = 2.5 D), and average E_T^N values of 0.3 to 0.5. These solvents do not act as hydrogen-bond donors since their C—H bonds are not sufficiently polarized. However, they are usually good EPD solvents and hence cation solvators due to the presence of lone electron pairs. Among the most important dipolar aprotic solvents are acetone, acetonitrile [75], benzonitrile, *N*,*N*-dimethylacetamide [76, 77], *N*,*N*-dimethylformamide [76–78], dimethylsulfone [79], dimethyl sulfoxide [80–84], hexamethylphosphoric triamide [85], 1-methylpyrrolidin-2-one [86], nitrobenzene, nitromethane [87], cyclic carbonates such as propylene carbonate (4-methyl-1,3-dioxol-2-one) [88], sulfolane (tetrahydrothiophene-1,1-dioxide) [89, 90, 90a], 1,1,3,3-tetramethylurea [91, 91a]; and tetrasubstituted cyclic ureas such as 3,4,5,6-tetrahydro-1,3-dimethyl-pyrimidin-2-(1*H*)-one (dimethyl propylene urea, DMPU) [133]. The latter is a suitable substitute for the carcinogenic hexamethylphosphoric triamide (*cf.* Table A-14) [134].

Protic solvents contain hydrogen atoms bound to electronegative elements (F—H, —O—H, —N—H, *etc.*) and are, therefore, hydrogen-bond donors *i.e.* HBD solvents (*cf.* Section 2.2.5). With the exception of acetic acid (and its homologues), the relative permittivities are usually larger than 15, and the E_T^N values lie between 0.5 and 1.0, indicating that these solvents are strongly polar. To this class of solvents belong water, ammonia, alcohols, carboxylic acids, and primary amides.

It is emphasized that this classification is not rigid. There are several solvents that cannot be unequivocally assigned to any one of these three groups, *e.g.* ethers, carbox-

* Although widely used, the term *dipolar aprotic* solvent is in fact rather misleading. Solvents referred to as dipolar *aprotic* are in fact *not* aprotic. In reactions where strong bases are employed, their protic character can be recognized. In dimethyl sulfoxide solution, the pK_a values are, for CH_3NO_2 17.2, CH_3CN 31.3, CH_3SOCH_3 35, 1-methyl-2-pyrrolidone *ca.* 35, HMPT *ca.* 45 or above [135]. It has therefore been recommended by Bordwell *et al.* [135] that the designation *dipolar aprotic* for these solvents be replaced by *dipolar nonhydroxylic* or better still by *dipolar non-HBD* solvents. The abbreviations *HBD* (hydrogen-bond donor) and *HBA* (hydrogen-bond acceptor) refer to donation and acceptance of the proton, and not to the electron pair involved in hydrogen bonding (*cf.* Section 2.2.5).



Fig. 3-4. An extension of Parker's classification of organic solvents [73] (concerning the E_T^N values, see Section 7.4).

ylic esters, primary and secondary amines, and *N*-monosubstituted amides such as *N*-methylacetamide [91b]. The choice of $\varepsilon_r = 15$ as the borderline is arbitrary but practical, since, in solvents with smaller ε_r values, ion association occurs, so that the free solvated ions can no longer be observed (*cf.* Section 2.6). This division of solvents into three classes has mainly heuristic value. Its usefulness is due to the fact that special prominence is given to the dipolar aprotic solvents with their extraordinary specific ion solvation [73, 92–96].

Protic solvents are particularly good anion solvators due to their hydrogen-bonding ability [136]. This tendency becomes more pronounced, the higher the charge density (*i.e.* ratio of charge to volume) of the anion to be solvated, and hence the greater its hardness according to the HSAB principle. It should be noted that the stronger the solvation, the more the nucleophilic reactivity of the anion will be decreased. Therefore, in protic solvents, the strongest nucleophiles will be those with lower or more diffused charge density, *i.e.*, "soft anions" (*cf.* Section 5.4.1).

In contrast, in dipolar aprotic solvents, anion solvation occurs mainly by ion-dipole and ion-induced dipole forces. The latter are important for large, polarizable, soft anions, with low charge density, in soft dipolar aprotic solvents. Therefore, although these solvents tend to be poor anion solvators, they are usually better, the larger and softer the anion. This has the consequence that the reactivity of anions is exceptionally high in dipolar aprotic solvents, and the rate constants of S_N2 reactions can increase by several powers of ten when the solvent is changed from protic to dipolar aprotic (*cf.* Section 5.4.2).

The observation that protic solvents are far better anion solvators than dipolar aprotic solvents, and that the reverse is true for cation solvation, has led to extremely valuable rules for the selection of solvents for specific reactions [73, 92–97].

3.5 Classification of Solvents using Multivariate Statistical Methods

Chemical experience suggests that more than three solvent classes, as recognized by Parker [73] (*cf.* Section 3.4 and Fig. 3-4), may be necessary to classify solute/solvent interactions for a wide range of organic solvents. Multivariate statistical methods have therefore been used in the classification and selection of organic solvents [102, 138–143]. Compilations of their physicochemical constants (*e.g.* boiling points, molar volumes, heats of vapourization, dipole moments, relative permittivities, molar refractions, *etc.*) and sometimes, additionally, empirical parameters of solvent polarity (*cf.* Chapter 7), are used as basic data sets. The extraction of chemical information contained in such a data set, *i.e.* the detection of the relative importance of individual variables in determining the data structure, can be done by two statistical methods: multiple linear regression analysis (MRA) [144] and factor (FA) or principal component analysis (PCA) [145]. This kind of analysis is part of the relatively new research field of *chemometrics**) [146, 147].

---

* *Chemometrics* has been defined as the application of mathematical and statistical methods to chemical measurements, in particular in providing maximum chemical information through the analysis of chemical data. Because of the enormous increase in generating analytical data, analytical chemists were among the first to use chemometrical methods extensively [148]. The first paper mentioning the name *chemometrics* was from Wold and was published in 1972: S. Wold, Kem. Tidskr. *84*, 34 (1972).

In MRA, a dependent variable Y is described in terms of a series of explanatory variables $X_1 \ldots X_n$, as given in Eq. (3-14).

$$Y = a_1 \cdot X_1 + a_2 \cdot X_2 + \cdots + a_n \cdot X_n + b \tag{3-14}$$

It is assumed that all the explanatory variables are independent of each other and truly additive as well as relevant to the problem under study [144]. MRA has been widely used to establish linear Gibbs energy (LGE) relationships [144, 149, 150]. The Hammett equation is an example of the simplest form of MRA, namely bivariate statistical analysis. For applications of MRA to solvent effects on chemical reactions, see Chapter 7.7.

The other statistical method used for seeking regularities in physicochemical data, FA, was first developed and used in psychometrics [145]. FA may be described as a mathematical method for seeking the simplest existing linear structure within a given set of multidimensional data. Starting with a matrix of such experimental data (descriptors), it is possible to extract, using sophisticated statistical methods, the minimum number of underlying, non-measurable variables (factors or principal components) necessary to describe this whole data set in multiple regression equations. After the number of factors (or components) have been found and their magnitudes have been calculated for particular solvents, often a physical or chemical meaning emerges. Although these factors are purely mathematical constructs and do not necessarily embody a direct physical significance, the advantage of FA is that an otherwise hidden physical or chemical interpretation will emerge. The two methods, FA and PCA, are coincident when PCA is used after normalizing the data set. The mathematics of fitting FA and PCA to a matrix of chemical data is well described in the literature [145, 151], and the capabilities of PCA in different fields of pattern recognition have been reviewed [152].

A clear geometrical description of solvent classification using FA or PCA can be given as follows. Common descriptors used for the classification of solvents are their physicochemical constants and empirical parameters of solvent polarity (see Chapter 7). Each descriptor defines a coordinate axis in a coordinate system. If m descriptors are used as the basic data set, they will therefore define an m-dimensional space in which each solvent can be described by a point (with coordinates equal to the m descriptors). The whole set of different solvents will then define a swarm of points in the m-dimensional descriptor space. If only three descriptors are involved, say boiling point, dipole moment, and relative permittivity, a simple right-angled three-dimensional coordinate system, as given in Fig. 3-5, results with t_{bp}, μ, and ε_r plotted along the x, y, and z axes, respectively.

FA/PCA now constitutes a projection of this swarm of points down to a space of lower dimensions in such a way that the first component vector (factor F_1) describes the direction through the swarm showing the largest variation in the data. The second component (factor F_2) shows the next largest variation, and so on. The supposition that the components (factors) should be independent of one another means that their vectors must be at right angles to one another, *i.e.* mutually orthogonal. To the extent that the solvent points in Fig. 3-5 fall into the plane defined by F_1 and F_2, the position of an individual solvent now needs only two coordinates, instead of the original three, for its localization. Thus, the intrinsic dimensionality of the three sets of solvent property data is reduced to two.



Fig. 3-5. Geometrical representation of FA. The three-dimensional property space is defined by three solvent descriptors (*e.g.* t_{bp}, μ, and ε_r) and filled with 30 solvent points, some of them already lying in the plane defined by the two factors F_1 and F_2 (according to [139] and [142]).

The coordinates of each solvent point are (i) the factor (or principal component) scores F, and (ii) the factor (or principal component) loadings L. They give the information necessary to reconstitute the original physical properties D of any solvent according to Eq. (3-15).

$$D = F_1 \cdot L_1 + F_2 \cdot L_2 + \cdots + F_n \cdot L_n \tag{3-15}$$

Eigenvectors and eigenvalues are the products of calculation at the beginning. They characterise the property of the square matrix (correlation or covariance) derived from the initial data matrix, and they allow calculation of the factor scores F and factor loadings L, respectively.

The advantage of this empirical model is that the systematic variation in the solvent data can now be described using fewer variables than in the original data set. Eventually, an attempt is made to explain the factors F_1 and F_2, which themselves define a new coordinate system, by considering an underlying physical or chemical meaning (*e.g.* polarity, polarizability, or Lewis acidity/basicity of the solvent molecule), thus leading finally to a new solvent classification.

Martin *et al.* [102] were the first to apply FA to solvent classification. A factor analysis for 18 organic solvents with 18 physicochemical parameters led to a solvent classification similar to Parker's classification [73], which was mainly based on chemical intuition (*cf.* Fig. 3-4).

Using PCA, Cramer [139] found that more than 95% of the variances in six physical properties (activity coefficient, partition coefficient, boiling point, molar refractivity, molar volume, and molar vaporization enthalpy) of 114 pure liquids can be explained in terms of only two parameters, which are characteristic of the solvent molecule. These two factors are correlated with the molecular bulk and cohesiveness of the individual solvent molecules, the interaction of which depends mainly upon nonspecific, weak intermolecular forces. This is closely related to nonspecific, weak solute/solvent interactions. With these factors, experimental values of 18 common physical properties

for 139 additional liquids of diverse structure have been predicted with surprising accuracy [139].

FA of data matrices containing 35 physicochemical constants and empirical parameters of solvent polarity (*cf.* Chapter 7) for 85 solvents has been carried out by Svoboda *et al.* [140]. An orthogonal set of four parameters was extracted from these data, which could be correlated to solvent polarity as expressed by the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, to solvent polarizability as expressed by the refractive index function $(n^2 - 1)/(n^2 + 1)$, as well as to the solvent Lewis acidity and basicity. Thus, four solvent parameters are generally needed for the quantitative empirical description of solvent effects on chemical reactions and light absorptions: two are needed to describe the nonspecific solvation of polar and dispersion character, and two to describe specific solvation of electrophilic and nucleophilic character. For correlations of solvent effects using only one empirical solvent parameter, the best parameters are the $E_T(30)$ values, which are derived from the UV/Vis absorption of a solvatochromic dye (*cf.* Sections 6.2.1 and 7.4).

Elguero *et al.* [141] have reduced Palm's analogous tetraparametric model for the multiple correlation of solvent effects [*cf.* Eq. (7-48) in Section 7.7] to a triparametric one, with two factors explaining 94% of the data variance given in an original set of four descriptors [Y, P, E, and B of Eq. (7-48) in Section 7.7] for 51 solvents.

A data matrix of eight common descriptors of solvent properties for 82 solvents was analysed with PCA by Carlson *et al.* [142]. The eight descriptors were melting point, boiling point, density, relative permittivity, dipole moment, refractive index, $E_T(30)$ [*cf.* Eq. (7-27) in Section 7.4], and lg P (*i.e.* the logarithm of the equilibrium partition coefficient of a solvent between 1-octanol and water at 25 °C [153]). Using a two-component model of the whole data set [one component explained principally by ε_r, μ, and $E_T(30)$, the other strongly correlated to the refractive index], different strategies for a systematic selection of solvents for chemical reactions were proposed. This has been applied to the Willgerodt-Kindler reaction between acetophenone and sulfur in the presence of morpholine, in order to reveal the influence of different solvents on the optimum reaction conditions (other variables: amount of sulfur/ketone, amount of morpholine/ketone, and reaction temperature) [165].

Application of PCA to a set of five thermodynamic and spectroscopic basicity-dependent properties of 22 organic non-HBD solvents (related to hydrogen-bonding, proton-transfer, and interaction with hard and soft Lewis acids) by Maria *et al.* [143] has led to the interesting result that just two factors are sufficient to account for about 95% of the total variance of the solvent data. Physical significance has been given to these two principal factors by correlating them with intrinsic gas-phase affinities of the solvent molecules toward the proton and the potassium cation. A blend of electrostatic and charge-transfer (or electron-delocalization) character can be attributed to the first factor F_1. The second factor F_2 corresponds to an essentially electrostatic character. A third factor F_3, of marginal importance only, arises in part from steric hindrance to acid/base complexation. Thus, the inherent dimensionality of the condensed-phase basicity of organic non-HBD molecules commonly used as solvents is essentially reduced to two [143].

Further useful attempts at the classification of organic solvents by means of chemometrical methods have been made by Zalewski *et al.* [242] and by Abraham *et al.* [243], using various sets of physicochemical solvent data.

The most ambitious approach to a general classification of solvents by PCA has been that of Chastrette *et al.* [138]. His classification is based on the representation of 83 solvents as points in an eight-dimensional property space, using the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, molar refraction $V_m \cdot (n^2 - 1)/(n^2 - 2)$, Hildebrand's δ parameter [*cf.* Eq. (2-1) in Section 2.1], refractive index, boiling point, dipole moment, and the HOMO and LUMO energies as solvent descriptors. Five descriptors are the properties of bulk solvents, whereas the last three (μ, HOMO, and LUMO) are molecular properties. The calculated HOMO and LUMO energies of the solvents are included in the set of basic variables in order to take into account Lewis acid/base interactions between solute and solvent. Because some of the eight descriptors are linearly correlated to each other, the spatial dimensionality needed to describe the solvent classification should be lower than eight.

Indeed, it was possible to reduce the original eight-dimensional space by suppressing five principal components, providing an easily visualised three-dimensional solvent property space, with only an 18% loss of information. This subspace is defined by the principal components F_1 (strongly correlated with the molar refraction, refractive index, and HOMO energy), F_2 (strongly correlated with the Kirkwood function, dipole moment, and boiling point), and F_3 (strongly correlated with the LUMO energy). Therefore, F_1 can be interpreted as an index of the polarizability of the solvent, F_2 represents the polarity of the solvent, and F_3 can be explained by the electron affinity and Lewis acidity of the solvent. The Lewis basicity of the solvent seems to be included in F_1.

The 83 organic solvents have been grouped into *nine classes* from the clustering of their principal component values, using a nonhierarchical multivariate taxonomy to progressively classify solvents by means of the discriminating power of the eight descriptors (*cf.* Fig. 3-6).

Classes (1)...(3) comprise dipolar aprotic*) solvents. The first class (AD) contains the usual protic solvents having a relative low dipolarity ($\mu \leq ca.\ 12 \cdot 10^{-30}$ Cm). More dipolar aprotic solvents ($\mu \geq ca.\ 12 \cdot 10^{-30}$ Cm) are found in the second class (AHD). A third class (AHDP) contains only two members, differing from the second by their high polarizability.

Classes (4)...(6) include apolar aprotic*) solvents. In classes (4) and (5), ARA and ARP, are found aromatic apolar ($\mu \approx 0 \ldots 4 \cdot 10^{-30}$ Cm) and aromatic relatively dipolar solvents ($\mu \approx 4 \ldots 10 \cdot 10^{-30}$ Cm). Class (6), called EPD, consists of solvents that are typical electron pair donors ($\mu \approx 4 \ldots 10 \cdot 10^{-30}$ Cm).

Protic or HBD solvents are found in classes (7) and (8). These two classes of hydrogen-bonding solvents (HB and HBSA) are clearly separated by the degree of their intermolecular association. If a total of ten solvent classes are established, water would then constitute a subclass of class HBSA.

Lastly, class (9), labelled MISC (from miscellaneous), consists of four solvents, which have only their high polarizabilities in common.

---

* Bordwell *et al.* [135] have pointed out that solvents referred to as *dipolar aprotic* are in fact not aprotic. In reactions employing strong bases their protic character can be recognized. Therefore, instead of *dipolar aprotic* the designation *dipolar nonhydroxylic* or better *dipolar non-HBD* solvents is strongly recommended. *Cf.* Section 2.2.5 and 3.4 (footnote). In order to avoid confusion, the nomenclature proposed by Chastrette *et al.* [138] is retained in Fig. 3-6.



Fig. 3-6. Chastrette's classification of organic solvents [138].

It is remarkable that this overall solvent classification, obtained entirely using statistical methods (PCA), correlates strongly with the chemist's intuition! Some of the solvent classes of Parker's scheme (*cf.* Fig. 3-4) are reproduced in Fig. 3-6.

However, certain limitations are obvious. Some solvents, marked by (!) in Fig. 3-6, are not expected to be classed as found. Trifluoroacetic acid is classified as AD; benzyl alcohol, 1-octanol, and tetrachloromethane appear in the class ARP; *n*-hexane and cyclohexane in class EPD; tetrahydrofuran and 1,2-dimethoxyethane in the class HB.

The reasons for some of these irregularities may be associated with the choice of HOMO and LUMO for multifunctional solvents. For heteroatom-containing solvents such as benzyl alcohol, the HOMO of the π-electrons or the high-lying heteroatom lone-pair HOMO are available for solute/solvent interactions. The possible regiospecific interaction of the solvent frontier orbitals with solute molecules could mean that a solvent has two points in the descriptor space and hence two places in the resulting classification scheme, depending on the particular solute/solvent system under consideration. According to the classification in Fig. 3-6, benzyl alcohol is treated as an aromatic solvent, rather than as an alcohol.

This duality in the descriptor choice demonstrates the fallibility of searching for a unique, universally valid classification of all organic solvents. Further work has to be done in order to clarify this point.

Another problem that has been tackled by multivariate statistical methods is the characterization of the solvation capability of organic solvents based on empirical parameters of solvent polarity (see Chapter 7). Since such empirical parameters of solvent polarity are derived from carefully selected, strongly solvent-dependent reference processes, they are molecular-microscopic parameters. The polarity of solvents thus defined cannot be described by macroscopic, bulk solvent characteristics such as relative permittivities, refractive indices, etc., or functions thereof. For the quantitative correlation of solvent-dependent processes with solvent polarities, a large variety of empirical parameters of solvent polarity have been introduced (see Chapter 7). While some solvent polarity parameters are defined to describe an individual, more specific solute/solvent interaction, others do not separate specific solute/solvent interactions and are referred to as general solvent polarity scales. Consequently, single- and multi-parameter correlation equations have been developed for the description of all kinds of solvent effects, and the question arises as to how many empirical parameters are really necessary for the correlation analysis of solvent-dependent processes such as chemical equilibria, reaction rates, or absorption spectra.

By a quantitative structure-property relationship (QSPR) analysis of a total of 45 different empirical solvent scales and 350 solvents, the direct calculation of predicted values of solvent parameters for any scale and for any previously unmeasured solvent was possible using the CODESSA program (*i.e. co*mprehensive *de*scriptors for *s*tructural and *s*tatistical *a*nalysis) developed by Katritzky *et al.* [244]. The QSPR models for each of the solvent scales were constructed using only theoretical descriptors, derived solely from the molecular solvent structure. This QSPR study enabled classification of the various solvent polarity scales and ultimately allowed a unified PCA treatment of these scales. This PCA treatment, carried out with 40 solvent scales as variables (each having 40 data points for 40 solvents as objects), allowed a rational classification and grouping

of both solvents and solvent scales [245]. Surprisingly, it emerged that only three major components, orthogonal (=unrelated) to each other, are sufficient to determine solvent polarity. For 36 of the solvent scales, an average of 88% of the variance is described by the first three principal components, which can be roughly related to the relative permittivities and dipolarities of the solvents, as well as to their electrophilicities (acidities) and nucleophilicities (basicities). The solvents included in this study also showed a clear clustering into five groups: (i) highly polar, strong HBD (*e.g.* formamide), (ii) polar, HBD (water, alcohols), (iii) dipolar non-HBD (DMF, DMSO), (iv) less dipolar, non-HBD (ethers, dichloromethane), and (v) apolar, non-HBD (*n*-hexane, tetrachloromethane) (*cf.* Fig. 3-6!) [245].

Another statistical treatment of a set of 32 solvent parameter scales for 45 solvents using the program SMIRC (*s*election of a set of *m*inimally *i*nterr*e*lated *c*olumns) has been carried out by Palm *et al.* [246], who, incidentally, introduced the first multi(four)-parameter equation for the correlation analysis of solvent effects in 1971 [*cf.* Eq. (7-50) in Chapter 7]. The minimum sufficient set of residual descriptors for the multilinear description of solvent effects consists of nine solvent parameter scales. This set of nine (purified) descriptors has been successfully applied to an extended set of 359 different solvent-dependent processes; for more details, see reference [246].

4 Solvent Effects on the Position of Homogeneous Chemical Equilibria

4.1 General Remarks

An equilibrium is homogeneous when all components are either exclusively in the gas phase or exclusively in solution. For gas-phase equilibria, the ratio of the product concentrations for end and starting materials is constant at a given temperature (law of mass action of Guldberg and Waage, 1867). When the reaction partners are dissolved, the standard molar Gibbs energy of solvation, ΔG°_{solv}, is liberated due to the intermolecular interactions between solvent and solute. In general, this quantity is different for starting and end products. Thus, a displacement of the equilibrium can take place when going from the gas phase to solution [1–7]. An unchanged equilibrium constant can only be expected when ΔG°_{solv} is accidently the same for starting and end products.

The effect of the medium on the position of equilibrium can be considered from two points of view: (a) comparison of the gas-phase and solution equilibrium constants, and (b) comparison of the equilibrium constants for different solvents. Unfortunately, few equilibrium reactions have been studied both in the gas and liquid phases [5, 6]. These are primarily non-ionic reactions where the interaction between reacting molecules and solvent is relatively small (*e.g.* the Diels-Alder dimerization of cyclopentadiene). In this chapter, therefore, equilibria which have been examined in solvents of different polarity will be the main topic considered (except for acid-base reactions described in Section 4.2.2).

Let us consider a simple isomerization reaction A $\rightleftharpoons$ B in the solvents I and II, whose abilities to solvate A and B are different. This corresponds to the Gibbs energy diagram shown in Fig. 4-1.

From Fig. 4-1, Eq. (4-1) can be immediately derived,

$$\Delta G^\circ(\mathrm{II}) + \Delta G^\circ_t(\mathrm{A}, \mathrm{I} \rightarrow \mathrm{II}) = \Delta G^\circ_t(\mathrm{B}, \mathrm{I} \rightarrow \mathrm{II}) + \Delta G^\circ(\mathrm{I}) \tag{4-1}$$

which, on rearrangement, leads to Eq. (4-2) [102]:

$$\begin{aligned} \Delta G^\circ(\mathrm{II}) - \Delta G^\circ(\mathrm{I}) &= \Delta\Delta G^\circ(\mathrm{I} \rightarrow \mathrm{II}) \\ &= \Delta G^\circ_t(\mathrm{B}, \mathrm{I} \rightarrow \mathrm{II}) - \Delta G^\circ_t(\mathrm{A}, \mathrm{I} \rightarrow \mathrm{II}) \end{aligned} \tag{4-2}$$

Since, for equilibria, the logarithm of the equilibrium constant is proportional to the standard molar Gibbs energy change, ΔG°, according to Eq. (4-3),

$$\Delta G^\circ = -R \cdot T \cdot \ln K \tag{4-3}$$

it follows from Eqs. (4-2) and (4-3) that the difference in the molar transfer Gibbs energies of educt A and product B, $\Delta\Delta G^\circ_t(\mathrm{I} \rightarrow \mathrm{II})$, determines the solvent effect on the position of this equilibrium. In the particular case of Fig. 4-1, $\Delta G^\circ_t(\mathrm{B}, \mathrm{I} \rightarrow \mathrm{II}) > \Delta G^\circ_t(\mathrm{A}, \mathrm{I} \rightarrow \mathrm{II})$, so that the equilibrium is displaced towards B when the solvent is changed from I to II.

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8



Fig. 4-1. One-dimensional Gibbs energy diagram for an equilibrium reaction A $\rightleftharpoons$ B in the solvents I and II. Ordinate: standard molar Gibbs energies of the reactants A and B in solvents I and II; Abscissa: not defined. $\Delta G^\circ(\mathrm{I})$ and $\Delta G^\circ(\mathrm{II})$: standard molar Gibbs energies of reaction in solvents I and II, respectively; $\Delta G_t^\circ(\mathrm{A}, \mathrm{I} \rightarrow \mathrm{II})$ and $\Delta G_t^\circ(\mathrm{B}, \mathrm{I} \rightarrow \mathrm{II})$: standard molar Gibbs energies of transfer of the solutes A and B from solvent I to solvent II, respectively [$\Delta G_t^\circ(\mathrm{A}, \mathrm{I} \rightarrow \mathrm{II}) = G^\circ(\mathrm{A} \text{ in } \mathrm{I}) - G^\circ(\mathrm{A} \text{ in } \mathrm{II})$, and $\Delta G_t^\circ(\mathrm{B}, \mathrm{I} \rightarrow \mathrm{II}) = G^\circ(\mathrm{B} \text{ in } \mathrm{I}) - G^\circ(\mathrm{B} \text{ in } \mathrm{II})$], *cf.* Eq. (2-12a) in Section 2.3; $\neq$ = transition state.

The required standard molar Gibbs energies of transfer can be obtained from activity coefficient measurements, using Eq. (4-3a),

$$\Delta G_t^\circ(\mathrm{X}, \mathrm{I} \rightarrow \mathrm{II}) = -R \cdot T \cdot \ln(\gamma_\mathrm{I}/\gamma_\mathrm{II}) \tag{4-3a}$$

in which γ refers to activity coefficients of solute X in solvents I and II. Methods used to obtain these activity coefficients have included vapour pressure, solubility, and distribution coefficient measurements [103]. It should be stressed that solvent-transfer activity coefficients can be directly obtained only for neutral solutes. For ionophores, only the product of the γ values of the cation and anion can be determined. Values for single ions can be obtained indirectly by means of so-called extrathermodynamic assumptions; *cf.* Eq. (2-12b) and Table 2-9 in Section 2.3.

In studying solvent effects on equilibria, it is, in principle, not sufficient to investigate the ΔG° changes alone, because this term is determined by both an enthalpy and an entropy term according to Eq. (4-4).

$$\Delta G^\circ = \Delta H^\circ - T \cdot \Delta S^\circ \tag{4-4}$$

Transfer functions can also be defined for the thermodynamic state functions ΔH° and ΔS° [102]. The ease of calorimetric measurements has made the standard molar transfer enthalpy, $\Delta H_t^\circ(\mathrm{X}, \mathrm{I} \rightarrow \mathrm{II})$, readily available. If both transfer Gibbs energies and

transfer enthalpies are available, it should be possible to achieve a complete dissection of the effect of solvents on the various thermodynamic parameters.

Four types of reaction control can be recognized:

(a) Cooperative effects, with $\Delta\Delta H^\circ$ and $T \cdot \Delta\Delta S^\circ$ having opposite signs. Then these two terms will be additive;
(b) Enthalpy-controlled effects, in which the two terms are opposed, but the enthalpy term is larger;
(c) Entropy-controlled effects, in which the two terms are opposite but the $T \cdot \Delta\Delta S^\circ$ is larger; and
(d) Compensating effects, in which the two terms are opposed but nearly equal.

A change in temperature may interconvert categories (b), (c), and (d).

A quantitative description of the influence of the solvent on the position of chemical equilibria by means of physical or empirical parameters of solvent polarity is only possible in favourable and simple cases due to the complexity of intermolecular solute/solvent interactions. However, much progress has recently been made in theoretical calculations of solvation enthalpies of solutes that can participate as reaction partners in chemical equilibria; see the end of Section 2.3 and references [355–364] to Chapter 2. If the solvation enthalpies of all participants in a chemical equilibrium reaction carried out in solvents of different polarity are known, then the solvent influence on this equilibrium can be quantified. A compilation of about a hundred examples of the application of continuum solvation models to acid/base, tautomeric, conformational, and other equilibria can be found in reference [231].

In the following sections, the influence of solvents on chemical equilibria is treated on a more qualitative basis, using acid/base, tautomeric, and other equilibria as representative examples.

4.2 Solvent Effects on Acid/Base Equilibria [8–13]

4.2.1 Brønsted Acids and Bases in Solution

As already emphasized in Section 3.3.1, the ionization equilibrium of an acid, Eq. (4-5), or of a base, Eq. (4-6), is affected by a solvent change, not only because of the

$$HA^{z+1} + SH \rightleftharpoons SH_2^+ + A^z \tag{4-5}$$

$$A^z + SH \rightleftharpoons HA^{z+1} + S^- \tag{4-6}$$

acidity or basicity of the solvent, but also because of its relative permittivity and the ability of the solvent to solvate the various species of Eqs. (4-5) and (4-6). A change in relative permittivity or in solvating ability can thus influence the acidity of an acid HA or the basicity of a base A. Thus, for example, the acidity constant K_a for carboxylic acids is up to 10^6 times larger in water ($\varepsilon_r = 78.3$) than in absolute ethanol ($\varepsilon_r = 24.6$), although water is only 15 . . . 20 times stronger a base than ethanol.

Eq. (4-5) shows the reaction of an acid with an amphiprotic solvent to form a solvated proton and the conjugate base of the acid, at an infinite distance from each other. Part of the energy required for this reaction comes from the electrostatic interactions between these ions and can be estimated from simple electrostatic theory. The electrostatic work necessary for gradually increasing the charge of a spherical species HA of radius r_{HA} from zero up to its full charge $z_{HA} \cdot e$, immersed in a continuous medium of relative permittivity ε_r, is given by the Born equation (4-7) (ε_0 = permittivity of vacuum; z = number of elementary charges e; N_A = Avogadro constant) [14].

$$\Delta G^{\circ}_{\text{electrostatic}} = \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{N_A \cdot z_{HA}^2 \cdot e^2}{2 \cdot \varepsilon_r \cdot r_{HA}} \tag{4-7}$$

The Born equation gives the electrostatic Gibbs free energy of an ion immersed in a medium of relative permittivity ε_r. However, it does not include the energy expended by solvents in forming the cavities to accommodate the ions. This energy is generally small compared to the large Born energy, which amounts to as much as ca. −1000 kJ/mol for the transfer of monovalent cations or anions (with radius r = 140 pm) from the gas phase ($\varepsilon_r = 1$) to water ($\varepsilon_r = 78$).

Application of this equation to acid/base reaction (4-5) leads to a net change in Gibbs energy per mol as shown in Eq. (4-8), if only pure electrostatic interactions are considered [8].

$$\Delta G^{\circ}_{\text{electrostatic}} = \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{N_A \cdot e^2}{2 \cdot \varepsilon_r} \cdot \left(\frac{1}{r_{SH_2^{\oplus}}} + \frac{z_A^2}{r_A} - \frac{z_{HA}^2}{r_{HA}} \right) \tag{4-8}$$

Thus, the Gibbs energy difference $\Delta\Delta G$ for the ionization of a mol of HA in solvent 1 and solvent 2 with relative permittivities $\varepsilon_{r(1)}$ and $\varepsilon_{r(2)}$, respectively, provided that the radii of the reactants are the same in both solvents, is given by Eq. (4-9).

$$\begin{aligned} \Delta\Delta G^{\circ} &= (\Delta G^{\circ}_{\text{electrostatic}})_2 - (\Delta G^{\circ}_{\text{electrostatic}})_1 \\ &= \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{N_A \cdot e^2}{2} \cdot \left(\frac{1}{r_{SH_2^{\oplus}}} + \frac{z_A^2}{r_A} - \frac{z_{HA}^2}{r_{HA}} \right) \cdot \left(\frac{1}{\varepsilon_{r(2)}} - \frac{1}{\varepsilon_{r(1)}} \right) \end{aligned} \tag{4-9}$$

Because the Gibbs energy of reaction, ΔG, is related to the equilibrium constant K_a, according to Eq. (4-3), Eq. (4-9) can be converted into Eq. (4-10).

$$\ln \frac{(K_a)_2}{(K_a)_1} = -\frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{N_A \cdot e^2}{2 \cdot RT} \cdot \left(\frac{1}{r_{SH_2^{\oplus}}} + \frac{z_A^2}{r_A} - \frac{z_{HA}^2}{r_{HA}} \right) \cdot \left(\frac{1}{\varepsilon_{r(2)}} - \frac{1}{\varepsilon_{r(1)}} \right) \tag{4-10}^{*)}$$

* A modification of Eq. (4-10) has been developed by Izmailov by expressing the specific and nonspecific solute/solvent interactions by means of a conventional division of the ΔG values into electrostatic and nonelectrostatic ones, in order to obtain a more complete equation that also contains the constants of complex formation between solutes and solvent [318].

Eq. (4-10) can be used only for solvents of equal acid and base strength, because only the effect of the solvent relative permittivity on the degree of ionization is considered. Under these conditions, Eq. (4-10) predicts that the logarithm of the ionization constant K_a of HA should be inversely proportional to the relative permittivity of the solvent in which HA is dissolved. However, one has to take into account the fact that the relative permittivities near solute ions can differ considerably because of the effect of dielectric saturation, which hinders the precise calculation of electrostatic interactions. Because of these restrictions, Eq. (4-10) can be expected to yield only semiquantitative results. Nevertheless, it allows us to predict qualitatively how the charge type of an acid affects the ionization constant in solvents of different relative permittivities.

If the acid HA in Eq. (4-5) has a charge of +1 ($z = 0$; *e.g.* NH_4^+), the right-hand side of Eq. (4-10) involves only the difference between two reciprocal radii ($z_A = 0$), and is often very nearly zero. Therefore, a change in the relative permittivity should in this case have only a small, if any, influence on the ionization equilibrium of an acid such as NH_4^+. This is plausible, because in this acid/base reaction (4-5) no charges are created or destroyed at all. Also, there is no electrostatic attraction between the positively charged species HA and the neutral conjugate base A. Indeed, the cationic acid NH_4^+ is only about 1/10 as strong in ethanol as it is in water according to the lower basicity of the solvent ethanol compared with water (*cf.* Table 4-1).

On the other hand, if HA is an uncharged acid ($z = -1$; *e.g.* $CH_3—CO_2H$), the right-hand side of Eq. (4-10) involves the sum of two reciprocal radii ($z_{HA} = 0$) and a strong influence of the relative permittivity on the ionization equilibrium is expected. Because in acid/base reactions of this charge type, neutral molecules are converted into anions and cations, which attract each other, reaction (4-5) will shift to the right with an increase in relative permittivity of the solvent in which HA is dissolved. Ionization increases when ε_r increases. This rule is qualitatively verifiable for water and alcohols as

Table 4-1. Solvent influence on the acid/base equilibrium $HA^{z+1} + SH \rightleftharpoons SH_2^+ + A^z$ for various charge types[a].

z	Acid-base charge type	pK_a change with increasing basicity of SH	pK_a change with increasing ε_r of SH	Examples	$\Delta pK_a = (pK_a)_{H_2O} - (pK_a)_{C_2H_5OH}$[a,b]
0	$HA^{\oplus}/A^0$	decrease	no or small effect	$NH_4^{\oplus}/NH_3$	−1.2
				$H_5C_6NH_3^{\oplus}/H_5C_6NH_2$	−0.4
−1	$HA^0/A^{\ominus}$	decrease	decrease	$H_3C—CO_2H/H_3C—CO_2^{\ominus}$	−5.7
				Picric acid/Picric anion	−3.3
−2	$HA^{\ominus}/A^{2\ominus}$	decrease	decrease	$HO_2C—(CH_2)_2—CO_2^{\ominus}/{}^{\ominus}O_2C—(CH_2)_2—CO_2^{\ominus}$	−5.8[d]
				$HO_2C—CH=CH—CO_2^{\ominus}/{}^{\ominus}O_2C—CH=CH—CO_2^{\ominus}$[c]	−5.5[d]

[a] See reference [11] for more examples.
[b] Relative permittivities: $\varepsilon_r = 78.4$ (H_2O), $\varepsilon_r = 32.7$ (CH_3OH), and $\varepsilon_r = 24.6$ (C_2H_5OH) at 25 °C.
[c] Fumaric acid.
[d] $\Delta pK_a = (pK_a)_{H_2O} - (pK_a)_{CH_3OH}$ in this case.

solvents. In fact, acetic acid is only about 10^{-6} times as strong an acid in ethanol (low ε_r) as in water (high ε_r) (*cf.* Table 4-1).

In the case of an anionic acid ($z = -2$; *e.g.* HSO_4^-), the acid/base equilibrium (4-5) will be shifted to the right with an increase in relative permittivity of the solvent in which HA is dissolved. Here, the ionization will increase much more quickly with an increase in relative permittivity than in the preceding case.

Table 4-1 shows the predicted solvent influence on the acidity constants K_a for various charge types of acid/base pairs.

Analogous rules apply to the basicity constants K_b according to Eq. (4-6). For example, for a pair A^0/HA^+ ($z = 0$; *e.g.* $NH_3 + ROH$), K_b increases as the relative permittivity of the solvent increases.

Because it is practically impossible to find solvents that differ only in their relative permittivities, and not in their basicity or acidity, the predictions based on Eq. (4-10) are usually not in good agreement with the experimental results. In addition, the solvation capability or polarity of a solvent is not described by its relative permittivity alone. Besides the purely electrostatic Coulomb interactions, there exist other specific and unspecific interaction forces, such as ion-dipole, dipole-dipole, hydrogen bonding, ion-pair formation, *etc.* Also, the model used to describe the coulombic effects on acid/base ionization constants ignores the actual shape and size of the individual ions. For example, in contrast to the carboxylic acids the acidity constant of picric acid increases by a factor of 1500 simply for a solvent change from ethanol to water (*cf.* Table 4-1). Because the negative charge of the picric anion is delocalized over a large molecule, the solvation enthalpy of this anion will be lower than that of the carboxylate anions. This means that its stability does not change with the solvation capability of the solvent to the same extent as for the carboxylate anions, where the negative charge is more or less localized on only two oxygen atoms.

Using activity coefficients, it is possible to examine in greater detail the effect of solvents on acid and base strengths. Equilibrium constants expressed in terms of concentration are solvent-dependent. Solvent-independent, so-called thermodynamic equilibrium constants are obtained when the concentration terms are replaced by concentration-based activity terms. The thermodynamic equilibrium constant K for the reaction $HA \rightleftharpoons H^{\oplus} + A^{\ominus}$ is given by the following equation:

$$K = \frac{a_{H^+} \cdot a_{A^-}}{a_{HA}} = \frac{[H^+]\gamma_{H^+}[A^-]\gamma_{A^-}}{[HA]\gamma_{HA}} = K_a \cdot \frac{\gamma_{H^+} \cdot \gamma_{A^-}}{\gamma_{HA}} \tag{4-11}$$

In highly dilute aqueous solution, the activity coefficients approach the value 1. That is, in aqueous solution, K and K_a become practically equal at infinite dilution. If the equilibrium constant, expressed in concentration terms, is denoted by K_{SH} for the solvent SH, and K_{H_2O} is the value of K_a in water at infinite dilution, then it follows that:

$$K = K_{SH} \cdot \frac{\gamma_{H^+} \cdot \gamma_{A^-}}{\gamma_{HA}} = K_{H_2O} \tag{4-12}$$

where the values of γ are the appropriate activity coefficients in the solvent SH. Analogously, for another acid HB with an acid/base equilibrium $HB \rightleftharpoons H^{\oplus} + B^{\ominus}$, Eq. (4-13) can be written:

$$K' = K'_{SH} \cdot \frac{\gamma_{H^+} \cdot \gamma_{B^-}}{\gamma_{HB}} = K'_{H_2O} \tag{4-13}$$

Division of Eq. (4-12) by Eq. (4-13) gives Eq. (4-14)

$$\frac{K_{SH} \cdot \gamma_{HB} \cdot \gamma_{A^-}}{K'_{SH} \cdot \gamma_{HA} \cdot \gamma_{B^-}} = \frac{K_{H_2O}}{K'_{H_2O}} \tag{4-14}$$

which can be rearranged to Eq. (4-15)

$$K_{SH} = K'_{SH} \cdot \frac{K_{H_2O}}{K'_{H_2O}} \cdot \frac{\gamma_{HA} \cdot \gamma_{B^-}}{\gamma_{HB} \cdot \gamma_{A^-}} \tag{4-15}$$

The acidity constant of a given acid in any solvent SH can then be calculated from Eq. (4-15) when its acidity constant in water at infinite dilution, the acidity constant of another acid HB in water and in solvent SH, and the activity term are all known. The values γ_{HA} and γ_{HB} can be obtained from solubilities, partial pressures, distribution coefficients, *etc.* The ratio $\gamma_{B^-}/\gamma_{A^-}$ can be determined potentiometrically, or from the solubility of salts. The values of K_{SH} calculated in this way correspond satisfactorily with the measured values.

At this point, we can only refer to the various acidity scales for different series of solvents, such as those of Hammett [15] and Grunwald [16]. The acidity functions are introduced in order to obtain expressions that are not affected by the relative permittivity and that allow a quantitative comparison of acidity in different solvents. It should be stated, however, that there exists no single scale of acidity or basicity that is universally valid in all types of solvents and applicable to both equilibrium and kinetic situations [17, 109].

Medium effects on acid/base equilibria in aqueous solutions of strong acids have been analyzed not only in terms of Hammett acidity functions, but also in terms of linear Gibbs energy relationships, first developed by Bunnett *et al.* [225]; see [226] for a review.

The ionisation constants of many acidic organic compounds determined in water [110a] and in twelve of the most popular dipolar non-HBD solvents [110b] have been compiled, as have the methods of determination [111] and prediction [112] of pK_a values. Particular attention has been paid to C—H acidic compounds [113]. Whereas the ionisation constants of Brønsted acids and bases for aqueous solutions are well known, the corresponding pK_a values for nonaqueous solutions are comparatively scarce.

4.2.2 Gas-Phase Acidities and Basicities [18–21, 114–118]

Acidities and basicities of organic compounds are expected to be different in the gas phase and in solution. Whereas in the gas phase acidity and basicity are intrinsic properties of the individual molecules, in the liquid state these properties belong to the phase

as a whole due to the interaction between solute and solvent molecules. Solution-phase acidities and basicities reflect solvent properties as well as the intrinsic proton-donating and -accepting power of the solute. Therefore, acidities and basicities measured under solvent-free conditions must be available before the interplay between solute properties and solvent effects can be studied. Efforts in this direction have not made much progress until recently due to the lack of methods for measurements of gas-phase acidities and basicities.

Gas-phase acidities and basicities for many organic compounds are now available, primarily due to the development within the past decades of three new experimental techniques: pulsed high-pressure (*i.e.* 0.1 ... 1300 Pa) mass spectrometry (HPMS) [22, 23, 118], the flowing afterglow (FA) technique with a fast-flowing gas like helium in the pressure range of *ca.* $10^{-1} \ldots 10^{-2}$ Pa [119], and pulsed electron beam, trapped ion cell, ion cyclotron resonance (ICR) spectrometry, carried out at *ca.* $10^{-6} \ldots 10^{-3}$ Pa [24–26, 115].

The development of these ion-molecule equilibrium measurements has completely changed the status of acid/base reactions (and of other reactions; *cf.* Section 5.2) in the gas phase. It is now possible to compare the complex and poorly understood situation in solution with the simple state in the gas phase. It is also possible to determine the acidity of all acids in the gas phase, from the weakest such as methane to the strongest. In solution, however, due to the levelling effect of the solvent or solubility problems, only a certain range of acids can be measured in a given solvent.

Using these methods, relative intrinsic acidities and basicities of molecules in the gas phase have been determined by measuring equilibrium constants, $K = [A^{\ominus}] \cdot [BH]/[HA] \cdot [B^{\ominus}]$, for proton-transfer reactions such as

$$(HA)_{gas} + (B^{\ominus})_{gas} \rightleftharpoons (A^{\ominus})_{gas} + (BH)_{gas} \tag{4-16}$$

Similar measurements for proton-transfer reactions such as (S = solvent molecules)

$$(HA)_{gas} + (B^{\ominus} \cdot S_n)_{gas} \rightleftharpoons (A^{\ominus} \cdot S_n)_{gas} + (BH)_{gas} \tag{4-17}$$

have allowed the scrutiny of the solvent influence as a function of stepwise solvation of the participating ions by sequential addition of solvent molecules ($n = 0$ up to $4 \ldots 9$), thus bridging the gap between the gas-phase and solution reaction.

The standard molar Gibbs energy change for reaction (4-16), $\Delta G^{\circ} = -RT \cdot \ln K$, is then a measure of the relative acidity of HA and BH (or of the relative basicity of $B^{\ominus}$ and $A^{\ominus}$). Series of acids and bases have been studied to establish a scale of relative acidities in the same manner as pK_a values are determined in solution.

In addition, an absolute intrinsic acidity or basicity scale as the case may be, corresponding to the reaction (4-18),

$$(HA)_{gas} \rightleftharpoons (H^{\oplus})_{gas} + (A^{\ominus})_{gas} \tag{4-18}$$

can be established by incorporating certain standard reactions such as $H_2 \rightleftharpoons H^{\oplus} + H^{\ominus}$ and $HF \rightleftharpoons H^{\oplus} + F^{\ominus}$, for which ΔH° and ΔG° can be calculated from available data, into the relative scale. These absolute gas-phase acidities are conveniently expressed

in terms of the proton affinity, PA, of the anion $A^{\ominus}$, defined as the standard molar enthalpy change ΔH° for the gaseous deprotonation reaction (4-18): $PA(A^{\ominus}) = -\Delta H^{\circ}$. For most simple cases, the change in ΔS° for this reaction is about the same. Thus, ΔH° (and ΔG°) varies in about the same way as the proton affinity does.

From consideration of the following thermodynamic cycle,

$H\text{-}A \longrightarrow H^{\bullet} + A^{\bullet}$	D_{HA} (homolytic dissociation energy of the H—A bond)
$A^{\bullet} + e^{\ominus} \longrightarrow A^{\ominus}$	$-EA_A$ (electron affinity of radical A)
$H^{\bullet} \longrightarrow H^{\oplus} + e^{\ominus}$	IP_H (ionization potential of atomic hydrogen, *i.e.* 1312 kJ/mol)
$H\text{-}A \longrightarrow H^{\oplus} + A^{\ominus}$	

the proton affinity of $A^{\ominus}$ is given by Eq. (4-19):

$$PA = D_{HA} - EA_A + IP_H \tag{4-19}$$

Since IP_H is a constant for every reaction, for the sake of simplicity this term is usually omitted; the acidities are then expressed as $D_{HA} - EA_A$. The acid strength increases as this difference decreases.

Scales of gas-phase proton affinities and gas-phase acidities for numerous organic compounds, including super bases and super acids [232], can be found in references [115, 120, 232, 233]; for comparisons of gas-phase and solution acidities, see references [124, 234–238].

In most cases, the gas-phase acidity orders differ dramatically from those observed in solution since the Gibbs energies of solvation (*ca.* 200–600 kJ/mol; *cf.* Table 2-8 in Section 2.3) are much larger than the intrinsic acidity differences for most pairs of compounds. Thus, the relative acidities in solution are often dictated by the differential Gibbs energies of solvation rather than by the intrinsic properties of the solute molecules.

The solution and gas-phase acidities of C—H acids are of particular interest because of the wide structural variations that are possible in this class of compounds [113, 123]. A qualitative ordering of a selection of C—H acids (and some O—H acids for comparison) gives the following sequence of increasing acidity in the gas phase [120]:

$$CH_4 < H_2O < CH_3OH < C_6H_5CH_3 < HC{\equiv}CH < CH_3SOCH_3 < CH_3CN$$
$$< CH_3COCH_3 < CH_3CHO < C_6H_5COCH_3 < CH_3NO_2 < \text{Cyclopentadiene}$$
$$< \text{Fluorene} < CH_3CO_2H < CH_2(CN)_2$$

Surprisingly enough, toluene is more acidic than water in the gas phase, but *ca.* 20 orders of magnitude less acidic in solution. Thus, in the gas phase, the reaction of $HO^{\ominus}$ with toluene proceeds rapidly, with the release of energy, to yield charge-delocalized $C_6H_5CH_2^{\ominus}$ and H_2O. In aqueous solution, the reaction proceeds in the opposite direction, converting the benzyl anion into toluene by water, and the latter giving charge-localized $HO^{\ominus}$.

Contrary to the behaviour in solution, malononitrile is a stronger acid than acetic acid in the gas phase, while fluorene, which is almost 10^5 times less acidic than cyclopentadiene in solution, becomes the stronger of the two in the gas phase. The reason for the reversed cyclopentadiene/fluorene acidity order is the better solvation of the smaller cyclopentadienide ion relative to the large fluorenide ion with the more delocalized negative charge, on going from the gas phase to solution. Whereas the larger, more charge-delocalized anion is preferably produced in the gas phase, in solution, the smaller anion with the higher charge density, *i.e.* the better solvated anion, is favoured.

A study of some C—H acids in dimethyl sulfoxide solution led to an acidity order that almost paralleled that found in the gas phase, whereas in protic solvents the order was different [116, 124, 125]. This result highlights the importance of specific solute/solvent interactions such as hydrogen bonding in comparing acid/base equilibria measured in the gas phase and in solution.

Some features of the solution pK_a scale are retained in the gas phase. For instance, the acidity of hydrogen halides increases in the order HF < HCl < HBr < HI, both in aqueous solution and in the gas phase [120].

The discovery of Brauman and Blair in 1968 [34] that the acidities of aliphatic alcohols are completely reversed on going from bulk solution to the gas phase was a landmark in the interpretation of solvent and substituent effects on acid/base equilibria. The gas-phase acidity of alcohols increases in the following order [34, 125, 126]:

$$\begin{aligned}&H\text{-}OH < CH_3\text{-}OH < CH_3CH_2\text{-}OH < CH_3CH_2CH_2\text{-}OH < (CH_3)_2CH\text{-}OH\\ &< CH_3CH_2CH_2CH_2\text{-}OH < (CH_3)_3C\text{-}OH < Me_3CCH_2\text{-}OH < C_6H_5CH_2\text{-}OH\\ &< (Me_3C)_2CH\text{-}OH < CH_3\text{-}SH < CH_3CH_2\text{-}SH \ll C_6H_5\text{-}OH\end{aligned}$$

In the gas phase, tertiary alcohols are more acidic than secondary alcohols, and these in turn are stronger acids than primary alcohols. In other words, the anion $R—CH_2O^{\ominus}$ is a stronger base than both $R_2CH—O^{\ominus}$ and $R_3C—O^{\ominus}$. This is in striking contrast to the solution behaviour, where introduction of alkyl groups at the OH-bearing carbon atom causes a significant increase in basicity of alkoxide anions [35, 127].

Prior to the availability of gas-phase data, the solution order of acidities was taken as evidence of anion destabilization through electron donation by methyl groups. Now it is clear that the liquid-phase ordering is due entirely to differential solvation of the reactants of Eq. (4-18). The effects of alkyl groups on gas-phase acidities have been considered in terms of inductive and polarization effects. It is now established that the stabilization of anions by alkyl substitution is due mostly to the role played by polarization forces between the negatively charged centre and the alkyl groups. A charged atom in an isolated ion in the gas-phase has only its attached alkyl groups to interact with, whereas in solution the polarizability of the surrounding solvent molecules is an additional factor in stabilizing the ion. Since polarizability generally increases with molecular volume, all alkyl groups are more polarizable than the hydrogen atom, and can stabilize a nearby charge whether that charge is negative or positive. Hence, the gas-phase acidities (and basicities) of alcohols should increase the greater the number and size of the alkyl groups. The reversed acidity order of alcohols obtained in bulk solution ($RCH_2—OH > R_2CH—OH > R_3C—OH$) can be explained by the assumption that

stabilization of the alkoxide ions through hydrogen bonding should be much better with the sterically less hindered $RCH_2—O^{\ominus}$ than with $R_2CH—O^{\ominus}$ and $R_3C—O^{\ominus}$ [128]. Since the Gibbs energies of solvation of the alkoxide ions are large compared to the difference in ionization energies according to Eq. (4-18), a reversal of the acidity order easily occurs in going from the gas phase to solution. Thus, the solution acidity order of alcohols is an artefact, and does not represent any intrinsic property of the alcohol molecules.

By quantitatively measuring the equilibrium given in Eq. (4-20), it has been shown [129] that the reversal of the relative acidities of methanol and ethanol on going from the

$$CH_3O^{\ominus}\cdots HOCH_3 + C_2H_5OH \rightleftharpoons C_2H_5O^{\ominus}\cdots HOCH_3 + CH_3OH \qquad (4\text{-}20)$$

gas phase to bulk solvent is almost half completed with the first molecule of solvation. The Gibbs energy of solvation of $CH_3O^{\ominus}$ using one molecule of CH_3OH is approximately 71 kJ/mol (17 kcal/mol) [130]. Thus, the first molecule of solvent causes the behaviour of the alkoxide ion to become already "solution-like". In other cases [*cf.* Eq. (4-17)], more solvent molecules are necessary for the reactivity to approach that of the bulk solvated form. Obviously, the first few solvent molecules can contribute most of the total solvation energy. These results also show that the solvating abilities of isolated solvent molecules can be very different from their performance as bulk liquids. For example, the extremely low gas-phase acidity and basicity of "monomeric" water is completely different from its acid/base behaviour as "polymeric" water, *i.e.* as the classical amphiprotic solvent.

Other well-studied cases of acidity order changes induced by differential solvation are substituted phenols [131] and halo-substituted carboxylic acids [34a, 122, 132, 239]. For a comparison of the acid/base behaviour of oxygen-versus sulfur acids and bases (*e.g.* R—OH/R—SH) in the gas phase and in solution, see reference [214].

By comparing the gas-phase and solution acidities of substituted phenols, it has been found that the solvent not only plays a dominant role in controlling the phenol acidity, but also in modifying the effects of substituents on this acidity. Modifications due to differential substituent solvation can significantly change the order of substituent effects on phenol acidity [131]. An analogous solvent influence on the substituent effects on gas-phase and solution basicity was found in the series of 4-substituted pyridines [33].

Generally, it has been found that substituent effects on acidity or basicity are significantly attenuated by the transfer of the substituted reactant from the gas phase into solution [235, 237, 240, 241]. Substituent effects are smaller in solution and this attenuation is different for different classes of substituted organic compounds [241].

The gas-phase acidity order of haloacetic acids $X—CH_2—CO_2H$ is: $H < F < Cl < Br < I$, *i.e.* the order is reversed compared to that in aqueous solution [34a, 122, 132, 239]. Thus, the well-known aqueous acidity order is not caused by the increasing inductive substituent effect ($I < Br < Cl < F$), as is generally assumed, but rather by solvation effects.

α-Amino acids such as glycine, which are known to exist as zwitterions in the crystalline state and in aqueous solution, are not zwitterionic in the gas phase [133]. By measuring the gas-phase acidity and basicity of glycine, it has been found that in the gas phase it exists as a non-ionic molecule $H_2N—CH_2—CO_2H$! Although the zwitter-

ionic form of α-amino acids predominates in dipolar non-HBD solvents, the ratio of the zwitterionic to the uncharged form is, at 2 . . . 40 in dimethyl sulfoxide, much smaller than the corresponding values of $10^4 \ldots 10^5$ in aqueous solution [219]. This large difference can be explained by the better solvation of the carboxylate group in water compared to dimethyl sulfoxide, with solvation of the ammonium group being similar in the two solvents [219]. The reaction path for the proton transfer from the neutral to the zwitterionic form on going from the gas to the aqueous phase does not correspond to a direct intramolecular proton transfer, but probably to a proton transfer mediated by the water molecules forming the first hydration shell [242].

Structurally related to α-amino acids is sulfamic acid, which can exist as neutral ($H_2N—SO_3H$) and zwitterionic species ($^+H_3N—SO_3^-$). According to high level *ab initio* MO calculations, the neutral form is slightly more stable in the gas phase (by ca. 2 kJ/mol), whereas in polar media the zwitterion is predicted to be strongly favoured, in agreement with measurements in aqueous and DMSO solution [261].

A historically interesting example of solvent influence on basicity that has puzzled chemists for a long time will be used to conclude this section.

It has long been known that the proton-acceptor abilities of alkylamines in aqueous solution, as expressed by their pK_b values, are in the order $NH_3 < RNH_2 < R_2NH > R_3N$(!) [27]. The unexpected reduced base strength of tertiary alkylamines is found for all the common alkyl groups. However, if the basicities of methylamines are determined in the gas phase, they increase monotonically from ammonia to trimethylamine, as expected from theoretical considerations: $NH_3 < CH_3NH_2 < (CH_3)_2NH < (CH_3)_3N$ [28–30, 115]. Therefore, the "basicity anomaly" of tertiary alkylamines in aqueous solution must be related to the differential solvation of the reacting species of the corresponding acid/base equilibrium.

The reaction between the ammonium ion, $NH_4^{\oplus}$, and trimethylamine, $(CH_3)_3N$, analogous to Eq. (4-16), has been studied by pulsed ICR mass spectrometry [115]. The Gibbs energy diagram in Fig. 4-2 describes what happens to the reactants on going from



Fig. 4-2. One-dimensional Gibbs energy diagram for the acid/base equilibrium reaction between the ammonium ion and trimethylamine in the gas phase (top) and in aqueous solution (bottom) [115].

the gas phase to aqueous solution. In the gas phase, the products are more stable than the educts by 92 kJ/mol. In aqueous solution, the Gibbs energy of reaction falls to only 3 kJ/mol because of preferential solvation of the $NH_4^{\oplus}$ ion by hydrogen bonding. This means that at equilibrium in aqueous solution the concentration of the ammonium ion is higher by a factor of 10^{15} than it is in the gas phase.

Why is trimethylamine a much stronger base than ammonia in the gas phase? The increase in base strength with increasing number of alkyl substituents at the amine nitrogen atom can be considered in terms of inductive and polarization effects. The electric field of the positive charge permeates the space around the alkylammonium ion and distorts the electron clouds of the alkyl groups. Both the inductive and polarization effect make $Me_3NH^{\oplus}$ a more stable ion than $NH_4^{\oplus}$. In particular, the replacement of hydrogen atoms by the larger, more polarizable alkyl groups stabilizes charged centres. The observation that the gas-phase acidity of alcohols increases with alkyl substitution (see above) shows that alkyl groups can stabilize a charge of either sign, withdrawing or donating electrons as needed.

What happens to the same reaction in aqueous solution? Whereas the neutral reactants, ammonia and trimethylamine, are hydrated about equally well, the ammonium ion is hydrated much more strongly than is $Me_3NH^{\oplus}$. As shown by Eqs. (4-21) and (4-22), solvation through hydrogen bonding will tend to increase the base strength of all amines



(4-21)



(4-22)

in aqueous solution because the positively charged ammonium ions will be better solvated than the uncharged amines [31]. However, the solvation through hydrogen bonding will decrease with increasing alkyl substitution; *cf.* Eq. (4-22). The ammonium ion can be stabilized by four hydrogen bonds, whereas $Me_3NH^{\oplus}$ has only one acidic hydrogen atom.

The apparent basicity "anomaly" of alkylamines can now be understood in terms of two opposing influences, one base-strengthening (due to increasing alkylation of the amine), and the other base-weakening (because of reduced solvation of the ammonium ions with increasing alkylation). For solvation effects on the basicities of alkyl amines in solvents other than water, see reference [243].

Why are aniline ($pK_b = 9.4$) and pyridine ($pK_b = 8.8$) so much less basic than ammonia ($pK_b = 4.8$) in aqueous solution? For a long time, students of organic chemistry have been given reasons in terms of lone-pair delocalization and sp^2/sp^3 hybridization of the nitrogen atom. From gas-phase studies, it is now clear that aniline and

pyridine are inherently much stronger bases than ammonia [116]*). Therefore, the reversed basicity order obtained in aqueous solution is caused by differential hydration! However, if more suitable reference compounds are used, cyclohexylamine ($pK_b = 3.3$) for aniline and piperidine ($pK_b = 2.9$) for pyridine, then aromatic amines are indeed less basic than saturated ones in aqueous solution. Thus, according to Arnett [213], "the right idea has been promoted over the years but for the wrong reasons, a not infrequent occurrence in chemistry".

Another acid/base anomaly, the anomalous relationship between rates and equilibria for the proton-transfer reactions of nitroalkanes such as $H_3C—NO_2/H_3C—CH_2—NO_2/(H_3C)_2CH—NO_2$ found in water, does not exist in the gas phase [244]. Only in aqueous solution is the rate of proton abstraction by HO^- unexpectedly slower for the more acidic nitro compound.

Some general rules for solvation effects on acid/base equilibria as transferred from the gas phase to solution have been collected by Arnett [213].

4.3 Solvent Effects on Tautomeric Equilibria

4.3.1 Solvent Effects on Keto/Enol Equilibria [36–43, 134]

In general, 1,3-dicarbonyl compounds, which include β-dialdehydes, β-ketoaldehydes, β-diketones, and β-ketocarboxylic esters, can exist in solution or as the pure compound in three tautomeric forms**): the diketo form *(4a)*, the *cis*-enolic *(4b)*, and the *trans*-enolic form *(4c)*.



(4b) *(4a)* *(4c)*

Open-chain 1,3-dicarbonyl compounds are observed in the *trans*-enolic form only in rare cases [41] (for examples, see references [44, 45]). When the *trans*-enolic form is excluded, the keto/enol equilibrium constant K_T is given by Eq. (4-23).

$$K_T = \frac{[\text{enol}]}{[\text{diketo}]} \tag{4-23}$$

---

* The much greater gas-phase base strength of aniline (and cyclohexylamine) compared with ammonia is due to the polarizability of the large carbocyclic residue. However, aniline is less basic than cyclohexylamine in the gas phase as well as in aqueous solution. The nitrogen lone-pair in aniline, unlike that in cyclohexylamine, is conjugated with the aromatic π-system of the benzene ring and is thus, to some extent, delocalized. Protonation of the aniline nitrogen atom localizes this electron pair and causes some loss of delocalization energy.

** If $R \neq R'$, two *cis*-enolic and two *trans*-enolic forms can exist.

Table 4-2. Equilibrium constants and mole fractions of enol tautomers of ethyl acetoacetate *(4a)* ($R = CH_3$, $R' = OEt$; K_T), acetylacetone *(4a)* ($R = R' = CH_3$; K'_T), and 5,5-dimethylcyclohexane-1,3-dione *(5a)* (K''_T), determined ^1H NMR spectroscopically at *ca.* 20 °C at solute concentrations of *ca.* $10^{-3} \ldots 10^{-2}$ mol/L, *i.e.* under conditions unperturbed by self-association of the solute [134], values for pure solutes excepted [47].

Solvents (deuterated)	K_T	x(enol) / (cmol · mol^{-1})	K'_T	x(enol) / (cmol · mol^{-1})	K''_T	x(enol) / (cmol · mol^{-1})
Gas phase[a)]	0.74	42.5	11.7	92	–	–
Cyclohexane	1.65	62	42	98	–	–
Tetrahydrofuran	0.40	29	7.2	88	–	–
Toluene	0.39	28	10	91	0.08	7
Tetrachloromethane	0.29	22.5	29	97	–	–
Benzene	0.26	21	14.7	94	0.12	11
Ethanol	0.14	12	5.8	85	169	99.4
1,4-Dioxane	0.13	11.5	4.8	83	2.8	74
Acetone	0.13	11.5	–	–	4.2	81
Pyridine	0.10	9	3.7	79	–	–
Trichloromethane	0.09	8	5.94	86	0.05	5
Dichloromethane	0.09	8	4.2	81	–	–
Pure solute[b)]	0.081	7.5	4.3	81	–	–
Methanol	0.07	6.5	2.9	74	148	99.3
Water	0.07	6.5	0.23	19	19	95
Dimethyl sulfoxide	0.05	5	2.0	67	94	99.0

[a)] Values at 40 °C; see reference [38].
[b)] Values at 33 °C; see reference [47].

In solution, open-chain 1,3-dicarbonyl compounds enolize practically exclusively to the *cis*-enolic form *(4b)*, which is stabilized by intramolecular hydrogen bonding. In contrast, cyclic 1,3-dicarbonyl compounds (*e.g.* cycloalkane-1,3-diones [46]), can give either *trans*-enols (for small rings) or *cis*-enols (for large rings). As the diketo form is usually more dipolar than the chelated *cis*-enolic form, the keto/enol ratio often depends on solvent polarity. This will be discussed in more detail for the cases of ethyl acetoacetate and acetylacetone [47–50, 134, 135].

The equilibrium constants, measured by ^1H NMR spectroscopy, of ethyl acetoacetate and acetylacetone [47, 48, 134] (Table 4-2) indicate a higher enol content for these *cis*-enolizing 1,3-dicarbonyl compounds in apolar aprotic than in dipolar protic or dipolar aprotic solvents.

The ratios obtained in the apolar aprotic solvents approach the gas-phase values [38]. In principle, on dissolution of β-dicarbonyl compounds in solvents of low polarity, the percentage of the *cis*-enolic form increases, whereas polar solvents displace the equilibrium towards the diketo form. Although at first sight it is surprising that increasing solvent polarity diminishes the enol content, this is understandable in terms of intramolecular chelation of the enol. The enol form is the least polar of the two tautomers because intramolecular hydrogen bonding helps reduce the dipole-dipole repulsion of the carbonyl groups; this is unreduced in the diketo form. Furthermore, the enol stabilization due to the intramolecular hydrogen bonding will be more pro-



Fig. 4-3. Effect of solvent and concentration on the keto/enol equilibrium of acetylacetone in four solvents of different polarity at 37 ± 2 °C: CCl_4 (○), $CHCl_3$ (△), CH_2Cl_2 (●), and $HCON(CH_3)_2$ (▲) [50].

nounced when intermolecular hydrogen bonding with the solvent does not compete. Thus, a change to a more polar solvent, with a tendency towards intermolecular hydrogen bonding (EPD solvents), is generally associated with a decline in enol content.

In agreement with this, the enol content also depends strongly on the initial concentration of the 1,3-dicarbonyl compound; see Fig. 4-3 [50].

As the dipolar 1,3-dicarbonyl compound acetylacetone is progressively diluted with apolar solvents, the enol content increases. Conversely, progressive dilution with a dipolar aprotic EPD solvent such as, *N*,*N*-dimethylformamide reduces the enol content of the acetylacetone solution.

The addition of comparatively less polar alcohols to solutions of acetylacetone in water shifts its keto/enol equilibrium in favour of the less polar *cis*-enolic form *(4b)*, which has been quantitatively rationalized in terms of so-called pairwise solute/solvent interactions [245]. The keto/enol equilibrium of ethyl acetoacetate and acetylacetone has also been studied in polar supercritical fluids such as CHF_3 ($\mu = 1.65$ D) and $CClF_3$ ($\mu = 0.50$ D) [246]. In polar trifluoromethane, the dipolar keto form was found to be favoured, although the change in the equilibrium constant with increasing sc-fluid density (*i.e.* increasing pressure) was quite minor. For *ab initio* calculations of the relative stabilities of various enols of acetylacetone in the gas phase, and theoretical calculations of keto/enol equilibria in aqueous solutions, see references [247] and [248], respectively.

In contrast to the *cis*-enolizing 1,3-dicarbonyl compounds, the *trans*-enolizing cycloalkane-1,3-diones with four- to six-membered rings show exactly the opposite dependence on solvent polarity [46]. In these compounds, intramolecular hydrogen bonding is excluded on steric grounds. For example, 5,5-dimethyl cyclohexane-1,3-dione *(5a,b)* is 95% enolized in aqueous solution [51]. However, in dilute solution in toluene, an apolar solvent, it is only 7% enolized [52, 134]; $x(\text{enol}) = 7$ cmol/mol; *cf.* Table 4-2.

Analogously, the keto/enol equilibrium of *trans*-enolizing 2-phenylindane-1,3-dione lies far on the side of the diketo form in apolar chloroform (>95 cmol/mol), whereas the *trans*-enol form is favoured in the polar HBA solvent dimethyl sulfoxide (ca. 77 cmol/mol) [249].



(5a) (5b)



(6a) (6b)

Another example is the β-ketonitrile *(6a,b)*. Because of the linearity of the cyano group, a cyclic structure with an intramolecular hydrogen bond is impossible. As predicted, it is found that the enol content is greater in polar than in apolar solvents [53].

In general, for the protomer pairs in which the enol cannot form an intramolecular hydrogen bond, such as *(5a)* ⇌ *(5c)*, the tautomeric equilibrium seems to be controlled almost completely by the hydrogen-bond acceptor property (Lewis basicity) of the solvent. EPD solvents enhance the enol content strongly; *cf. (5a)* in Table 4-2.

For protomer pairs such as *(4a)* ⇌ *(4b)*, in which intramolecular hydrogen bonding is possible, the solute/solvent effect due to dipolarity/polarizability interactions dominates, although differential stabilization of the tautomers by hydrogen bonding remains significant. If there exists a substantial difference in the permanent dipole moments of the two tautomers, and both can donate a hydrogen bond to the solvent, the dipole/dipole solute/solvent interactions will dominate [134]. Similar results have been obtained for acetoacetic acid itself, for which the enol mole fraction ranges from less than 2 cmol/mol in D_2O to 49 cmol/mol in CCl_4 [136].

Open-chain β-keto carboxylic esters with two mesityl substituents, such as methyl 3-hydroxy-2,3-dimesityl-2-propenoate, exist in solution only as the (*Z*)-isomer *(4b)* and the (*E*)-isomer *(4c)*; no keto form *(4a)* has been observed [223]. The (*Z*)-form predominates in nonpolar solvents such as cyclohexane (90 cmol/mol) and benzene (87 cmol/mol). Increasing solvent polarity shifts this (*Z*)/(*E*) equilibrium in favour of the more polar (*E*)-isomer, up to 76 cmol/mol (*E*)-form in ethanol [223]. The introduction of mesityl substituents stabilizes enols of simple monocarbonyl compounds such as 2,2-dimesitylethenol, $Mes_2C{=}CH{-}OH$ [42, 224].

The influence of solvents on tautomeric equilibria has been related to the solubilities of the two tautomers. Formally, the dissolution of a solute to give a saturated solution may be regarded as an equilibrium:

$$\text{crystals} + \text{solvent} \rightleftharpoons \text{dilute saturated solution}$$

The energy of the solid phase is independent of the solvent, and thus differences in solubility on going from one solvent to another will be a measure of the solvent effect on the Gibbs energy of the dissolved compound [1, 2]. Table 4-3 shows that the variation in

Table 4-3. Keto/enol equilibrium of 3-benzoyl camphor in five solvents at 0 °C [37].

Solvents	K_T [a]	$\frac{S_{enol}}{S_{diketo}}$ [b]	G
Diethyl ether	6.81	6.39	1.06
Ethyl acetate	1.98	1.81	1.09
Ethanol	1.67	1.57	1.06
Methanol	0.87	0.75	1.15
Acetone	0.85	0.80	1.06

[a] K_T = [enol]/[diketo].
[b] Enol solubility/diketo solubility.

the equilibrium constants for the keto/enol tautomerization of 3-benzoyl camphor *(7a,b)*, in a series of solvents, is consistent with the solubilities of the two tautomers.



(7a) *(7b)*

The van't Hoff-Dimroth relationship (4-24) [37, 54] states that two interconvertible isomers are in equilibrium when the ratio of their concentrations is proportional to that of their solubilities S in the relevant solvent. In Eq. (4-24), G is a solvent-independent constant, characteristic of the 1,3-dicarbonyl compound. Therefore, the equilibrium constant, K_T equals G multiplied by the solubility ratio.

$$K_T = \frac{[\text{enol}]}{[\text{diketo}]} = G \cdot \frac{S_{enol}}{S_{diketo}} \tag{4-24}$$

From this, it follows that the concentration of an enol will be at a maximum in a solvent in which it has the highest relative solubility.

The applicability of Eq. (4-24) was demonstrated by O. Dimroth using 3-benzoyl camphor *(7a,b)* as an example [37]. This was particularly suitable, since both the diketo and the enol form are separately isolable, and the solubility of each is readily determined due to the slowness of their tautomerization. Although the ratio of the S terms varies by a factor of 8, the value of G is constant within an error of $\pm 5\%$. This is understandable, since for each 1,3-dicarbonyl compound, the same characteristic difference between diketo and enol forms should, in principle, be observed.

The solubility is a reflection of how well the diketo and enol forms are solvated. Deviations from Eq. (4-24) are due mainly to the fact that the equation holds only for dilute solutions. At high concentrations, the solubility is no longer determined by solvent-solute interactions alone, but self-solvation (*i.e.* interactions between the dis-

solved molecules themselves) also plays a role. Thus, as already stated, the enol content of a solution also depends on the concentration of the 1,3-dicarbonyl compound [55].

From the thermodynamic point of view, the keto/enol equilibrium is determined by the change in Gibbs energy, ΔG°, which, in turn, is comprised of the enthalpy ΔH° and entropy ΔS° of enolization; *cf.* Eq. (4-4). Therefore, the position of equilibrium in solution will be determined by the differences $\Delta H^\circ_{\text{solv}}$ and $\Delta S^\circ_{\text{solv}}$ between the keto and enol forms, according to Eq. (4-25).

$$\Delta G^\circ_{\text{keto-enol}} = \Delta H^\circ_{\text{keto-enol}} \pm \Delta H^\circ_{\text{solv}} - T \cdot \Delta S^\circ_{\text{keto-enol}} - T \cdot \Delta S^\circ_{\text{solv}} \tag{4-25}$$

The values of $\Delta H_{\text{keto-enol}}$ and $T \cdot \Delta S_{\text{keto-enol}}$ can be determined from the temperature dependence of the equilibria in the gas phase [38]. However, the magnitudes and signs of the solvation terms are not well known. ΔH_{solv} will be either negative or positive when changing from a nonpolar to a polar solvent depending on which of the tautomers is the more polar. The resulting change in the equilibrium can be compensated by the corresponding entropy change. In a nonpolar solvent, stronger solvation leads to a higher degree of order of the solvent molecules and, hence, to a decrease in the entropy. In contrast, polar solvent molecules show a high degree of order even in the absence of a dipolar solute. The thermodynamic parameters for the enolization reaction of acetyl-acetone in seven solvents have been determined. The molar Gibbs energy of reaction varies from −0.25 kJ/mol in methanol to −9.1 kJ/mol in cyclohexane at 25 °C [134a].

Since these complications have prevented a quantitative estimation of the enthalpies and entropies of keto and enol forms in different solvents, a quantitative relationship between ΔG or K_T, and the solvating ability or polarity of the solvent (expressed in physical characteristics such as the relative permittivity ε_r, the dipole moment μ, the refraction index n, *etc.*) has not been possible either. Such equations would have the form of Eq. (4-26):

$$\Delta G^\circ = f(\varepsilon_r, \mu, n, \ldots) \quad \text{or} \quad K_T = f(\varepsilon_r, \mu, n, \ldots) \tag{4-26}$$

For purely electrostatic solute/solvent interactions, the Kirkwood equation, Eq. (4-27) [56], is applicable, which relates the standard molar Gibbs free energy of transfer of spherical dipolar molecules of radius r and dipole moment μ from the gas phase ($\varepsilon_r = 1$) to a continuous medium of relative permittivity ε_r.

$$\Delta G^\circ = -\frac{N_A}{4\pi \cdot \varepsilon_0} \cdot \frac{\mu^2}{r^3} \cdot \frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \tag{4-27}$$

Application of this equation to keto/enol equilibria gives Eq. (4-28), derived by Powling and Bernstein [57].

$$(\Delta H^\circ)_{\text{solution}} = (\Delta H^\circ)_{\text{gas}} - \frac{N_A}{4\pi \cdot \varepsilon_0} \cdot \left(\frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \cdot \frac{\varrho}{M_r} \right) \cdot (\mu_1^2 - \mu_2^2) \tag{4-28}$$

M_r is the molar mass and ϱ is the density of the solvent, μ_1 and μ_2 are the dipole moments of the least and more stable isomers, respectively. The proposed linear function between the enthalpy of tautomerization and the solvent quantity $[(\varepsilon_r - 1)/(2\varepsilon_r + 1)]\varrho/M_r$ gives, in the case of ethyl acetoacetate and acetylacetone, approximate straight lines (omitting the alcohols), but with considerable scattering of the points [47]. This scattering is not surprising in view of the rather concentrated solutions used, the specific interactions expected in hydrogen-bonding solvents, and the possible entropy effects in these systems.

A wide variety of different theoretical (*e.g.* Kirkwood function) and empirical (*cf.* Chapter 7) parameters of solvent polarity have successfully been tested using multivariate statistical methods in order to model the solvent-induced changes in keto/enol equilibria [134].

Most simple monocarbonyl compounds are enolized to such a small extent, that it is difficult to determine reliably their enol content in solution [58, 137]. The enol content of acetone, for example, is about $6 \cdot 10^{-8}$ cmol/mol at equilibrium [137]. A remarkable example of a solvent-dependent keto/enol equilibrium of a monocarbonyl compound is 2-hydroxy-7-isopropyl-1,4-dimethylazulene (2-hydroxy-guaiazulene) *(8a,b)* [59].



(8a) *(8b)*

Whereas in chloroform and water, no enol form *(8b)* is detectable, the enol content is 5 cmol/mol in cyclohexane, 20 cmol/mol in methanol, 55 cmol/mol in 1,4-dioxane, and 95 cmol/mol in dimethyl sulfoxide. Apparently, the enol is stabilized in solvents that can act as hydrogen-bond acceptors, while the keto form is favoured in protic solvents acting as hydrogen-bond donors. 6-Hydroxy-4,8-dimethylazulene behaves similarly [59a]. In polar solvents such as acetonitrile or dimethyl sulfoxide, the enolic azulenoid structure is exclusively observed, whereas in less polar solvents, dichloromethane or chloroform, a keto/enol equilibrium in a ratio of about 3:1 is detectable by ^{1}H NMR measurements [59a].

In the case of the tautomerization between 9-anthrone, *(9a)*, and 9-anthranol, *(9b)*, the equilibrium lies practically completely on the side of the keto form *(9a)* in the gas phase as well as in inert solvents such as *iso*-octane and benzene; *e.g.* in benzene at 20 °C, the enol content is 0.25 cmol/mol [60, 134].



(9a) *(9b)*

Addition of increasing amounts of triethylamine to a benzene solution of *(9a)* leads to a gradual shift of the equilibrium towards the enol form *(9b)*. This can be interpreted in terms of hydrogen-bond formation between 9-anthranol and triethylamine. In hydrogen-bond accepting solvents such as *N*,*N*-dimethylformamide (enol content 56.5 cmol/mol at 20 °C), pyridine (58 cmol/mol), and dimethyl sulfoxide (61.5 cmol/mol), the anthranol content increases further [61, 134].

Particularly well-studied tautomeric keto/enol equilibria are those of 3-pyridinyl 2-picolyl ketone [138] and *t*-butyl 2-picolyl ketone [139].



(10a) *(10b)*

In both cases, the tautomeric equilibria are shifted to the more dipolar keto form *(10a)* with increasing solvent polarity. For instance, *t*-butyl 2-picolyl ketone (R = *t*-butyl) is the only tautomer observed in 2,2,2-trifluoroethanol [139], and 3-pyridinyl 2-picolyl ketone (R = 3-pyridinyl) dominates in water with a mole fraction of 60 cmol/mol [138].

Another illustrative example of a solvent-dependent keto/enol tautomerism of a monocarbonyl compound is given by 4-[(pent-1-ylthio)acetyl]-*N*-(4′-chlorobenzyl)-pyridinium chloride. In the HBA solvent dimethyl sulfoxide, the enol predominates (84 cmol/mol), while in the HBD solvent water it exists as the ketone (47 cmol/mol) and its hydrate (40 cmol/mol) [250]. Recrystallization of this pyridinium salt from a DMSO/ethyl acetate mixture yields the pure enolic form, with one molecule of DMSO per molecule of enol present in the crystal lattice.

Further examples of solvent-dependent keto/enol equilibria can be found in reference [41].

4.3.2 Solvent Effects on Other Tautomeric Equilibria [62–64, 140]

Solvent effects similar to those described for the keto/enol equilibria can also be found for other tautomerisms, *e.g.* lactim/lactam, azo/hydrazone, ring/chain equilibria, *etc.* [62–64]. The pecularities arising here can only be illustrated by means of a few representative examples.

One of the classic studies of *lactim/lactam tautomerism* is the determination of the 2-hydroxypyridine *(11a)* ⇌ 2-pyridone *(11b)* equilibrium [63–65, 141–145, 251–255].



(11a) *(11b)* *(11b′)*

Table 4-4. Gas-phase and solution equilibrium constants $K_T = [NH]/[OH]$ of 2- and 4-hydroxypyridine at 25 . . . 30 °C unless otherwise stated [65, 67].

Solvents	2-Hydroxypyridine K_T	4-Hydroxypyridine K_T
Vapour[a)]	0.4 ± 0.25 (by UV)[c)] 0.5 ± 0.3 (by IR)[d)]	<0.1[d)]
Cyclohexane[b)]	1.7	–
Trichloromethane[b)]	6.0	1.3
Acetonitrile[b)]	148	4.6
Water[a)]	910	1900

a) Reference [65].
b) Reference [67].
c) At 130 °C.
d) At 250 °C.

IR and UV/Vis [65a], mass spectrometric [65b], photoelectron [65c], microwave [65d], as well as low-temperature matrix-isolation IR spectroscopic measurements [65e] reveal that 2- and 4-hydroxypyridine (as well as 2- and 4-mercaptopyridine [65f]) exist in the gas phase and in inert matrices (N_2, Ar) under equilibrium conditions mainly in the lactim (hydroxy or mercapto) form, in contrast to the situation in solution. While in nonpolar solvents such as cyclohexane and chloroform both tautomers exist in comparable amounts, the tautomeric equilibrium is shifted entirely in favour of the lactam (oxo or thioxo) form in polar solvents such as water, as well as in the crystalline state [66, 67, 141–145, 251–255]. Supercritical-fluid 1,1-difluoroethane can be used to adjust the tautomeric constant $K_T = [(11b)]/[(11a)]$ isothermally over a continuum from gas-phase values to those measured in polar solvents, simply by increasing the pressure [254]. The gas-phase and solution equilibrium constants of 2- and 4-hydroxypyridine are given in Table 4-4.

The gas-phase equilibrium constants differ from those in aqueous solution by as much as 10^4! The large differences between the stabilities of the tautomeric forms in the gas phase and in solution once more reveal the dominant influence of solvation on relative molecular stabilities.

By considering the equilibrium $(11a) \rightleftharpoons (11b)$ in solvents of varying polarity, it has been found that increasing solvent polarity shifts the equilibrium towards the pyridone-form. This form is more dipolar than the hydroxy-form due to the contribution of the charge-separated mesomeric form $(11b')$. Whereas the solvent has little influence on the ground-state electronic structure of the lactim form, significant effects on the molecular geometry, charge distribution, and vibrational frequencies have been found for the lactam form $(11b)$, which are readily understood in terms of increasing importance of the dipolar mesomeric structure $(11b')$ in polar solvents [251]. Furthermore, the hydrogen-bonding ability of the solvent plays an important role since hydrogen-bond donors tend to stabilize the oxoform, whereas hydrogen-bond acceptors stabilize the hydroxy form. For example, the oxo form of 6-chloro-4-methyl-2-hydroxypyridine predominates in water, while in various other solvents the hydroxy form predominates: 67 cmol/mol in methanol, 56 cmol/mol in chloroform, 96 cmol/mol in dimethyl sulfoxide, and 95 cmol/mol in cyclohexane [66].

The precise determination of the protomeric equilibrium constants K_T for 2(4)-hydroxypyridine $\rightleftharpoons$ 2(4)-pyridone is rather difficult because of self-association even in highly diluted solutions of the tautomers in nonpolar solvents such as cyclohexane. Self-associated tautomers may have K_T values which are substantially different from those of the unassociated isomers [142, 255].

A quantitative model for the differential solvation of the hydroxypyridine/pyridone tautomer pair in terms of reaction-field and hydrogen-bonding effects, using multivariate regression analysis, has been given by Beak *et al.* [141]. *Ab initio* and other calculations correctly predict the greater stability of 2-hydroxypyridine *(11a)* as compared to 2-pyridone *(11b)* in the gas-phase [144, 251–253]. The calculated free energy difference for the lactim/lactam pair *(11a)* and *(11b)* in the gas phase at 25 °C is $\Delta G = 2.9$ kJ/mol in favour of *(11a)*, in good agreement with the experimental value [252].

In contrast to mono-hydroxypyridines and mono-hydroxypyrimidines, the lactim-lactam equilibria of uracils are not found to be markedly influenced by solvent polarity [143].

In the vapour phase, both 2- and 4-hydroxyquinoline exist as the NH forms, *i.e.* as 2- and 4-quinolones, in contrast to the results found for 2- and 4-hydroxypyridines [145].

The lactim/lactam tautomerism of hydroxamic acids and their O-alkyl and O-acyl derivatives has also been studied [146]. Hydroxamic acids exist in the solid state and in polar solvents as the lactam tautomer only, whereas in nonpolar solvents the hydroximic tautomer is also present. Further analogous solvent-dependent lactim/lactam equilibria have been observed for certain Schiff bases (prepared from anilines and 2(4)-hydroxybenzaldehyde [256] or 2-hydroxynaphthaldehyde [257]), for 3-hydroxypyrazole [258], and for 3-methyl-1-phenylpyrazolin-5-one [259].

Di-(2-quinolyl)methane exists in a solvent-dependent tautomeric equilibrium between a colourless form *(12a)* and a coloured, hydrogen-bonded form *(12b)* in solution [68].



(12a) *(12b)*

The thermodynamic data for the transformation of the two forms have been estimated and are given in Table 4-5.

In hydrogen-bond donor solvents such as alcohols and trichloromethane, the tautomeric equilibrium is shifted in favor of the colourless form *(12a)* more than in other solvents. This is obviously due to the formation of hydrogen bonds between *(12a)* and these protic solvents. In aprotic solvents, ΔH° is negative and the reaction is exothermic. Since, however, all ΔG° values are positive, the negative value of ΔH° must be over-compensated by a positive entropy change; *cf.* Eq. (4-4).

Table 4-5. Thermodynamic data for the tautomeric conversion *(12a)* → *(12b)* of di(2-quinolyl)-methane at 20 °C [68].

Solvents	ΔG° [a] / (kJ · mol^{-1})	ΔH° [a] / (kJ · mol^{-1})	$T \cdot \Delta S^\circ$ [a] / (kJ · mol^{-1})
Ethanol	+12.6	+9.6	−2.9
Trichloromethane	+9.6	+8.4	−1.3
tert-Butanol	+8.8	+6.7	−2.1
Benzene	+6.3	−0.4	−6.7
Tetrachloromethane	+5.4	−1.7	−7.1
N,*N*-Dimethylformamide	+4.6	−3.3	−7.9
n-Heptane	+4.2	−6.3	−10.5
Carbon disulfide	+2.9	−10.0	−13.0

[a] The energy which is added to the system is considered to be positive, and that given up by the system to be negative.

This entropy decrease for the formation of the coloured form *(12b)* may be due to the planarization of the molecule (formation of the N—H · · · N bridge) and the fixation of the di-*cis*-form *(12b)*, while in *(12a)* there is free rotation about the central C—C bond. In protic solvents, the colourless form *(12a)* is already stabilized by hydrogen bonding and $T \cdot \Delta S^\circ$ is small. Therefore, the ΔG° values are greater in these solvents and ΔH° is positive [68].

The ketimine *(13a)*, prepared from desoxybenzoin and aniline, is also subject to a solvent-dependent tautomerism called *imine/enamine tautomerism*. The enamine content of a solution of *(13a)* increases in the order tetrachloromethane (31 cmol/mol at 35 °C), [D_5]pyridine (47.5 cmol/mol at 55 °C), and [D_6]dimethyl sulfoxide (67 cmol/mol at 55 °C) [69]. Hydrogen-bond acceptor solvents favour the enamine form *(13b)* due to hydrogen-bonding, whereas in less polar and apolar solvents the equilibrium is shifted towards the imine form *(13a)* [69].



(13a) *(13b)*

Other remarkable cases of solvent-dependent imine/enamine tautomerism have been reported by Ahlbrecht *et al.* [147], Scheffold *et al.* [69a], and Pérez-Ossorio *et al.* [211].

Compounds capable of a solvent-dependent *amino/imino tautomerism* are 3-methylcytosine *(14a,b)* and 1-alkyladenines [69b]. It has been shown by IR and UV/Vis spectroscopy that, in all cases, the imino forms such as *(14b)* predominate in nonpolar media (*e.g.* 1,4-dioxane). However, the content of the amino form *(14a)* increases with increasing solvent polarity, and in aqueous solution the amino form predominates [69b]. Further interesting examples of solvent-dependent tautomeric amino/imino equilibria

are given in references [148, 149]. For a review on the solvent-dependent tautomerism of unsymmetrical acyclic and cyclic amidines, X—NH—CR=N—Y, see reference [260].



(14a) *(14b)*

An extreme case is the *nitrone/hydroxylamine tautomerism* between 2-methylindolenine-*N*-oxide *(15a)* and 2-methyl-*N*-hydroxyindole *(15b)* [70]. The position of this equilibrium depends strongly on the proton-accepting and donating abilities of the solvent: in pyridine and acetonitrile there exists 0 cmol/mol of form *(15a)*, 33 cmol/mol in tetrachloromethane, but 100 cmol/mol in phenol. Thus, it is possible to observe either *(15a)* (in phenol) or *(15b)* (in pyridine) depending on the solvent used.



(15a) *(15b)*

The position of the *N*-oxide/*N*-hydroxy equilibrium of 1-hydroxybenzotriazole, which is similar in structure to *(15b)*, is also solvent-dependent: approximately 6, 11, 18, and 26 cmol/mol of the *N*-oxide form is present at equilibrium in dimethyl sulfoxide, acetone, formamide, and methanol, respectively [218].

Because azo dyes are of commercial importance as colouring materials, the *azo/hydrazone tautomerism* of hydroxy-substituted azo compounds has been intensively studied [71, 228]. In the case of 4-phenylazo-1-naphthol *(16a)*, an increase in the solvent polarity displaces the tautomeric equilibrium towards the more dipolar quinone hydrazone form *(16b)* [72–74, 74a, 74b, 150–154]. In addition, the NH and OH groups of both tautomers are capable of forming hydrogen bonds with suitable solvents. Due to



(16a) *(16b)*

the stronger hydrogen-bond donor ability of the OH group compared with that of the NH group, it will be to a different degree. Thus, the formation of hydrogen bonds with HBA solvents such as pyridine should mainly stabilize the azo form, whereas the basic imino group in the hydrazone form should be more stabilized in HBD solvents such as chloroform or acetic acid. From UV/Vis spectroscopic measurements, the following order of increasing proportions of the hydrazone form *(16b)* has been found in solution

(mole fractions in cmol/mol): pyridine (15) < acetone (30) < ethanol (31) < methanol (40) < benzene (56) < trichloromethane (79) < acetic acid (89) [151]. That is, the azo form *(16a)* is indeed stabilized in pyridine, acetone, ethanol, and methanol, whereas the hydrazone form dominates in trichloromethane and acetic acid. This is relative to the equilibrium in benzene, which is used as reference solvent. In *N*,*N*-dimethylformamide and dimethyl sulfoxide solutions, the azo dye *(16)* is converted into its mesomeric anion, due to the high basicities and high relative permittivities of these solvents [151]. Similar results have been obtained with 4-*alkyl*azo-1-naphthols [152]. Quantum-chemical calculations have shown that the azo form *(16a)* should be the more stable isomer in the gas phase [153, 154].

Relatively independently of solvent polarity, 4-nitrosophenol *(17a)* exists in solution mainly in the 1,4-benzoquinone monoxime form *(17b)* [75, 76]: *ca.* 83 cmol/mol monoxime in 95 cl/L aqueous ethanol [75], *ca.* 86 cmol/mol in 1,4-dioxane, and *ca.* 75 cmol/mol in acetone (at 20 °C) [76].



(17a) *(17b)*

In the case of 2-nitrosophenols such as 2,4-dialkyl-6-nitrosophenols, the tautomeric equilibrium is shifted towards the 1,2-benzoquinone monoxime form with increasing polarity of the solvent [76a].

The reversible tautomeric equilibrium between phosphane oxides and ylides with a P—OH bond, *e.g.* $R_2P(=O)—CHR'_2 \rightleftharpoons R_2P(—OH)=CR'_2$ ($R = C_6H_5$; $R' = 4\text{-}Cl—C_6H_4—SO_2$), has been found to be solvent-dependent [155]. The more dipolar phosphane oxide dominates in polar solvents (90 cmol/mol in dichloromethane at 25 °C), whereas in HBA solvents such as tetrahydrofuran the P—OH form is favoured (*ca.* 54 cmol/mol).

Finally, four different examples of solvent-dependent *ring/chain tautomerism* should be mentioned [77, 210]. The equilibrium between phthalaldehydic acid *(18a)* and phthalide *(18b)*, which in decalin lies in favor of *(18b)* (90 cmol/mol phthalide at 20 °C), is strongly shifted towards the open-chain form *(18a)* in the hydrogen-bond accepting solvent dimethyl sulfoxide (only 5 cmol/mol phthalide) [78].



(18a) *(18b)*

In the case of 5-hydroxy-2-pentanone, there is a slight preference for the open-

chain form *(19b)* over the cyclic hemiketal *(19a)* (in [D_{12}]cyclohexane 55 cmol/mol and in [D_6]dimethyl sulfoxide 61 cmol/mol *(19b)*) in most organic solvents [79].



(19a) *(19b)*

An increase in solvent polarity further favours the open-chain tautomer; in water there is no evidence for any cyclic form [79]. This is remarkable in view of the fact that the furanose/pyranose equilibria of sugars, which are interconverted through the open-chain form, are also solvent-dependent [80, 81, 159]. Arabinose, for example, in [D_5]pyridine consists of 66 cmol/mol pyranose form (α:β = 33:33) and 34 cmol/mol furanose form (α:β = 21:13), compared with 95.5 cmol/mol pyranose form (α:β = 60:35.5) and 4.5 cmol/mol furanose form (α:β = 2.5:2.0) in deuterium oxide as solvent [80, 159].

In solution, the hydroxylaminomethylation product of 2-naphthol and acetaldehyde exhibits an equilibrium between the hydroxynitrone form *(20a)* and the cyclic hydroxylamine form *(20b)*, the position of which depends on the solvent: 29 cmol/mol hydroxynitrone form in [D_6]dimethyl sulfoxide, and 94 cmol/mol in [D_4]methanol [80a].



(20a) *(20b)*

The ring/chain tautomeric equilibrium between (2-hydroxyphenylimino)phosphorane *(21a)* and 1,3,2-benzoxazaphospholine *(21b)* has been studied in thirteen solvents by NMR spectroscopy [80b]. This equilibrium is shifted towards the ring-form *(21b)* in hydrogen-bond acceptor solvents (*e.g.* tris-*n*-propylamine, dimethyl sulfoxide), compared to inert solvents such as benzene or acetone. Therefore, depending on substituents and solvents, it is possible to prepare either iminophosphoranes or benzoxazaphospholines [80b].



(21a) *(21b)*

Further remarkable examples of solvent-dependent ring/chain tautomeric equilibria can be found in references [156–158, 210].

The positions of *metallotropic tautomeric equilibria* can also be strongly solvent-dependent [160–163]. Metallotropic transformations of the σ,σ-type are related to prototropic tautomeric equilibria whereby the mobile hydrogen atom is replaced by an organometallic group. Metallotropic reactions also include π,π- and σ,π-transitions depending on the nature of the bond formed by the metal [160]. Two examples of σ,σ-type metallotropic equilibria will conclude this Section.

The metallotropic C → O transition of the trimethylstannyl group of liquid trimethyl-phenacyltin *(22a)* to give the O-isomer *(22b)* has been found to be solvent-dependent [161].



(22a) *(22b)*

As expected, the relative concentration of the more dipolar C-isomer *(22a)* increases with increasing solvent polarity (concentration of *(22a)* in cmol/mol): C_6H_{12} (74) < pure liquid (78) < C_6H_6 (81) < $ClCH_2CH_2Cl$ (83) < $CHCl_3$ (95) < CH_3COCH_3 (>99) [161].

Dynamic exchange of the trimethylstannyl substituent in cyclopenta-1,3-dienyl-trimethylstannane *(23a)* (and in cyclonona-1,3,5,7-tetraenyl-trimethylstannane) proceeds in less polar, weakly coordinating solvents such as tetrahydrofuran, 1,2-dimethoxyethane, trichloromethane, and dichloromethane through an intramolecular, orbital-symmetry controlled sigmatropic reaction. Addition of EPD solvents such as



(23a) *(23b)* *(23c)*

hexamethylphosphoric triamide or *N*,*N*-dimethylformamide to the tetrahydrofuran solution of *(23a)* shifts the equilibrium in favour of the ion pair *(23c)*, thus facilitating an intermolecular, dissociative mechanism for the substituent exchange [162].

In general, polar but weakly coordinating solvents will facilitate intramolecular metallotropic processes. Solvents exhibiting both high polarity and high coordinating capacity (EPD solvents), capable of inducing the heterolysis of the carbon-metal bond, should accelerate metallotropic processes through an intermolecular dissociative mechanism.

4.4 Solvent Effects on Other Equilibria

Not only tautomeric equilibria are subject to considerable solvent effects. Other equilibria, such as rotational and conformational equilibria [81–83], *cis*/*trans* (or *E*/*Z*) isomerization, valence isomerization [84], ionization, dissociation, and association [85] (some of which are considered in Section 2.6), complex equilibria [86, 163, 262, 263], acid/base equilibria [264, 265] *etc.*, are also strongly affected by the medium. Only a small number of representative examples will be considered in this Section in order to give an idea of how solvents can affect these different kinds of equilibria.

4.4.1 Solvent Effects on Brønsted Acid/Base Equilibria [8–13, 104–108, 264, 265]

Sections 3.3.1 and 4.2.1 dealt with Brønsted acid/base equilibria in which the solvent itself is involved in the chemical reaction as either an acid or a base. This Section describes some examples of solvent effects on proton-transfer (PT) reactions in which the solvent does not intervene directly as a reaction partner. New interest in the investigation of such acid/base equilibria in non-aqueous solvents has been generated by the pioneering work of Barrow *et al.* [164]. He studied the acid/base reactions between carboxylic acids and amines in tetra- and trichloromethane. A more recent compilation of Brønsted acid/base equilibrium constants, determined in up to twelve dipolar aprotic solvents, demonstrates the appreciable solvent influence on acid ionization constants [264]. For example, the pK_a value of benzoic acid varies from 4.2 in water, 11.0 in dimethyl sulfoxide, 12.3 in *N*,*N*-dimethylformamide, up to 20.7 in acetonitrile, that is by about 16 powers of ten [264].

According to Eq. (4-29), protons can be transferred from Brønsted acids A—H to bases |B *via* the hydrogen-bonded covalent and ionic complexes *(a)* and *(b)*, depending on both the relative acidity and basic strength of A—H and |B, respectively, and the solvation capability of the surrounding medium [265, 266]. Eq. (4-29) is simplified because not only 1:1 complexes but 1:2 and higher complexes can be formed in solution.

$$A\text{-}H + |B \underset{}{\overset{\text{Association}}{\rightleftharpoons}} \underset{(a)}{A\text{-}H\cdots B} \overset{\text{Proton Transfer}}{\rightleftharpoons} \underset{(b)}{A^{\ominus}\cdots H\text{-}B^{\oplus}} \overset{\text{Dissociation}}{\rightleftharpoons} A^{\ominus} + H\text{-}B^{\oplus} \qquad (4\text{-}29)$$

The solvent can influence all three steps of Eq. (4-29): the association, the proton-transfer, and the dissociation step. The main factor which determines the position of the acid/base equilibrium given in Eq. (4-29) is the differential solvation of the covalent and the ionic hydrogen-bonded complexes *(a)* and *(b)*. Both the hydrogen-bonded complex *(a)* and the proton-transfer ion pair *(b)* have been observed in the systems 4-nitrophenol/triethylamine [165], picric acid/triethylamine [166], chloro-substituted phenols/*N*-methylpiperidine or *n*-octylamine [167], and trifluoroacetic acid/pyridine [168]. With increasing solvent polarity, the proton-transfer equilibrium *(a)* $\rightleftharpoons$ *(b)* is shifted in favour of the ionic structure *(b)*. Thermodynamic parameters ($K, \Delta H^{\circ}$) for the formation of hydrogen-bonded complexes of phenol with various bases in different solvents can be found in reference [209]. The strengths of hydrogen bonds between solutes (protonated amines and phenolate ions) in aqueous solution have been studied [220]. Formation of such solute/solute hydrogen bonds in water as solvent requires that

competition from hydrogen-bonding of the HBD and HBA molecules to 55 M water must be overcome.

A simple example of an *intra*molecular proton-transfer reaction (thus avoiding the association and dissociation step) is given by the Mannich base *(24)* [169, 170], which can be considered as an analogue of the corresponding intermolecular complexes between phenols and amines [163]. UV/Vis and IR spectroscopic measurements show that this proton-transfer equilibrium is shifted to the right-hand side with increasing solvent polarity (concentration of *(24b)* in cmol/mol): CCl_4 (0) < $CHCl_3$ (15) < CH_2Cl_2 (26) < $ClCH_2CH_2Cl$ (30) < CH_3CN (40) [170]. A linear relationship exists between ln K_{PT} and Onsager's reaction field parameter $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, demonstrating the presence of nonspecific solute/solvent interactions only in these non-HBD solvents.



(24a) *(24b)*

Another rather simple example is the acid/base reaction between tropolone and triethylamine, which has been studied using IR and ^1H NMR spectroscopy in various solvents [171].



In non-HBD solvents such as *n*-heptane, tetrachloromethane, diethyl ether, deuterio-trichloromethane, and dimethyl sulfoxide, tropolone transfers its proton to triethylamine to give an ion pair, which is in equilibrium with the non-associated reactants. There is no formation of a hydrogen-bonded complex between tropolone and triethylamine because of the fact that tropolone itself is intramolecularly hydrogen-bonded. The extent of the ion pair formation increases with solvent polarity. In polar HBD solvents such as ethanol, methanol, and water, this proton-transfer equilibrium is shifted completely towards the formation of triethylammonium tropolonate [171].

A peculiar example of a solvent-dependent regiospecific proton-transfer equilibrium is found for 4-amino-5-methylacridine *(25)* [172]. In aqueous hydrochloric acid, the ring nitrogen atom of *(25)* is protonated to give *(25a)*, whereas in ethanolic hydrochloric acid the primary amino group preferentially accepts the proton to give *(25b)*. Without a methyl group in the 5-position (*i.e.* with 4-aminoacridine) only the ring nitrogen atom is protonated in both solvents.



(25a) *(25)* *(25b)*

Obviously, the 5-methyl substituent sterically impedes the ethanol solvation of the $NH^{\oplus}$ form *(25a)*, thus favouring reaction at the more exposed 4-amino group. In water, protonation of the ring nitrogen atom and solvation of the resulting $NH^{\oplus}$ form *(25a)* by the smaller water molecules can take place despite the methyl group [172].

The regioselectivity of protonation of various types of amides has been studied theoretically and experimentally in the gas phase and in water, showing that the site of protonation (acid residue or amide nitrogen) is often altered on going from the gas phase to aqueous solution [267].

4.4.2 Solvent Effects on Lewis Acid/Base Equilibria [106–108, 173, 174, 268]

Sections 2.2.6 and 2.6 dealt with Lewis acid/base equilibria in which principally the solvent itself is involved in the chemical reaction, either as a Lewis acid (EPA solvents) or as a Lewis base (EPD solvents). This Section includes some examples of solvent-dependent Lewis acid/base equilibria in which the solvent is not directly involved as the reaction partner, but as the surrounding and interacting medium.

Formally analogous to Eq. (4-29), Eq. (4-30) describes in a simplified manner the reaction between a Lewis acid $A^{\oplus}$ and a Lewis base $|B^{\ominus}$, via tight ion pairs (which can sometimes be considered as EPD/EPA complexes), to give the covalent ionogen A—B [*cf.* also Eq. (2-13) in Section 2.6].

$$\underset{\text{Free Ions}}{A^{\oplus} + |B^{\ominus}} \underset{\text{Dissociation}}{\overset{\text{Association}}{\rightleftharpoons}} \underset{\text{Tight Ion Pair}}{\underset{(a)}{A^{\oplus}|B^{\ominus}}} \underset{\text{Ionization}}{\overset{\text{Electron transfer}}{\rightleftharpoons}} \underset{\text{Ionogen}}{\underset{(b)}{A\text{-}B}} \qquad (4\text{-}30)$$

The position of this equilibrium depends on the electrophilicity or nucleophilicity of $A^{\oplus}$ and $|B^{\ominus}$, respectively, as well as the solvation capability of the surrounding medium. The solvent can influence the association as well as the electron-transfer step (or in the reverse reaction the ionization and dissociation step). The position of the Lewis acid/base equilibrium given in Eq. (4-30) will depend mainly on the differential solvation of the ionic and covalent species *(a)* and *(b)*.

A simple example of an *intra*molecular Lewis acid/base reaction (thus avoiding the association step) is the xanthene dye rhodamine B, which exists in solution either in the red-coloured zwitterionic form *(26a)* or as the colourless lactonic form *(26b)* [175, 221, 222]. Solutions of rhodamine B in non-HBD solvents such as dimethyl sulfoxide,



λ_{max} = 543 nm (in EtOH)

(26a) *(26b)*

N,*N*-dimethylformamide, 1,4-dioxane, pyridine, and hexamethylphosphoric triamide are entirely colourless, indicating complete conversion into the inner lactone *(26b)*. Protic

solvents stabilize the zwitterion and shift the equilibrium toward the highly coloured zwitterionic form (concentration of *(26a)* in cmol/mol): $(CH_3)_3COH$ (1.6) < *n*-C_3H_7OH (65.2) < C_2H_5OH (70.6) < H_2O (81.5) < $HCONH_2$ (88.5) < CH_3OH (89.2) < CF_3CH_2OH (94.6) [221]. The conversion of *(26b)* to *(26a)* is sensitive to the presence of hydroxy groups to such an extent that a white piece of cellulose turns red on contact with a colourless solution of *(26b)* [175]. Addition of acids to solutions of rhodamine B in either non-HBD or protic solvents produces an intensely coloured cation ($\lambda_{max} = 553$ nm in ethanol [222]) by protonation of the carboxylate group of *(26a)* [221, 222]. The application of increasing external pressure to alcoholic solutions of rhodamine B shifts its equilibrium in favour of the zwitterion *(26a)*, which causes a larger "electrostriction" of the surrounding solvation shell than lactone *(26b)* (see Section 5.5.11) [269]. Rational design of the rhodamine B structure by fine tuning of the substituents in the 3,6-positions of the xanthylium ring can lead to compounds that form the coloured form only under certain conditions [270].

Analogous solvent-dependent intramolecular Lewis acid/base equilibria between lactonic and zwitterionic forms have also been found for the xanthene dyes fluorescein and eosin [176]. Structurally related colourless 3,3-diarylphthalides form highly coloured triarylcarbenium ions by interaction with acidic compounds such as phenols. This acid/base reaction has been used to design optochemical sensors suitable for the detection of vapours of polar solvents in the air or in waste water [271].

The colourless spiropyran *(27b)* is another important example of an intramolecular Lewis acid/base equilibrium. In solution, it is in equilibrium with the coloured zwitterionic species *(27a)* [99].



λ = 532 nm (in EtOH) λ = 330 nm (in EtOH)

(27a) *(27b)*

The thermodynamic data presented in Table 4-6 reveal that the change in equilibrium constant is in the direction anticipated, with more ionization occurring as the

Table 4-6. Thermodynamic data for the intramolecular Lewis acid/base reaction *(27a)* ⇌ *(27b)* at 25 °C [99a].

Solvents	$K \cdot 10^{5}$ [a]	ΔG° / (kJ · mol^{-1})	ΔH° / (kJ · mol^{-1})	$T \cdot \Delta S^\circ$ / (kJ · mol^{-1})
Benzene	4.1	25.1	18.4	−6.7
Trichloromethane	9.8	22.6	2.2	−20.0
Acetone	134	16.3	7.5	−8.8
Ethanol	843	11.7	8.8	−3.0

[a] $K = [(27a)]/[(27b)]$.

solvent becomes more polar. Accordingly, the Gibbs energy change decreases monotonically with increasing solvent polarity.

The negative entropy changes observed in all solvents are a result of an ordering of solvent molecules in the environment of the zwitterionic form. Since polar solvents are *per se* more structured than apolar solvents, proportionally less negative entropy changes are obtained in more polar solvents such as ethanol. The rate of the spiropyran/merocyanine interconversion is also solvent-dependent, as is the position of the visible absorption band of *(27a)*, which, as is typical for a merocyanine, exhibits a pronounced negative solvatochromism (see Section 6.2) [99c, 99d].

A further illustrative example is the *inter*molecular Lewis acid/base reaction between tropylium and isothiocyanate ions via tight ion pairs *(28a)* to give 7-isothiocyanatocycloheptatriene *(28b)* [177].



(28a) *(28b)*

In solution, both the ionic and covalent forms of tropylium isothiocyanate have been directly observed by low-temperature ^{13}C and ^{1}H NMR spectroscopy. In deuteriotrichloromethane and in diethyl ether, the covalent form *(28b)* is exclusively present below −10 °C. With increasing temperature and with increasing solvent polarity (addition of CD_3CN to the solution of *(28b)* in $CDCl_3$), the relative concentration of the ionic *(28a)* increases. In pure acetonitrile, the ionic form *(28a)* dominates. Evidence that the ionic form is a tight ion pair in these solvents is given by NMR and UV/Vis spectra (*e.g.* the occurrence of a charge-transfer absorption between isothiocyanate and tropylium ion). Accordingly, the Gibbs energy of activation for the random migration of the isothiocyanato group around the cycloheptatriene ring decreases with increasing solvent polarity [177]. That is, the stabilization of the ionic *(28a)* by polar solvents corresponds to a similar stabilization of the preceding dipolar activated complex.

Analogous results have been obtained for the Lewis acid/base equilibrium between ionic tropylium azide and covalent 7-azidocycloheptatriene [178]. Again, in less polar solvents such as deuterio-trichloromethane and even $[D_6]$acetone, no ionization to give the tropylium and azide ions could be detected. Dipolar liquid sulfur dioxide, however, induces complete ionization at low temperature (−70 °C).

An interesting example of a Lewis acid/base reaction between neutral reactants is the formation of tris(*n*-butyl)phosphonium-dithiocarboxylate, $(n\text{-}Bu)_3P^+—CS_2^-$, from tris(*n*-butyl)phosphane and carbon disulfide in solution. As expected, this equilibrium is strongly shifted in favour of the dipolar zwitterion with increasing solvent polarity (diethyl ether → dimethyl sulfoxide) [272, 273].

The examples mentioned above are characterized by heterolysis of C—O, C—N, or C—P bonds. Finally, a solvent-dependent Lewis acid/base reaction between carbocations and carbanions, produced by heterolysis of a weak C—C bond, is presented (*cf.* also Section 2.6).

A large variety of different combinations of charge-delocalized carbenium ions with carbanions has been investigated in order to find a well-balanced equilibrium

mixture of free ions or ion-pairs and a neutral covalent product with clean C—C bond formation in solution [179]. The equilibrium between (4-nitrophenyl)malononitrile anions and trianisylmethyl cations [179] or triphenylcyclopropenium cations [180] was finally found to be the most useful. The Lewis acid/base equilibrium between the (4-nitrophenyl)malononitrile anion and the triphenylcyclopropenium cation has already been discussed in Section 2.6 as an example of a solvent-dependent ionization reaction. The first-mentioned reaction can be represented as:



(29a) *(29b)*

A [D_2]dichloromethane solution at room temperature contains 70 cmol/mol ionic *(29a)* and 30 cmol/mol covalent *(29b)*. With decreasing solvent polarity, the equilibrium is shifted toward the right-hand side (concentration of *(29b)* in cmol/mol): CD_2Cl_2 (30) < CD_3CN (55) < CD_3COCD_3 (85) < $(CD_2)_4O$ (100). In [D_8]tetrahydrofuran solution, only the covalent *(29b)* is present [179].

4.4.3 Solvent Effects on Conformational Equilibria [81–83, 181–184]

Changing the medium has a particular effect on various conformational and rotational equilibria [83, 181–184]. Because the Gibbs energy differences between conformational isomers are almost always very small (*ca.* 0 . . . 13 kJ/mol) and the solvation enthalpies of dipolar solutes are at least as large and often much larger than this, the medium can affect conformational equilibria very considerably. It is often found that one conformer or rotamer is predominant in one medium but not in another. This has led to the long-established rule that the conformer (rotamer) of higher dipole moment is more favored in media of high relative permittivity [83].

For example, the standard molar Gibbs energy for the *rotational* equilibrium *(30a)* ⇌ *(30b)* of chloroacetaldehyde is strongly solvent-dependent, as shown in Table 4-7 [87].

(30a) *(30b)*

Inspection of Table 4-7 reveals a substantial increase in the more dipolar rotamer *(30b)* as the polarity of the solvent increases. In saturated hydrocarbon solvents (the

Table 4-7. Solvent dependence of the relative rotamer population (mole fraction of *(30b)*) and the standard molar Gibbs energy differences between rotamers of chloroacetaldehyde at 36 °C [87].

Solvents	$\frac{x\ (30b)}{(\mathrm{cmol \cdot mol^{-1}})}$	$\frac{\Delta G^\circ}{(\mathrm{kJ \cdot mol^{-1}})}$ for *(30a)* ⇌ *(30b)*
trans-Decalin	44	−1.26
Cyclohexane	45	−1.30
Tetrachloromethane	47	−1.46
Trichloromethane	55	−2.34
Benzene	58	−2.68
Dichloromethane	61	−2.97
Acetone	72	−4.18
Acetonitrile	76	−4.60
N,*N*-Dimethylformamide	79	−5.23
Dimethyl sulfoxide	84	−6.07
Formamide	85	−6.28

least polar solvents used), *(30b)* is only favored by *ca.* 1.3 kJ/mol, whereas in formamide (the most polar solvent used) it is favored by *ca.* 6 kJ/mol. In view of the higher dipole moment of *(30b)* over *(30a)*, this appears reasonable [87].

A second well-studied example of a solvent-dependent *rotational* equilibrium is that of 1,2-dichloroethane, which exists preferentially in the apolar *trans* (or antiperiplanar) form ($\mu = 0$) and the dipolar *gauche* (or synclinal) form ($\mu \approx 2.6$ D). According to thorough experimental measurements and theoretical calculations [274–279], the *trans* form is the more stable conformer in the gas phase (ca. 79%). In solution, the population of the *trans* conformer decreases with increasing solvent polarity (ca. 26% in acetonitrile) in favour of the more dipolar *gauche* conformer, as expected. Interestingly, even hydrocarbons [276] and noble gases as cryogenic solvents (liquefied Xe, Kr, and Ar) [277] shift this equilibrium slightly in favour of the *gauche* conformer because of polarization and weak dipole/induced dipole solute/solvent interactions. Thus, noble gas solutions do not behave as "pseudo gas phase" model solutions, as is often assumed; they can significantly shift rotational equilibria if the conformers have different dipolarities, as shown in this case [277].

Another remarkable example is the medium effect on the rotational equilibrium of ethoxycarbonylmethylene triphenylphosphorane *(31a)* ⇌ *(31b)*. As the polarity of the solvent increases, the equilibrium shifts in the direction of the *s-trans*-isomer *(31b)*, as shown by the equilibrium constants presented in Table 4-8.

In the *s-cis* rotamer *(31a)* the $P^{\oplus}/O^{\ominus}$ attraction is maximized. Increasing relative permittivity of the solvent reduces the attraction of the opposite charges, resulting in an increase in the amount of *s-trans* rotamer *(31b)*. An exception is found with trichloromethane as solvent ($\varepsilon_r = 4.9$). Here, the *s-trans* isomer is stabilized more effectively than in the more polar acetonitrile ($\varepsilon_r = 35.9$) and nitromethane ($\varepsilon_r = 35.9$). This is obviously due to the fact that trichloromethane associates with the negative oxygen atom through hydrogen bonding. Addition of a HBD solvent such as methanol to the trichloromethane solution enhances the *s-cis* → *s-trans* conversion to an even greater extent.

Table 4-8. Equilibrium constants for the *s-cis*/*s-trans*-isomerization reaction *(31a)* ⇌ *(31b)* of ethoxycarbonylmethylene triphenylphosphorane in various solvents at −10 . . . 0 °C [95].

Solvents (deuterated)	$K = [s\text{-}cis]/[s\text{-}trans]$
Tetrachloromethane	6.2
Benzene	5.9
Nitromethane	2.4
Acetonitrile	2.3
Trichloromethane	1.8
Trichloromethane/Methanol (5:1)	0.83
Trichloromethane/LiBr	0.57



(31a) *(31b)*

Addition of the extremely polar lithium bromide favours the formation of the *s-trans* isomer even more [95]. Analogous results were obtained with formylbenzylidene triphenylphosphorane [96].

In this context, the solvent influence on the C—N rotational barrier in *N*,*N*-dimethylformamide, $Me_2N—CH{=}O \leftrightarrow Me_2N^+{=}CH—O^-$, is noteworthy [280]. For this rotation, the Gibbs free energy of activation in the gas phase ($\Delta G^{\neq} = 81$ kJ/mol) is much smaller than in polar HBD solvents such as water ($\Delta G^{\neq} = 92$ kJ/mol). Thus, the rate of amide bond rotation decreases as the polarity and the HBD ability of the solvent increases. This can be attributed to the change in dipole moment on rotation, whereby a polar solvent stabilizes the ground state with the higher dipole moment ($\mu = 3.8$ D) in preference to the less dipolar activated complex [280].

Further well-studied examples of solvent-dependent rotational equilibria have been those of furfural [185], the *N*,*N*-dimethylamides of furoic and thenoic acids [186], benzil monoimines [187], and methyl 2-, 3-, and 4-fluorobenzoates [188].

Alicyclic compounds exhibiting *conformational* isomerism are also subject to considerable medium effects [80–83, 182, 184].

One interesting example is (+)-*trans*-2-chloro-5-methylcyclohexanone *(32)*. The sign of the Cotton effect in its ORD spectrum is reversed on transferring it from water to *n*-heptane (molar optical rotation at $\lambda = 330$ nm and 25 °C; $[\Phi]_D = +382$ and -1486, respectively) [189]. This can be ascribed to a diaxial/diequatorial conformational equilibrium *(32a)* ⇌ *(32b)*, which lies more to the left-hand side in *n*-heptane than in water.



(32a) *(32b)*

Due to the nearly parallel C═O and C—Cl dipole vectors in *(32b)*, the *ee* isomer must have the larger net dipole moment and is therefore better solvated in polar solvents. In nonpolar solvents, the electrostatic repulsion between the two equatorial C═O and C—Cl dipoles in *(32b)* is unfavourable, and the molecule escapes from this situation to give the *aa* isomer *(32a)*, even at the cost of nonbonded axial repulsions [189].

The majority of results obtained for other cyclohexanone derivatives, such as 4-methoxycyclohexanone [94a], a bridged 4-oxacyclohexanone [94b], 2- and 4-halocyclohexanones [190], 2-chloro- [191] and 2-bromocyclohexanone [281] have been similar. The conformational Gibbs energy difference $\Delta G^\circ(e \rightarrow a)$ for 4-methoxycyclohexanone has been determined in thirty-four solvents. It shows a marked sensitivity to solvent change, the axial conformer being the more stable in all but the most dipolar HBD solvents [94a]. It is interesting to note that although the equatorial conformer is the least dipolar one, it is stabilized by polar solvents more effectively than the axial isomer. This is due to the larger quadrupole moment of the equatorial conformer [197].

In the case of diaxial/diequatorial equilibria of *trans*-1,2-dihalocyclohexanes (with and without a 4-*tert*-butyl group), ΔG° ($aa \rightarrow ee$) also shows a pronounced dependence on the medium, varying from about 4 kJ/mol in apolar solvents to about -2 kJ/mol in polar solvents [90–94, 192–194]. This is mainly due to the very different dipole moments of the two conformers. In general, the more dipolar diequatorial isomer is favoured in polar solvents.

Another remarkable example is provided by the observation of the strong influence of solvents on the conformation of phencyclidine *(33)*, a drug developed as an anaesthetic, but later withdrawn because of its psychotomimetic effects [195].



(33a) *(33b)*

Table 4-9. Solvent dependence of standard molar Gibbs energy differences between *cis*/*trans*-isomers of 2-isopropyl-5-methoxy-1,3-dioxane at 25 °C [89].

Solvents	$\Delta G^{\circ}_{OCH_3}/(kJ \cdot mol^{-1})$ for *(34a)* ⇌ *(34b)*[a]
n-Hexane	−4.44
Cyclohexane	−4.31
Tetrachloromethane	−3.77
1,3,5-Trimethylbenzene	−3.64
tert-Butylbenzene	−3.47
Diethyl ether	−3.47
Toluene	−2.97
Tetrahydrofuran	−2.72
Benzene	−2.47
1,1,1-Trichloroethane	−2.43
Acetone	−1.42
Nitrobenzene	−0.84
Deuterio-trichloromethane	−0.79
Trichloromethane	−0.67
Dichloromethane	−0.42
Methanol	−0.13
Acetonitrile	+0.04

[a] $K = [trans]/[cis]$.

In dipolar non-HBD solvents such as $[D_6]$acetone and $[D_3]$acetonitrile, the equilibrium is more to the left-hand side than in the less polar $[D_2]$dichloromethane. Upon transfer from CD_2Cl_2 to an HBD solvent (CD_2Cl_2/CD_3OD, 1:2, cl/L), the equilibrium is shifted substantially to the right-hand side. The ratio *(33b)*/*(33a)* varies from 99:1 in CD_2Cl_2/CD_3OD at −80 °C to 1:1 in CD_3COCD_3/CD_3CN at room temperature. Obviously, conformer *(33b)* with the piperidine ring in the equatorial position is stabilized by hydrogen-bonding in HBD solvents; similar stabilization of the axial piperidine ring in *(33a)* is sterically inhibited. Consequently, it can be expected that the structure of this drug is subject to change on passing through a cell membrane [195].

A thoroughly examined case, namely that of a heterocycloalkane, which involves an axial/equatorial conformational change shall conclude this Section.

The position of the acid-catalyzed equilibrium between *cis*- *(34a)* and *trans*-2-isopropyl-5-methoxy-1,3-dioxane *(34b)* has been determined in seventeen different solvents; *cf.* Table 4-9 [89].



(34a)
axial *cis* isomer
$\mu = 9.5 \cdot 10^{-30}$ C m

(34b)
equatorial *trans* isomer
$\mu = 4.3 \cdot 10^{-30}$ C m

Inspection of Table 4-9 reveals that the axial *cis* isomer *(34a)*, which is the conformer with the higher dipole moment, becomes more favoured as the solvent polarity increases. In the most polar solvent studied, acetonitrile, ΔG° is nearly zero. Benzene, toluene, trichloromethane, dichloromethane, and methanol are seen to behave as more polar solvents than their relative permittivities would lead one to predict. The deviation for trichloromethane was particularly difficult to explain (for a full discussion, see reference [89]). In general, good correlations between $\Delta G^\circ_{OCH_3}$ values and other solvent-dependent phenomena such as absorption maxima of solvatochromic dyes, rate constants of reactions involving dipolar activated complexes, *etc.*, were obtained [89]. Because of this, it was recommended that the solvent scale obtained should be used as an empirical scale of solvent polarity, useful for the prediction of medium effects on other solvent-dependent reaction rates or equilibria [89] (*cf.* Section 7.2).

The so-called *anomeric effect*, *i.e.* that polar substituents X attached to a carbon α to a heteroatom Y (Y = O, N) in a six-membered ring preferentially reside in the axial position, has been shown to be solvent-dependent [82, 83, 217, 282–286]. In general, the position of an anomeric equilibrium shifts in favour of the equatorial anomer with increasing solvent polarity. The anomeric effect is thought to be the result of either molecular orbital interactions, which stabilize the axial conformer, or electrostatic interactions, which destabilize the equatorial conformer [82, 282].

The MO explanation for the anomeric effect considers the n–σ^* overlap between the lone-pair of Y and the vacant σ^* orbital of the C—X bond. This stabilizing interaction is more effective when X is axial and thus the axial conformer is favoured. The electrostatic explanation invokes the destabilizing interaction between the dipole moment of the C—X bond and the dipole moment resulting from the C—Y bond and the lone-pairs of Y. Such dipole/dipole interactions are minimized when X is axial and again the axial conformer is preferred in the gas phase or in nonpolar solvents. It is not so easy to distinguish between the relative importance of each interaction. However, the observation that the axial preference is diminished by increasing solvent polarity is best explained by the electrostatic interaction model [82, 282–284]. The unfavourable electrostatic dipole/dipole repulsion in the equatorial anomer decreases with increasing solvent polarity, and hence the equilibrium shifts towards the equatorial conformer in polar solvents. This solvent-dependent anomeric effect has been particularly well studied with 4,6-dimethyl-2-methoxytetrahydropyran [283, 284] and 2-methoxy-1,3-dimethylhexahydropyrimidine [282].

Another well-known example is D-glucopyranose, the anomeric equilibrium mixture of which is made up of 36% of the axial α anomer and 64% of the equatorial β anomer at 20 °C in water as a polar solvent ($\varepsilon_r = 78$), with $\Delta G(\alpha \rightarrow \beta) = -1.4$ kJ/mol [82]. This is in agreement with free-energy simulations with an empirical force field [285] and quantum-chemical calculations [286], showing that the preference of D-glucopyranose for the β-anomer in water is mainly due to electrostatic solvation and hydrogen-bonding effects, which stabilize the more dipolar β-anomer better than the α-anomer.

Numerous attempts have been made to calculate relative conformer energies in solution, using physical properties of solutes and solvents, in order to derive theoretical procedures or models with predictive ability [83, 88, 182, 188, 190, 192, 196–198, 274–281]. The methods used include quantum-chemical calculations (*e.g.* [198]), statistical

mechanics and molecular dynamics calculations (*e.g.* [182]), direct dipole-dipole methods (*e.g.* [83]), and reaction field methods based on Onsager's theory [199] of dipolar molecules in the condensed phase (*e.g.* [83, 88, 188, 190, 194, 197]). In general, a quantitative description of solvent effects on conformational equilibria can be given on the basis of these methods, except in cases where specific solute/solvent interactions occur.

According to the reaction field method, the electrostatic stabilization of a solute molecule, located in the centre of a spherical cavity, with dipole moment μ and radius r in a solvent modeled as a uniform dielectric with relative permittivity ε_r, can be expressed as in Eq. (4-31) (ε_0 = permittivity of vacuum).

$$\begin{aligned}\Delta G^\circ_{solv} &= G^\circ_{vapour} - G^\circ_{solution} \\ &= -\frac{N_A}{4\pi \cdot \varepsilon_0} \cdot \frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \cdot \frac{\mu^2}{r^3}\end{aligned} \tag{4-31}$$

The difference in the Gibbs energy of solvation, ΔG°_{solv}, for two species in equilibrium, A $\rightleftharpoons$ B, is then given by Eq. (4-32), assuming that they have the same size.

$$\Delta\Delta G^\circ_{solv} = -\frac{N_A}{4\pi \cdot \varepsilon_0} \cdot \frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \cdot \left(\frac{\mu_A^2}{r^3} - \frac{\mu_B^2}{r^3}\right) \tag{4-32}$$

Qualitatively, Eq. (4-32) predicts that the more dipolar isomer will be preferentially stabilized in more polar media. Quantitatively, the expression significantly overestimates the solvent effects obtained experimentally for conformational equilibria [182]. Further modifications are necessary, *e.g.* adjustment for back-polarization of the solute by its own reaction field, inclusion of the effect of the solute's quadrupole moment on the reaction field [197], *etc.* Specific solute-solvent interactions, such as those with HBD solvents, cannot be treated with this reaction field theory. For a more detailed discussion, see references [83, 182].

A purely empirical correlation between the $\Delta\Delta G^\circ_{solv}$ of equilibria such as A $\rightleftharpoons$ B and solvent polarity has been given by the parabolic Eq. (4–33), where $X = (\varepsilon_r - 1)/(2\varepsilon_r + 1)$ and, in the majority of cases, $C = 0.5 = \lim_{\varepsilon_r \to \infty} X$ [196].

$$\Delta\Delta G^\circ_{solv} = A + B \cdot \sqrt{C - X} \tag{4-33}$$

If $X = 0$, the A parameter should be regarded as the extreme $\Delta\Delta G^\circ_{solv}$ value for solvents with infinitely large polarity, and the B parameter could be considered as a measure of the susceptibility of the equilibrium to changes of solvent polarity. Eq. (4-33) has been successfully applied to various conformational and tautomeric equilibria [196].

4.4.4 Solvent Effects on *cis/trans* or *E/Z* Isomerization Equilibria [82, 200]

One of the simplest examples of a *cis/trans* or *E/Z* isomerization equilibrium is represented by *trans- (35a)* and *cis*-1,2-dichloroethene *(35b)*. In the gas phase at 185 °C,

the equilibrium mixture contains 63.5 cmol/mol of the thermodynamically more stable *cis*-isomer *(35b)* [201].



(35a) $\mu = 0$ C m

(35b) $\mu = 6.3 \cdot 10^{-30}$ C m

Due to the high activation barrier for *cis*/*trans* isomerization reactions at carbon-carbon double bonds (*ca.* 260–270 kJ/mol [82]), it is often impossible to measure directly the non-catalyzed thermal equilibration reaction in solution. For 1,2-dichloroethene, however, the relative stability of its *cis* and *trans* isomers in various solvents has been determined by means of calorimetric measurements of heats of solution [202]. Surprisingly, these measurements show a quite similar solvent effect on both diastereomers, even though the *cis* isomer is a dipolar molecule and the *trans* isomer is not. Therefore, the position of this *cis*/*trans* equilibrium should not be very solvent-dependent.

One reason for this at first sight unexpected result is the fact that probably 70 . . . 90% of the solute/solvent interaction term is caused by London dispersion forces, which are more or less equal for the *cis* and *trans* isomers. Another important reason is that one has to take into account higher electric moments: the *trans* isomer has a quadrupole moment, and the *cis* isomer also has moments of a higher order than two. Calculations of solute/solvent interactions of both diastereomers using a reaction field model led to the conclusion that the quadrupolar contribution of the *trans* isomer is comparable to the dipolar contribution of the *cis* isomer. It has been pointed out that the neglect of solute/solvent interactions implying higher electric moments than the dipole moment can lead to completely false conclusions [202].

Unlike those for *cis*/*trans* isomers, the activation barriers separating *s-cis* and *s-trans* isomers are usually small (*ca.* 40 . . . 50 kJ/mol); the Gibbs energy differences for *s-cis* and *s-trans* isomers are also small (*ca.* 4 . . . 20 kJ/mol). For example, a more easily measurable *cis*/*trans* isomerization reaction can be carried out with 3-*tert*-butylaminopropenal *(36)* [203].



(36a) ***E-s-E*** form

(36b) ***Z-s-Z*** form

According to its ^{13}C and ^{1}H NMR spectra, this vinylogous amide exists as the *E-s-E* form in polar solvents such as [D_4]methanol, and as a mixture of *Z-s-Z* and *E-s-E* isomers in nonpolar solvents such as deuterio-trichloromethane (30 cmol/mol *(36a)* and 70 cmol/mol *(36b)*). As expected, the more dipolar *E-s-E* form is stabilized in polar solvents (dipole moment of the related *E-s-E* 3-dimethylaminopropenal $21 \cdot 10^{-30}$ Cm).

Experimental and theoretical studies on the solvent influence on molecular geometries and *cis/trans* isomerization processes of other so-called push-pull ethenes, $R_2N—CH=CH—A \leftrightarrow R_2N^+=CH—CH=A^-$ (with $A = NO_2$, CHO, CN, *etc.*), have been collected in references [287, 288]. The barriers to isomerization about the C=C bonds of these acceptor-substituted enamines are considerably smaller than those for simple ethenes such as 2-butene ($E_a = 259$ kJ/mol [82]), owing to a significant contribution of the mesomeric zwitterionic structure to the electronic ground state. Increasing solvent polarity increases the contribution of this dipolar mesomeric structure, and hence leads to a decrease in the barrier to C=C isomerization and a simultaneous increase in the barrier to rotation about the C—N bond [288]. The calculated sequence of solvent stabilization for such acceptor-substituted enamines is: (activated complex for C=C rotation) $\gg E$ form $> Z$ form $>$ (activated complex for C—N rotation). Obviously, the activated complex for isomerization about the C=C bond corresponds to a full zwitterion with maximal solvent stabilization [288].

Finally, a *cis/trans* isomerization process of a heterocycloalkane is presented. The *cis/trans* isomer ratios of 2,3-dibenzoyl-1-benzylaziridine *(37)* have been determined by means of ^{1}H NMR measurements [97] ($R = CH_2C_6H_5$).



(37a) $\mu = 8.7 \cdot 10^{-30}$ C m

(37b) $\mu = 15.8 \cdot 10^{-30}$ C m

This isomerization reaction passes through *cis/trans*-isomeric open-chain azomethine-ylides, which arise from conrotatory ring-opening at the C—C bond [98]. The equilibrium constants range from 5.25 in dimethyl sulfoxide to 0.32 in *tert*-butanol, as shown in Table 4-10, and approximately parallel the polarities of the solvents used. The more dipolar *cis*-aziridine *(37b)* is the more stable isomer in polar solvents; the reverse is true for the less dipolar *trans*-aziridine *(37a)*. The isomer ratio is 1.63 in methanol (corresponding to 62 cmol/mol *cis* isomer); this is in close agreement with the separately determined solubility ratio of 1.50 (corresponding to 60 cmol/mol *cis* isomer). This close agreement of the ratios of the solubilities of the two isomers with their equi-

Table 4-10. Equilibrium constants and mole fractions of *(37b)* in various solvents for the base-catalyzed *trans/cis*-isomerization *(37a)* $\rightleftharpoons$ *(37b)* of 2,3-dibenzoyl-1-benzylaziridine at 33 °C [97].

Solvents	x *(37b)*/(cmol · mol^{-1})	$K = [cis]/[trans]$
tert-Butanol	24	0.32
Ethanol	45	0.82
Methanol	62	1.63
Dimethyl sulfoxide	84	5.25

librium constants is a modern example of a long recognized phenomenon [37]; *cf.* Table 4-3 and Eq. (4-24).

4.4.5 Solvent Effects on Valence Isomerization Equilibria [84]

Valence isomerization reactions interconvert so-called valence isomers by simple reorganization of some of the bonding electrons, without any atom migration. Since both valence isomers may have different structures and thus different physical properties (*e.g.* different dipole moments) it is to be expected that solvents should influence the equilibrium between non-degenerate isomers.

For example, ^{1}H NMR measurements showed that the azido/tetrazole equilibrium of thiazolo[2,3-*e*]tetrazole *(38b)* is considerably affected by the medium [100].



(38a) *(38b)*

Whereas in the gas phase and in nonpolar solvents such as tetrachloromethane and benzene, the 2-azidothiazole *(38a)* is the more stable isomer, in polar solvents such as dimethyl sulfoxide and hexamethylphosphoric triamide the bicyclic valence isomer *(38b)* is the dominant species [100]. This result is in line with the fact that the dipole moment of phenyl azide ($\mu = 5.2 \cdot 10^{-30}$ Cm) is smaller than that of cyclic 1,2,3-benzotriazole ($\mu = 13.7 \cdot 10^{-30}$ Cm). Similar results have been obtained for the valence isomers of 3-azidopyrazine-1-oxide [204].

Another remarkable example is 2,4,6-tri-*tert*-butyl-*N*-thiosulfinylaniline *(39a)*, which is in equilibrium with its valence isomer *(39b)* in solution, while in the solid state only *(39b)* exists [205]. The equilibrium ratio *(39b)*/*(39a)* is subject to a considerable

(39a) *(39b)*

$\mu \approx 5 \cdot 10^{-30}$ C m $\mu \approx 10 \cdot 10^{-30}$ C m

solvent effect: the mole fraction of *(39b)* in *n*-hexane solution is 86 cmol/mol and in acetonitrile solution 96 cmol/mol. Polar solvents enhance the preference for the cyclic isomer *(39b)*, mainly due to its larger dipole moment [205]. This is noteworthy because the aromaticity of the benzene ring is destroyed in *(39b)*.

In the case of the oxepin/benzene oxide valence isomerization *(40a)* $\rightleftharpoons$ *(40b)*, it has been found by UV/Vis measurements that, with *iso*-octane as the solvent, only about 30 cmol/mol benzene oxide is present, whereas in water/methanol (85:15, cL/L)

the benzene oxide portion rises to about 90 cmol/mol [101]. Increasing solvent polarity



(40a) *(40b)*

shifts this equilibrium towards the more dipolar benzene oxide isomer, in agreement with quantumchemical MNDO calculations [206]. These calculations have shown that in the gas phase *(40a)* is more stable than *(40b)*, but that *(40b)* is more stabilized in polar media.

The behaviour of the corresponding hexafluoro derivative of *(40b)* in solution is somewhat different: solutions of hexafluorobenzene oxide in nonpolar solvents are stable at room temperature. In acetonitrile or acetone, the hexafluorobenzene oxide rearranges spontaneously to hexafluorocyclohexa-2,4-dienone and not to hexafluorooxepin. However, ^{19}F NMR measurements in chlorobenzene at various low temperatures indicate the existence of a dynamic equilibrium between perfluorobenzene oxide and perfluorooxepin [289].

A further example is the reversible valence isomerization equilibrium between the dipolar 8,8-diformylheptafulvene *(41a)* and the less dipolar 8a*H*-cyclohepta[*b*]furan-3-carbaldehyde *(41b)* [212].



(41a) *(41b)*

$\mu = 17.5 \cdot 10^{-30}$ C m (5.2 D)

In deuterium oxide, only *(41a)* could be detected by ^1H NMR spectroscopy. In less polar solvents, such as perdeuterated acetonitrile, acetone, and benzene, the mole fraction of *(41a)* decreases to $x = 69$, 58, and 54 cmol/mol, respectively. Eventually, in tetrachloromethane, the less dipolar *(41b)* predominates; $x(41b) = 63$ cmol/mol [212].

Finally, the allylcarbinyl/cyclopropylcarbinyl anion rearrangement *(42a)* $\rightleftharpoons$ *(42b)* should be mentioned as a striking example of a solvent-dependent valence isomerization [207].

The deep-red lithium cyclopropyl-diphenylmethanide *(42b)*, which is stable in tetrahydrofuran solution, opens completely to the colourless lithium (3,3-diphenylprop-2-enyl)methanide in diethyl ether. The *retro* rearrangement to *(42b)* can be achieved simply by adding tetrahydrofuran to the solution of *(42a)* in diethyl ether [207]. Obviously, tetrahydrofuran, which is the better cation-solvating EPD solvent, makes the carbon-lithium bond in *(42a)* more ionic, thus favouring the ring closure to *(42b)*, which has a more delocalized negative charge.



(42a) (42b)

A closely related example, the equilibrium between tight and solvent-separated ion pairs of lithium 10-phenylnonafulvene-10-oxide, has already been given in Section 2.6 (formulas *(2b)* and *(3b)*). Depending on the solvent-influenced association with the lithium cation, the anion exists either as the aromatic benzoyl [9]annulene anion or as the olefinic nonafulvene oxide anion [208].

The first metallaquinone (*i.e.* a compound in which one of the 1,4-benzoquinone oxygen atoms is replaced by ruthenium) has recently been synthesized and exhibits another interesting type of solvent-dependent valence isomerization [290]. This metallaquinone interconverts between a quinoid-Ru(0) carbene form in nonpolar solvents (reddish-orange solution in benzene) and a zwitterionic Ru(II)-phenolate form in polar solvents (yellow solution in methanol).

4.4.6 Solvent Effects on Electron-Transfer Equilibria

Examples of solvent effects on electron-transfer equilibria between organic species (*i.e.* redox and disproportionation reactions) are rather scarce.

A nice example is the disproportionation reaction of the 1-ethyl-4-(methoxycarbonyl)pyridinyl radical *(43)*, which leads to the ion pair *(43a)/(43b)* [215, 216].



(43) (43a) (43b)

Solvent	$HCON(CH_3)_2$	CH_3CN	C_2H_5OH	$HCONH_2$	H_2O
$K=\frac{[\text{ion pair}]}{[\text{radical}]}$	$4\cdot10^{-13}$	10^{-13}	10^{-8}	10^{-6}	$5\cdot10^{-5}$

As expected, an increase in solvent polarity shifts this equilibrium from the left to the right-hand side due to the better solvation of the ion pair as compared to the neutral radical [215].

From equilibrium constants for transfer from water to vapour, determined by dynamic vapour pressure measurements, the hydrophilic character of 1,4-benzoquinone and 1,4-hydroquinone has been estimated [227]. 1,4-Benzoquinone is about 3.2 orders of magnitude less strongly solvated by water than is 1,4-hydroquinone. Because 1,4-hydroquinone is so much more strongly solvated than 1,4-benzoquinone, its reducing

power in water is less by *ca.* 18 kJ/mol (0.2 V) than it would be in a medium of unit relative permittivity. Therefore, the redox potentials of biologically important hydroquinone/quinone systems should be strongly affected by the surrounding medium, particularly if the corresponding electron-transfer reaction is part of an electron transport chain embedded in mitochondrial inner membranes [227].

The redox properties of tetraphenylporphyrin have been studied in dimethyl sulfoxide (DMSO) and more polar water/DMSO mixtures. Increasing solvent polarity (*i.e.* increasing water content) shifts the reduction potentials to more positive values and makes the reduction step more favourable, since a better solvated anion is formed from a neutral species [291].

An interesting solvent-dependent, reversible electron-transfer equilibrium between a pair of resonance-stabilized carbocations and carbanions and the corresponding carbon radicals has been described by Arnett *et al.* [229]. Addition of bis(4-dimethyl-



aminophenyl) phenylcarbenium tetrafluoroborate (malachite green) to sodium tris(4-nitrophenyl) methanide leads, in tetrahydrofuran, to the corresponding triphenylmethyl radicals by single-electron transfer, whereas in sulfolane as solvent only traces of the radicals could be detected by ESR spectroscopy. That is, in the more polar solvent sulfolane, both the carbocation and carbanion are stabilized by electrostatic ion/solvent interactions, whereas in the less polar tetrahydrofuran single-electron transfer from the carbanion to the carbocation occurs readily to produce trityl radicals. Dilution of the radical solution in tetrahydrofuran with sulfolane leads to a sharp decrease of the ESR signal and to an increase of the carbanion absorption in the UV/Vis spectrum, in accordance with a reversible electron-transfer equilibrium [229].

Another hydrocarbon salt, composed of the tri(cyclopropyl)cyclopropenylium cation and Kuhn's anion, which can exist in all three types of elementary organic species (*i.e.* as an ionic, radical, and covalent compound) in a solution equilibrium, depending on the solvent, has already been mentioned in Section 2.6 [292].

Comprehensive reviews on the solvent influence on electron- and charge-transfer reactions between organic electron-donor and acceptor compounds, resp., [230], as well as on the redox properties of predominately inorganic redox couples [293] can be found in the literature.

4.4.7 Solvent Effects on Host/Guest Complexation Equilibria [294–299]

The formation of supramolecular host/guest complexes between two or more neutral or charged species, which are held together only by comparatively weak, noncovalent intermolecular forces, has been thoroughly investigated during the last decades, not least because of its paramount importance with regard to the chemical molecular recognition found in biochemical processes [294–299]. For example, the reversible formation of such complexes is responsible for the selective recognition and high catalytic activity of enzymes, usually observed in aqueous media. However, solvents other than water are known to have a profound influence on the stabilities of all kinds of host/guest complexes [300–306].

Host/guest complex formation in solution can be simply described by the following equation,

$$(\text{Host})_S + (\text{Guest})_S \rightleftharpoons (\text{Host/Guest complex})_S + S$$

where the first term represents solvated hosts such as acyclic and cyclic polyethers (*e.g.* podands, coronands, cryptands; see Section 5.5.5), cyclophanes, calixarenes, cyclodextrins, *etc.*, and the second term represents guests such as solvated metal ions, organic ions, neutral dipolar molecules, dyes, *etc*. The standard Gibbs free energy for host/guest complexation depends on the change in surface area as the two solvent cavities containing the host and guest collapse into a single solvent cavity containing the complex, which is usually formed in a 1:1 stoichiometry. This reaction is associated with (a) some desolvation of both the host and guest molecules in order for them to come closer together, and (b) simultaneous solvation of the newly formed complex, which is presumably accompanied by the release of solvent molecules S from the host and guest solvation shells into the bulk solvent, which then experience solvent/solvent interactions. Accordingly, by means of a thermodynamic cycle, the Gibbs free energy of complex formation in solution, $\Delta G(\text{host} + \text{guest})_S$, is given by Eq. (4-34),

$$\Delta G(\text{host} + \text{guest})_S = \Delta G(\text{host} + \text{guest})_G - \Delta G(\text{host})_S - \Delta G(\text{guest})_S + \Delta G(\text{host/guest complex})_S \quad (4\text{-}34)$$

and thus depends on the Gibbs free energies of (from left to right) complex formation in the gas phase, desolvation of the host and guest, as well as of solvation of the host/guest complex.

In addition, the uncomplexed, usually rather large guest molecules may have different conformations in different solvents (*e.g.* uncomplexed crown ethers may be present in oxygen-in and oxygen-out conformations depending on the solvent [295]). According to Cram [294], the principle of preorganization of the binding sites prior to complexation includes not only geometrical changes of the host, but also the minimization of solvation energy changes during complexation.

Overall, the solvent influence on such host/guest equilibria is obviously a rather complex phenomenon and has to be carefully analysed for each individual solvent-dependent complex formation. A few representative examples of solvent-dependent host/guest complexation reactions are given in the following.

The 1:1 host/guest complexation between the cyclic polyether [18]crown-6 and potassium thiocyanate [*cf.* Eq. (5-128) in Section 5.5.5] has been studied calorimetrically in fourteen solvents of different polarity at 25 °C [300]. The complex formation (or association) constant K_a increases by a factor of $>10^4$ (!) on going from water to propylene carbonate (PC), with K_a increasing in the solvent order $H_2O < HMPT < DMSO < DMF < Me_2CHOH < MeCN < Me_2C{=}O < MeOH < PC$. Surprisingly, a satisfactory linear correlation was found between lg K_a and the standard molar Gibbs energy of transfer of potassium cations from water to the solvents given in this sequence, $\Delta G_t^\circ(K^+, W \rightarrow S)$; *cf.* Table 2-9 in Section 2.3. Thus, the position of the complexation equilibrium between the neutral crown ether and a cationic guest is mainly determined by the desolvation energy of the metal cation, $\Delta G(\text{host})_S$ in Eq. (4-34), which is largest for the good cation solvators water, HMPT, DMSO, and DMF. This finding is in line with the much larger solvation energies obtained for charged species as compared with those of electroneutral ligands. It is also consistent with the perfect fit between the hole size of the coronand ($d = 260$ pm) and the guest diameter ($d = 266$ pm) [295]; there is practically no room left for solvent molecules inside the complex. An analysis of the thermodynamic parameters shows that the host/guest complex formation is accompanied by a considerable decrease in entropy in all solvents [300]. Obviously, the entropy increase associated with the host desolvation is more than counterbalanced by the entropy decrease caused by the host/guest complexation. The release of solvent molecules from the guest solvation shell into the bulk of highly structured solvents seems to be less important in this respect.

Analogous dramatic solvent-dependent host/guest complexation constants have been found for the inclusion of silver(I) cations (which are similar in size to K^+, but softer) into the cyclic polyether 1,10-diaza-[18]crown-6 [301]. The K_a values increase by nearly eight orders of magnitude in the solvent sequence $DMSO < H_2O \approx CH_3CN < DMF < MeOH < Me_2C{=}O < CH_3NO_2 < PC$. They are smallest in EPD solvents such as DMSO, H_2O, and CH_3CN, which are known to solvate Ag^+ ions strongly. Again, the major factor in controlling the variation in the complex formation constants K_a is the desolvation of the silver(I) cations. The HBD solvents water and methanol form hydrogen bonds to the basic nitrogen atoms of the aza-crown ether, thereby leading to somewhat lower K_a values than one might expect [301].

Not only metal cations but also organic cations such as arenediazonium ions can be complexed by polyethers. Benzenediazonium tetrafluoroborate forms 1:1 host/guest complexes with acyclic (*e.g.* pentaglyme) and cyclic polyethers (*e.g.* crown ethers) [302]. This solvent-dependent complexation increases the stability of the benzenediazonium

ion in solution and determines the first-order rate constant and the outcome of its thermal decomposition, with a change from the heterolytic (in the absence of complexing agents) to the homolytic dediazoniation mechanism; *cf.* Eq. (5-149) in Section 5.5.7.

Other useful hosts for a great variety of guests are cyclic oligosaccharides such as α-cyclodextrin, composed of six α-(1 → 4)-linked D-glucopyranose units forming an *endo*-hydrophobic/*exo*-hydrophilic host with a cavity of diameter 47–60 pm [295]. The binding constants for the 1:1 host/guest complexation between α-cyclodextrin and the azo dye methyl orange ($Me_2N—C_6H_4—N{=}N—C_6H_4—SO_3H$) have been measured in water ($K_a/L \cdot mol^{-1} = 682$) and in seven water/solvent mixtures ($K_a \ll 682$), demonstrating the destabilizing effect of organic solvents on complex formation. In water, there is a hydrophobic driving force for the inclusion of the organic dye molecule into the *endo*-hydrophobic cavity of α-cyclodextrin, which is reduced upon addition of organic solvents such as dimethyl sulfoxide, acetonitrile, or acetone; this is accompanied by a decreased solvent stabilization of the *exo*-hydrophilic host/guest complex by these organic solvents as compared to water.

Cyclophanes, having at least two benzenoid rings disubstituted by closed chains of carbon atoms (usually methylene groups), constitute another class of excellent hosts with apolar hydrophobic cavities [295, 296, 299, 304]. A particularly well-studied example is the solvent-dependent 1:1 complex formation between a macrobicyclic cyclophane with two tertiary cryptand nitrogen atoms and pyrene (as well as other polycyclic arenes), as shown in the following equation [305]:



The inclusion of planar pyrene into the macrobicyclic cyclophane (which displays solubility in nonpolar *and* polar solvents) has been studied in a total of eighteen solvents of differing polarities as a kind of model reaction for molecular recognition, mimicking both the highly polar biological aqueous phases and the less polar lipid membranes. On going from water, the most polar, to carbon disulfide, the least polar of the eighteen solvents, the Gibbs free energy of host/guest complexation decreases from $\Delta G^\circ = -39.4$ kJ/mol to $\Delta G^\circ = -5.4$ kJ/mol ($\Delta\Delta G^\circ = 34.0$ kJ/mol), which corresponds to a change in K_a ($L \cdot mol^{-1}$) from 6.0×10^6 (H_2O) to 9×10^0 (CS_2) at 30 °C [305]. Calorimetric measurements of the enthalpies of formation of the host/pyrene complex show that the complex formation is enthalpy-driven in all solvents studied, with mostly unfavourable complexation entropies (*i.e.* entropy decrease).

A linear correlation has been found between the solvent-dependent ΔG° values and the empirical parameter of solvent polarity, $E_T(30)$ (see Section 7.4). Thus, the host/guest binding strength increases steadily on going from nonpolar solvents to water, thus shifting the complexation equilibrium more and more to the right-hand side with increasing solvent polarity.

All solvent molecules are small enough to easily enter and exit the large, highly preorganized host cavity, leading to a complete equilibrium solvation of the host. The host/pyrene complex is stabilized by dispersion interactions and local dipole/induced dipole interactions and, in all solvents, adopts the geometry shown, with exclusion of solvent molecules from the host interior. Consequently, the large differences in binding strength result entirely from solvation effects. Computer simulations have shown that the major contribution to the experimentally observed solvent effect is due to the different free energies of cavitation for water and less polar, less structured solvents. The stronger cohesive interactions of water facilitate the transfer of pyrene from the solution into the cyclophane cavity, whereas the free energy needed for the desolvation of the cyclophane cavity and subsequent pyrene complexation depends only slightly on the nature of the solvent. Molecules of solvents with high cohesive pressure interact more favourably with bulk solvent molecules than with the apolar surfaces of the free host and guest. Therefore, free energy is gained upon the release of surface-solvating solvent molecules into the bulk during the host/guest complexation step. Furthermore, upon complexation, the less favourable dispersion interactions between solvent molecules of low polarizability (*e.g.* water) and the highly polarizable arene surface of the guest molecule pyrene are replaced by the more favourable dispersion interactions between the complementary surfaces of host and guest [299, 305].

Similar results have been reported for the host/guest complex formation between a tetracationic cyclophane host [=cyclobis(paraquat-*p*-phenylene)] and the planar heteroarene indole. The binding constants K_a, determined in six solvents of different polarities, increase with increasing solvent polarity, being *ca.* 100-fold greater in water than in acetone [306]. Further examples of solvent-dependent cyclophane/guest complexation reactions can be found in reference [299].

Solvents can also act as a kind of template and increase the selectivity in reactions leading to more than one product. A more recent example is the synthesis of so-called glycolurils by acylation of cyclic hydrazines (tetrahydro-1,2-diazines) with active tetracarboxylic esters, leading to three diastereomeric *C*-, *S*-, and *W*-shaped products in a solvent-dependent ratio [307]. The *C*-shaped isomer is self-complementary in hydrogen-

bond donor and acceptor sites and readily forms dimeric capsules in solvents that compete poorly for hydrogen bonds. While in the HBA solvent DMSO only monomeric species are found, in [D_6]benzene H-bonded dimers with the encapsulation of benzene as guest molecules are formed ($K_a \approx 3 \cdot 10^{10}$). Interestingly, in solvents that favour the formation of a dimeric capsule, the glycoluril-forming reaction between hydrazine and ester shows a higher selectivity in favour of the *C*-shaped isomer, whereas in DMSO only the statistically expected distribution of *C*-, *S*-, and *W*-shaped isomers is found. Obviously, in the reaction sequence forming the glycolurils, benzene acts as a template, directing the reaction predominantly towards the formation of the *C*-shaped isomer, which forms H-bonded dimers with encapsuled benzene molecules. Thus, in other words, the solvent controls the covalent bond formation through molecular recognition within the monomeric intermediate in the glycoluril synthesis. The intermediate with the best solvated surface, with optimally filled niches, is formed preferentially [307].

Biochemical reactions within or outside of living cells are usually catalysed by macromolecular proteins called *enzymes*, which combine high catalytic power with high specificity, *i.e.* high chemo-, regio-, enantio-, and diastereoselectivity. The rate acceleration and high selectivity of enzymatic reactions stems from a further kind of specific host/guest interaction between the active site of the large enzyme and the smaller substrate, referred to as the 'lock-and-key' interaction by Emil Fischer in 1894 [308]. To this end, the enzyme must have, or must be able to assume (by induced fit), a shape complementary to that of the substrate.

According to the Michaelis-Menten mechanism of enzyme-catalysed reactions in solution, the enzyme E binds reversibly to the substrate S in a pre-equilibrium to yield an intermediate solvated host/guest complex $(E \cdot S)_S$, which then converts the substrate to the solvated products P, according to the following equation:

$$(E)_S + (S)_S \rightleftharpoons (E \cdot S)_S \rightarrow (E)_S + (P)_S$$

In contrast to non-enzymatic chemical reactions, enzymatic reactions occur under rather mild physiological conditions, that is to say at moderate temperatures, reasonable pH values, and in *aqueous* solution. Evolution has taken place in an environment with water as one of the major components. Therefore, biochemical reactions in living cells occur in a dilute aqueous medium.

Surprisingly, it has been shown that many enzymatic reactions can advantageously be carried out in nonaqueous solutions, that is, in carefully selected organic solvents containing little or no added water, sometimes with dramatic changes in enzyme specificity as a function of solvent polarity [309–312].

An illustrative example of a change in *chemoselectivity* is the inversion of substrate specificity of the serine protease *Subtilisin Carlsberg* in the transesterification reaction of ethyl esters of *N*-acetyl-L-serine and *N*-acetyl-L-phenylalanine with 1-propanol, measured in twenty anhydrous organic solvents. The enzyme-catalysed reaction with the serine substrate is strongly favoured in dichloromethane, while the reaction with the phenylalanine substrate is preferred in *t*-butylamine, with a 68-fold change in substrate specificity [313].

An example of a solvent-dependent change in *regioselectivity* is the lipase-catalysed partial transesterification of the diester derived from 2-octyl-1,4-

dihydroxybenzene and butyric acid with 1-butanol, to give butyl butyrate and two isomeric 1,4-dihydroxybenzene monoesters. By two alternative pathways, either the monoester with the hydroxy group *ortho* to the octyl group or that with the hydroxy group *meta* to the octyl group is formed preferentially, depending on the solvent used. Thus, the regioselectivity of the lipase from *Pseudomonas cepacia* is reversed on going from toluene to acetonitrile as the reaction medium [314].

A striking example of a complete reversal of enzyme *enantioselectivity* upon a change of solvent is given by the transesterification reaction of *N*-acetyl-(L or D)-phenylalanine 2-chloroethyl ester with 1-propanol, catalysed by *Aspergillus oryzae* protease in eighteen anhydrous organic solvents [315]. Whereas in hydrophilic polar solvents such as acetonitrile, *N*,*N*-dimethylformamide, and pyridine, the L-enantiomer of the ester is much more reactive than its D counterpart, the opposite holds true for hydrophobic apolar solvents such as toluene, octane, and tetrachloromethane. The enantioselectivity (given by the ratio of initial rates v_L/v_D) changes from 7.1 in acetonitrile to 0.19 in tetrachloromethane, *i.e.* by a factor of 37. This result can be explained in terms of the variation in solvent hydrophobicity (as measured by lg $P_{o/w}$; see Section 7.2), without invoking solvent-induced conformational changes of the enzyme [315].

Another remarkable example of a solvent-induced change in *enantioselectivity* is the partial hydrolysis of achiral 4-aryl-substituted 1,4-dihydro-2,6-dimethyl-3,5-pyridine dicarboxylic diesters, catalysed by a lipase from *Pseudomonas sp.*, leading either to the (*R*)- or (*S*)-configured monoesters. Whereas in water-saturated di-*i*-propyl ether, the (*S*)-monoesters are obtained with *ee* values up to 99%, the (*R*)-monoesters are formed in water-saturated cyclohexane with *ee* values of 88–91% [316].

Many other examples of the use of enzymes in organic solvents can be found in some excellent reviews [309–312].

Attempts have been made to correlate the influence of solvents on enzyme activity, stability, and selectivity with physicochemical solvent characteristics such as relative permittivity, dipole moment, water miscibility, and hydrophobicity, as well as empirical parameters of solvent polarity. However, no rationale of general validity has been found, except the simple rule that nonpolar hydrophobic solvents are generally better than polar hydrophilic ones. The best correlations are often obtained with the logarithm of the 1-octanol/water partition coefficient, lg $P_{O/W}$, a quantitative measure of the solvent's hydrophobicity (*cf.* Section 7.2).

In the selection of suitable organic solvents for enzyme-catalysed reactions, care has to be taken that the removal of water does not denature or deactivate the protein. This leads to the question as to how much water is needed to preserve the native conformation of the enzyme-forming protein. Certainly not the entire 55.5 mol/L water that surrounds it in an aqueous solution. It seems that only a few monolayers of water around the enzyme molecule are sufficient to preserve its biological activity. As long as this water (*i.e.* constitutional water inside the protein molecule and interfacial water located on the protein surface) is available around the enzyme molecule, the bulk water can evidently be replaced by other solvents without adversely affecting the enzyme's activity. It has been found that, although enzymic proteins are denatured in water/organic solvent mixtures, their secondary structure remains essentially intact in the corresponding pure organic solvents [317].

Nonaqueous biocatalytic reactions represent significant recent progress in enzyme

biotechnology. Some advantages of biocatalysis in organic solvents as opposed to water are: (a) better solubility of organic (mostly hydrophobic) substrates and products, (b) enhanced thermostability of enzymes in organic solvents, (c) the ability to carry out new reactions that are impossible in water, (d) suppression of undesirable side reactions caused by water, (e) insolubility of enzymes in organic solvents, which permits their easy recovery and reusability, and (f) as already mentioned, the possibility of controlling enzyme specificity [309–312]. A more detailed treatment of biocatalysis in organic solvents and its practical applications is beyond the scope of this book and the reader is referred to recent reviews [311, 312].

5 Solvent Effects on the Rates of Homogeneous Chemical Reactions

5.1 General Remarks [1–29, 452–463]

A change of solvent can considerably change both the rate and order of a homogeneous chemical reaction. As early as 1890, Menschutkin demonstrated in his classical study on the quaternization of triethylamine with iodoethane in 23 solvents, that the rate of reaction varied remarkably depending on the choice of solvent. Compared to the rate in *n*-hexane, the rate was found to be four times faster in diethyl ether, 36 times faster in benzene, 280 times faster in methanol, and 742 times faster in benzyl alcohol [30]. Thus, through judicious choice of solvent, decisive acceleration or deceleration of a chemical reaction can be achieved. This can be of great practical importance both in the laboratory and in the chemical industry. In some extreme cases, rate accelerations by a factor of up to *ca.* 10^9 (!) can be achieved solely by a solvent change [31]. Therefore, it is very important to establish rules and theories, enabling a rational selection of solvent and design for chemical synthesis.

The dependence of the reaction rate on the medium can, in principle, be approached from two points of view: (a) comparison of the rates of reaction in the gas phase and in solution, and (b) comparison of the rates of reaction in different solvents.

Until recently, only very few reactions which occur in solution have also been sufficiently examined in the gas phase [32]. Therefore, the comparison of solvent effects has been essentially limited to method (b). Reactions which have been studied according to method (a) are mainly non-ionic reactions, *i.e.* reactions without any charge separation or charge dispersion during the activation process, such as, for example, pericyclic reactions in which neutral reactants produce neutral products. Reactions that follow ionic mechanisms with considerable charge separation or charge dispersion during activation, such as, for example, proton-transfer or S_N2 ion-molecule reactions with charged species as reactants, have until recently only been investigated using method (b).

Three new experimental techniques, developed within the past decades, now make it possible to study ionic reactions in the gas phase as well. These are pulsed ion-cyclotron-resonance (ICR) mass spectrometry, pulsed high-pressure mass spectrometry (HPMS), and the flowing afterglow (FA) technique [469–478; see also the references given in Section 4.2.2]. Although their approaches are quite independent, the results obtained for acid/base and other ionic reactions agree within an experimental error of 0.4 . . . 1.3 kJ/mol (0.1 . . . 0.3 kcal/mol) and are considered as reliable as those obtained in solution.

In solution, ions are produced by the heterolysis of covalent bonds in ionogens. This ionization reaction is favored by solvents due to their cooperative EPD and EPA properties (*cf.* Section 2.6). In the gas phase, however, ionization of neutral molecules to form free ions is rarely observed because this reaction is very endothermic. For example, in order to ionize gaseous H—Cl into $H^{\oplus}$ and $Cl^{\ominus}$, an energy of 1393 kJ/mol (333 kcal/mol) must be provided. This considerably exceeds the 428 kJ/mol (102 kcal/mol) needed to homolytically cleave H—Cl into hydrogen and chlorine atoms. Thus, for the creation of isolated ions in the gas phase, energy must be supplied by some means other than solvation with EPD/EPA solvents. The most widely used method is ionization by elec-

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

tron impact of gaseous molecules, as employed in mass spectrometry. However, whereas ions in solution are stabilized by their solvation shells, gaseous ions are immediately destroyed when they strike a solid surface. Therefore, special care has to be taken to restrict the motion of gaseous ions and to hold them long enough so that their chemical reactivity can be studied. Several methods have been developed for storing gaseous ions. In ICR mass spectrometry, a static magnetic trap is used [469].

The study of reactions of isolated ions and molecules in the gas phase without interference from solvents has led to very surprising results. Gas-phase studies of proton-transfer and nucleophilic substitution reactions permit the measurement of the intrinsic properties of the bare reactants and make it possible to distinguish these genuine properties from effects attributable to solvation. Furthermore, these studies provide a direct comparison of gas-phase and solution reactivities of ionic reactants. It has long been assumed that solvation retards the rates of ion-molecule reactions. Now, using these new techniques, the dramatic results obtained make it possible to show the extent of this retardation. For example, in a typical S_N2 ion-molecule reaction in the gas phase, the substrates react about 10^{10} times faster than when they are dissolved in acetone, and about 10^{15} (!) times faster than in water (*cf.* Table 5-2 in Section 5.2).

Clearly, the effect of the solvent on a chemical reaction is much larger than previously assumed. In solution, the behaviour of ions and molecules is dictated mainly by the solvent and only to a lesser extent by their intrinsic properties. This will be elaborated on in subsequent Sections. A comparison of gas-phase reactivities and solution reactivities is given in Section 5.2.

There are, in principle, two ways in which solvents can affect the reaction rates of homogeneous chemical reactions: through static, or equilibrium, solvent effects and through dynamic, or frictional, solvent effects [463, 465, 466].

The static influence of solvents on rate constants can be understood in terms of transition-state theory. According to this theory, solvents can modify the Gibbs energy of activation (as well as the corresponding activation enthalpies, activation entropies, and activation volumes) by differential solvation of the reactants and the activated complex. Reaction rates are very sensitive to barrier heights. For example, a change of only 8.4 kJ/mol (2 kcal/mol) in an activation barrier can alter the reaction rate at room temperature by a factor of 31. Here, it is implicitly assumed that the required reorientational relaxation of solvent molecules during the activation process is sufficiently fast and that the activated complex is in thermal equilibrium with the solvent due to the frequent collisions of the reacting system with the surrounding solvent molecules. The Hughes–Ingold rules of solvent effects on reaction rates (*cf.* Section 5.3.1) are based on equilibrium solvation of the activated complex.

This equilibrium hypothesis is, however, not necessarily valid for rapid chemical reactions. This brings us to the second way in which solvents can influence reaction rates, namely through dynamic or frictional effects. For broad-barrier reactions in strongly dipolar, slowly relaxing solvents, non-equilibrium solvation of the activated complex can occur and the solvent reorientation may also influence the reaction rate. In the case of slow solvent relaxation, significant dynamic contributions to the experimentally determined activation parameters, which are completely absent in conventional transition-state theory, can exist. In the extreme case, solvent reorientation becomes rate-limiting and the transition-state theory breaks down. In this situation, rate con-

stants will depend on the solvent dynamics, and will vary with friction, *i.e.* with some measure of the coupling of the solvent such as density, internal pressure, or viscosity. In the opposite regime of reactions, *i.e.* those with sharp barriers and weakly dipolar, rapidly relaxing solvents, reaction rates are fairly well described by transition-state theory. At present, it seems to be difficult to quantitatively separate the influence of the dynamic and static solvent effects on the reaction rate. The importance of these less familiar frictional solvent effects on reaction rates has only recently been stressed [463, 465, 466], although the first theoretical treatment of such dynamic solvent effects was given by Kramers as early as 1940 [479].

Because the transition-state theory is essential for even a qualitative understanding of solvent effects on reaction rates, some important features of this theory are outlined below.

Arrhenius' classical theory of reaction kinetics is based on the assumption that the starting materials (reactants) have to overcome an energy barrier, the activation energy, in order to be transformed into the products. This picture has been developed and made more explicit in the theory of absolute reaction rates [2–5, 7, 8, 11, 24, 464–466, 770, 771]. The influence of solvent on reaction rates is best treated by means of this theory – also known as transition-state theory, developed almost simultaneously in 1935 by Eyring as well as Evans and Polanyi [464].

Consider a reaction between the starting compounds A and B, and suppose that during the course of the reaction these two first form an activated complex, which then decomposes to the end products, C and D. The reaction can then be described as follows:

$$\underset{\text{Reactants}}{A + B} \rightleftharpoons \underset{\substack{\text{Activated}\\ \text{Complex}}}{(AB)^{\neq}} \rightarrow \underset{\text{Products}}{C + D} \tag{5-1}$$

The theory of absolute reaction rates is based on the following assumptions:

(a) The reactants are envisaged as being in quasi-equilibrium with the activated complex. The corresponding quasi-thermodynamic equilibrium constant is given by Eq. (5-2) (with a = activities, [ ] = molar concentrations, and γ = activity coefficients).

$$K^{\neq} = \frac{a_{(AB)^{\neq}}}{a_A \cdot a_B} = \frac{[(AB)^{\neq}]}{[A] \cdot [B]} \cdot \frac{\gamma^{\neq}}{\gamma_A \cdot \gamma_B} \tag{5-2}$$

The use of activity coefficients takes into account deviations from ideal behaviour in solution. The activity coefficients usually refer to the standard state of infinite dilution for the solutes: $\lim_{c \to 0} \gamma = 1$.

(b) The formation of the products C and D does not significantly affect the equilibrium between reactants and activated complex.

(c) The activated complex has all the properties of a normal molecule (with N atoms) except that one vibrational degree, of the $3N$–6 vibrational degrees of freedom, is



Fig. 5-1. One-dimensional Gibbs energy diagram for reaction (5-1) in solution. Ordinate: relative standard molar Gibbs energies of reactants, activated complex, and products; Abscissa*): not defined, expresses only the sequence of reactants, activated complex, and products as they occur in the chemical reaction. ΔG°: standard molar Gibbs energy of the reaction; $\Delta G^{\neq}$: standard molar Gibbs energy of activation for the reaction from the left to the right.

transformed into a translational degree of freedom, which leads to the decomposition of the activated complex.

(d) The activated complex exists at the top of an energy barrier as shown in Fig. 5-1. The activated complex represents that point in the progress of the reaction at which the reformation of reactants is as likely as the formation of products. The region in the neighbourhood of this maximum is also called the "transition state"**).

(e) The net reaction rate is determined by the rate at which the activated complex passes over the energy barrier in the direction of product formation. The probability of a forward passage through the activated complex is given by a transmission coefficient, which is usually assumed to approach unity.

---

* The so-called *reaction coordinate* – often used as abscissa in such diagrams – is essentially a molecular-microscopic quantity. In the case of a unimolecular reaction it is the internuclear distance; in the case of a bimolecular reaction it represents a translational degree of freedom and in the transition state a normal mode of vibration that leads to the decomposition of the activated complex. ΔG° and $\Delta G^{\neq}$ are, however, macroscopic thermodynamic state functions, which are experimentally available only at the maxima and minima and not at configurations between them. Therefore, in order to avoid the indiscriminate mixing of macroscopic and molecular-microscopic quantities, the one-dimensional presentation of Fig. 5-1 is preferable to the smooth curves often used and called Gibbs energy "profiles". In Fig. 5-1, the reaction coordinate has no meaning. The abscissa expresses only the sequence of reactants, activated complex, reaction intermediates, and products as they occur in chemical reactions. *Cf.* also Fig. 4-1 in Chapter 4.

** The terms *activated complex* and *transition state* are often wrongly regarded as synonymous. *Activated complex* refers to the real molecular entity at the point of maximum Gibbs energy on the reaction path, whereas *transition state* describes only the set of states or energy levels at this point. The transition state has no physical existence [*cf.* K. J. Laidler, J. Chem. Educ. *65*, 540 (1988)]. In order to avoid the unnecessary confusion of state with structure (*i.e.* energy space with geometric space), it has been proposed that the configuration of atoms in the geometrical 3D space at the potential energy saddle point should be called the *critical transition structure* (CTS) [*cf.* S. H. Bauer and C. F. Wilcox, J. Chem. Educ. *72*, 13 (1995)]. Using ultrafast pulsed laser techniques with femtosecond resolution (1 fs = 10^{-15} s), it is now even possible to study directly some properties of *activated complexes* as molecular species [*cf.* A. H. Zewail, J. Phys. Chem. A *104*, 5660 (2000)].

(f) The change in the Gibbs energy of activation in going from the gas phase to solution or from one solvent to another is evaluated as the relative modification in Gibbs energy by differential solvation of the reactants and the activated complex, whereby it is implicitly assumed that in solution reactants and activated complex are in thermal equilibrium with the solvent. This equilibrium hypothesis will not be valid for rapid chemical reactions with slow reorientational relaxation of the solvent molecules. For broad-barrier reactions in slowly relaxing solvents, solvent reorientation becomes rate limiting and transition-state theory may break down [463, 465, 466].

Because the reaction rate is assumed to be proportional to the concentration of the activated complex, the specific rate constant k is proportional to $K^{\neq}$. By statistical calculations, the proportionality factor can be estimated as $k_B \cdot T/h$. Provided that each activated complex passing the transition state actually becomes product (transmission coefficient close to unity), and that the activity coefficients are close to unity, the specific rate constant of the elementary reaction is given by Eq. (5-3),

$$k = \frac{k_B \cdot T}{h} \cdot K^{\neq} \cdot (1\ \text{mol} \cdot \text{L}^{-1})^{1-n} = \frac{R \cdot T}{N_A \cdot h} \cdot K^{\neq} \cdot (1\ \text{mol} \cdot \text{L}^{-1})^{1-n} \tag{5-3}$$

where k is the rate constant including units with the dimensions $(\text{mol} \cdot \text{L}^{-1})^{1-n} \cdot (\text{time})^{-1}$, k_B the Boltzmann constant, h the Planck constant, T the absolute temperature, R the gas constant, N_A the Avogadro constant, $K^{\neq}$ the quasi-thermodynamic equilibrium constant related to a hypothetical unit molar concentration standard state for the postulated equilibrium between reactant(s) and activated complex, and n the molecularity and kinetic order of the elementary reaction (usually 1 or 2).

This simple equation predicts that the rate constant for any chemical reaction consists of an equilibrium constant multiplied by a universal frequency factor, $k_B \cdot T/h$, which varies only with the temperature.

Since $K^{\neq}$ represents an equilibrium constant, it is possible to define quantities such as $\Delta G^{\neq}$, $\Delta H^{\neq}$, and $\Delta S^{\neq}$ according to Eqs. (5-4) and (5-5), which are called the Gibbs energy, enthalpy, and entropy of activation, respectively.

$$\Delta G^{\neq} = -R \cdot T \cdot \ln K^{\neq} \tag{5-4}$$

$$\Delta G^{\neq} = \Delta H^{\neq} - T \cdot \Delta S^{\neq} \tag{5-5}$$

From Eqs. (5-3), (5-4), and (5-5), it follows that the specific rate constant for the standard state ($\gamma = 1$) can also be given by Eq. (5-6).

$$\begin{aligned} k &= \frac{R \cdot T}{N_A \cdot h} \cdot e^{-\Delta G^{\neq}/(R \cdot T)} \cdot (1\ \text{mol} \cdot \text{L}^{-1})^{1-n} \\ &= \frac{R \cdot T}{N_A \cdot h} \cdot e^{-\Delta H^{\neq}/(R \cdot T)} \cdot e^{\Delta S^{\neq}/R} \cdot (1\ \text{mol} \cdot \text{L}^{-1})^{1-n} \end{aligned} \tag{5-6}$$

Eq. (5-6) is assumed to be valid for reactions in solution, although it is probably applicable only to an ideal gas system. According to this equation, the smaller the value of $\Delta G^{\neq}$, *i.e.* the difference between the Gibbs energies of reactants and activated complex, the greater the reaction rate of a chemical reaction. The fastest reaction (with $\Delta G^{\neq} = 0$) is given by $R \cdot T/N_A \cdot h$, which corresponds to the frequency for the passage of the activated complex through the transition state. At 25 °C, this value is 6×10^{12} s^{-1}, a typical value for the frequency of molecular vibrations.

In principle, when studying solvent effects on reaction rates, it is not sufficient to investigate only the $\Delta G^{\neq}$ change because, according to Eq. (5-5), this term is determined by both an enthalpy and an entropy term. There are four types of reaction rate control [41]:

(a) Cooperative effects, with $\Delta\Delta H^{\neq}$ and $T \cdot \Delta\Delta S^{\neq}$ having opposite signs. Then these two terms will be additive;
(b) Enthalpy-controlled reactions, for which the two terms are opposed (*i.e.* $\Delta\Delta H^{\neq}$ and $T \cdot \Delta\Delta S^{\neq}$ have equal signs), but the activation enthalpy term is larger;
(c) Entropy-controlled reactions, for which the two terms oppose, but the $T \cdot \Delta\Delta S^{\neq}$ term is larger; and
(d) Compensating effects, where the two terms are opposed, but nearly equal.

If reaction (5-1) takes place in solution, then the initial reactants, as well as the activated complex, will be solvated to differing extents according to the solvating power of the solvent used. This differential solvation can retard or accelerate a reaction in the manner described in Fig. 5-2.

$\Delta G_I^{\neq}$ in Fig. (5-2a) and (5-2b) represents the Gibbs energy of activation for a given chemical reaction in an ideal solvent I. In such a case, neither the reactants nor the activated complex are solvated. If in another solvent II only the activated complex is solvated, then $\Delta G_{II}^{\neq}$ in Fig. (5-2a) results. The Gibbs energy of activation is reduced by the Gibbs energy of transfer, $\Delta G_{I\to II}^{\neq}$, with a consequent rate acceleration. If on the other hand, only the initial reactants are solvated, as happens in solvent III, then $\Delta G_{III}^{\neq}$ in Fig. (5-2b) results. The Gibbs energy of activation is increased by the Gibbs energy of transfer, $\Delta G_{I\to III}^{R}$, with a consequent rate deceleration. The solvation of the products does not have any influence on the reaction rate. Because in reality the initial reactants as well as the activated complex are solvated, but usually to a different extent, the difference of both Gibbs transfer energies determines the reaction rate in solution.

A dissection, therefore, of the reaction rate solvent effects into initial and transition state contributions would lead to more direct information concerning the nature of the solvent involvement [453, 467].

Fig. 5-3 shows the Gibbs energy diagram for a chemical reaction carried out in two different solvents I and II. The standard molar Gibbs energy of the reactants R in solvent I is designated G_I^R and in solvent II as G_{II}^R. The difference in Gibbs energies between the two solvents, $(G_I^R - G_{II}^R)$, is termed the Gibbs energy of transfer, $\Delta G_{I\to II}^R$. Similarly, for the activated complex one obtains $\Delta G_{I\to II}^{\neq} = G_I^{\neq} - G_{II}^{\neq}$.

The difference in Gibbs energies of activation for the reaction in the two solvents is designated $\Delta\Delta G^{\neq}$, and from Fig. 5-3 it follows that



Fig. 5-2. One-dimensional Gibbs energy diagram for a chemical reaction in three different solvents I, II, and III (*cf.* Fig. 5-1). *(a)* Reaction with non-solvated (solvent I) and solvated (solvent II) activated complex (preferential solvation of the activated complex); *(b)* Reaction with non-solvated (solvent I) and solvated (solvent III) reactants (preferential solvation of the reactants). $\Delta G_{\mathrm{I}}^{\neq}$, $\Delta G_{\mathrm{II}}^{\neq}$, and $\Delta G_{\mathrm{III}}^{\neq}$: standard molar Gibbs energies of activation in solvents I, II, and III, respectively; $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\neq}$ and $\Delta G_{\mathrm{I}\rightarrow\mathrm{III}}^{\mathrm{R}}$: standard molar Gibbs energies of the transfer of the activated complex and the reactants R from solvent I to solvents II and III, respectively.



Fig. 5-3. One-dimensional Gibbs energy diagram for a chemical reaction in two different solvents I and II (*cf.* Figs. 5-1 and 5-2). $\Delta G_{\mathrm{I}}^{\neq}$ and $\Delta G_{\mathrm{II}}^{\neq}$: standard molar Gibbs energies of activation in solvents I and II; $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{R}}$ and $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\neq}$: standard molar Gibbs energies of transfer of the reactants R and the activated complex from solvent I to solvent II, respectively.

$$\Delta\Delta G^{\neq} = \Delta G_{\mathrm{I}}^{\neq} - \Delta G_{\mathrm{II}}^{\neq} = (G_{\mathrm{I}}^{\neq} - G_{\mathrm{I}}^{\mathrm{R}}) - (G_{\mathrm{II}}^{\neq} - G_{\mathrm{II}}^{\mathrm{R}}) \tag{5-7}$$

this can be simplified to give

$$\Delta\Delta G^{\neq} = \Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\neq} - \Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{R}} \tag{5-8}$$

From Eq. (5-8), it is apparent that $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\neq}$ can be evaluated from the measurable Gibbs energies of transfer of the reactants, $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{R}}$, in conjunction with the measured kinetic activation parameters, $\Delta\Delta G^{\neq}$.

The required Gibbs transfer energies, $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{R}}$, can be obtained from activity coefficient measurements according to Eq. (5-8a),

$$\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{i}} = -R \cdot T \cdot \ln(\gamma_{\mathrm{I}}/\gamma_{\mathrm{II}}) \tag{5-8a}$$

where γ refers to the activity coefficients of solute i in solvents I and II. The γ values can be obtained from vapour pressure, solubility, and distribution coefficient measurements [23, 467].

Consequently, by combining the thermodynamic and kinetic measurements, values for $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\neq}$ and $\Delta G_{\mathrm{I}\rightarrow\mathrm{II}}^{\mathrm{R}}$ can be obtained. Both terms can be either positive (destabilization), negative (stabilization), or zero (no solvent effect). When both terms have the same sign, a balancing situation occurs, when they are of opposite sign, a reinforcing situation. The effect of the solvent on the rate can be expected to be largest in the reinforcing situation and smallest in the balancing situation [467].

Because of the complicated interactions between solvents and solutes, the prediction of solvent effects on reaction rates, and the correlation of these effects with intrinsic solvent properties, is very difficult. Nevertheless, many authors have tried to establish – empirically or theoretically – correlations between rate constants or Gibbs energies of activation and characteristic solvent parameters such as relative permittivity, ε_{r}, dipole moment, μ, refractive index, n, solubility parameter, δ, empirical solvent polarity parameters, *etc.*, as schematically shown by Eq. (5-9).

$$\lg k \quad \text{or} \quad \Delta G^{\neq} = f(\varepsilon_{\mathrm{r}}, \mu, n, \delta, \ldots) \tag{5-9}$$

An early attempt in this direction was O. Dimroth's effort in correlating rate constants k with the solubility S of the reactants in the solvents used [42]. While investigating the intramolecular rearrangement of 5-hydroxy-1-phenyl-1,2,3-triazole-4-carboxylic esters in various solvents, he found, in agreement with a rule formulated by van't Hoff [43], that the rate constants are inversely proportional to the solubility of the rearranging isomers; *cf.* Eq. (5-10).

$$k = const \cdot \frac{1}{S} \tag{5-10}$$

Relationships such as this have only a very limited application [2]. More modern attempts to correlate the solvent influence on reaction rates with physical and empirical parameters of the solvents can be found in Sections 5.4 and 7.6, respectively.

The following Sections deal first with gas-phase reactivities and then with the most important qualitative and quantitative general relationships between reaction rates and solvent properties. The next Section (5.2) begins with a comparison of reactions in the gas phase and in solution, thus demonstrating the huge changes in reaction rates obtained on going from the gas phase to solution. The following Section (5.3) deals with the qualitative Hughes–Ingold rules (and their limitations) concerning solvent effects on substitution and elimination reactions, based on a sub-classification of these reactions according to the different charge-types of the initial reactants [16]. Thereafter, further analogous examples of solvent effects on organic reactions, using a classification of reactions according to Kosower [15] and Reichardt [468], are given. Attempts at establishing quantitative relationships between reaction rates and physical solvent properties are then reviewed (Section 5.4), which is followed by a treatment of some specific solvation effects on reaction rates (Section 5.5).

5.2 Gas-Phase Reactivities

In the gas phase, bond fission is invariably homolytic and complications from solvents are absent. Reactants follow their reaction route in splendid isolation which depends only on the intrinsic (inherent) properties of the reactant molecules. On the other hand, bond fission in solution is generally heterolytic due to the EPD/EPA properties of the solvents. The ubiquitous solvent molecules perturb the reactants in their reaction course, sometimes to such an extent that the solvent is entirely responsible for the observed rate constants.

Accordingly, in the gas phase, reactions without charge separation or charge dispersion are preferred, *e.g.* radical-producing and pericyclic reactions. In solution, on the other hand, reactions involving charge separation and charge dispersion can also be carried out, *e.g.* ionization and S_N2 ion-molecule reactions. Reactions can only occur in the gas phase *and* in solution when the intermolecular interactions between reactants and solvents are so weak that the non-ionic gas-phase reaction mechanism is virtually unchanged on transferring the reactants to solution.

An example of such a reaction, which has the same mechanism in the gas phase and in solution, is the bimolecular Diels–Alder cycloaddition reaction of cyclopentadiene to give *endo*-dicyclopentadiene – a reaction between neutral reactants to give a neutral product. As the Arrhenius activation energies and the rate constants of this reaction in Table 5-1 show, there is only a slight variation on going from the gas phase to solution. The rate constants vary by only a factor of *ca.* 3, accompanied by a corresponding small change in the activation energies.

The homolytic thermolyses of di-*t*-butyl peroxide [36, 37] and diacetyl peroxide [38, 39] have roughly the same rate constants and activation energies both in the gas phase and in solution in a variety of solvents.

On the contrary, reactions involving charge separation or charge dispersion during the activation process depend strongly on the reaction medium, due to the strong intermolecular interactions between the ionic and dipolar reactants and the solvent. Examples include S_N1 and S_N2 reactions, elimination reactions, isomerizations involving polar and charged groups, as well as proton- and electron-transfer reactions.

Table 5-1. Comparison of the activation energies and specific rate constants for the bimolecular Diels–Alder cycloaddition reaction of cyclopentadiene giving *endo*-dicyclopentadiene in the gas phase and in solution at 20 °C [3, 33, 34].

Medium	E_a / (kJ · mol^{-1})	k_2 / (L · mol^{-1} · s^{-1})	k_2^{rel}
Gas phase	69.9	$6.9 \cdot 10^{-7}$	1.2
Ethanol	68.6	$19 \cdot 10^{-7}$	3.4
Nitrobenzene	63.2	$13 \cdot 10^{-7}$	2.3
Paraffin oil	72.8	$9.8 \cdot 10^{-7}$	1.8
Carbon disulfide	70.7	$9.3 \cdot 10^{-7}$	1.7
Tetrachloromethane	71.5	$7.9 \cdot 10^{-7}$	1.4
Benzene	68.6	$6.6 \cdot 10^{-7}$	1.2
Neat liquid	67.8	$5.6 \cdot 10^{-7}$	*1.0*

A typical example of such reactions is the exothermic S_N2 nucleophilic displacement reaction $Cl^{\ominus} + CH_3—Br \rightarrow Cl—CH_3 + Br^{\ominus}$. Table 5-2 provides a comparison of Arrhenius activation energies and specific rate constants for this Finkelstein reaction in both the gas phase and solution. The new techniques described above (*cf.* Sections 4.2.2 and 5.1) have made it possible to determine the rate constant of this ion-molecule reaction in the absence of any solvent molecules in the gas phase. The result is surprising: on going from a protic solvent to a non-HBD solvent and then further to the gas phase, the ratio of the rate constants is approximately $1:10^5:10^{15}$! The activation energy of this S_N2 reaction in water is about ten times larger than in the gas phase. The suppression of the S_N2 rate constant in aqueous solution by up to 15 orders of magnitude demonstrates the vital role of the solvent.

A detailed molecular-level understanding of the role of solvation on the nature of S_N2 reaction pathways has been revealed only during the last decades. Fig. 5-4 compares the gas-phase S_N2 enthalpy diagram with two minima, first proposed by Brauman *et al.* [474], with the more familiar single transition-state diagram obtained in solution.

Table 5-2. Comparison of the activation energies and specific rate constants for the bimolecular S_N2 ion-molecule reaction $Cl^{\ominus} + CH_3—Br \rightarrow Cl—CH_3 + Br^{\ominus}$ in the gas phase and in solution at 25 °C [480].

Medium	E_a / (kJ · mol^{-1})	k_2 / (cm^3 · molecule^{-1} · s^{-1})	k_2^{rel}
Gas phase (at 24 °C)	11	$2.1 \cdot 10^{-11}$	$2.5 \cdot 10^{15}$
Acetone[a)]	66	$5.5 \cdot 10^{-21}$	$6.6 \cdot 10^{5}$
N,*N*-Dimethylformamide[a)]	75	$8.3 \cdot 10^{-22}$	$1.0 \cdot 10^{5}$
Methanol[b)]		$1.0 \cdot 10^{-26}$	1.2
Water[c)]	103	$8.3 \cdot 10^{-27}$	*1.0*

[a)] D. Cook and A. J. Parker, J. Chem. Soc., Part B *1968*, 142.
[b)] R. Alexander, E. C. F. Ko, A. J. Parker, and T. J. Broxton, J. Am. Chem. Soc. *90*, 5049 (1968).
[c)] R. H. Bathgate and E. A. Moelwyn-Hughes, J. Chem. Soc. *1959*, 2642.



Fig. 5-4. Schematic one-dimensional enthalpy diagram for the exothermic bimolecular Finkelstein reaction $Cl^{\ominus} + CH_3—Br \rightarrow Cl—CH_3 + Br^{\ominus}$ in the gas phase and in aqueous solution [469, 474, 476]. Ordinate: standard molar enthalpies of *(a)* the reactants, *(b, d)* loose ion-molecule clusters held together by ion-dipole and ion-induced dipole forces, *(c)* the activated complex, and *(e)* the products. Abscissa: not defined, expresses only the sequence of *(a)* ... *(e)* as they occur in the chemical reaction.

The unexpected gas-phase double-minimum diagram can be best explained as follows: As the reactants approach one another, long-range ion-dipole and ion-induced dipole interactions first produce loose ion-molecule association complexes or clusters. This is related to a decrease in enthalpy prior to any chemical barrier produced by orbital overlap between the reactants. For reasons of symmetry, an analogous drop in enthalpy must exist on the product side. Because the neutral reactant and product molecules will, in general, have different dipole moments and polarizabilities, the two minima will also be different. Only in the case of degenerate identity S_N2 reactions ($X^{\ominus} + CH_3—X \rightarrow X—CH_3 + X^{\ominus}$) will the enthalpy of the two minima be equal.

Overall, the gaseous S_N2 reaction shown in Fig. 5-4 consists of a collision step, which generates reactant ion-dipole complexes *(b)*, followed by a chemical activation step forming activated complexes *(c)* that contain pentacoordinated carbon atoms. Once the transition state has been surmounted, product ion-dipole complexes *(d)* are formed, the dissociation of which yields the products *(e)*.

During the activation process, the enthalpy increases up to the central transition state level. The height of this central, intrinsic barrier is the major factor responsible for the inherent reactivities of the various reactant combinations. It is important to note that the enthalpy of the activated complex is less than that of the reactants. In other words, the enthalpy barrier for the formation of the activated complex is smaller than the lowering in enthalpy due to the attraction between the reactant ion and the dipolar reactant molecule.

This S_N2 reaction model, containing pre- and post-ion/dipole clusters that are lower in enthalpy than both the initial reactants and final products, and separated by an

(b) An increase in solvent polarity results in a decrease in the rates of those reactions in which the charge density is lower in the activated complex than in the initial reactant molecule(s).

(c) A change in solvent polarity will have a negligible effect on the rates of those reactions that involve little or no change in the charge density on going from reactant(s) to the activated complex.

In other words, a change to a more polar solvent will increase or decrease the reaction rate depending on whether the activated reaction complex is more or less dipolar than the initial reactants. In this respect, the term "solvent polarity" was used synonymously with the power to solvate solute charges. It was assumed to increase with the dipole moment of the solvent molecules, and to decrease with increased thickness of shielding of the dipole charges.

For example, the reaction between two equally charged ions results in an increase in charge density during the activation process. Therefore, the reaction rate will increase with increasing solvent polarity. On the other hand, a reaction between oppositely charged ions will be slower in polar solvents, which are good ion solvators, because in this case a reduction of charge density occurs in going from the reactants to the activated complex. Furthermore, reactions where charge is either created or destroyed during the activation process should be affected to a greater extent by a solvent polarity change than reactions in which there is only charge dispersal. Thus, the rates of substitution reactions involving charge creation or destruction are altered by a factor of $10^3 \ldots 10^6$ in going from water to ethanol, whereas S_N reactions involving charge dispersal are increased by only a factor of $3 \ldots 10$ when the solvent is changed from ethanol to water.

These Hughes–Ingold rules can be used for making qualitative predictions about the effect of solvent polarity on the rates of all heterolytic reactions of known mechanisms. For nucleophilic substitution reactions of types (5-11) and (5-12)

$$R\text{-}X \overset{S_N1}{\rightleftharpoons} [R^{\delta\oplus}\cdots X^{\delta\ominus}]^{\ddagger} \longrightarrow R^{\oplus} + :X^{\ominus} \xrightarrow{+:Y^{\ominus}} R\text{-}Y + :X^{\ominus} \qquad (5\text{-}11)$$

$$Y:^{\ominus} + R\text{-}X \overset{S_N2}{\rightleftharpoons} [Y^{\delta\ominus}\cdots R\cdots X^{\delta\ominus}]^{\ddagger} \longrightarrow Y\text{-}R + :X^{\ominus} \qquad (5\text{-}12)^*$$

---

* S_N2 reactions of the type shown in Eq. (5-12) have been commonly represented as occurring by attack of a lone electron pair from $Y^{\ominus}$ at the backside of R with simultaneous displacement of $X^{\ominus}$. However, this representation is misleading because electrons shift singly, not in pairs! Therefore, synchronous S_N2 displacement reactions are better represented in terms of $Y:^{\ominus} + R \cdot\cdot X \rightarrow Y \cdot\cdot R + :X^{\ominus}$, where a single electron from the donor anion $Y:^{\ominus}$ pairs with one electron from R, and an electron from R is transferred to X, accompanied by cleavage of the R—X bond [14, 483]. In certain cases, this process can merge with a single-electron transfer (SET) pathway, wherein the product Y—R is formed by coupling of an intermediate geminate radical pair according to

$$Y:^{\ominus} + R \cdot\cdot X \xrightarrow{SET} Y^{\odot} + R \cdot\cdot X^{\odot\ominus} \longrightarrow Y \cdot\cdot R + :X^{\ominus}$$

That is, the act of shifting the single electron from Y to X may occur either with or without free-radical formation. Usually, the concerted non-radicaloid process is energetically favoured. For a more detailed discussion of the various mechanisms of nucleophilic substitution reactions in aliphatic compounds and their solvent dependence, see references [14, 483, 782–785].

Table 5-4. Predicted solvent effects on rates of nucleophilic substitution reactions [16, 44–46].

Reaction type	Initial reactants	Activated complex	Charge alteration during activation	Effect of increased solvent polarity on rate[a)]
(a) S_N1	R—X	$R^{\delta+}\ldots X^{\delta-}$	Separation of unlike charges	Large increase
(b) S_N1	$R—X^+$	$R^{\delta+}\ldots X^{\delta+}$	Dispersal of charge	Small decrease
(c) S_N2	Y + R—X	$Y^{\delta+}\ldots R\ldots X^{\delta-}$	Separation of unlike charges	Large increase
(d) S_N2	$Y^- + R—X$	$Y^{\delta-}\ldots R\ldots X^{\delta-}$	Dispersal of charge	Small decrease
(e) S_N2	$Y + R—X^+$	$Y^{\delta+}\ldots R\ldots X^{\delta+}$	Dispersal of charge	Small decrease
(f) S_N2	$Y^- + R—X^+$	$Y^{\delta-}\ldots R\ldots X^{\delta+}$	Destruction of charge	Large decrease

[a)] The terms "large" and "small" arise from the theory that the effect of the dispersal of charge should be notably smaller than the effect of its creation or destruction and have therefore only relative significance.

the predictions are compiled in Table 5-4. The middle three columns show, for each charge type and mechanism, what happens to the charges on going from the initial to the transition state. The last column shows the predicted solvent effects. The conclusions derived from Table 5-4 were experimentally confirmed in a large number of substitution reactions. Some typical examples shall serve to demonstrate the application of these rules.

The solvolysis of 2-chloro-2-methylpropane is 335000 times faster in water than in the less polar solvent ethanol [40]; *cf.* reaction type (a) in Table 5-4.



$\mu = 2.9 \cdot 10^{-30}$ C m $\quad \mu \approx 27 \cdot 10^{-30}$ C m (5-13)

Solvent	C_2H_5OH	CH_3OH	$HCONH_2$	HCO_2H	H_2O
k_1^{rel}*)	1	9	430	12200	335000

Comparison of the solvolysis rate constants of 2-chloro-2-methylpropane obtained in water and in benzene solution reveals a rate acceleration of *ca.* 10^{11} with increasing solvent polarity [47]**). The solvolysis rate of 1-bromoadamantane in

* Here, and in the following examples, only the relative rate constants, k^{rel}, with respect to the "slowest solvent" are given. The selected solvents are ordered from left to right with increasing polarity.

** In the S_N1 solvolysis reaction of 2-chloro-2-methylpropane, leading mainly to *t*-butanol and *t*-butyl ethers together with some *i*-butene, the term *solvolysis* is normally restricted to the reaction in water and other HBD solvents. In non-HBD solvents, however, the only reaction product is *i*-butene. For convenience, the term *solvolysis* is often used in the literature to cover both types of reaction, solvolysis and dehydrohalogenation of 2-chloro-2-methylpropane, because the solvent-dependent rate-determining step of both reactions, S_N1 and E1, is the same. For a detailed review on the heterolysis of tertiary haloalkanes in the gas phase and in solution, see G. F. Dvorko, E. A. Ponomareva, and N. I. Kulik, Usp. Khim. *53*, 948 (1984); Russ. Chem. Rev. *53*, 547 (1984).

ethanol/water increases by a factor of 4900 when the volume fraction of water goes from 10 to 60 cL/L [48].

Rearrangement reactions, where the first step corresponds to the ionization of a tertiary haloalkane, also proceed faster with increasing solvent polarity, as shown by the Wagner–Meerwein rearrangement of 3-chloro-2,2,3-trimethylnorbornane to 2-*exo*-chlorobornane [49].



(5-14)

Solvent	Petroleum ether	C_6H_5Cl	$C_6H_5NO_2$	CH_3CN	CH_3NO_2
k_1^{rel}	1	20	65	200	610

In agreement with the predictions for reaction type (b) in Table 5-4, the thermolysis of triethylsulfonium bromide takes place more slowly in polar solvents, *e.g.* alcohols, than in less polar solvents, *e.g.* acetone [50].



(5-15)

Solvent	CH_3COCH_3	$CHCl_3$	$C_6H_5NO_2$	i-$C_5H_{11}OH$	$C_6H_5CH_2OH$
k_1^{rel}	290	>230	180	4.5	1

Solvolysis rates of *t*-butyl-dimethylsulfonium salts decrease with increasing solvent polarity [710]. Analogously, the rate of solvolysis of triethyloxonium salts in ethanol/water mixtures decreases with increasing water content [490]. Solvolysis rates of *N*-*t*-alkylpyridinium salts such as 1-(1-methyl-1-phenylethyl)pyridinium perchlorate are almost independent of solvent polarity, whereas *N*-*sec*-alkylpyridinium salts exhibit small decreases in rate with increasing solvent polarity [710]. The non-creation of charge in the activation process of these type (b) reactions much reduces the influence of solvent polarity on rate; *cf.* Table 5-4.

An example of reaction type (c) in Table 5-4 is the well-known Menschutkin reaction [30] between tertiary amines and primary haloalkanes yielding quaternary ammonium salts. Its solvent dependence was studied very thoroughly by a number of investigators [51–65, 491–496, 786–789]. For instance, the reaction of tri-*n*-propylamine with iodomethane at 20 °C is 120 times faster in diethyl ether, 13000 times faster in chloroform, and 110000 times faster in nitromethane than in *n*-hexane [60]. It has been estimated that the activated complex of this Menschutkin reaction should have a dipole moment of *ca.* $29 \cdot 10^{-30}$ Cm (8.7 D) [23, 64], which is much larger than the dipole moments of the reactant molecules (tris-*n*-propylamine $2.3 \cdot 10^{-30}$ Cm = 0.70 D; iodomethane $5.5 \cdot 10^{-30}$ Cm = 1.64 D) [64].

Table 5-5. Absolute and relative rate constants, Gibbs activation energies, activation enthalpies, and activation entropies of the Menschutkin reaction between triethylamine and iodoethane in twelve solvents at 50 °C [59].

Solvents	$k_2 \cdot 10^5$ / (L · mol^{-1} · s^{-1})	k_2^{rel}	$\Delta G^{\neq}$ / (kJ · mol^{-1})	$\Delta H^{\neq}$ / (kJ · mol^{-1})	$\Delta S^{\neq}$ / (J · mol^{-1} · K^{-1})
1,1,1-Trichloroethane	1.80	1	102.6	52.3	−156
Chlorocyclohexane	3.09	2	101.6	54.4	−146
Chlorobenzene	9.30	5	98.2	46.9	−159
1,1-Dichloroethane	11.9	7	97.1	48.1	−151
trichloromethane	15.4	9	96.2	49.0	−146
1,2-Dichlorobenzene	18.1	10	96.7	49.4	−146
Acetone	31.7	18	94.1	47.3	−145
Cyclohexanone	33.7	19	94.8	51.5	−134
1,2-Dichloroethane	42.6	24	93.4	45.2	−149
Propionitrile	59.6	33	92.3	48.5	−135
Benzonitrile	76.5	43	92.6	49.0	−135
Nitrobenzene	93.4	52	92.0	49.4	−133

Activation parameter data as a function of solvent have been published for the reaction of triethylamine with iodoethane [59]. The reaction rates obtained in dipolar aprotic solvents together with the activation parameters are given in Table 5-5 [59].



(5-16)

A solvent change from 1,1,1-trichloroethane to nitrobenzene causes a 52-fold rate acceleration for reaction (5-16), which corresponds to a decrease in $\Delta G^{\neq}$ of 10.6 kJ/mol. Bimolecular reactions such as (5-16), which produce charge during the activation process, usually show large negative entropies of activation. A negative entropy of activation indicates a greater degree of ordering in the transition state than in the initial state, due to an increase in solvation during the activation process. The $\Delta S^{\neq}$ values of Table 5-5 show that the largest decrease of activation entropy is obtained in the less polar solvents. This observation can be rationalized because polar solvents will have some structure corresponding to the orientation of the dipolar solvent molecules due to intermolecular solvent–solvent interactions. In less polar solvents, however, which have only a small or no dipole moment, the solvent molecules will be relatively unoriented and consequently have a higher entropy. Thus, nonpolar solvents will have a greater entropy loss as a result of increased solvation during the activation process. Consequently, reactions proceeding through dipolar activated complexes should have a larger negative entropy of activation in less polar solvents than in polar solvents [226].

The Menschutkin-type S_N2 reaction of triphenylphosphane with iodomethane has been studied in thirteen solvents [500]. In propylene carbonate, this reaction is 245 times faster than in di-*i*-propyl ether. In agreement with the highly dipolar activated complex, large solvent-dependent negative activation volumes, $\Delta V^{\neq}$, have been obtained as a

result of charge-induced reorientation of the surrounding solvent molecules during the activation process; *cf.* also Section 5.5.11.

For many physical organic chemists, the Menschutkin reaction was a kind of "guinea pig", which has been extensively used for the study of solvent effects on chemical reactivity. A comprehensive review of this reaction has been given by Abboud *et al.* [786]. More recent theoretical treatments of the solvent influence on Menschutkin reactions can be found in references [787–789].

Another remarkable example of reaction type (c) in Table 5-4, somewhat related to the Menschutkin reaction, is the S_N2 reaction between tris(*n*-butyl)phosphane and carbon disulfide, to give zwitterionic tris(*n*-butyl)phosphonium dithiocarboxylate via a corresponding dipolar activated complex [790, 791].

$$(n\text{-Bu})_3\text{P} + \text{CS}_2 \underset{k_1}{\overset{k_2}{\rightleftharpoons}} (n\text{-Bu})_3\text{P}^+\text{—CS}_2^- \qquad (5\text{-}16a)$$

The rate constants of both the forward (k_2) and the reverse reaction (k_1) are solvent-dependent, albeit to a different extent. In going from less polar diethyl ether to polar dimethyl sulfoxide, the forward reaction is accelerated by a factor of just 26, while the reverse reaction is decelerated to a larger extent. In other words, an inverted solvent change dimethyl sulfoxide → diethyl ether accelerates the reverse reaction 685-fold. This suggests that the activated complex of both reactions is located closer to the initial weakly dipolar reactants than to the zwitterionic adduct.

An example of reaction type (d) in Table 5-4 is the Finkelstein halide exchange reaction between iodomethane and radioactive labeled iodide ion. The rate constant for this reaction decreases by about 10^4 on going from less polar acetone to water as solvent [66].

$$\overset{*}{\text{I}}^{\ominus} + \text{CH}_3\text{-I} \underset{25\,^\circ\text{C}}{\overset{k_2}{\rightleftharpoons}} \left[\overset{*}{\text{I}}^{\delta\ominus}\cdots\text{CH}_3\cdots\text{I}^{\delta\ominus}\right]^{\ddagger} \longrightarrow \overset{*}{\text{I}}\text{-CH}_3 + \text{I}^{\ominus} \qquad (5\text{-}17)$$

Solvent	CH_3COCH_3	C_2H_5OH	$(CH_2OH)_2$	CH_3OH	H_2O
k_1^{rel}	13000	44	17	16	1

Other examples of this type of reaction are: S_N2 reactions between azide ion and 1-bromobutane [67], bromide ion and methyl tosylate [68], and bromide ion and iodoethane [497]. In changing the medium from non-HBD solvents (HMPT, 1-methylpyrrolidin-2-one) to methanol, the second-order rate constants decrease by a factor of $2 \cdot 10^5$ [67], $9 \cdot 10^4$ [68], and $1 \cdot 10^5$ [497], respectively. The large decrease in these rate constants in going from the less to the more polar solvent is not only governed by the difference in solvent polarity, as measured by dipole moment or relative permittivity, but also by the fact that the less polar solvents are dipolar aprotic and the more polar solvents are protic (*cf.* Section 5.5.2).

The second-order reaction between trimethylamine and the trimethylsulfonium ion gives the predicted rate decrease with increasing solvent polarity [69]; *cf.* reaction type (e) in Table 5-4.



(5-18)

Solvent	CH_3NO_2	C_2H_5OH	CH_3OH	H_2O
k_2^{rel}	119	10	6	1

Obviously, the initial reactants are more strongly solvated than the activated complex with its dispersed positive charge.

Finally, as an example of reaction type (f) in Table 5-4, the alkaline hydrolysis of the trimethylsulfonium ion demonstrates the predicted large rate decrease by increasing the proportion of water in an aqueous ethanolic reaction medium [70].



(5-19)

Solvent	Ethanol/water			
H_2O in cL/L	0	20	40	100
k_2^{rel}	19600	480	40	1

Similar findings were obtained for the alkaline decomposition of triarylsulfonium halides with ethoxide ion in aqueous ethanol at 120 °C [71]. The results indicate that the decomposition rate is decreased about 10^6-fold by increasing the water content from 2.3 to 98.2 cmol/mol [71].

It should be mentioned that a solvent change affects not only the reaction rate, but also the reaction mechanism (see Section 5.5.7). The reaction mechanism for some haloalkanes changes from S_N1 to S_N2 when the solvent is changed from aqueous ethanol to acetone. On the other hand, reactions of halomethanes, which proceed in aqueous ethanol by an S_N2 mechanism, can become S_N1 in more strongly ionizing solvents such as formic acid. For a comparison of solvent effects on nucleophilic substitution reactions at primary, secondary, and tertiary carbon atoms, see references [72, 784].

The mechanisms of nucleophilic substitution and β-elimination reactions are closely parallel with regard to the rate-determining step. The two monomolecular reactions (S_N1 and E_1) have a common rate-controlling step, whereas in the two bimolecular reactions (S_N2 and E_2), the electron transfers from the reagent to the leaving group are similar. However, they pass through a larger chain of carbon atoms in elimination than in substitution reactions. Therefore, similar rules for solvent effects on monomolecular, (5-20), and bimolecular, (5-21), β-elimination reactions of different charge-type have been obtained by Hughes and Ingold [16, 44] (*cf.* Table 5-6).

(5-20)



(5-21)

Table 5-6. Predicted solvent effects on rates of β-elimination reactions [16, 44, 73, 74].

Reaction type	Initial reactants	Activated complex	Charge alteration during activation	Effect of increased solvent polarity on rate[a)]
(a) E_1			Separation of unlike charges	Large increase
(b) E_1			Dispersal of charge	Small decrease
(c) E_2			Separation of unlike charges	Large increase
(d) E_2			Dispersal of charge	Small decrease
(e) E_2			Dispersal of charge	Small decrease
(f) E_2			Destruction of charge	Large decrease

[a)] The terms "large" and "small" arise from the theory that the effect of the dispersal of charge should be notably smaller than the effect of its creation or destruction and have therefore only relative significance.

The changes undergone by the charges during the activation process are shown in the three middle columns of Table 5-6. The last column shows the predicted solvent effects.

Owing to the similarity of solvent effects for E_1 and S_N1 reactions due to the common rate-controlling step, only some examples of E_2 reactions are mentioned. Typical examples of the reaction types (c) to (f) in Table 5-6 are shown in Eqs. (5-22) to (5-25).



$$(CH_3)_3N + (CH_3)_3C\text{-}Br \longrightarrow (CH_3)_3\overset{\oplus}{N}\text{-}H + H_2C{=}C(CH_3)_2 + Br^{\ominus} \quad (5\text{-}22)$$



$$HO^{\ominus} + CH_3\text{-}CH_2\text{-}Br \longrightarrow H_2O + H_2C{=}CH_2 + Br^{\ominus} \quad (5\text{-}23)$$



$$H_2O + O_2N\text{-}C_6H_4\text{-}CH_2\text{-}CH_2\text{-}\overset{\oplus}{N}(CH_3)_3 \longrightarrow H_3O^{\oplus} + O_2N\text{-}C_6H_4\text{-}CH{=}CH_2 + N(CH_3)_3 \quad (5\text{-}24)$$



$$C_2H_5O^{\ominus} + CH_3\text{-}CH_2\text{-}\overset{\oplus}{S}(CH_3)_2 \longrightarrow C_2H_5OH + H_2C{=}CH_2 + S(CH_3)_2 \quad (5\text{-}25)$$

In Eqs. (5-23) and (5-24), there is no net change of charge after reaction, but for Eq. (5-22) charge is created and for Eq. (5-25) it is destroyed. Further examples of observed effects for solvent changes on rates of mono- and bimolecular eliminations are given by Hughes and Ingold [16, 44]. In most cases studied, haloalkanes and *'onium* salts in ethanol/water mixtures, the observed solvent effects are in the expected direction.

According to the Hughes–Ingold rules, in E_2 and S_N2 reactions the dispersal of charge is spread over more atoms than in E_1 and S_N1 reactions, so increase in solvent

polarity will generally favor the E_1 and S_N1 mechanisms over the E_2 and S_N2 mechanisms. Thus, solvent change can alter not only the rate but also the reaction mechanism.

The competition between β-elimination and substitution reaction determines the proportion of alkene produced. The ratio of elimination to substitution is affected by the solvent as well as other factors (concentration and strength of the attacking base,



temperature, structure of substrate and attacking base). Looking at the activated complexes for S_N2 and E_2 reactions, it can be seen that the charge is more widely dispersed in the elimination transition state than in that of substitution.

From this it follows that both reactions are decelerated with increasing solvent polarity, but due to the larger charge dispersal on activation in E_2 reactions, solvent stabilization of the activated complex of elimination is less than that for substitution. Therefore, E_2 eliminations are more decelerated than S_N2 substitutions with increasing solvent polarity, and the alkene yield should fall slightly on making this solvent change. Table 5-7 gives some examples of solvent effects on the proportion in which the alkene is formed in β-elimination reactions.

Although the solvent effects are small, the alkene formation diminishes as predicted with increasing water content (corresponding to increased solvent polarity). The S_N2/E_2 reaction of 2-phenylpropyl tosylate with sodium cyanide (in hexamethylphosphoric triamide and in *N*,*N*-dimethylformamide as solvents at 100 °C) gives α-methylstyrene (elimination product) and 1-cyano-2-phenylpropane (substitution product) [75]. It has been found, in accordance with the predictions of the Hughes–Ingold rules, that the elimination/substitution ratio decreases as the polarity of the solvents (measured by the relative permittivity) increases [75]. Theoretical investigations of the

Table 5-7. Observed solvent effects on the proportion of alkene formed in mono- and bimolecular β-eliminations in ethanol/water mixtures [16, 44].

		Water in ethanol in cL/L			
		0	20	40	100
(a) *Bimolecular reactions*	($E_2 + S_N2$)				
$HO^{\ominus} + i\text{-}C_3H_7\text{—}Br$	(55 °C)	71	59	54	–
$HO^{\ominus} + (H_5C_2)_3S^{\oplus}$	(100 °C)	–	100	100	86
(b) *Monomolecular reactions*	($E_1 + S_N1$)				
$t\text{-}C_4H_9Br$	(25 °C)	19.0	12.6	–	–
$t\text{-}C_5H_{11}\text{—}Br$	(25 °C)	36.3	26.2	–	–
$t\text{-}C_5H_{11}\text{—}S(CH_3)_2^{\oplus}$	(50 °C)	64.4	47.8	–	–

dichotomic S_N2/E_2 reaction between F^- and CH_3CH_2F in the presence of n HF (n = 0–4) as a 'polar solvent' using the density-functional method have confirmed the experimental results: by microsolvation, the S_N2 activated complex is much more stabilized and becomes lower in energy than the *anti*-E_2 activated complex [854].

Analogously, solvent effects on alkene formation in S_N1 and E_1 reactions can be predicted [16, 44]. Owing to the fact that the first step in both reactions, the heterolysis of the C—X bond, is exactly the same, we have to consider the activated complexes which lead to either the alkene or substitution product.



Both reactions are decelerated with increasing solvent polarity. However, due to the greater dispersal of charge in the transition state of the E_1 reaction, the β-elimination is again more decelerated than the substitution reaction. This results in decreasing alkene formation with increasing solvent polarity. Similar results are obtained for reactions in which Y: represents a neutral base (*e.g.* H_2O). Some observed effects of solvent changes on alkene formation in E_1 and S_N1 reactions are given in Table 5-7. Experimental results are relatively scarce. Nevertheless, the predicted rate trends for all charge-types of *'onium* salts and haloalkanes in protic media are obtained [16, 44]. Newer studies of the competition between substitution and elimination in solvolyses of various 2-halo-2-methylpropanes in acetic acid, ethanol, and water have shown that decreasing solvent polarity favours alkene formation [76].

A further refinement of the Hughes–Ingold rules has been given by Westaway regarding the influence of solvents on the structure of S_N2 activated complexes [498]; *cf.* Eq. (5-12) and Table 5-4. His *solvation rule for S_N2 reactions* states that a change in solvent will not lead to a change in the structure of an S_N2 transition state if both the attacking nucleophile, Y, and the leaving group, X, have the same charge as in reaction types (d) and (e) in Table 5-4 (called *type I* S_N2 reactions). A solvent change will, however, lead to a change in the structure of the S_N2 transition state when the nucleophile and leaving group have opposite charges, as in reaction types (c) and (f) in Table 5-4 (called *type II* S_N2 reactions).

If the activated complex of *type I* S_N2 reactions is transferred from the gas phase to solution, the solvent will interact nearly equally with the two partially charged groups Y and X, thus lowering the Gibbs energy of the activated complex. The Gibbs energy of the activated complex will vary from solvent to solvent and the rate constant changes with solvent, depending on the interaction between Y and X and the solvent. However, the relative charge density of Y and X will not be altered by a solvent change, and thus the structure of the activated complex will not be changed even though its Gibbs energy has been changed.

On the other hand, in activated complexes of *type II* S_N2 reactions, the charges on Y and X are of opposite sign, and thus the nucleophile Y and the leaving group X will

interact differently with the solvent. For instance, in protic solvents, the negatively charged group will be preferentially solvated *via* hydrogen-bond interactions, whereas a much weaker ion-dipole type interaction will occur at the positively charged group. As a result, the difference in solute/solvent interactions at both ends of the activated complex will be large enough to cause a shift in charge density between Y and X along the Y . . . C . . . X bond axis of the S_N2 transition state. In this case, not only the Gibbs energy but also the structure of the activated complex will be altered from solvent to solvent.

In other words, whether or not an S_N2 reaction has a *tight* or *loose* activated complex will not only depend upon the nature of the reactants Y and R–X, in solution it will also be affected by the nature of the solvent. Better solvation of the activated complex of a *type II* S_N2 reaction by solvents with improved EPD/EPA properties will lead to a loosening of the activated complex. Transferring this activated complex from solution to the gas phase, with subsequent loss of the charge-separation stabilizing solvation, will therefore increase its tightness; *cf.* also [499].

This *solvation rule for S_N2 reactions* can be useful in predicting the influence of a change in solvent on the structure of activated complexes. It is in agreement with studies involving leaving group heavy atom and secondary α-deuterium kinetic isotope effects, as well as theoretical calculations of solvent effects on transition-state structures. Possible limitations of this solvation rule have been discussed; see [498] and relevant references cited therein.

Although the Hughes–Ingold theory of solvent effects was first illustrated in the field of nucleophilic aliphatic substitutions and β-eliminations, it should be applicable to all other heterolytic reactions in solutions, where the activation processes are connected with creation, dispersal, or destruction of charge. Using Kosower's classification [15, 468], in the following Sections the solvent effects on other organic reactions will be discussed to give the reader a feeling for the expected solvent influence, and to help him or her in finding the right solvent for a particular reaction under study.

5.3.2 Solvent Effects on Dipolar Transition State Reactions

Reactions involving dipolar activated complexes are those in which, compared to the initial state, there are considerable differences in charge distribution. In addition to S_N1/S_N2 and E_1/E_2 reactions as described in Section 5.3.1, solvent effects on the following reactions involving dipolar activated complexes have been studied: aromatic nucleophilic (S_NAr) and aromatic electrophilic (S_EAr) reactions; aliphatic electrophilic substitution (S_E1/S_E2) reactions; aliphatic electrophilic (A_E) and aliphatic nucleophilic addition (A_N) reactions; cycloaddition and cycloreversion reactions; aldol addition reactions; as well as rearrangement, fragmentation, and isomerization reactions. A selection of typical and particularly instructive examples, taken from the vast number of examples given in the literature, shall demonstrate the usefulness of the simple qualitative Hughes–Ingold approach.

The simplest example of an aromatic nucleophilic substitution is the S_NAr reaction between 1-chloro-2,4-dinitrobenzene and piperidine, the two-step mechanism of which, given in Eq. (5-26), is now fully established [501–503]. It involves formation of a Meisenheimer-type zwitterionic intermediate *via* a dipolar activated complex, followed

by spontaneous or base-catalysed elimination of HCl to give the reaction product. Regarding this last step, the solvent can modify the relative rates of the first and second step as well as the reaction order.



(5-26)

Solvent	*c*-C_6H_{12}	C_6H_6	C_6H_5Cl	CH_3COCH_3	$HCON(CH_3)_2$	CH_3SOCH_3
k_2^{rel}	1	2	5	13	29	50

In agreement with the separation of unlike charges during the activation process, an increase in rate by up to a factor of 50 with increasing solvent polarity has been found for reaction (5-26), carried out in thirteen non-HBD solvents [503]. The absence of base-catalysis suggests that specific solvent effects are negligible in non-HBD solvents. In protic solvents, however, specific solvation of the piperidine nucleophile leads to a diminution in rate with increasing HBD ability of the hydroxylic solvents [503].

As an example of a solvent-dependent electrophilic substitution reaction, the azo coupling (S_EAr) reaction of 4-nitrobenzenediazonium tetrafluoroborate with *N*,*N*-dimethylaniline is given in Eq. (5-27) [504]. According to the two-step arenium ion mechanism, the activation process of the rate-limiting first step is connected with the dispersion of the positive charge. This should lead to a decrease in rate with increasing solvent polarity.



(5-27)

Solvent	CH_3NO_2	CH_3CN	CH_3COCH_3	$HCON(CH_3)_2$	CH_3SOCH_3	HMPT
k_2^{rel}	4748	509	110	7	4	1
DN	2.7	14.1	17.0	26.6	29.8	38.8

The solvent-induced change in rate is, however, much larger than expected from the relatively small difference in polarity between nitromethane and hexamethylphosphoric triamide. This, together with the correlation between rate decrease and increase in the solvent donor number *DN* (*cf.* Table 2-3 in Section 2.2.6), suggests that specific solvation and stabilization of the diazonium ion by EPD solvents play a dominant role in the reaction (5-27). Very likely, formation of an EPD/EPA complex between the reactants in a rapid preequilibrium step precedes the rate-controlling first step [504, 792].

The addition of bases (*e.g.* tertiary amines) can induce proton abstraction from the arenium ion intermediate in the second reaction step. Investigation of base-catalysed azo coupling reactions with *N*,*N*-dialkylanilines in different organic solvents have shown that the solvent itself can abstract the proton from the arenium ion intermediate. In solvents of relatively high basicity (CH_3OH, CH_3SOCH_3), neither base-catalysis nor a kinetic isotope effect was observed. In less basic solvents (CH_3NO_2), however, the reaction is strongly accelerated by base and exhibits a substantial deuterium isotope effect [505]. In other words, a change in solvent can also induce a change in the rate-limiting step of azo coupling reactions.

The influence of the reaction medium on azo coupling reactions has been reviewed by Zollinger *et al.* [506].

The S_E2 cleavage of tetramethyltin with iodine is an aliphatic electrophilic substitution reaction which is subject to strong solvent dependence [507–509]. The rate of iodinolysis of $(CH_3)_4Sn$ in acetonitrile at 25 °C is more than 10^5 times faster than that in tetrachloromethane (carried out at 50 °C owing to its slow rate in this solvent) [509].



$$I^{\ominus} + I\text{-}CH_3 + {}^{\oplus}Sn(CH_3)_3 \longrightarrow I\text{-}CH_3 + I\text{-}Sn(CH_3)_3 \qquad (5\text{-}28)$$

Solvent	CCl_4	C_6H_5Cl	CH_2Cl_2	CH_3COCH_3	C_2H_5OH	CH_3CN
k_2^{rel}	1[a)]	7.4	252	739	47800	274000

[a)] at 50 °C.

The fact that the second-order rate constant is strongly solvent-dependent is in agreement with a highly dipolar activated complex as shown schematically in Eq. (5-28) by the S_E2(back) mechanism. Neither reactant has a dipole moment; the degree of charge separation in the activated complex is estimated to be between 0.8 and 1 [509]. A fast preequilibrium formation of an EPD/EPA complex between the reactants, followed by a rate-limiting iodinolysis of the alkyl metal by electron-transfer from the alkyl metal donor to the iodine moiety to give an intermediate radical ion pair, $(CH_3)_4Sn^{\oplus\cdot}$, $I_2^{\ominus\cdot}$, is also consistent with the observed solvent-dependence [509]. In S_E2 reactions, solvents not only affect the reaction rates, but also the dichotomy between S_E2(front), S_E2(back), and S_Ei mechanisms, as well as selectivity $S(R/CH_3)$ in the cleavage of unsymmetrical alkyl metals such as $(CH_3)_nSnR_{4-n}$ [508].

Solvent effects on the rates of organotin and organomercury alkyl exchange reactions have been studied and reviewed by Petrosyan [510]. It should be mentioned that

the solvent influence on the reactivity of organometallic compounds was first studied by Ziegler *et al.* in 1929/30. He showed that organolithium compounds react faster in diethyl ether than in hydrocarbons such as benzene and cyclohexane [511].

Now, some addition reactions will be considered, the solvent dependences of which have been reviewed [77, 78]. Addition of uncharged electrophiles (*e.g.* Br_2, ArS—Cl, NO—Cl, R—CO_3H) to carbon–carbon multiple bonds leads to the development of a small, usually dispersed charge in the activated complex. In more polar solvents, this is accompanied by a slight rate acceleration. In reactions with substantial charge development in the activated complex, larger rate accelerations with increasing solvent polarity are observed.

A strongly solvent-dependent electrophilic reaction is the addition of halogens to alkenes [79–81] and alkynes [81a]. In a rapid equilibrium, a loose transitory EPD/EPA complex (1:1) between halogen and alkene is formed [512]. This is followed by the rate-determining step, which involves an S_N1-like unimolecular ionization to form a halonium intermediate which can be either symmetrical or unsymmetrical. This then reacts with a nucleophile Nu: to give the products; *cf.* Eq. (5-29).



(5-29)

Solvent	CCl_4	1,4-Dioxane	CH_3CO_2H	CH_3OH	H_2O
k_2^{rel}	1	51	4860	$1.6 \cdot 10^5$	$1.1 \cdot 10^{10}$

It has been shown that in the case of bromine addition to 1-pentene in solvents of different polarity, the overall rate constant varies by a factor of 10^{10} (!) [81]. This dramatic solvent effect has been taken – together with other findings – as strong evidence for the so-called Ad_EC1-mechanism, which involves considerable charge separation in the activation step. It has also been demonstrated that protic solvents enhance this addition by a specific electrophilic solvation of the anionic part of the activated complex [81]. It appears that in the rate-determining step of alkene brominations there is also a small specific nucleophilic solvent contribution [513]. In addition it should be mentioned that, not only the rate, but also the stereospecifity of this halogen addition is strongly solvent-dependent (*cf.* Section 5.5.7) [79, 81].

The rate of the analogous bromine addition to 1-hexyne is *ca.* 10^4 times faster in methanol/water (50:50 cL/L) than in acetic acid [81a].

In the same manner, the second-order rate constant for the reaction of cyclohexene with chloro-2,4-dinitrophenylsulfane increases with increasing solvent polarity [82].



(5-30)



Solvent	CCl_4	$CHCl_3$	CH_3CO_2H	$ClCH_2CH_2Cl$	$C_6H_5NO_2$
k_2^{rel}	1	605	1370	1380	2800

Neutral reactants form a dipolar activated complex, which reacts through a thiiranium intermediate [83, 515] to give the products. Similar solvent effects have been observed in the addition of chloro-phenylsulfane to alkenes [514].

A reaction corresponding to Eq. (5-30) is the addition of nitrosyl chloride to alkenes such as cyclohexene or styrene [84, 85]. The reaction seems to be faster in polar solvents (*e.g.* nitrobenzene and trichloromethane) than in less polar solvents (*e.g.* toluene and tetrachloromethane). This is consistent with the view that the reaction involves an electrophilic attack of $NO^{\delta\oplus}—Cl^{\delta\ominus}$. The reaction was, however, also found to be very slow in diethyl ether. Presumably, this is due to strong bonding of the NO^+ cation to the ether oxygen atom [84].

The epoxidation of alkenes with peroxycarboxylic acids gives the corresponding oxirane by an electrophilic 1,1-addition mechanism as outlined in Eq. (5-31) [77, 86, 86a, 87, 516].



(5-31)

Solvent	$(C_2H_5)_2O$	1,4-Dioxane	n-C_6H_{14}	C_6H_6	CH_2Cl_2	$CHCl_3$
k_2^{rel} for R = C_6H_5 and cyclohexene [86a]	1	2.5	6.2	40	58	122

In the reactions of cyclohexene with peroxybenzoic acid (R = C_6H_5) [86a] and of trimethylethene with peroxyacetic acid (R = CH_3) [87], a moderate increase in rate with increasing solvent polarity was observed. These experimental results and recent DFT calculations [86] are in agreement with the asynchronous formation of a slightly dipolar spiro-geometrical activated complex, as proposed in Eq. (5-31). The highest rates for these reactions are obtained in non-HBD halogen-containing solvents (CH_2Cl_2, $CHCl_3$). The slowest rates are found in solvents capable of intermolecular association with the peroxycarboxylic acid (ethers). The strength of intermolecular peracid–solvent interaction increases with increasing solvent basicity [78, 87].

The addition of peroxyacetic acid to the heteronuclear C═O double bond of cyclohexanone (*i.e.* the first step of the Baeyer–Villiger oxidation reaction) also exhibits a small solvent dependence [517].

Nucleophilic additions to carbon–carbon triple bonds are also subject to solvent influence, as shown by the example given in Eq. (5-32) [88].



(5-32)

Solvent	*c*-C_6H_{12}	$(C_2H_5)_2O$	C_6H_6	CH_3OH	1,4-Dioxane	CH_3CN
k_2^{rel}	1	2	10	65	73	865

The second-order rate constant for the reaction between methoxycarbonylacetylene and piperidine increases with increasing solvent polarity. This can be attributed to the increased solvation of the strongly dipolar activated complex, which is formed from neutral molecules [88]. Analogous solvent effects have been observed for the nucleophilic addition of aziridine to 3-dimethylaminopropynal [89] and the addition of diethylamine to β-alkoxyvinyl methyl ketones [793].

Concerted $[_\pi 2_s + _\pi 2_s]$ cycloadditions are, in principle, forbidden by orbital symmetry [90]. This restriction is bypassed when these reactions occur *via* zwitterions or biradicals, or by the symmetry-allowed $[_\pi 2_a + _\pi 2_s]$ process. Since cycloadditions proceeding through zwitterionic intermediates or dipolar activated complexes should be affected by solvent polarity, the investigation of the solvent effects on rates can be of considerable value when considering potential models for the activated complex and the reaction mechanism [91–93]. The possible solvent effects on one-step and two-step cycloaddition reactions are shown schematically in Fig. 5-6 [92]*).

In the case of one-step cycloaddition reactions involving an activated complex with a different dipolarity than the reactants, an increase in solvent polarity should enhance the reaction rate (*cf.* Fig. 5-6a). However, since two-step cycloadditions are consecutive reactions, the solvent effect depends on the relative size of $\Delta G_I^{\neq}$ and $\Delta G_I^{\neq\prime}$ or of $\Delta G_{II}^{\neq}$ and $\Delta G_{II}^{\neq\prime}$ (*cf.* Fig. 5-6b). If the formation of the zwitterionic intermediate is irreversible, and $\Delta G^{\neq} \gg \Delta G^{\neq\prime}$, then the first step is rate-determining in all solvents. Consequently, there is a rate acceleration with increasing solvent polarity. When $\Delta G^{\neq} \ll \Delta G^{\neq\prime}$, this behaviour is reversed. If ever $\Delta G^{\neq} \approx \Delta G^{\neq\prime}$, then only relatively

* Some definitions may be of importance for this and further discussions. A *two-step* reaction is one which takes place in two distinct kinetic steps, *via* a stable intermediate. A *one-step*, *concerted* reaction is one which takes place in a single kinetic step. A *synchronous* reaction is a concerted reaction in which all the bond-making and bond-breaking are parallel processes, having proceeded to the same extent in the activated complex. A *two-stage* reaction is a concerted but non-synchronous reaction. That is, some of the bond-making and bond-breaking processes take place mainly in the first half of the reaction (between reactants and activated complex), while the rest take place in the second half of the reaction (between activated complex and products). *Cf.* [116] and the references cited therein.



Fig. 5-6. Gibbs energy diagram for *(a)* one-step cycloaddition reactions proceeding via a dipolar activated complex, and *(b)* two-step cycloaddition reactions proceeding via a zwitterionic intermediate, in both nonpolar (solvent I) and polar solvents (solvent II) [92]. For the sake of simplicity, no notice is taken of the different solvation of the initial reactants.

small, positive or negative, solvent effects will be observed. If the formation of the zwitterionic intermediate is reversible, then a rate increase with increasing solvent polarity will always be obtained. Under these circumstances, the distinction between a one-step and a two-step mechanism can be made only (if at all) when the solvent effects are large. Additionally, the activated complexes of the first stage of two-step reactions and of one-step reactions presumably differ only slightly in structure and dipolarity. Provided that the first step in the two-step reaction is rate-determining, the expected solvent effects for both reactions should be similar in direction and extent [92]. Three typical examples shall demonstrate solvent effects on [2 + 2] cycloadditions; for more examples, see reference [92].

A prototype cycloaddition involving a zwitterionic intermediate is the reaction of *n*-butyl vinyl ether with tetracyanoethene, as shown in Eq. (5-33) [94–98].

The influence of the solvent on the cycloaddition rate constants for electron-rich alkenes is of an unusual magnitude, as shown by $k_2(CH_3CN)/k_2(C_6H_{12}) = 2600$ for *n*-butyl vinyl ether, 10800 for 1-ethoxybutene, and 29000 for 1-methoxy-4-propenylbenzene (anethole) [94]. These ratios correspond to a $\Delta\Delta G^{\neq}$ of up to 23 kJ/mol (5.5 kcal/mol) for the increase in solvation during activation. The formation of the zwitterionic intermediate is assumed to be both reversible and rate-controlling. The dipole moments of reactants and adduct, as well as the calculated dipole moments of the activated complex and the zwitterion, reveal a notable increase in charge separation, corresponding to the large solvent effect obtained*). The dipole moment of the activated

* The calculation of the dipole moment of the activated complex was based on an electrostatic model of Kirkwood, Laidler and Eyring (*cf.* Section 5.4.3) [2, 11]. The value for the dipole moment of the zwitterion was estimated as the vector sum of the partial moments [94].



(5-33)

complex is larger than expected for the transition state of a concerted one-step cycloaddition reaction. The transition state dipole moment is two-thirds of the zwitterion dipole moment, which is reasonable because the charge separation is not yet completed in the transition state for zwitterion formation. Apart from solvent dependence of the rate constants, additional evidence for a two-step mechanism and formation of a zwitterionic intermediate in reaction (5-33) can be obtained by examining the effect of structural variation of the enol ether on the rate [94], the activation parameter [94], and the solvent dependence of the activation volume [95, 96]. The intermediate 1,4-dipole can also be trapped using alcohols or 1,4-dipolarophiles [97, 98]. Further agreement with the two-step mechanism is the lack of solvent-dependent stereospecificity for the cycloaddition reaction [94]. Rotation of the assumed zwitterionic intermediate competes with the cyclization to give the cyclobutane product. Similarly large solvent effects on rate have been observed for tetracyanoethene additions to ethyl propenyl thioether [518] and verbenene, a *trans*-fixed 1,3-diene [519]. The [2 + 2] cycloaddition of *cis*- and *trans*-1,2-bis(trifluoromethyl)-1,2-dicyanoethene to *tert*-butyl vinyl thioether also proceeds *via* a zwitterionic intermediate, corresponding to a rate acceleration of $k_2(CH_3CN)/k_2(CCl_4) = 2160$ [99]. Further 1,4-dipolar zwitterionic intermediates have been postulated for the thermal [2 + 2] cycloadditions of tetracyanoethene to styrene [99a] and 1,1-diarylbutadienes [520], as well as for the cycloaddition of tris(methoxycarbonyl)ethene to 4-(dimethylamino)styrene [521] on the basis of the rate changes observed upon altering the solvent polarity.

In contrast to reaction (5-33), the rate for the cycloaddition reaction of diphenylketene to *n*-butyl vinyl ether shows a much smaller solvent dependence; *cf.* Eq. (5-34) [100].

$\mu = 5.9 \cdot 10^{-30}$ C m (1.76 D)



(5-34)

$\mu = 4.2 \cdot 10^{-30}$ C m (1.25 D) $\mu = 10.1 \cdot 10^{-30}$ C m (3.20 D)

Solvent	*c*-C_6H_{12}	C_6H_5Cl	CH_3COCH_3	C_6H_5CN	CH_3CN
k_2^{rel}	1	13	43	63	163

The smaller rate increase with increasing solvent polarity, the large negative activation entropy, as well as electronic and steric substituent effects and the stereospecificity of this reaction, are all in agreement with a concerted, but non-synchronous one-step mechanism. The unequal bond formation in the activated complex creates partial charges, which are stabilized by increasingly polar solvents. Reaction (5-34) should be solvent-independent if the dipole moment of the activated complex approaches $10.1 \cdot 10^{-30}$ Cm (*i.e.* sum of the dipole moments of the reactants), a figure which corresponds to the dipole moment of the product. The solvent dependence obtained shows that there must be a larger charge separation in the activated complex than in the adduct [100]. A similar solvent effect was reported for the dimerization reaction of dimethyl ketene ($k_2(CH_3CN)/k_2(CCl_4) = 29$), in agreement with a concerted one-step mechanism with charge separation in the activation step [101].

In the reaction of dimethylketene with enamines such as *N*-isobutenylpyrrolidine, the two-step process *via* a zwitterion (k_I) competes successfully with the concerted one-step mechanism (k_C), leading to cyclobutanones and 2:1 adducts, respectively [102]. In contrast to the activated complex formed from vinyl ethers, there is superior stabilization of the positive charge in the zwitterion, $>C^{\delta\oplus}\cdots NR_2$ *vs.* $>C^{\delta\oplus}\cdots OR$; *cf.* Eqs. (5-33) and (5-34). The experimental results for the cycloaddition of dimethylketene to *N*-isobutenyl-pyrrolidine show that k_I depends to a much higher extent on solvent polarity than k_C, the rate constant of the concerted reaction. In acetonitrile, the zwitterionic 1,4-dipole is produced 560 times faster than in cyclohexane, while the same solvent change accelerates the concerted reaction only 36-fold [102].

Graf [103] originally proposed a two-step mechanism for the cycloaddition of chlorosulfonyl isocyanate to alkenes. This leads to the 1,4-dipole shown in Eq. (5-35), which can then ring close to give a β-lactam (and as by-product an unsaturated amide *via* a proton shift from R^1 or R^2 to $N^{\ominus}$). Moriconi [104], on the other hand, has proposed a nearly concerted, thermally allowed $[_\pi 2_s + _\pi 2_a]$ cycloaddition, probably initiated by a π-complex formation, and proceeding through the dipolar activated complex shown in the lower part of Eq. (5-35).

Among the evidence cited in favour of the two-step mechanism (for a review, see reference [105]) is the marked increase in reaction rate produced by polar solvents



(5-35)

Solvent	n-C_6H_{14}	$(C_2H_5)_2O$	$CHCl_3$	CH_2Cl_2	$C_6H_5NO_2$	CH_3NO_2
k_2^{rel}	1	31	250	1700	5000	18800

for the reaction of chlorosulfonyl isocyanate with 2-ethyl-1-hexene ($R^1 = C_2H_5$; $R^2 = n$-C_4H_9) [106, 107]. *Ab initio* MO calculations have shown that the [2 + 2] cycloaddition between *cis*- or *trans*-alkenes and isocyanates to yield *β*-lactams takes place by a concerted mechanism in the gas phase, with retention of the alkene configuration in the product. However, increasing solvent polarity contributes to the asynchronicity of the reaction to such an extent that the mechanism changes from a concerted to a two-step process involving a zwitterionic intermediate, with a consequent loss of alkene stereospecificity, as observed in the reaction between chlorosulfonyl isocyanates and vinyl ethers [794].

Not only thermal [2 + 2] cycloaddition reactions but also the corresponding [2 + 2] cycloreversions are subject to large solvent effects. A good example is the thermal decarboxylation of the *β*-lactone *trans*-3-*t*-butyl-4-phenyloxetan-2-one, as described in Eq. (5-36).



(5-36)

$\mu \approx 14 \cdot 10^{-30}$ C m (4.2 D) $\mu \approx 27 \cdot 10^{-30}$ C m (8.2 D) [524] $\mu \approx 0$ C m

Solvent	Decalin	$C_6H_5OCH_3$	C_5H_5N	Propylene carbonate	$HCONHC_6H_5$
k_2^{rel}	1	4	14	98	438

The substantial rate increase with increasing solvent polarity is consistent with a two-step route *via* a highly dipolar activated complex [522]. This mechanism is also



Fig. 5-7. Schematic Gibbs energy diagram for a general aldol addition of enolates to carbonyl compounds in both *(a)* nonpolar solvents, and *(b)* polar solvents, according to Heathcock [525].

supported by the negative volume of activation, $\Delta V^{\neq} = -28$ cm^3/mol, although the volume of reaction is $+52$ cm^3/mol! [523]. The thermal decarboxylation of 3,3-difluoro-4,4-dimethyloxetan-2-one, which could be studied in solution *and* in the gas phase, is even 3600 times faster in acetonitrile than in the gas phase [524a]. *Ab initio* MO calculations have shown that the mechanism in the gas phase is obviously quite different from that in solution. Experimental and computational kinetic data point to a concerted mechanism in the gas phase (via biradicaloid or non-Woodward–Hoffmann topological activated complexes), whereas even in moderately polar solvents zwitterionic intermediates must be involved [524a,b].

One very important addition reaction is the aldol addition of enolate ions to carbonyl double bonds to give β-hydroxy aldehydes (aldols) or β-hydroxy ketones (ketols). According to Fig. 5-7, the negative charge is localized on one oxygen atom in both the reactants and products of the aldol addition. In the activated complex, however, the negative charge is shared between two oxygen atoms. In polar solvents, reactants and products are expected to be more stabilized by solvation than the activated complex with its dispersed charge. Thus, both aldol addition and retro-aldol reaction should be faster in nonpolar than in polar solvents [525]. Enolate, activated complex, and aldolate are considered as monomeric species, which is certainly an oversimplification. Nevertheless, the arguments regarding charge delocalization during the activation process should still apply for oligomers, and the conclusion that nonpolar solvents should accelerate aldol additions (and aldol reversals) is still valid.

Simple, clear-cut examples of aldol reactions exhibiting such solvent effects are scarce. Heathcock *et al.* [526] have reported that the *erythro* $\rightarrow$ *threo* equilibration of lithium aldolates (*via* retro-aldol reaction) is much faster in pentane than in tetrahydrofuran or diethyl ether.



(5-37)

Thus, the *erythro* lithium aldolates given in Eq. (5-37) (R = Me, Et, *n*-Pr, *n*-Bu) equilibrate to their *threo* counterparts in less than two hours at 25 °C in pentane ($t_{1/2} = 45$ min for the aldolate with R = CH_3), whereas in diethyl ether the rate of equilibration is much slower ($t_{1/2} = 8$ hours for the aldolate with R = CH_3) [525, 526].

We shall conclude this Section with four other examples of rearrangements, fragmentations, and isomerizations for solvent-dependent reactions involving dipolar activated complexes.



(5-38)

Solvent	Methylcyclohexane	C_6H_6	1,4-Dioxane	CH_3CN	C_2H_5OH	$CHF_2CF_2CH_2OH$
k_1^{rel}	332	121	73	25	11	1

A study of the solvent effects on the rate of thermal racemization of chiral allyl sulfoxides has revealed that polar solvents significantly decelerate the racemization [108]. The reaction proceeds by way of a reversible and concerted rearrangement: achiral allyl sulfenates are formed as intermediates and an intramolecular α, γ-shift of the allyl group between the sulfoxide oxygen and sulfur termini occurs as shown in Eq. (5-38).

The proposed mechanism is in accordance with the observed solvent dependence of the reaction. Whereas the dipolar sulfoxide is expected to be more strongly solvated with an increase in solvent polarity, the less dipolar sulfenate should be relatively insensitive to such a solvent change. Stabilization of the sulfoxide, relative to the less dipolar activated complex (which should be similar to the sulfenate intermediate), increases the enthalpy of activation, $\Delta H^{\neq}$. This is reflected in the necessity of breaking increasingly strong solute–solvent interactions. On the other hand, because desolvation on activation is expected to increase the degrees of freedom in the system, a more positive $\Delta S^{\neq}$ is expected to work in the opposite direction and effect a compensating increase in k_1 with

increasing solvent polarity. These suppositions are confirmed by experiment [108]. The sulfenate–sulfoxide equilibrium is shifted towards the more dipolar sulfoxide with increasing solvent polarity, as measured by the corresponding equilibrium constants [108]. Monte Carlo simulations of the solvent influence on this [2,3]sigmatropic rearrangement are essentially in agreement with this mechanism, indicating that the activated complex is more product-like than reactant-like [795].

The amine-catalyzed thermal decomposition of *tert*-butyl peroxyformate [110] represents an example of a solvent-dependent heterolytic fragmentation reaction (for a review, see reference [109]). This decomposition reaction occurs *via* a radical transition state in the absence of bases, with $\Delta H^{\neq} = 159$ kJ/mol and $\Delta S^{\neq} = +63$ J/K · mol in chlorobenzene, according to Eq. (5-39a). However, in the presence of pyridine, the reaction rate increases uniformly with increasing solvent polarity. This indicates a change in reaction mechanism from a radical to a heterolytic fragmentation pathway as shown in Eq. (5-39b).

The activation parameters obtained for reaction (5-39b) in chlorobenzene are in accordance with a dipolar activated complex involving considerable charge separation:



(5-39a)

(5-39b)

Solvent	n-C_7H_{16}	CCl_4	C_6H_6	$CHCl_3$	CH_3NO_2	$C_6H_5NO_2$
k_2^{rel} for Eq. (5-39b)	1	4	14	30	110	140

$\Delta H^{\neq} = 64$ kJ/mol; $\Delta S^{\neq} = -96$ J/K · mol [110]. Thus, an increase in charge separation in the activation step should lead to more strongly orientated solvent molecules around the dipolar activated complex, as evidenced by the larger negative entropy of activation for reaction (5-39b). The solvent-dependence of the amine-catalyzed fragmentation of 2-*tert*-butylperoxy-2-methylpropanoic acid can be explained in a similar manner [111].

Push-pull substituted azobenzenes are nice examples of solvent-dependent (*E*)/(*Z*)-isomerization reactions [527–530, 561, 729]. For instance, the thermal *cis*-to-*trans* isomerization of 4-(dimethylamino)-4′-nitroazobenzene exhibits a rate enhancement of *ca.* 10^5 on changing the solvent from *n*-hexane to formamide [528].

Two mechanisms have been proposed for this isomerization reaction; *cf.* Eq. (5-40). One involves rotation around the N=N double bond and includes complete π-bond rupture to give a highly dipolar activated complex. The other involves inversion at one (or both) of the azo-nitrogen atoms with simultaneous $sp^2 \rightarrow sp$ rehybridization during the activation process. The π-bond remains intact.



(5-40)

Solvent	n-C_6H_{14}	C_6H_6	CH_2Cl_2	CH_3COCH_3	CH_3OH	$HCONH_2$
k_1^{rel}	1	5	450	4760	$2.5 \cdot 10^4$	$8.2 \cdot 10^5$
$\Delta V^{\neq}/(cm^3 \cdot mol^{-1})$	$-3^{a)}$	$-22^{a)}$	−29	−30	−27	

a) At 40 °C.

This huge solvent effect was first interpreted as an indication of the dipolar character of the activated complex, thus supporting the rotation mechanism. The isomerization rate is also considerably accelerated by external pressure in benzene and in polar solvents, but is little affected in *n*-hexane; *cf.* the negative activation volumes, $\Delta V^{\neq}$, due to electrostriction during the activation process (see Section 5.5.11 for definitions). From this, it was concluded that the isomerization mechanism changes from inversion in *n*-hexane to rotation in benzene and polar solvents [528]. That is, in the absence of solvational stabilization of the rotational activated complex, the inversion route becomes dominant (*n*-hexane; gas-phase). Other observations, however, indicate that a modification of this dual mechanism seems to be necessary. Taking into account that the inversional activated complex may also have some dipolar character due to resonance interactions of its push-pull substituents, the observed solvent, pressure, and substituent effects on rates can also be explained by means of a pure inversion mechanism [561, 729]. However, the two-route isomerization mechanism of push-pull substituted azobenzenes, *i.e.* inversion in nonpolar aprotic solvents and rotation in polar protic ones, was unequivocally proven by concave-up Arrhenius plots for reactions in solvents of medium polarity [527–529]. A change in mechanism from inversion to rotation has also been observed in supercritical (sc) carbon dioxide as solvent with small amounts of cosolvents added (1–5 cmol/mol). While in sc-CO_2/methanol the inversion mechanism dominates, the rotation mechanism is operative in more polar sc-CO_2/hexafluoro-*i*-propanol [530]. Unsubstituted and push-push substituted (*Z*)-azobenzenes, as well as (*Z*)-*N*-arylazomethines (*e.g.* *N*-benzylideneanilines), generally isomerize by way of the less solvent-dependent inversion mechanism.

We shall conclude this Section with an example of a reaction that undergoes an extreme rate acceleration with an increase in solvent polarity. Thermolysis of α-chlorobenzyl methyl ether in a series of non-nucleophilic, non-HBD solvents shows rate variations up to 10^5, encompassing a $\Delta G^{\neq}$ range of 30 kJ/mol (7 kcal/mol) [112]. This dramatic solvent effect is best explained by a mechanism involving ionization of the C—Cl bond to form an ion pair, followed by a nucleophilic attack by $Cl^{\ominus}$ on CH_3 to give an aldehyde and chloromethane; *cf.* Eq. (5-41).



(5-41)

Solvent	CCl_4	C_6H_5Cl	$CHCl_3$	$C_6H_5NO_2$	CH_3CN
k_1^{rel}	1	36	750	22000	166000

The experimental activation entropies are all negative, showing a higher degree of solvent organization in the transition state than in the initial state.

The thermal decomposition of alkyl chlorosulfites (R–O–SO–Cl → R–Cl + SO_2), the prototype of an S_Ni reaction, is also strongly accelerated in polar solvents: 353-fold upon a solvent change from *i*-octane to 1,4-dioxane [796a]. This is in agreement with a mechanism involving ionization either (a) of the S–Cl bond to yield a sulfinyl cation (R–O–SO^+) and Cl^- (with R = primary alkyl), or (b) of the R–O bond to give a carbenium ion and a chlorosulfinyl anion ($^-$O–SO–Cl) (with R = tertiary alkyl) [796b]. The generation of both ion pairs is facilitated in polar solvents, according to reaction type (a) in Table 5-4.

5.3.3 Solvent Effects on Isopolar Transition State Reactions

Reactions involving isopolar activated complexes (neither dipolar nor radical in nature), normally exhibit only small solvent rate effects. This is because charge distribution in the activated complexes and the reactants is very similar. Owing to the different polarizabilities of the initial and transition state molecules, there are changes in the solute/solvent dispersion interactions leading to small rate changes. Many reactions involving isopolar transition states are electrocyclic, sigmatropic, cheletropic, or cycloaddition reactions, all of which are representative of a larger class of concerted reactions known as "pericyclic reactions". Woodward and Hoffmann have suggested that the course of these reactions is controlled by the symmetry of the molecular orbitals of the reactants and products [90, 113]. Changing either the substituent or the medium usually has little effect on the rates of pericyclic reactions. This fact has been used by some chemists as a criterion for establishing the mechanism of such reactions. Examples illustrating the lack of solvent sensitivity of pericyclic and related reactions are presented. Unfortunately,

because no solvent effects are expected for these reactions, thorough investigations in a large variety of solvents are lacking.

Some concerted $[m+n]$ cycloaddition reactions*), however, have been well examined [90–93], *e.g.* Diels–Alder reactions [114–116] and 1,3-dipolar cycloaddition reactions [117–120, 541–543]. Both of these reactions have been established as concerted $[4+2]$ cycloaddition reactions.

The solvent effect on the bimolecular rate constant of a Diels–Alder reaction is usually quite small. As a rule, in going from nonpolar to polar solvents, the rate constant increases only by a factor of about 3 . . . 20 [34, 35, 121–130, 531–537].

Typical examples are the cycloaddition of isoprene and maleic anhydride as shown in Eq. (5-42) [127], and the dimerization of cyclopentadiene (*cf.* Table 5-1 in Section 5.2) [33, 34].



(5-42)

Solvent	$(i\text{-}C_3H_7)_2O$	C_6H_6	C_6H_5Cl	CH_3NO_2	$C_6H_5NO_2$	*o*-Dichlorobenzene
k_2^{rel}	1	3.5	5.0	6.6	11	13

It is clear from the above figures that the activated complex is not much more dipolar than the initial state. Neutral reactants produce a neutral product in a single, concerted, often synchronous step. Although dienes and dienophiles may have a dipole moment, these dipoles are usually incorporated unchanged into the product, and there is no reason to believe that they increase or decrease greatly during the reaction. The small solvent effects observed are in agreement with this picture of the Diels–Alder reaction.

Further examples of Diels–Alder cycloaddition reactions with small or negligible rate solvent effects can be found in the literature [531–535]. The thermolysis of 7-oxabicyclo[2.2.1]hept-5-ene derivatives is an example of a solvent-independent *retro*-Diels–Alder reaction [537]. For some theoretical treatments of the solvent influence on Diels–Alder cycloaddition reactions, which, in general, confirm their small solvent-dependence, see references [536, 797–799].

Nevertheless, there are some examples known with larger, although still moderate, solvent effects [124, 125, 129, 538–540]. Over a range of solvents from *o*-xylene to trichloromethane, the reaction rates for the addition of tetracyanoethene to anthracene have been found to increase by a factor of 70 [125]. In the case of the reaction between cyclopentadiene and acrolein, changing the solvent from ethyl acetate to acetic acid causes a 35-fold acceleration in rate [129]. A strongly dipolar activated complex is unlikely, as reflected by this small sensitivity to solvent. These data are more consistent with the following mechanism: first the diene and dienophile form an EPD/EPA com-

* $[m+n]$-Cycloaddition reactions involve the addition of a molecule (or fragment) with m π-electrons to one with n π-electrons to produce a new ring in a concerted, but not necessarily synchronous, orbital-symmetry controlled process [90].

plex, this is then converted directly into the adduct *via* an electron-rich, polarizable activated complex. Some Diels–Alder reactions experience a significant change in activation enthalpy when the solvent is changed. When the relative solvation enthalpies for the reactants are determined in EPD and EPA solvents by calorimetry, it is found that in EPD solvents the reactants are stabilized, whereas the more electronegative EPA solvents stabilize the electron-rich activated complex [128]; see however references [130, 800] for a criticism of this approach.

Another explanation for enhanced solvent effects in Diels–Alder reactions in HBD solvents is that carbonyl-containing reactants and activated complexes can form R–O–H $\cdots$ O=C hydrogen bonds to a different extent, depending on the polarization of the C=O group, which is larger in the activated complex according to C^+–O^-. For example, the Diels–Alder reaction between cyclopentadiene and butenone was reported to be twelve times faster in methanol than in *i*-octane, but in water the rate is 730 times faster than in *i*-octane ($\Delta\Delta G^{\neq} \approx -16$ kJ/mol) [801]. This remarkable rate acceleration in water cannot explained simply in terms of solvent polarity differences. *Ab initio* MO calculations on the cycloaddition of cyclopentadiene and butenone have shown that during the activation process the hydrogen bond between butenone and water is enhanced by ca. 8 kJ/mol [797]. In addition to this H-bond effect, a hydrophobic effect is certainly operative in water. In the activated complex for Diels–Alder reactions, two hydrocarbon surfaces must come together, and this aggregation is facilitated in water [801]; see also Sections 2.2.7 and 5.5.8. However, there are limitations associated with the use of water as a solvent for Diels–Alder reactions because the vast majority of organic compounds are insoluble in water, and water precludes the use of water-sensitive reactants. Further examples of water-mediated Diels–Alder reactions can be found in a recent review [802].

The particular influence of water as solvent on Diels–Alder reactions can be altered by the addition of ionophores (e.g. LiCl, NaCl, $LiClO_4$, *etc.*) [801]. For the 'cyclopentadiene + butenone' reaction, the rate increases 2.5-fold on addition of LiCl (4.86 M), while a marginal decrease is observed on addition of guanidinium chloride. Obviously, lithium chloride increases the hydrophobic effect, while guanidinium chloride decreases it. The effect of aqueous salt solutions can also be understood in terms of structure-making (rate-enhancing) and structure-breaking (rate-decreasing) agents [802].

Much more impressive rate accelerations for several Diels–Alder (and other) reactions have been observed by employing solutions of lithium perchlorate (up to 5 M) in diethyl ether (LPDE solutions) [802–806]. The dramatic rate accelerations found for Diels–Alder reactions in LPDE solutions appear to stem from Lewis acid catalysis by the coordinative unsaturated Li^+ ion (see the end of Section 3.1). The Lewis acid catalysis by LPDE is applicable to those Diels–Alder reactions in which the lithium cation can coordinate with suitable functional groups in the reactants (*e.g.* $Li^+ \cdots O{=}C$). Addition of lithium-specific crown ethers (*e.g.* [12]crown-4) leads to a loss of the catalytic activity of the Li^+. For a recent extensive review of salt effects on Diels–Alder reactions, see reference [802].

Rather large rate enhancements with increasing solvent polarity have also been found for *intramolecular* Diels–Alder reactions such as the cyclization of 2-furfuryl methyl fumarate shown in Eq. (5-42a) to yield lactone D. This first-order reaction is about 3200 times faster in dimethyl sulfoxide than in toluene as solvent [807].



(5-42a)

In order to form the activated complex required for the formation of product D, rotational changes of the less dipolar *anti*-form A to the more dipolar *syn*-conformer B are necessary, to give an activated complex C with more parallel bond dipoles, which is thus more dipolar and better solvated than the reactant molecule. In agreement with this explanation is the observation that the reverse *retro*-Diels–Alder reaction exhibits no large solvent effect, since the activated complex C is quite similar to the reactant D [807].

A very subtle solvent effect has been observed in the Diels–Alder addition of methyl acrylate to cyclopentadiene [124]. The polarity of the solvent determines the ratio of *endo* to *exo* product in this kinetically controlled cycloaddition reaction, as shown in Eq. (5-43). The more polar solvents favour *endo* addition.



(5-43)

Solvent	$(C_2H_5)_3N$	$(CH_3OCH_2)_2$	CH_3CN	CH_3CO_2H	CH_3OH
$\lg \frac{[endo]}{[exo]}$	0.445	0.543	0.692	0.823	0.845

Consideration of the dipolarity of the two activated complexes can explain the observed trend. If the reactants are pictured as lying in roughly parallel planes, the dipole moments for the *exo* orientation are seen to be nearly opposite in direction, whereas for the *endo* orientation they are parallel. Therefore, the net dipole moment for the *endo* transition state is greater than that for the *exo*. Thus, the solvation of the *endo* activated complex will be more pronounced as the polarity of the solvent increases. This leads to a lowering of the activation enthalpy and preferential formation of the *endo* adduct. The logarithm of the *endo*/*exo* product ratio in various solvents has been used to define an empirical solvent polarity scale [124] (*cf.* Section 7.3). Analogous solvent-dependent *endo*/*exo* product ratios have been obtained in [4 + 2]cycloadditions of cyclopentadiene to other acrylic acid derivatives [560]. Theoretical calculations on *exo*/*endo* structures for activated complexes of [4 + 2]cycloadditions have shown that the observed *endo* preference in polar solvents is due to the influence of the medium, and that secondary orbital interactions are not involved [808]. The solvent has the decisive influence on the *exo*/*endo* selectivity.

As measured by the criteria of stereospecificity, regioselectivity, kinetic isotope effects, and solvent effects [117–120, 541–543], 1,3-dipolar cycloaddition reactions represent orbital symmetry-allowed [$_{\pi}4_{s} + {}_{\pi}2_{s}$] cycloadditions, which usually follow concerted pathways*). Diels–Alder reactions and 1,3-dipolar cycloadditions resemble each other, as demonstrated by the small solvent effects on their bimolecular rate constants. In going from nonpolar to polar solvents, the rate constants of 1,3-dipolar cycloadditions change only by a factor of 2 . . . 10 [120, 131–134].

The cycloaddition of *N*-methyl-*C*-phenylnitrone to ethyl acrylate yielding a substituted isoxazolidine serves to illustrate this point [132]. In solvents of increasing polarity, this reaction, shown in Eq. (5-44), exhibits only a 5.6-fold rate deceleration [132].

Similarly small rate factors were obtained for 1,3-dipolar cycloadditions between diphenyl diazomethane and dimethyl fumarate [131], 2,4,6-trimethylbenzenecarbonitrile oxide and tetracyanoethene or acrylonitrile [811], phenyl azide and enamines [133], diazomethane and aromatic anils [134], azomethine imines and dimethyl acetylenedicarboxylate [134a], diazo dimethyl malonate and diethylaminopropyne [544] or *N*-(1-cyclohexenyl)pyrrolidine [545], and *N*-methyl-*C*-phenylnitrone and thioketones [812]. Huisgen has written comprehensive reviews on solvent polarity and rates of 1,3-dipolar cycloaddition reactions [541, 542]. The observed small solvent effects can be easily explained by the fact that the concerted, but non-synchronous, bond formation in the activated complex may lead to the destruction or creation of partial charges, connected

* Firestone postulated that many 1,3-dipolar cycloadditions occur by a two-step mechanism with a discrete spin-paired diradical intermediate [118], but his arguments were criticized by Huisgen [119]. Distinction between concerted and stepwise-diradical mechanisms cannot be made on the basis of the negligible small solvent effects obtained for most 1,3-dipolar cycloaddition reactions. References [90c, 543] give a fair discussion of this controversial point. Using thiocarbonyl ylides, $R_2C{=}S^{+}{-}CH_2^{-}$, and dimethyl dicyanofumarate as reaction partners, the first nonstereospecific, two-step 1,3-dipolar cycloaddition with ring-opened, equilibrating zwitterions as intermediates has been reported by Huisgen *et al.* [809]; for further examples, see reference [810]. When steric hindrance to bond formation at one end of the 1,3-dipole is particularly severe, then a stepwise, more solvent-dependent cycloaddition can obviously compete with the concerted process.

$\mu = 11.8 \cdot 10^{-30}$ C m (3.55 D)



(5-44)

$\mu = 5.9 \cdot 10^{-30}$ C m (1.76 D) $\mu = 8.3 \cdot 10^{-30}$ C m (2.48 D)

Solvent	$C_6H_5CH_3$	1,4-Dioxane	CH_3COCH_3	CH_3NO_2	CH_3CN	C_2H_5OH
k_2^{rel}	5.6	3.2	2.2	1.9	1.9	1

with the disappearance of the formal charges of the 1,3-dipole. A concerted mechanism, involving a small charge imbalance in the transition state, is characteristic of most 1,3-dipolar cycloaddition reactions. In the case of cycloadditions between diazomethane and certain aromatic anils to give substituted 1,2,3-triazolines, a specific protic/dipolar aprotic solvent effect was postulated, involving better solvation of the activated complex relative to the reactants in dipolar non-HBD solvents such as *N*,*N*-dimethylformamide [134].

Likewise, 1,3-dipolar (and other) cycloreversions have been studied in a variety of solvents [546–549]. As expected, only rather small or negligible solvent effects on rate have been observed.

Other examples of reactions closely related to the Diels–Alder cycloaddition reaction are the ene reactions between alkenes with allylic hydrogen atoms (ene) and compounds with a double bond (enophile) [135, 136], and the dye-sensitized photooxygenation of allylic alkenes by singlet oxygen to give allylic hydroperoxides with a shifted double bond [137–139].

The observed range of solvent effects (less than a factor of four) for the ene reaction between 3-phenyl-1-*p*-tolylpropene-(1) and diethyl azodicarboxylate, given in Eq. (5-45), is best explained by a concerted mechanism involving an isopolar six-centre transition state [136].



(5-45)

R = 4-CH_3-C_6H_4

Solvent	*c*-C_6H_{12}	1,4-Dioxane	$HCON(CH_3)_2$	CH_3CN	$(ClCH_2)_2$	$C_6H_5NO_2$
k_2^{rel}	1	1.1	2.0	2.2	3.1	3.9

The addition of singlet oxygen to 2-methyl-2-pentene occurs *via* a concerted "ene-type" mechanism as shown in Eq. (5-46)*). This is entirely consistent with the small solvent effect observed for this reaction [138]. When the solvent is changed from methanol to carbon disulfide, the rate changes by a factor of seven**). Thus, it would appear that the activated complex does not involve much charge separation [138].



(5-46)

Solvent	CS_2	$CH_3CO_2C_2H_5$	CH_3SOCH_3	CH_3COCH_3	C_6H_6	CH_3OH
β-value*)	0.022	0.04	0.07	0.08	0.10	0.16

In the photo sensitized oxygenation reactions of alkenes, not only the influence of the solvent on the reaction rate but also the effect of solvent on product distribution (*i.e.* from competing hydroperoxide, 1,2-dioxetane, and *endo*-peroxide formation) is rather small [550, 551].

In contrast to strongly solvent-dependent [2 + 2]cycloaddition reactions, which proceed through a 1,4-dipolar zwitterionic intermediate by a two-step mechanism or through a dipolar activated complex by a one-step mechanism (*cf.* Section 5.3.2, and Eqs. (5-33) to (5-35) [92, 94–107], [2 + 2]cycloadditions are also known that exhibit concerted, nearly synchronous bond formation without significant charge separation on activation in the transition state. An example is given in Eq. (5-47). Since the rate constant for this diphenylketene/styrene addition is practically independent of solvent polarity [140], it can be classed as concerted.



(5-47)

Solvent	C_6H_5Br	*o*-Dichlorobenzene	$(ClCH_2CH_2)_2O$	$HCON(CH_3)_2$
k_2^{rel}	1.2	1.1	1.1	1

The rates of [2 + 2]cycloaddition reactions between di-*tert*-butylthioketene and azomethines [141], thiobenzophenones and keteneimines [552] as well as [2 + 2 + 2]

* Since there exists a second group of allylic hydrogen atoms at the other end of the 2-methyl-2-pentene molecule, two different allylic hydroperoxides are obtained in the solvent-independent ratio of *ca.* 1:1 [138].

** Only the ratio $\beta = k_d/k_A$ of the rate constant for decay of singlet oxygen to its ground-state triplet (designated k_d) and the specific rate constant for the product-forming step (designated k_A) is readily determined experimentally [137, 138]. Therefore, the β values provide an inverse measure of the reactivity of the allylic alkene.

cycloadditions between quadricyclane and, for example, acrylonitrile [553] also reveal no significant solvent dependence.

Ab initio MO calculations for the uncatalyzed [2 + 2]cycloaddition reaction between ketenes (*e.g.* chloroketene) and carbonyl compounds (*e.g.* formaldehyde) to yield oxetan-2-ones (β-lactones) predict a relatively synchronous, concerted process, with a $[{}_{\pi}2_s + ({}_{\pi}2_s + {}_{\pi}2_s)]$ arrangement of the reactants in the activated complex and a negligible solvent effect [813].

Another remarkable example of a [2 + 2]cycloaddition reaction is the Wittig reaction of alkylidenephosphoranes and carbonyl compounds [142, 143, 554]. The solvent dependence of some Wittig reactions has been studied [144–148]. The relatively small inverse solvent effect found for the "salt-free" Wittig reaction between 4-nitrobenzaldehyde and the resonance-stabilized ylide benzoylmethylene-triphenylphosphorane – a 58-fold rate decrease with increasing solvent polarity [146–148] – is in agreement with a concerted, but not necessarily synchronous formation of C—C and P—O bonds in the rate-controlling first step, leading to a non-ionic cyclic oxaphosphetane intermediate such as that described in Eq. (5-48). Because C—C bond

$\mu = 8.0 \cdot 10^{-30}$ C m (2.41 D) [150]



$\mu = 18.2 \cdot 10^{-30}$ C m (5.45 D) [149] $\mu^{\neq} < 26 \cdot 10^{-30}$ C m (7.9 D)



(5-48)

Solvent	CCl_4	n-C_6H_{14}	CH_3OH	CH_3CN	C_6H_6	$(HOCH_2)_2$	$HCON(CH_3)_2$
k_2^{rel}	58	16	9.5	9	3.3	1.6	1

formation is more advanced than P—O bond formation in the activation process [554], small partial charges are generated. However, according to the inverse solvent effect, the activated complex must be less dipolar than the reactants. That is, the dipole moment of the activated complex should definitely be less than the sum of the reactant dipole moments. This certainly rules out the formation of a zwitterionic phosphonium betaine in

the first, rate-determining step. In addition, the somewhat solvent-dependent activation volumes are – with values of $\Delta V^{\neq} = -20 \ldots -30$ cm^3/mol – insufficiently negative for a fully zwitterionogenic activation step [555].

According to Bestmann [554], the initially formed 1,2-oxaphosphetane with an apical-located oxygen atom, undergoes a ligand rearrangement process (pseudorotation) thus bringing the bond necessary for alkene formation into the apical position. After this conversion, depending on the reaction conditions, alkene formation takes place either directly by an asynchronous cycloreversion of the oxaphosphetane, or by a preferred cleavage of the C–P bond to give equilibrating zwitterions which decompose to the products. The substituent R^2 determines the lifetime of this zwitterion and consequently the configuration of the alkenes formed. If R^2 is an electron-donating group, fast phosphane oxide elimination occurs and the (*Z*)-alkene is formed. Electron-withdrawing groups R^2 [*e.g.* $R^2 = COC_6H_5$ as in Eq. (5-48)] stabilize the betaine which now isomerizes to the thermodynamically more stable conformer yielding (*E*)-alkenes *via* phosphane oxide elimination.

Protic solvents shift the alkene (*E*)/(*Z*) ratio in the direction of the (*E*)-form. The alkene (*E*)/(*Z*) ratio of salt-free Wittig reactions is thus influenced not only by the electronic character of R^2, but also by the solvent and the stereochemistry of the formation of the 1,2-oxaphosphetane in the first rate-determining step. According to Eq. (5-48), the thermodynamically less stable (*Z*)-1,2-oxaphosphetane is formed in the first activation step. A conformational analysis of the activated complex leading to the 1,2-oxaphosphetane intermediate provides a reasonable explanation for this unexpected *cis*-selectivity [143, 556].

A complete analysis of the solvent influence on the Wittig reaction given in Eq. (5-48), based on Gibbs energies of transfer, $\Delta G_t^{\circ}(X, CH_3OH \rightarrow S)$, from methanol to thirteen other solvents for reactants, activated complex, and products has been given [148]. As confirmed by the kinetic results, this Wittig reaction is an example of a balanced reaction type [467] with similar solvation effects for both the initial and transition state, as well as for the final state. The activated complex appears to be reactant-like and product-like, as expected for an activated complex similar to a cyclic intermediate, without localized charges on specific ring atoms [148]. A more detailed discussion of the solvent (and salt) influence on reaction rates and (*E*)/(*Z*) alkene ratios of Wittig reactions can be found in reference [143].

Sigmatropic reactions, *i.e.* reactions involving migration of a σ-bond, flanked by one or more conjugated systems, to a new position within the system, are also pericyclic reactions [90, 113]. Solvent studies of such reactions have been carried out, *e.g.* the effect on rate of solvent polarity has been examined for some [3,3]sigmatropic reactions such as the Cope rearrangement of substituted 1,5-hexadienes [151, 152, 154] and the *ortho*-Claisen rearrangement of allyl aryl ethers [153, 154, 814]. Cope and Claisen rearrangements show little response to variation of the polarity of the solvent, in accordance with an isopolar activated complex. Some [1,3]- [557] and [1,5]sigmatropic reactions [558] can also be considered as isopolar transition state reactions involving rather small solvent effects on rate.

As an example, the rate of rearrangement of 4,4-dicyano-5-ethylhepta-1,5-diene changes only by a factor of 3.8, even when the solvent is changed drastically by going from cyclohexane to dimethyl sulfoxide (Eq. (5-49)) [151].



(5-49)

Solvent	*c*-C_6H_{12}	1,4-Dioxane	C_2H_5OH	CH_3SOCH_3
k_1^{rel}	1	2.0	2.6	3.8

It can be concluded, therefore, that there is negligible charge separation on activation, even when the compounds have two nitrile groups, which are ideal for stabilizing a potential carbanion in an ionic mechanism. Similar results were obtained for the Cope rearrangement of diethyl allylisopropenylmalonate [152]. It should be mentioned, however, that some 1,5-hexadienes, containing radical-stabilizing substituents in the 2,5-positions, seem to react *via* a diradicaloid pathway [155]. The analogous azo-Cope rearrangement of arylazo (α,α-dimethylallyl)malononitriles into *N*-(γ,γ-dimethylallyl)-aryl-hydrazonomalononitriles, involving simultaneous carbon–carbon bond cleavage and carbon–nitrogen bond formation, exhibits a small but significant rate increase with increasing solvent polarity (a factor of 19 on changing from CCl_4 to Me_2SO). This result suggests a concerted mechanism *via* an activated complex of low dipolarity [593].

An interesting case is the fast degenerate Cope rearrangement of the donor/acceptor-substituted 2,6-dicyano-4,8-diphenylbarbaralane, which has been studied in 26 solvents of different polarity [815]. Such degenerate Cope rearrangements are characterized by rather small activation energies in going from the reactant with a classical localized electronic structure to the activated complex with a delocalized electronic structure, corresponding to a flat double-minimum potential. The solvent-dependent thermochromism of solutions of this barbaralane suggests that dipolar polarizable solvents such as DMSO, HMPT, and DMPU stabilize the polarizable delocalized activated complex more than the less polarizable reactants. This leads to an inversion of the energetic order of reactant and activated complex, corresponding to a single-minimum potential with the former activated complex becoming the more stable species, which can be observed UV/Vis spectrophotometrically ($\lambda_{max} = 460$ nm). In most other solvents (*e.g.* cyclohexane, di-*n*-butyl ether, benzene), the normal order as obtained in the gas phase is maintained. Evidently, it is possible to visualize activated complexes not only by means of femtosecond spectroscopy [815], but also through the solvent-dependent thermochromism of barbaralanes (and semibullvalenes).

For the *ortho*-Claisen rearrangement of allyl *p*-tolyl ether shown in Eq. (5-50), the rate enhancement with increasing solvent polarity is modest, in agreement with a cyclic process involving concerted bond-making and -breaking on activation [153].



(5-50)

Solvent	*n*-$C_{14}H_{30}$	$(n\text{-}C_4H_9)_2O$	Sulfolane	$(CH_2OH)_2$	C_6H_5OH
k_1^{kel}	1	1.1	3.6	22	34

For this reaction, the rate was found to vary by a factor of 34 in going from the least polar (*n*-tetradecane) to the most polar solvent (phenol), and by a factor of 102 in going from the gas phase to the most polar solvent [153]. The small solvent and substituent effects observed suggest that a slightly dipolar activated complex must be formed during the Claisen rearrangement of this allyl aryl ether [153].

A similar slightly dipolar activated complex has been postulated for an aliphatic pendant to the aromatic Claisen rearrangement, that is for the [3,3]sigmatropic rearrangement of alkoxyallyl vinyl ethers [767]. For example, the rate of rearrangement of 6-methoxyallyl vinyl ether to 3-methoxy-4-pentenal is increased 3 . . . 68 fold upon changing the solvent from benzene to acetonitrile and methanol, respectively [767]. The first-order rate constant for the [3,3]sigmatropic rearrangement of a similar alkoxyallyl vinyl ether, methyl and sodium 8-vinyloxy-9-decenoate, increases by a factor of 56 in going from cyclohexane to trifluoroethanol [816]. A comparison with solvolytic rate data (*Y* values of Winstein; see Section 7.3) shows that the activated complexes of such [3,3]sigmatropic rearrangements are substantially less dipolar than those of S_N1 solvolysis reactions. However, in water, the rearrangement of the sodium salt is about 214 times faster than in cyclohexane. This result suggests that both hydrophobic destabilization of the reactant and H-bond stabilization of the activated complex (which contains a developing carbonyl group) are responsible for this enhanced rate acceleration [797, 814]. For a *retro*-Claisen rearrangement, a 12-fold rate increase was found for the solvent change cyclohexane → trifluoroethanol, again indicating a concerted reaction with a small partial charge separation in the activated complex [817]. A more detailed discussion of the solvent influence on Claisen rearrangements can be found in reference [814].

A reaction similar to the Claisen reaction is the [3,3]sigmatropic rearrangement of an allyl thiobenzoate into an allyl thiolbenzoate, as shown in Eq. (5-51) [156].



(5-51)

Solvent	*c*-C_6H_{12}	C_6H_5Cl	CH_3COCH_3	CH_3CN	CH_3CO_2H
k_1^{rel}	1	1.9	2.3	4.9	4.8

From the first-order rate constants obtained in different solvents (in sealed ampoules), it is apparent that this isomerization is not very sensitive to the polarity of the medium, in accordance with an isopolar, six-membered activated complex [156]. A similar small solvent effect has been observed for the [3,3]sigmatropic rearrangement of allyl S-methyl xanthate to allyl methyl dithiol carbonate [559].

Finally, the solvent dependence results for two electrocyclic reactions are mentioned. Electrocyclic reactions are generally defined as reactions involving the concerted cyclization of an n π-electron system to an $(n-2)$ $\pi + 2\sigma$-electron system, or the reverse process [90, 113, 818].

The conrotatory cyclization of *all-cis*-deca-2,4,6,8-tetraene to *trans*-7,8-dimethylcycloocta-1,3,5-triene has been studied in solvents of different polarity [157]. In agreement with a synchronous conrotatory ring closure *via* an isopolar activated complex, the solvent effect is negligible as shown by the relative first-order rate constants in Eq. (5-52).



(5-52)

Solvent	$CDCl_3$	Pyridine	CH_3CN
k_1^{rel}	1.1	1.2	1

In the case of 2-methyl-4,4-diphenylcyclobutenone, the reverse process, an electrocyclic ring opening, has also been examined in different solvents [158]; *cf.* Eq. (5-53).



(5-53)

Solvent	*c*-C_6H_{12}	C_6H_6	$CH_3CO_2C_2H_5$	C_2H_5OH	CH_3OH
k_1^{rel}	3.0	2.3	1.4	1.1	1

The small inverse dependence of the first-order rate constant on solvent polarity is in agreement with a concerted electrocyclic ring cleavage through an isopolar activated complex to vinylketene, which is converted into the corresponding ester in alcoholic solvents [158].

Desimoni *et al.* have thoroughly studied the solvent dependence of the electrocyclic ring closure of 1,2,6-triphenylhexa-1,3,5-triene (to give 1,5,6-triphenylcyclohexa-1,3-diene) and the ring opening of dimethyl 3,4-dimethyl-1,2-diphenylcyclobutene-*cis*-3,4-dicarboxylate (to yield dimethyl 2,5-dimethyl-3,4-diphenylmuconate) in 15 different solvents [819]. Both electrocyclic reactions are influenced very little by the solvent, with small rate accelerations of 1.2 and 1.8, respectively, on going from cyclohexane to methanol.

The solvent-dependence of cheletropic reactions*) has also been investigated [158a]. The thermolysis of 3-methyl-2,5-dihydrothiophene-1,1-dioxide appears to involve a concerted fission of the two σ-bonds. This is in accordance with the very small solvent effect on rate observed in six solvents of different polarity [158a].

* Cheletropic reactions are defined as processes in which two σ-bonds that terminate at a single atom are made or broken in a concerted fashion.



(5-54)

Solvent	n-$C_{10}H_{22}$	Triethylene glycol	$(H_5C_6)_2O$	H_5C_6—CO—C_6H_5	Sulfolane
k_1^{rel}	1	1.3	1.6	1.8	1.8

From these results, it is clear that this reaction meets the requirements for a pericyclic reaction involving an isopolar activated complex.

Another more recent example of a cheletropic reaction, studied in various solvents, is the addition of aryl-halocarbenes (generated photolytically from diazirines) to tetramethylethene to give the corresponding cyclopropane derivatives [820]. The addition of chlorophenylcarbene is only about three times faster in ethyl acetate than in pentane, as befits an isopolar activated complex.

The thermal rearrangement of allyl(silylmethyl) ethers, an example of a dyotropic reaction*), in which the silyl and allyl groups exchange their positions, exhibits only a very small solvent dependence; this is as expected for a concerted reaction according to Eq. (5-55) [158b]:



(5-55)

R + R = 2,2′-Biphenylylene

Solvent	Decalin	Benzene	o-Dichlorobenzene	Propylene carbonate
k_1^{rel}	1	1.1	2.8	5.1

Finally, it should be mentioned that no strict limit between reactions with dipolar and isopolar activated complexes exists. Some borderline cases with significant but relatively small charge separation in going from the initial to the transition state, with correspondingly small solvent rate effects, have been mentioned in this Section.

5.3.4 Solvent Effects on Free-Radical Transition State Reactions

According to Kosower, a third category of reactions involving free-radical activated complexes may be defined. These complexes are formed either through the creation of unpaired electrons during radical pair formation or atom-transfer reactions [15, 468] (*cf.*

* Dyotropic rearrangements are uncatalyzed, intramolecular pericyclic reactions in which two σ-bonds simultaneously migrate [158b].

Section 5.3). These two different types of free-radical reactions have been investigated for solvent effects (for reviews, see references [159–166]). Reactions in which the medium affects the reactivity of free radicals have received greater attention than radical-forming reactions. The solvent often affects the course of free-radical reactions by participating as a reactant. This is evidenced by the incorporation of the solvent into the products of the reaction. This Section, however, will be limited to those solvent effects which are truly medium effects.

Free radicals may be generated by oxidation, reduction, or by homolytic cleavage of one or more covalent bonds, such as C—C bonds (*e.g.* dimers of triarylmethyl radicals), N—N bonds (*e.g.* tetrasubstituted hydrazines), O—O bonds (*e.g.* hydroperoxides, dialkyl and diacyl peroxides, peroxycarboxylic esters), C—N bonds (*e.g.* dialkyl azo compounds), and N—O bonds (as in the thermolysis of nitrogen pentoxide O_2N—O—NO_2). Two typical examples, which have been investigated in different solvents, are given in Eqs. (5-56) and (5-57); *cf.* also reaction (5-39a) in Section 5.3.2.



(5-56)



(5-57)

The dissociation rate of the dimer of the triphenylmethyl radical*) in 28 solvents was studied by Ziegler *et al.* [167]. The decomposition rate of azobisisobutyronitrile in 36 solvents was measured by different authors [183–185, 562]. Despite the great variety of solvents, the rate constants vary only by a factor of 2 . . . 4. This behaviour is typical for reactions involving isopolar transition states and often indicates, but does not prove, a radical-forming reaction. The lack of any marked solvent effects in most free-radical forming reactions will become more apparent after an examination of some further reactions presented in Table 5-8.

Generally, it can be said that radical-forming reactions are usually not very sensitive to medium effects, because activated complexes which produce neutral radicals normally exhibit no charge separation. Even in the case of 4-methoxyphenylazo-2-methylpropanedinitrile (*cf.* Table 5-8, last entry), which is predisposed for an ionic decomposition, there is only a modest 26-fold rate acceleration in going from *n*-decane to methanol as solvent. This rate factor can be easily explained by assuming that the cleavage of the two C—N bonds of the azo group is concerted but non-synchronous [189].

---

* Ziegler *et al.* undertook their experiments with a compound which they believed to be hexaphenylethane [167]. In 1968, it was shown that the dimer of the triphenylmethyl radical is not hexaphenylethane, but 1-diphenylmethylene-4-triphenylmethyl-2,5-cyclohexadiene [168, 169] in accordance with a proposal made by Jacobson in 1905 [170].

Table 5-8. Solvent influence on rates of monomolecular decomposition of various free-radical initiators [164].

Initiators	Number of solvents	Temperature °C	Range of variation of k_1/s^{-1}	$k_1^{rel.a)}$	References
	28	0	$(1.1 \ldots 4.2) \cdot 10^{-3}$	3.9	[167]
	12	125	$(1.5 \ldots 3.5) \cdot 10^{-5}$	2.3	[172]
$(H_3C)_3C-O-O-C(CH_3)_3$	6	75	$(1.9 \ldots 5.9) \cdot 10^{-2}$	3.1	[171]
$(H_3C)_2\underset{\displaystyle C_6H_5}{\overset{\displaystyle |}{C}}-O-O-\underset{\displaystyle C_6H_5}{\overset{\displaystyle |}{C}}(CH_3)_2$	5	140	$(2.2 \ldots 2.8) \cdot 10^{-4}$	1.3	[173]
$H_3C-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-O-O-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-CH_3$	15	85.2	$(1.0 \ldots 1.7) \cdot 10^{-4}$	1.7	[174–176]
$H_5C_6-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-O-O-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-C_6H_5$	40	80	$(1.8 \ldots 40) \cdot 10^{-5}$	22	[177–179]
$(H_3C)_2CH-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-O-O-\overset{\displaystyle O}{\overset{\displaystyle ||}{C}}-CH(CH_3)_2$	16	40	$(3.1 \ldots 68) \cdot 10^{-5}$	22	[180, 181]
$H_5C_6-\underset{\displaystyle CH_3}{\overset{\displaystyle |}{CH}}-N=N-\underset{\displaystyle CH_3}{\overset{\displaystyle |}{CH}}-C_6H_5$	7	97.3	$(3.2 \ldots 4.1) \cdot 10^{-5}$	1.3	[182]
$(H_3C)_2\underset{\displaystyle CN}{\overset{\displaystyle |}{C}}-N=N-\underset{\displaystyle CN}{\overset{\displaystyle |}{C}}(CH_3)_2$	36	67	$(1.0 \ldots 2.0) \cdot 10^{-4}$	2.0	[183–185, 562]
$(H_5C_6)_3C-N=N-C_6H_5$	7	25	$(1.9 \ldots 4.4) \cdot 10^{-6}$	2.3	[186–188]
$(NC)_2\underset{\displaystyle CH_3}{\overset{\displaystyle |}{C}}-N=N-C_6H_4-p-OCH_3$	13	85	$(3.8 \ldots 100) \cdot 10^{-5}$	26	[189]

a) $k_1^{rel} = k_1$ (fastest solvent)/k_1 (slowest solvent).

Even the mesolytic scission of C—C bonds in radical cations is practically solvent-independent. The radical cations obtained by oxidation of 1,2-dimethyl-1-(4-dimethylaminophenyl)-2-phenylethane undergo unimolecular cleavage of the central C—C bond, to yield benzyl cations and benzyl radicals, with Gibbs free energies (and rate constants) that remain constant in different solvents ($\Delta G^{\neq} \approx 69$–$72$ kJ/mol) [821].

Another interesting intramolecular, possibly radical-forming reaction is the thermolytic ring-opening of cyclopropanones *via* either a diradical ($\cdot H_2C—CO—CH_2 \cdot$) or a zwitterionic oxyallyl intermediate [$H_2C=C(O^-)–CH_2^+ \leftrightarrow {}^+H_2C–C(O^-)=CH_2$], to ultimately yield an allene oxide. The rate of racemization of enantiomerically enriched *trans*-2,3-di-*t*-butylcyclopropanone increases only by a factor of 12 on going from *i*-octane to acetonitrile as solvent [822]. This small solvent effect is incompatible with a zwitterionic intermediate and confirms, together with theoretical calculations [797, 823], the diradical nature of the oxyallyl intermediate and the preceding activated complex.

The decomposition of dibenzoyl [177–179] and diisobutyryl peroxide [180, 181] (*cf.* Table 5-8) shows relatively large enhancing effects, which may be explained by

the fact that the diacyl peroxide molecule contains two mutually repelling $C^{\delta\oplus} = O^{\delta\ominus}$ dipoles. The molecule should therefore prefer the conformation shown in Eq. (5-58), with only a small net dipole moment – analogous to glyoxal, which also possesses a *transoid* conformation of the two carbonyl groups [190]. Dissociation of the diacyl peroxide leads to two independent radical dipoles, with a comparatively greater net dipole



(5-58)

moment (for comparison, the dipole moment of methyl benzoate is $6.3 \cdot 10^{-30}$ Cm = 1.9 D). Since this dipole separation occurs on activation, the slightly better solvation of the activated complex with increasing solvent polarity leads to the 22-fold rate acceleration experimentally observed on changing the solvent from *iso*-octane to acetonitrile [180, 181]. An additional influence of the solvent on the reaction of diisobutyryl peroxide is that it favors a heterolytic product in a subsequent, product-determining step. It is believed that a common intermediate, resulting from the rate-controlling step, decomposes *via* homolytic and heterolytic pathways.

In the case of the thermolysis of unsymmetrical diacyl peroxides, R—CO—O_2—CO—Ar, with negatively substituted phenyl groups (*e.g.* Ar = 3-chlorophenyl), there is a moderate increase in reaction rate with increasing solvent polarity. They are generally considered to involve ion-pair intermediates (*e.g.* $R^{\oplus}{}^{\ominus}O_2C$—Ar), formed *via* dipolar activated complexes. A typical example is that of *endo*- and *exo*-(2-norbornyl)formyl 3-chlorobenzoyl peroxide: $k_1(CH_3CN)/k_1(\text{cyclohexane}) = 320$ for the *exo*-reactant [563].

The thermal decomposition of symmetrical dialkyl peroxides such as diisopropyl peroxide in solution has been shown to involve a competition between monomolecular homolysis (k_r) and an electrocyclic reaction yielding acetone and hydrogen (k_H); *cf.* Eq. (5-59) [564].



(5-59a)

(5-59b)

Whereas the activated complex for monomolecular homolysis [Eq. (5-59a)] has no dipolar character and k_r is nearly solvent-independent [$k_r(H_2O)/k_r(\text{toluene}) = 7$], k_H increases moderately with increasing solvent polarity [$k_H(H_2O)/k_H(\text{toluene}) = 59$]. This seems to be due to the development of dipolar character in the corresponding activated complex, which involves preformed dipolar acetone molecules [Eq. (5-59b)]. In the gas phase, the normal free-radical producing O–O homolysis is the preferred reaction route [564].

The rates for the thermolysis of cyclic peroxides are also only slightly solvent-dependent. For example, the first-order decomposition of acetone cyclic diperoxide (3,3,6,6-tetramethyl-1,2,4,5-tetroxane) increases only 21-fold on going from *n*-octane to acetic acid as solvent, corresponding to homolytic O–O bond rupture [824].

Azo compounds can exist in either the *cis* or *trans* form. It is reasonable to assume that the azoalkanes in Table 5-8 exhibit the *trans* configuration. Contrary to the small solvent effects obtained in the decomposition of *trans*-azoalkanes, the thermolysis of definite *cis*-azoalkanes reveals a significant solvent influence on rate. Thermolysis of aliphatic symmetrical *cis-tert*-azoalkanes can lead either to the corresponding *trans-tert*-azoalkanes, presumably *via* an inversion mechanism, or to *tert*-alkyl radicals and nitrogen by decomposition *via* a free-radical transition state [192]. An example of the first type of reaction is the $(Z)/(E)$ isomerization of [1,1′]azonorbornane. Its rate is virtually solvent-independent, which is consistent with a simple inversion mechanism [565, 566]. The second reaction type is represented by the thermal decomposition of *cis*-2,2′-dimethyl-[2,2′]azopropane, for which a substantial decrease in rate with increasing solvent polarity has been found [193]; *cf.* Eq. (5-60).



(5-60)

Solvent	*n*-C_5H_{12}	$(C_2H_5)_2O$	CH_3COCH_3	C_2H_5OH	CH_3OH
k_1^{rel}	65	31	8.4	2.1	1

Since *cis*-azoalkanes exhibit dipole moments of *ca.* $(7 \ldots 10) \cdot 10^{-30}$ Cm (2 . . . 3 D) [194], this solvent effect is best rationalized by assuming a decrease and final loss of the dipole moment during activation. Due to their dipole moments, *cis*-azoalkanes are more stabilized by polar solvents than the less dipolar activated complexes. The activation process corresponds to a synchronous, two-bond cleavage, probably accompanied by widening of the C—N═N bond angles [193]. A two-step, one-bond cleavage process *via* short-lived diazenyl radicals has been discussed [567], but this mechanism seems to be important only in the case of unsymmetrical azoalkanes, in particular arylazoalkanes [192].

Cyclic *cis*-azoalkanes (*e.g.* pyrazolines, pyridazines, and others) exhibit a similar solvent effect on rate [193, 549], but angle expansion is of course impossible in these compounds.

Another explanation of these solvent effects recognizes the fact that polar solvents have a higher internal pressure [*cf.* Eq. (3-6) in Section 3.2] [549]. Since the activation

volume, $\Delta V^{\neq}$, for loss of nitrogen is positive, *cis*-azoalkanes should decompose more slowly in polar solvents. This should also be true for *trans*-azoalkanes. These, however, exhibit a negligible solvent dependence of the decomposition rate; see reference [192] for further examples.

Analogous to *cis*-azoalkanes (1,2-diazenes), the thermolysis of 1,1-diazenes is also solvent-sensitive. The monomolecular decomposition rate of *N*-(2,2,5,5-tetramethylpyrrolidyl)nitrene decreases with increasing solvent polarity [568].



(5-61)

Solvent	*n*-C_6H_{12}	$(C_2H_5)_2O$	Tetrahydrofuran
k_1^{rel}	5.1	2.4	1.0

As shown by Eq. (5-61), the dipolarity of the solvated reactant molecule decreases during the activation process. Loss of nitrogen by simultaneous C—N bond-breaking produces a 1,4-biradical.

The effect of the medium on the thermolysis of peroxycarboxylic esters deserves particular mention. Some examples are compiled in Table 5-9. An interesting aspect of this reaction is that the peresters, depending on structure, substituents, and medium, can decompose by two different mechanisms given in Eqs. (5-62a) and (5-62b) [195, 196].



According to Eq. (5-62a), the perester may decompose by a concerted two-bond homolysis involving an isopolar or slightly dipolar activated complex, as established by Bartlett *et al.* [197]. The activated complex may have a slightly dipolar character resulting in a small rate increase with increasing solvent polarity (rate factors 2 . . . 11; *cf.* Table 5-9). This homolytic fragmentation, involving the so-called polar effect, is found

Table 5-9. Solvent influence on rates of monomolecular decomposition of various peroxycarboxylic esters R—CO—O—O—CR'_3 according to Eqs. (5-62a) and (5-62b).

No.	R	CR'_3	Number of solvents	Temperature °C	Slowest solvent	Fastest solvent	$k_1^{rel.}$ [a]	References
1	$H_5C_6-O-CH_2-$	$-C(CH_3)_3$	5	70	H_5C_6—C_2H_5	CH_3CN	1.6	[198]
2	$n-C_5H_{11}-$	$-C(CH_3)_3$	6	110	$CH_3(CH_2)_2CO_2H$	CH_3CO_2H	1.9	[199]
3	$H_5C_6-C(CH_3)_2-$	$-C(CH_3)_3$	3	50	n-$C_{12}H_{26}$	CH_3CN	3.5	[200]
4	$H_3CO-C_6H_4-CH_2-$	$-C(CH_3)_3$	3	70.4	c-C_6H_{12}	C_2H_5OH	3.8	[201]
5		$-C(CH_3)_3$	3	10.3	tetralin	CH_3OH	6.2	[202]
6		$-C(CH_3)_3$	9 [b]	95	C_6H_5Cl	n-C_4H_9OH	6.4	[203]
7		$-C(CH_3)_3$	6	30	H_5C_6—C_2H_5	H_5C_6—CH_2OH	11	[204]
8	$o-(H_5C_6)_2C=CH-C_6H_4-$	$-C(CH_3)_3$	2	90	c-C_6H_{12}	CH_3OH	62	[205]
9	$o-H_5C_6S-C_6H_4-$	$-C(CH_3)_3$	12	40	c-C_6H_{12}	CH_3OH	692	[206]
10	$H_5C_6-CH_2-$		7	60	H_5C_6—C_2H_5	CH_3OH	31	[207]
11	H_5C_6-		11 [c]	25	CH_3OH	CH_3OH/H_2O	3.5 [c]	[208, 209]
12		$-C(CH_3)_3$	6	100	c-C_6H_{12}	CH_3OH	670	[210, 211]

[a] $k_1^{rel.} = k_1$ (fastest solvent)/k_1 (slowest solvent).
[b] Besides the two pure solvents, seven chlorobenzene/n-butanol mixtures.
[c] The reaction rate increases in the order c-$C_6H_{12} < C_6H_6 < H_3C$—$CO$—$CH_3 < CHCl_3 < CH_3NO_2 < CH_3OH < CH_3CO_2H$ [208], but exact quantitative rate constants for this reaction are only available for eleven methanol/water mixtures up to 2.8 mmol/mol of water [209].

in reactions no. 1 . . . 7 in Table 5-9. The products can be best explained by postulating a homolytic fragmentation of the peresters to form intermediate alkoxy and alkyl radicals.

Another representative example of a homolytic perester fragmentation is given by the thermal decarboxylation of β-peroxylactones such as 4,4,5,5-tetramethyl-1,2-dioxolan-3-one [569]; *cf.* Eq. (5-63).

The observed rates vary only by a factor of three in the solvents used. This is consistent with a simple homolytic O—O bond cleavage leading to a short-lived 1,5-



(5-63)

Solvent	*c*-C_6H_{12}	CCl_4	C_6H_6	CH_3CN
k_1^{rel}	1.0	1.2	1.8	2.8

biradical, which subsequently decarboxylates with concurrent β-alkyl 1,2-migration to afford the rearranged pinacolone as the major product [569].

Reactions no. 8 [205] and no. 9 [206] of Table 5-9 show special behaviour. These reactions, which show larger medium effects, involve neighbouring group participation by the alkene or phenylthio groups in the homolytic cleavage of the O—O bond, as shown for the thermolysis of *tert*-butyl 2-(phenylthio)perbenzoate in Eq. (5-64) [206]. The



(5-64)

relatively large solvent and substituent effects observed for this anchimerically assisted reaction (rate factor 692), indicate that a large contribution to the transition state is made by a dipolar structure involving a five-membered ring. There is a concentration of positive charge on the neighbouring group involved in the displacement from oxygen and of negative charge on the oxygen leaving group. The activated complex is best described as a dipolar singlet, represented by neutral and ionic canonical mesomeric structures. Products isolated from the perester decomposition are compatible with the proposed bonding interaction in the activated complex leading to free-radical fragments. The radical nature of the intermediate – and consequently the homolytic character of the reaction – is confirmed by the products observed and by the fact that galvinoxyl solutions (a radical scavenger) are decolourized. The fraction of radicals trapped does not decrease with increasing solvent polarity. The rates of reaction (5-64), measured in various solvents, correlate well with the rates of another anchimerically assisted reaction [206], the ionization of 4-methoxyneophyl tosylate, used to establish a scale of solvent ionizing power (*cf.* Section 7.3).

In contrast to reactions no. 1 . . . 9 in Table 5-9, the solvent effects and products observed in reactions no. 11 [208, 209] and no. 12 [210, 211] strongly suggest that the predominant mode of decomposition of these two peresters involves heterolysis of the O—O bond and concurrent migration of the neighbouring alkyl group to the electron-deficient oxygen as described in Eq. (5-62b) [195, 196]. This ionic mechanism was first established by Criegee *et al.* [208]. Bartlett *et al.* [209] confirmed it using *trans*-9-decalyl-peroxybenzoate as substrate; *cf.* Eq. (5-65).



(5-65)

The strong rate acceleration observed in the thermal rearrangement of *trans*-9-decalyl-peroxybenzoate to 1-benzoyloxy-1,6-epoxycyclodecane with increasing solvent polarity (rate factor *ca.* 10^2) is in accordance with the postulated heterolytic O—O bond cleavage leading to a dipolar activated complex.

The decomposition of *trans*-9-decalyl-peroxyphenylacetate (no. 10 in Table 5-9) is an interesting borderline reaction. Depending on the reaction medium, either the heterocyclic Criegee mechanism or the homolytic fragmentation mechanism can be observed [207]. In alcohols, the decomposition occurs mainly by the heterolysis pathway, whereas in nonpolar solvents like ethylbenzene, the homolysis pathway predominates. In acetonitrile, both mechanisms compete, as evidenced by the product distribution. The thermolysis of this particular perester is thus an impressive example of the strong influence which solvents may have, not only on reaction rates but also on the mechanism of a chemical reaction (*cf.* Section 5.5.7 for further examples). In addition, it should be mentioned that peresters such as *tert*-butyl cycloheptatrieneperoxycarboxylate undergo heterolytic fragmentation if the reaction is acid-catalyzed [195, 212]. The base-catalyzed ionic fragmentation of *tert*-butyl peroxyformate has already been mentioned [110]; *cf.* Eq. (5-39b).

Apart from the above described radical-forming reactions, the influence of solvent on the reactions of the radicals themselves has also been thoroughly investigated [159–166]. The most important elementary reactions of radicals are atom transfer, combination, addition, disproportionation, and electron transfer, as listed in Table 5-10 [15, 213].

Most criteria for mechanisms depend upon intramolecular and extramolecular perturbation of the reacting system. A change in the medium is an extramolecular perturbation of the original system. The solvent effects produced by this perturbation can be predicted, as shown in Table 5-10, by taking into account whether or not the activated complex is dipolar or isopolar with the respect to the initial reactants. Only a small

Table 5-10. Expected substituent and solvent rate effects for elementary radical reactions [15, 213].

Reaction type	Reaction scheme	Intramolecular perturbation by substituents	Extramolecular perturbation by medium
Atom transfer	$R\odot + A\text{-}X \rightleftharpoons [R^{\delta\odot}\cdots A\cdots X^{\delta\odot}]^{\ddagger} \longrightarrow R\text{-}A + \odot X$	Modest	Modest
Combination	$R\odot + \odot R \rightleftharpoons [R^{\delta\odot}\cdots R^{\delta\odot}]^{\ddagger} \longrightarrow R\text{-}R$	Small	Small
Addition	$R\odot + CH_2{=}CH\text{-}R \rightleftharpoons [R^{\delta\odot}\cdots CH_2{=}\overset{\delta\odot}{C}H\text{-}R]^{\ddagger} \longrightarrow R\text{-}CH_2\text{-}\overset{\odot}{C}H\text{-}R$	Small	Small
Disproportionation	$R\odot + \underset{R}{\overset{\odot}{C}H_2}\text{-}CH_2 \rightleftharpoons [R^{\delta\odot}\cdots H\cdots \underset{R}{CH}{=}\overset{\delta\odot}{C}H_2]^{\ddagger} \longrightarrow R\text{-}H + \underset{R}{CH}{=}CH_2$	Small	Small
Electron transfer	$R\odot \longrightarrow R^{\oplus} + e^{\ominus}$ or $R\odot + e^{\ominus} \longrightarrow R^{\ominus}$	Large	Large

number of examples selected from the vast number of solvent-dependent free-radical reactions [159–166] shall be used to illustrate this point.

The most extensively studied reactions are those involving substitution between radicals (like halogen atoms, alkoxy or peroxy radicals, *etc.*) and a neutral molecule A—X (*cf.* first reaction in Table 5-10). In this atom-transfer reaction, the atom A, which is frequently a hydrogen atom, is transferred slowly from A—X to R·. In the isopolar activated complex of this reaction, there is no appreciable charge separation. In reactions of this type, therefore, only negligible solvent effects should be observed. However, there are also radical reactions known, in which a change in the polarity of the solvent can play an important role. For such processes, whose rate is clearly influenced by the medium, a certain charge separation during activation must be taken into account. The degree of separation of charges in an activated complex such as $[R^{\delta\ominus}\cdots A\cdots X^{\delta\oplus}]^{\neq}$ should depend on the electron affinity of the radical R· and the ionization potential of the molecule A—X.

Halogen abstraction by the stable free radical 1-ethyl-4-(methoxycarbonyl)pyridinyl (Py·) proceeds by the mechanism shown in Eq. (5-66) [214, 570]. The first step, which is rate-determining, is a transfer of the halogen atom to the pyridinyl radical.



(5-66)

Solvent	CH_2Cl_2	CH_3CN	$(CH_3)_2CHOH$	C_2H_5OH
k_2^{rel}	1.0	2.1	2.0	3.5

The negligible solvent effect of this radical reaction with dibromomethane demonstrates that the activated complex for bromine atom-transfer has the same charge separation as the initial reactants. The dipole moment expected for a molecule like the pyridinyl radical is probably $(0 \ldots 10) \cdot 10^{-30}$ Cm (0 ... 3 D). Dibromomethane has a modest dipole moment of $5 \cdot 10^{-30}$ Cm (1.5 D). Consequently, in view of the negligible solvent effect upon rate, the activated complex must also have a dipole moment between $(0 \ldots 10) \cdot 10^{-30}$ Cm [214, 570].

In contrast to the preceding atom-transfer reaction, the solvent-induced rate change for the reaction between 1-ethyl-4-(methoxycarbonyl)pyridinyl and 4-(halomethyl)-nitrobenzenes is so large that a change in mechanism must be involved [215, 570]. In changing the solvent from 2-methyltetrahydrofuran to acetonitrile, the relative rate constant for 4-(bromomethyl)-nitrobenzene increases by a factor of up to 14800. This is of the order expected for a reaction in which an ion pair is created from a pair of neutral molecules [*cf.* for example, reaction (5-16)]. It has been confirmed therefore that, according to scheme (5-67), an electron-transfer process is involved [215, 570].



(5-67)

Solvent	2-Methyltetrahydrofuran	CH_2Cl_2	CH_3COCH_3	$HCON(CH_3)_2$	CH_3CN
k_2^{rel} for X = Br	1	46	278	7400	14800

It has been established that for solvents in which specific solvation is not dominant, a small solvent effect implies an atom-transfer reaction and a large solvent effect suggests an electron-transfer reaction between neutral species. The high solvent sensitivity of electron-transfer reactions between neutral molecules should provide a useful test of their occurrence [215, 570]. From Table 5-11, it can be concluded that atom-transfer, according to Eq. (5-66), is the rate-limiting step in the reaction of pyridinyl radical with

Table 5-11. Rate constant solvent effects for the reaction of haloalkanes with 1-ethyl-4-(methoxycarbonyl)pyridinyl radicals at 25 °C [215, 570].

Haloalkane	k_2^{rel} [a]	$\Delta\Delta G^{\neq}/(kJ \cdot mol^{-1})$ [a]
Dibromomethane	*ca.* 0.5 [b]	−1.3
(Bromomethyl)benzene	30	8.4
(Chloromethyl)benzene	63	10.3
4-(Chloromethyl)-nitrobenzene	2900	19.7

[a] Rate constant for the reaction of pyridinyl radical in acetonitrile divided by that for the reaction in 1,2-dimethoxyethane and the corresponding change in Gibbs activation energy.
[b] $k_2^{rel} = k_2(i\text{-propanol})/k_2(\text{dichloromethane})$. This solvent polarity change is close to that used in the other cases.

(halomethyl)benzenes except for 4-(halomethyl)-nitrobenzenes, for which the solvent effect is compatible only with an electron-transfer mechanism according to Eq. (5-67) (for a more detailed discussion, see references [215, 570]).

Small but significant effects of solvent polarity were found in the autoxidation of a variety of alkenes and aralkyl hydrocarbons [216–220] (styrene [216, 218, 219], ethyl methyl ketone [217], cyclohexene [218], cumene [218, 219], tetralin [219], *etc.*). An extensive study on solvent effects in the azobisisobutyronitrile (AIBN)-initiated oxidation of tetralin in a great variety of solvents and binary solvent mixtures was made by Kamiya *et al.* [220].

Since solvent effects in radical oxidation (and autoxidation) reactions of organic compounds have been compiled and discussed in an excellent monograph by Emanuel, Zaikov, and Maizus [460], they are not discussed further here.

Another example of a solvent-dependent atom-transfer reaction is hydrogen abstraction by chlorine atoms during the photochemical chlorination of hydrocarbons with molecular chlorine; for an excellent review, see reference [571]. Russel reported that in the photochlorination of 2,3-dimethylbutane, according to reaction scheme (5-68), certain solvents do not have any effect on the selectivity of the reaction as measured by the rate ratio $k_2^{\text{tert}}/k_2^{\text{prim}}$, whereas other solvents increase this ratio significantly (*cf.* Table 5-12) [221]. The relative reactivity ratio $k_2^{\text{tert}}/k_2^{\text{prim}}$ of the tertiary hydrogen atoms, with respect to the primary hydrogens in 2,3-dimethylbutane, can be determined from the relative amounts of 2-chloro-2,3-dimethylbutane and 1-chloro-2,3-dimethylbutane



(5-68)

produced in the photochlorination of this hydrocarbon*). In the absence of any solvent, and in several aliphatic and cycloaliphatic solvents, the ratio $k_2^{\text{tert}}/k_2^{\text{prim}}$ varies from 3.3 to 9.1. In other solvents, particularly aromatic solvents and carbon disulfide, this ratio becomes comparatively large, ranging from 10 to 33 (*cf.* Table 5-12). Since the respective activated complexes leading to the primary and the tertiary haloalkane should not differ in their dipolarity, the solvation of the initial reactants must cause this selectivity. The latter group of solvents includes carbon disulfide and benzene derivatives bearing substituents that increase the electron density in the aromatic ring relative to that of benzene itself. These π-EPD solvents are able to form a loose π complex with the electrophilic chlorine radical in a reversible reaction as shown in Eq. (5-69). As might be

* It is necessary to multiply this product ratio by six to correct for the fact that there are twelve primary hydrogen atoms and two tertiary hydrogen atoms in the molecule.

Table 5-12. Solvent effect on the selectivity of the photochemical chlorination of 2,3-dimethylbutane at 55 °C (solvent concentration 4.0 mol/L) [221].

Solvent	Relative Reactivities k_2^{tert}/k_2^{prim} [a]
2,3-Dimethylbutane	3.7[b]
Nitromethane	3.3
Tetrachloromethane	3.5
Cyclohexene	3.6
Trichloroethene	3.6
Propionitrile	4.0
Methyl acetate	4.3
tert-Butanol	4.8
Nitrobenzene	4.9
1,4-Dioxane	5.6
Di-*n*-butyl ether	7.2
N,*N*-Dimethylformamide	9.1
Chlorobenzene	10.2
Flurobenzene	10.3
Benzene	14.6
Toluene	15.4
Methoxybenzene	18.4
p-Xylene	18.6
1,3,5-Trimethylbenzene	25
Iodobenzene	31
Carbon disulfide	33[c]

[a] *Cf.* reaction scheme (5-68).
[b] Solvent concentration 7.6 mol/L.
[c] Value at 25 °C.

expected, the complexed and therefore less reactive chlorine atoms show greater selectivity as hydrogen abstractors. The differences in the relative amounts of tertiary and primary chloroalkanes formed become more pronounced when the chlorination is performed in such complexing solvents. On the other hand, electron-withdrawing substituents on the benzene ring (*e.g.* NO_2) decrease the selectivity relative to benzene. Even in the presence of benzene, a fairly good π-EPD solvent, there is still some attack on the primary hydrogen atoms of 2,3-dimethylbutane. More direct evidence for the existence of a chlorine atom/benzene π-complex in the photochlorination of 2,3-dimethylbutane in solution in the presence of benzene has been given by Ingold *et al.* [572]. The interaction of $Cl\cdot$ with carbon disulfide, which displays a remarkably high efficiency in increasing the selectivity of the chlorine atom (*cf.* Table 5-12), probably leads to the formation of a σ-complexed radical according to Eq. (5-70), which will certainly have less energy than free $Cl\cdot$ and act as a more selective hydrogen abstractor. The comparatively small increase in selectivity caused by solvents lacking a π-electron system, but having non-bonding electrons (*n*-EPD solvents like alcohols, ethers, and *N*,*N*-dimethylformamide) might result from a complexation of the electrophilic chlorine atom by the oxygen of these solvents.



(5-69)

π-complex



(5-70)

σ-complex

Studies of the influence of the solvent on relative selectivity in the photochlorination of 1,1-dichloroethane [573], 1-chlorobutane [574], and 2-chlorobutane [575] have shown that solvents can be divided into three classes. First, there are nonselective solvents, which are approximately as selective as the neat liquid hydrocarbon reactants (*e.g.* CCl_4, CH_2Cl_2, CH_3CN). Secondly, there are moderately selective perfluorohydrocarbon solvents (*e.g.* C_6F_{14}, C_8F_{18}, $C_{10}F_{20}O$), which exhibit selectivities similar to those in gas-phase chlorinations. Thirdly, there are solvents which lead to a greatly increased selectivity (*e.g.* C_6H_6, CS_2). The first group of solvents are relatively polarizable (as compared to the perfluorohydrocarbon solvents) and are thus able to stabilize the activated complex by solvation. This lowers the Gibbs activation energy and tends to have a levelling effect. In contrast, the inert perfluorohydrocarbons have no tendency to solvate the activated complex. Chlorinations in such solvents have a selectivity approaching that of the gas-phase reaction. The third group of solvents stabilize the chlorine atoms by specific solvation [*cf.* Eqs. (5-69) and (5-70)] and the resulting reactant/solvent complex has to be broken open before hydrogen abstraction can occur [573–575]. A detailed review on the photochlorination of alkanes in solution can be found in reference [825].

Another remarkable example of the solvent-influenced chemoselectivity of chlorine radicals is the photochlorination of cyclopropane to give, in competitive reactions, either 3-chloropropyl radicals (by S_H2 ring opening; k_C) or cyclopropyl radicals (by H abstraction; k_H) as intermediates. The rate constant ratio k_C/k_H was found to increase by a factor of 372 on going from the gas phase to 1,2-dichloroethane, presumably as a result of the internal pressure of the solvents used. Thus, solvents of high internal pressure favour the S_H2 process, while H-abstraction becomes more important in the gas phase and in solvents of low internal pressure [826].

The selectivity of free-radical side-chain bromination of toluene derivatives using *N*-bromosuccinimide and leading to mono- and dibromo-substituted toluenes has been studied in different solvents [577]. Surprisingly, yields and selectivities are much better in solvents such as methyl formate and dichloromethane than in the more commonly used tetrachloromethane.

Compared with chlorination, hydrogen abstraction reactions of alkoxy radicals are relatively insensitive to solvent effects [160, 222, 223]. The results of the AIBN-initiated radical chain chlorination of 2,3-dimethylbutane with *tert*-butyl hypochlorite indicate a solvent effect on *tert*-butoxy radical reactions of much smaller magnitude, but greater selectivity in aromatic solvents [222, 223]. The reduced solvent effect for this hydrogen abstraction reaction has been attributed to steric effects. Due to the bulky

methyl groups around the electrophilic oxygen atom, complex formation involving solvent molecules and activated complex is hindered.

tert-Alkoxy radicals involved in the reactions of dialkyl peroxides and alkyl hypochlorites are not only capable of abstracting a hydrogen from a hydrocarbon yielding an alkyl radical and *tert*-alkyl alcohol, they can also decompose into a ketone and an alkyl radical, which subsequently reacts with the hydrocarbon. Changes in the rates of hydrogen abstraction (k_a) and decomposition (also called β-scission; k_β) of the radical can be determined from the relative amounts of the *tert*-alkyl alcohol and ketone produced, as shown for the *tert*-butoxy radical in Scheme (5-71) [160].



(5-71)

The ratio k_a/k_β can be solvent-dependent, as has been shown for the reaction of *tert*-butyl hypochlorite with cyclohexane in different solvents [224].

Replacement of one methyl group in the *t*-butoxy radical of Eq. (5-71) by a phenyl group leads to the cumyloxy radical, $H_5C_6–C(CH_3)_2–O\cdot$, the reaction of which has recently been studied in solvents of different polarity [827]. Analogously, cumyloxy radicals, generated photolytically from dicumyl peroxide, undergo H-abstraction reactions with cyclohexane or the solvent (k_a) and decomposition reactions (β-scission; k_β) with rate constant ratios k_a/k_β of up to 6.5 on going from polar acetic acid to apolar tetrachloromethane. The results indicate that k_a is practically solvent-independent, while k_β increases with increasing solvent polarity. This is in agreement with the results obtained for *t*-butoxy radicals [224], suggesting a decrease of k_a/k_β with increasing solvent polarity for both radicals. The increase in k_β with increasing solvent polarity is attributable to better stabilization of the activated complex for β-scission because of increased solvation of the developing dipolar ketone product (acetone or acetophenone, respectively) [827].

A free-radical addition reaction, the solvent-dependence of which has been studied in thirty-nine solvents, is the addition of the 4-(dimethylamino)benzenethiyl radical to α-methylstyrene; *cf.* Eq. (5-72) [576].

The rate of addition decreases moderately with increasing solvent polarity; there is a 35-fold rate deceleration in going from cyclohexane to dimethyl sulfoxide. In polar solvents, the dipolar reactant thiyl radical is more stabilized than the less dipolar activated complex. The stabilization of the thiyl radical by solvation has been proven by its strong positive solvatochromism (*i.e.* bathochromic shift of λ_{max} with increasing solvent polarity) [576]. Similar solvent effects on rate have been observed in the addition of the 4-aminobenzenethiyl radical to styrene [577].





(5-72)

Solvent	*c*-C_6H_{12}	CCl_4	C_6H_6	C_2H_5OH	CH_3CN	CH_3SOCH_3
k_2^{rel}	35	19	6.8	2.8	1.7	1.0

Finally, a solvent-dependent disproportionation reaction is discussed. The 2,6-di-*tert*-butyl-4-isopropylphenoxy radical disproportionates to the corresponding quinone methide and the parent phenol in a slightly solvent-dependent reaction, according to Eq. (5-73) [225]. The enthalpies of activation vary from 21 to 32 kJ/mol in going from cyclohexane to benzonitrile, increasing as the polarity of the solvent increases. Due to the compensating changes in the entropies of activation, the reaction rates are only slightly sensitive to changes in the medium. The formation of the activated complex may be regarded as the head-to-tail joining of two dipoles. Desolvation of one of the phenoxy radicals is essential for the formation of this activated complex. Thus, in a medium in which radicals are highly solvated, the enthalpy of activation should be relatively high in order to provide the necessary desolvation energy. Such cases should be accompanied by the largest entropy increase. The linear relationship observed between $\Delta H^{\neq}$ and $(\varepsilon_r - 1)/\varepsilon_r$ strongly suggests that the solvent effect in this reaction can mainly be attributed to dipolar interactions between radical and solvent molecules [225].



(5-73)

Solvent	C_6H_6	*c*-C_6H_{12}	C_6H_5Cl	$C_6H_5OCH_3$	C_6H_5CN
k_2^{rel}	1	1.8	1.4	1.5	2.6
$\Delta H^{\neq}/(kJ \cdot mol^{-1})$	24	21	27	28	32

Apart from the selection of reactions involving dipolar, isopolar, or free-radical activated complexes used to demonstrate the qualitative theory of solvent effects by Hughes and Ingold [16, 44] in the preceding sections, further illustrative examples can be found in the literature (*e.g.* [14, 15, 18, 21, 23, 26, 29, 460, 468]).

5.3.5 Limitations of the Hughes–Ingold Rules

The qualitative theory of solvent effects introduced by Hughes and Ingold in 1935 [16, 44] is expressed as a set of rules that take into account the changes in charge magnitude and in charge distribution that occur between reactants and activated complex, as well as of the dielectric characteristics of solvents that enable them to solvate charges (*cf.* Section 5.3.1). Although the countless successful applications of this qualitative theory testify to its widespread use, it does contain some inherent limitations.

One of these limitations is the assumption made by Hughes and Ingold that the contribution of entropy changes ($\Delta S^{\neq}$) to changes in Gibbs energy of activation ($\Delta G^{\neq}$) are negligible. This implies that enthalpy changes ($\Delta H^{\neq}$) dominate the Gibbs energy expression $\Delta G^{\neq} = \Delta H^{\neq} - \mathrm{T} \cdot \Delta S^{\neq}$. This assumption is necessary because an increase in solvation usually decreases the entropy of a given state. Decreases in the entropy of activation counteract increasing enthalpy changes, but fortunately these decreases are relatively small for most reactions. Therefore, most but not all chemical reactions are controlled by enthalpy changes. A number of reactions in which changes in the entropy of activation govern the Gibbs energy of activation were reported by Pearson [226]. As illustrated in Table 5-5 for the reaction of iodoethane and triethylamine (*cf.* Eq. (5-16) in Section 5.3.1 [59]), the rate constants increase by a factor of 52 with increasing solvent polarity, although the enthalpies of activation show only a relatively small change of $\Delta\Delta H^{\neq} = 2.9\ \mathrm{kJ \cdot mol^{-1}}$. The entropies of activation rise by $\Delta\Delta S^{\neq} = 23\ \mathrm{J \cdot mol^{-1} \cdot K^{-1}}$ for the same sequence of solvents. Polar solvents appear to reduce the loss of entropy on passing from the reactants to the activated complex of the reaction, and it is mainly this factor that causes the rates to increase with solvent polarity. An explanation in terms of solvent orientation around the dipolar activated complex of reaction (5-16) was given in Section 5.3.1.

Another example is the S_N1 solvolysis of 2-chloro-2-methylpropane (*cf.* Eq. (5-13) in Section 5.3.1 [40]), the rate data for which are given in Table 5-13. As the solvent is changed from ethanol to water, the rate of solvolysis increases by a factor of 335000, with a corresponding decrease in $\Delta G^{\neq}$. In going from ethanol to formic acid, the values of $\Delta S^{\neq}$ vary only from -13 to $-7\ \mathrm{J \cdot mol^{-1} \cdot K^{-1}}$. In the first five solvents, the $\Delta S^{\neq}$ values are relatively constant (*ca.* $-12\ \mathrm{J \cdot mol^{-1} \cdot K^{-1}}$), in agreement with domination of the overall solvent effects by $\Delta H^{\neq}$. Since $\Delta S^{\neq}$ is constant in these solvents, $\Delta H^{\neq}$ and $\Delta G^{\neq}$ must have a linear relationship. However, the large $\Delta S^{\neq}$ value for the reaction in water demonstrates that the $\Delta S^{\neq}$ changes are important in this case. This huge entropy effect is obviously due in part to the highly ordered structure of water.

Even in those cases where the rate constants, for a reaction in various solvents, are not significantly different, the activation parameters may indicate a significant amount of interaction between solute and solvent, as shown for the unimolecular decomposition of di-*tert*-butyl peroxide in Table 5-14 [172, 227]. The rate of decomposition of the per-

Table 5-13. Rate constants and activation parameters for the S_N1 solvolysis of 2-chloro-2-methylpropane in six solvents at 25 °C [40].

Solvents	ε_r	k_1 $(10^5 \cdot s^{-1})$	$k_1^{rel.}$	$\Delta G^{\neq}$ $(kJ \cdot mol^{-1})$	$\Delta H^{\neq}$ $(kJ \cdot mol^{-1})$	$\Delta S^{\neq}$ $(J \cdot mol^{-1} \cdot K^{-1})$
Ethanol	24.6	0.00860	1	113	109	−13
Acetic acid	6.2	0.0213	2.5	111	108	−10
Methanol	32.7	0.0753	9	108	104	−13
Formamide	109.5	3.72	430	98.3	93.6	−16
Formic Acid	58.5	105	12200	90.0	87.9	−7
Water	78.4	2880	335000	82.0	97.2	+51(!)

oxide does not vary by more than a factor of 2.3 in going from cyclohexane to acetonitrile, corresponding to a change in $\Delta G^{\neq}$ of only 4 $kJ \cdot mol^{-1}$. However, the activation parameters $\Delta H^{\neq}$ and $\Delta S^{\neq}$ show that the solvent apparently has a marked effect on the decomposition reaction ($\Delta\Delta H^{\neq} = 41$ $kJ \cdot mol^{-1}$; $\Delta\Delta S^{\neq} = 94$ $J \cdot mol^{-1} \cdot K^{-1}$). An evaluation of these activation parameters indicates that the compensating effects of the energy gained in solvating the radical-like activated complex, which would be expected to increase the decomposition rate, is counterbalanced by the decrease in entropy resulting from the more highly ordered arrangement of solvent molecules around the activated complex. The result is that the reaction rate is in the same range as that observed for the gas-phase reaction.

A second limitation of the Hughes–Ingold theory concerns the fact that the solvent is treated as dielectric continuum, characterized by one of the following: its relative permittivity, ε_r, the dipole moment, μ, or by its electrostatic factor, *EF*, defined as the product of ε_r and μ [27]. The term "solvent polarity" refers then to the ability of a solvent to interact electrostatically with solute molecules. It should be remembered, however, that solvents can also interact with solute molecules through specific intermolecular forces like hydrogen bonding or EPD/EPA complexation (*cf.* Section 2.2). For example, specific solvation of anionic solutes by protic solvents may reduce their nucleophilic reactivity, whereas in dipolar aprotic solvents solvation of anions is less,

Table 5-14. Rate constants and activation parameters for the decomposition of di-*tert*-butyl peroxide at 125 °C [172, 227].

Solvents	ε_r	k_1 $(10^{-5} \cdot s^{-1})$	$k_1^{rel.}$	$\Delta G^{\neq}$ $(kJ \cdot mol^{-1})$	$\Delta H^{\neq}$ $(kJ \cdot mol^{-1})$	$\Delta S^{\neq}$ $(J \cdot mol^{-1} \cdot K^{-1})$
Gas phase	1.00	–	–	136	159	59
Cyclohexane	2.0	1.52	1	136	171	88
Tetrahydrofuran	7.6	1.84	1.2	135	155	51
Benzene	2.3	1.99	1.3	135	148	33
Nitrobenzene	34.8	2.39	1.6	134	149	38
Acetic acid	6.2	2.98	2.0	132	140	19
Acetonitrile	35.9	3.47	2.3	132	130	−6

resulting in enormous rate accelerations [6], which cannot be explained by the simple electrostatic theory (*cf.* Section 5.5.2).

A third limitation of the Hughes–Ingold concept of solvent effects on reaction rates arises from the fact that it is based on static equilibrium transition-state solvation (*cf.* Section 5.1). That is, the reorientational relaxation of solvent molecules during the activation process is considered to be sufficiently fast so that the activated complex will be in thermal equilibrium with the surrounding solvent shell. However, this is not necessarily true in all cases, in particular not for very fast reactions. In such reactions, the rate will also depend on solvent reorientation rates and the standard transition-state theory will break down. Theoretical investigations of the S_N1 ionization reactions of $Me_3C–X$ (X = Cl, Br, I) in solution have shown that the calculated Gibbs energies of activation, $\Delta G^{\neq}$, decrease with increasing solvent polarity, in agreement with experiment and conventional expectations. However, a *decreasing* stabilization of the activated complexes with increasing solvent polarity has been found. That is, the activated complexes become *less* ionic in more polar solvents. This surprising result is at variance with the conventional Hughes–Ingold rules. Obviously, for S_N1 ionization reactions of tertiary haloalkanes, the conventional equilibrium reaction-rate methods cannot be applied. The S_N1 ionization process seems to be a non-equilibrium one [465]; *cf.* references [463, 465, 466] for further discussions.

It should be mentioned that an investigation of solvent rate effects is very often limited by the narrow range of solvents examined. When a particular reaction is studied in a small number of often very similar solvents, no far-reaching conclusions about the influence of solvent can be reached. As a rule, the reaction under consideration should be investigated in as many solvents of different polarity as possible. With a minimum of five different solvents, a good general picture of the reaction should be obtained. Very often, reactions have been carried out in a series of binary solvent mixtures such as alcohol/water. Solvents of such a similar structure will specifically solvate the solutes in a related manner throughout the whole solvent series, and a change of solvent mixture will mainly reflect the electrostatic influences of the solvents.

The Hughes–Ingold theory also neglects the changing solvent structure. Although solvent–solvent interactions are usually small compared to solute–solvent interactions, consideration should be given to solvent association when reactions are carried out in a highly structured solvent like water. The electrostatic theory ignores such solvent–solvent interactions.

The final limitation of the pure electrostatic theory is its inability to predict solvent effects for reactions involving isopolar transition states. Since no creation, destruction, or distribution of charge occurs on passing from the reactants to the activated complex of these reactions, their rates are expected to be solvent-independent. However, the observed rate constants usually vary with solvent, although the variations rarely exceed one order of magnitude (*cf.* Section 5.3.3). These solvent effects may be explained in terms of cohesive forces of a solvent acting on a solute, usually measured by the cohesive pressure of the solvent (*cf.* Section 5.4.2).

In spite of these limitations, the electrostatic Hughes–Ingold theory remains a good guide in predicting the solvent influence on chemical reactions, at least in a qualitative way. Exceptions can be safely assumed to involve strong specific solute–solvent interactions.

5.4 Quantitative Theories of Solvent Effects on Reaction Rates

5.4.1 General Remarks

Whereas the theoretical treatment of gas-phase reactions is comparatively simple, the calculation of rate constants for reactions in the liquid phase is very complicated. This is essentially due to the complexity of the many possible intermolecular solute-solvent interactions (*cf.* Section 2.2). When investigating solution-phase reaction kinetics, the problems to be faced include deciding which property of the solvent to use when setting up mathematical correlations with the reaction rates. Another problem is deciding which characteristics of the reacting molecules are to be considered when the effects of the solvent on their reactivity is determined. A quantitative allowance for the solvent effects on the rate constants k for elementary reactions involves establishing the following functions:

$$k = f(a, b, c, \ldots m, n, o, \ldots) \tag{5-74}$$

where a, b, and c are parameters characterizing the properties of the reactants, and m, n, and o are parameters characterizing the properties of the medium. The function thus obtained will then correctly describe the dependence of the rate constants on the medium. The problem then arises as to which reactant and solvent parameters are responsible for the observed dependences and must be included in Eq. (5-74).

When considering only pure electrostatic interactions leading to non-specific solvation, the solvent can be regarded as an isotropic continuum, with relative permittivity ε_r, and the reactants are characterized by both the magnitude and distribution of charge in the molecule. However, an analysis of solvent effects shows that not only non-specific solvation caused by electrostatic and dispersion forces, but also specific solvation caused by hydrogen bonding and EPD/EPA complexation, must be considered. The reaction kinetics for any particular reaction in a given solvent will be determined by the predominating type of solvation for that reaction.

The theoretical treatment of liquid-phase reaction kinetics is limited by the fact that no single universal theory on the liquid state exists at present. Problems which have yet to be sufficiently explained are: the precise character of interaction forces and energy transfer between reacting molecules, the changes in reactivity as a result of these interactions, and finally the role of the actual solvent structure. Despite some limitations, the absolute reaction rates theory is at present the only sufficiently developed theory for processing the kinetic patterns of chemical reactions in solution [2–5, 7, 8, 11, 24, 463–466]. According to this theory, the relative stabilization by solvation of the initial reactants and the activated complex must be considered (*cf.* Section 5.1).

Thus, the problem of making quantitative allowances for solvent-induced changes in the rate constant of a reaction $A + B \rightleftharpoons (AB)^{\neq} \rightarrow C + D$ (*cf.* Eq. (5-1) in Section 5.1) is reduced to the calculation of the difference between the partial Gibbs energies of solvation of the activated complex $(AB)^{\neq}$ and the initial reactants A and B as given in Eq. (5-75) [28].

$$\ln k = \ln k_0 - \frac{1}{R \cdot T}\left(\Delta G_{A,\,solv} + \Delta G_{B,\,solv} - \Delta G_{(AB)^{\neq},\,solv}\right) \tag{5-75}$$

k_0 is the rate constant of the reaction in a standard solvent or in the gas phase, k is the rate constant observed in the solvent under consideration. Some progress has been made in the calculation of initial-state and transition-state solvent effects on reaction rates using thermodynamic transfer functions [453, 467]; *cf.* also Section 5.5.3.

If all possible solute–solvent interactions are taken into account, then attempts to correlate the rate constant with the medium will generally lead to such complicated equations that their experimental verification is impossible. Therefore, equations correlating rate constants with medium properties are usually derived on the basis of more or less theoretically justified models, allowing for only a limited number of dominating interaction factors. If the model adopted correctly reflects the dominating solute–solvent interactions, then a good quantitative description of the experimental data using the derived theoretical equation may be obtained. It is convenient, therefore, to distinguish between the following four reaction types:

(a) Reactions between neutral, apolar molecules (*via* isopolar activated complexes);
(b) Reactions between neutral, dipolar molecules (*via* dipolar activated complexes);
(c) Reactions between neutral, dipolar molecules and ions (*via* dipolar and charged activated complexes); and
(d) Reactions between ions (*via* dipolar and/or charged activated complexes).

The following sections will deal only with the final results of these calculations based on suitable, but rather simple models, together with some examples, which shall illustrate the proposed dependences. More detailed discussions can be found in well-known monographs [2–5, 7, 8, 11, 12, 21, 24, 25, 28, 457, 459, 460]. During the last decades, theoretical chemists have developed an impressive variety of methods and computational strategies for describing and understanding the complex phenomena of solute solvation. These can also be applied to the solvation of reactants and activated complexes, which is of particular relevance to this Chapter. These theoretical treatments of solute–solvent interactions include, amongst others, quantum-chemical continuum models, supramolecular models, and semicontinuum quantum-chemical models; see the end of Section 2.3. For recent comprehensive reviews on these topics, see references [27d, 355–358] given in Chapter 2 and references [578, 579] given in this Chapter.

5.4.2 Reactions Between Neutral, Apolar Molecules

In any solution reaction, cavities in the solvent must be created to accommodate reactants, activated complex, and products. Thus, the ease with which solvent molecules can be separated from each other to form these cavities is an important factor in solute solubility (*cf.* Section 2.1). Furthermore, because solubility and reactivity are often related phenomena, the intermolecular forces between solvent molecules must also influence rates of reaction. The overall attractive forces between solvent molecules gives the solvent as a whole a cohesion which must be overcome before a cavity is created. The degree of cohesion may be estimated using the surface tension, but a more reliable estimate is obtained by considering the energy necessary to separate the solvent molecules. This is known as the *cohesive pressure c* (also called *cohesive energy density*) [228–

232]. The cohesive pressure is defined as the energy of vapourization, ΔU_v, per unit molar volume, V_m, as shown in Eq. (5-76); *cf.* also Eq. (3-5) in Section 3.2.

$$c = \frac{\Delta U_v}{V_m} = \frac{\Delta H_v - R \cdot T}{M_r \cdot \varrho^{-1}} \tag{5-76}$$

Values of c are calculated from experimentally determined enthalpies (heats) of vapourization of the solvent to a gas of zero pressure, ΔH_v, at a temperature T, as well as from the molecular mass M_r, the density of the solvent ϱ, and the gas constant, R. The cohesive pressure characterizes the amount of energy needed to separate molecules of a liquid and is therefore a measure of the attractive forces between solvent molecules. The cohesive pressure c is related to the *internal pressure* π, because cohesion is related to the pressure within a liquid; *cf.* Eq. (3-6) in Section 3.2 for the precise definition of π*).

In mixtures which are regular solutions**), the mutual solubility of the components depends on the cohesive pressure, hence Hildebrand termed the square root of c the *solubility parameter* δ, according to Eq. (5-77); *cf.* also Eq. (2-1) in Section 2.1 [228, 229, 231, 238].

$$\delta = \left(\frac{\Delta U_v}{V_m}\right)^{1/2} = \left(\frac{\Delta H_v - R \cdot T}{M_r \cdot \varrho^{-1}}\right)^{1/2} \tag{5-77}$$

A good solvent for a certain nonelectrolyte solute should have a δ value close to that of the solute (*cf.* Section A.1). Extensive compilations of δ values are given in references [231, 238]; a selection of δ values for various organic solvents is given in Table 3-3 in Section 3.2.

Assuming that it is only van der Waals forces which are acting in the solute/solvent system, and that the heat of mixing is responsible for all deviations from ideal behaviour, as well as the fact that the solute/solvent interaction energy is the geometric mean of solute/solute and solvent/solvent interactions, Hildebrand [228, 229] and Scatchard [230] were able to develop the following expression for the activity coefficient f_i of the nonelectrolyte solute i dissolved in a solvent s (mole fraction basis), referred to a standard state of pure liquid solute (not infinite dilution):

$$RT \cdot \ln f_i = V_{m,i} \cdot \phi_s^2 \cdot \left[\left(\frac{\Delta U_{v,i}}{V_{m,i}}\right)^{1/2} - \left(\frac{\Delta U_{v,s}}{V_{m,s}}\right)^{1/2}\right]^2 \tag{5-78}$$

---

* Since the internal pressure is actually defined in a slightly different way, values of internal pressure approach those of the cohesive pressure only for nonpolar and non-associated solvents (*cf.* Table 3-2 in Section 3.2) [228–232, 237]. Internal pressure is a measure of the instantaneous volume derivative of the cohesive pressure during isothermal expansion of a liquid (*cf.* Eq. (3-6) in Section 3.2). Because of the experimental difficulty in obtaining real internal pressures, it is usual to refer to $\Delta U_v / V_m$ as the internal pressure of a liquid.

** Regular solutions are characterized by a disordered distribution of solute and solvent molecules which is the same as in an ideal solution. In going from an ideal to a regular solution there is no change in entropy ($\Delta S = 0$), whereas the change in the activity coefficient f_i of the solute i is determined only by the enthalpy component of the Gibbs energy: $RT \cdot \ln f_i = \Delta G = \Delta H$ [228, 229].

where $V_{m,i}$ is the molar volume of solute i (as a pure liquid); $\Delta U_{v,i}$ is the molar energy of vapourization of the solute (as a pure liquid); $V_{m,s}$ and $\Delta U_{v,s}$ are the same quantities for the solvent s; ϕ_s is the volume fraction of the solvent s, equal to unity for a dilute solution.

Since reactions are usually carried out in dilute solution, Eq. (5-78) can be simplified to Eq. (5-79),

$$RT \cdot \ln f_i = V_{m,i} \cdot (\delta_i - \delta_s)^2 \tag{5-79}$$

where δ_i and δ_s are the solubility parameters of the reactant solutes and the solvent, respectively, according to Eq. (5-77) [5, 228–230].

The rate constant k of a bimolecular reaction $A + B \rightleftharpoons (AB)^{\neq} \rightarrow C + D$ can be expressed either as in Eq. (5-75) or in terms of the activity coefficients as shown in Eq. (5-80),

$$\ln k = \ln k_0 + \ln f_A + \ln f_B - \ln f_{\neq} \tag{5-80}$$

where k_0 is the rate constant in an ideal solution [5, 28].

By substituting Eq. (5-79) into Eq. (5-80), the following relationship for the rate of reaction between the nonpolar reactants A and B is obtained [5, 28]:

$$\ln k = \ln k_0 + \frac{1}{RT}[V_A(\delta_A - \delta_s)^2 + V_B(\delta_B - \delta_s)^2 - V_{\neq}(\delta_{\neq} - \delta_s)^2] \tag{5-81}$$

where V_A, V_B, and $V_{\neq}$ are the molar volumes of A, B, and the activated complex, respectively. Thus, the rate constant depends not only on the difference in molar volumes between reactants and activated complex [called the volume of activation, $\Delta V^{\neq} = V_{\neq} - (V_A + V_B)$], but also on the relative cohesive pressure of reactants, activated complex, and solvent. If the reactants have a greater solubility in the solvent than the activated complex, the rate is lower compared to the rate in an ideal solution. The reverse is true if the activated complex is more soluble than the reactants.

Eq. (5-81), which is difficult to handle, can be changed into the linear Eq. (5-82):

$$\begin{aligned} RT \cdot \ln(k/k_0) = {} & \delta_s^2(V_A + V_B - V_{\neq}) + (V_A \cdot \delta_A^2 + V_B \cdot \delta_B^2 - V_{\neq} \cdot \delta_{\neq}^2) \\ & + 2\delta_s(V_{\neq} \cdot \delta_{\neq} - V_A \cdot \delta_A - V_B \cdot \delta_B) \end{aligned} \tag{5-82}$$

A reasonable assumption in some cases is that $V_{\neq} = V_A + V_B$, thus the first term in Eq. (5-82) becomes zero. The second term is constant for all solvents if molar volumes and cohesion pressures of reactants and activated complex are the same in these solvents. Thus, under certain conditions, the third term is the most important. In gas-phase reactions, only the second term is left ($\delta_s = 0$ for the gas phase).

Intuitively, one would expect that the volume of the reactants, the volume of the activated complex, and the corresponding activation volume, as well as the internal

pressure of the solvent*) will affect the reaction rate [27]. As early as 1929, Richardson and Soper [233], and later Glasstone [234], put forward rules which consider the influence of cohesion of reactants, products, and the solvent on reaction rates. They observed that reactions in which the products possessed greater (or lower) cohesion than the reactants were generally accelerated (or decelerated) by solvents with high cohesion, whereas reactions in which reactants and products were of similar cohesion, the solvent had relatively little influence on the reaction velocity. Although the reaction products are not involved in the activation process, according to the transition state theory, the above observations are still valid, since the cohesion of the activated complex may be regarded as lying somewhere between the values for the reactants and the products [27]. It has been pointed out, however, that variations in rate, caused by the internal pressure or cohesive pressure of the solvents, should be small in most solvents and generally should not exceed an order of magnitude [235].

Use of Eqs. (5-81) or (5-82) to predict solvent effects on reaction rates between nonpolar solutes is limited by the fact that there are no experimental data available for the heats of vapourization and hence for δ_i of most reacting organic compounds. However, the δ_s values are known for most organic solvents [231, 232, 236, 238]. A good approximation for the evaluation of $\delta_{\neq}$ is given by Eq. (5-83) [25, 28]:

$$\delta_{\neq} = \left(\frac{V_A \delta_A^2 + V_B \delta_B^2}{V_{\neq}} \right)^{1/2} \tag{5-83}$$

Nevertheless, it is possible to obtain correlations between reaction rates and δ_s^2, as shown in Figs. 5-8 and 5-9 (for further examples, see references [25, 126]).

Fig. 5-8 shows that there is a rough correlation of $\lg(k/k_0)$ with δ_s^2 for the Diels–Alder dimerization of cyclopentadiene [35], as predicted by Eq. (5-81). The molar volumes of cyclopentadiene and the corresponding activated complex have been determined as 83.1 and 135 cm^3, respectively [126].

Fig. 5-9 shows that there is also a very rough, inverse correlation between $\lg(k/k_0)$ and δ_s^2 for the dissociation of the dimer of the triphenylmethyl radical [167]. It can be safely assumed that in this unimolecular reaction the molar volume of the activated complex is greater than the molar volume of the reactant, since a bond breaking must occur to some extent on activation.

The scatter of the points in the two figures is not surprising in view of the simplifications made in deriving Eq. (5-81). Furthermore, the solvent effects observed in these reactions are very small with a comparatively large experimental error, and the solvents used include such dipolar and associated liquids as nitromethane and alcohols.

In principle, it is possible to distinguish between reactions with negative activation volume $\Delta V^{\neq}$ $(=V_{\neq} - V_{\text{reactants}})$ and those with positive activation volume, since the activated complex for the former occupies less volume than the reactants, according to the schematic Eqs. (5-84) and (5-85) [239].

---

* How an external pressure may affect the rate of a reaction in solution is considered in Section 5.5.11. The activation volume of a reaction is usually obtained from external pressure measurements.



Fig. 5-8. Correlation of $\lg(k/k_0)$ [35] and the cohesive pressure δ_s^2 [238] in the Diels–Alder dimerization of cyclopentadiene at 40 °C (rate constants relative to acetone as 'slowest' solvent):
1) diethyl ether, 2) tetrachloromethane, 3) toluene, 4) tetrahydrofuran, 5) benzene, 6) trichloromethane, 7) chlorobenzene, 8) dichloromethane, 9) acetone, 10) 1,4-dioxane, 11) *tert*-butanol, 12) 1-butanol, 13) 1-propanol, and 14) ethanol. The point for methanol shows a large deviation and is not included in this figure.

Since in reaction (5-84) the reactants form a more compact activated complex, it is assumed that $(\delta_A + \delta_B) < \delta_{\neq}$. In the case of Eq. (5-85), $(\delta_{A-B}) > \delta_{\neq}$ because a loos-



(5-84)



(5-85)

ening of bonds occurs on activation. If the nonpolar reaction is of type (5-84), *e.g.* dimerization of cyclopentadiene, $\ln k$ will increase as δ_s increases, whereas for reactions of type (5-85), such as the unimolecular dissociation of the triphenylmethyl dimer, $\ln k$ will decrease as δ_s increases. Eq. (5-81) therefore predicts that the internal pressure (or the cohesive pressure) of solvents should influence reaction rates of nonpolar reactions in the same direction as external pressure (*cf.* also Section 5.5.11) [27, 232, 239].

The solvent influence on rates of bimolecular H-atom-transfer reactions $R\cdot + H{-}X \rightarrow R{-}H + \cdot X$ has been theoretically studied [580]. Rates for the model



Fig. 5-9. Correlation of $\lg(k/k_0)$ [167] and the cohesive pressure δ_s^2 [238] in the dissociation of 1-diphenylmethylene-4-triphenylmethyl-2,5-cyclohexadiene ("hexaphenylethane") at 0 °C; *cf.* Eq. (5-56) in Section 5.3.4 (rate constants relative to acetonitrile as 'slowest' standard solvent):
1) ethyl benzoate, 2) diethyl oxalate, 3) tetrachloromethane, 4) 1-bromopropane, 5) toluene, 6) 4-methyl-3-penten-2-one, 7) trichloromethane, 8) styrene, 9) propionic acid, 10) *N*,*N*-diethylacetamide, 11) carbon disulfide, 12) nitrobenzene, 13) aniline, 14) methyl salicylate, 15) pyridine, 16) ethyl cyanoacetate, 17) ethylene glycol monomethyl ether, 18) acetonitrile, 19) ethanol, and 20) nitromethane.

H-exchange reaction between $CH_3\cdot$ and CH_4 have been compared in the gas-phase and in rare gas solution (compressed Ar and Xe as solvents) over a wide range of internal pressures. Depending on the solvent and its internal pressure, relatively large rate enhancements have been calculated for this methyl/methane H-abstraction reaction. The major reason for this static solvent enhancement of rate in rare gas solutions seems to be the compact (tight) character of the activated complex, $[CH_3^{\delta\odot}\cdots H\cdots{}^{\delta\odot}CH_3]^{\neq}$ [580].

The concept of cohesive pressure (or internal pressure) is useful only for reactions between neutral, nonpolar molecules in nonpolar solvents, because in these cases other properties of the solvents, such as the solvation capability or solvent polarity, are neglected. For reactions between dipolar molecules or ions, the solvents interact with reactants and activated complex by unspecific and specific solvation so strongly that the contribution of the cohesive pressure terms of Eq. (5-81) to $\ln k$ is a minor one. It should be mentioned that cohesive pressure or internal pressure are not measures of solvent polarity. Solvent polarity reflects the ability of a solvent to interact with a solute, whereas cohesive pressure, as a structural parameter, represents the energy required to create a hole in a particular solvent to accommodate a solute molecule. Polarity and cohesive pressure are therefore complementary terms, and rates of reaction will depend

on both of them [27, 232]. The influence of solvent polarity on reaction rates will be discussed in the following Sections.

5.4.3 Reactions Between Neutral, Dipolar Molecules

In reactions between neutral, dipolar molecules, the electrostatic solute/solvent interactions such as dipole–dipole forces should above all determine the reaction rates in a manner as described qualitatively by the Hughes–Ingold theory [16, 44]. In order to describe electrostatic solvent effects, one considers the interaction between charged points separated by an isotropic dielectric continuum using functions of the relative permittivity. In rate equations involving dipolar reactants, the well-known equation of Kirkwood [240] is generally applicable. The Kirkwood expression relates the standard Gibbs free energy of transfer of a spherical dipolar molecule of radius r and dipole moment μ from the gas phase ($\varepsilon_r = 1$) to a continuous medium (with $\varepsilon_r > 1$) according to

$$\Delta G^{\circ}_{solv} = -\frac{N_A}{4\pi \cdot \varepsilon_0} \cdot \frac{\mu^2}{r^3} \cdot \frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \tag{5-86}$$

where ε_0 is the permittivity of vacuum and N_A is the Avogadro constant. This important formula relates the changes in Gibbs solvation energy of the dipolar solute both to the relative permittivity of the solvent and to the dipole moment and radius of the solute molecule, taking into account electrostatic forces only between solute and solvent molecules*).

Applying Kirkwood's formula to the transition-state theory for the bimolecular reaction $A + B \rightleftharpoons (AB)^{\neq} \rightarrow C + D$ and combining Eq. (5-86) with Eq. (5-75), one obtains an expression for the rate constant of a reaction between two dipolar molecules A and B with moments μ_A and μ_B to form an activated complex with dipole moment $\mu_{\neq}$ [2]:

$$\ln k = \ln k_0 - \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{N_A}{R \cdot T} \cdot \frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \cdot \left(\frac{\mu_A^2}{r_A^3} + \frac{\mu_B^2}{r_B^3} - \frac{\mu_{\neq}^2}{r_{\neq}^3} \right) \tag{5-87}$$

where k is the rate constant in the medium of relative permittivity ε_r, and k_0 is the rate constant in a condensed medium with $\varepsilon_r = 1$ Eq. (5-87) predicts that, if the activated complex is more dipolar than the reactants, the rate of the reaction increases with the relative permittivity of the medium.

Based on the same premises, but with some modification of the electrostatic model introduced by Kirkwood, Laidler, and Landskroener obtained another similar

* It should be noted that Kirkwood's formula does not appear in his first publication [240] in the form of Eq. (5-86). Nevertheless, it is this form of Kirkwood's formula which is widely known, representing only one of the terms of a more complex equation given in reference [240] with $n = 1$. Kirkwood's theory was further developed in papers by Kirkwood, Westheimer, and Tanford [241], Laidler and Landskroener [242], and Hiromi [243].

expression for the reaction of two dipole molecules in a medium as given by Eq. (5-88) [11, 242]:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{3 \cdot N_A}{8 \cdot R \cdot T} \cdot \left(\frac{2}{\varepsilon_r} - 1 \right) \cdot \left(\frac{\mu_A^2}{r_A^3} + \frac{\mu_B^2}{r_B^3} - \frac{\mu_{\neq}^2}{r_{\neq}^3} \right) \tag{5-88}$$

This equation predicts that a plot of $\ln(k/k_0)$ versus $1/\varepsilon_r$ should give a straight line, and gives an explicit expression for the slope s of this line in terms of the radii and dipole moments as shown by Eq. (5-89):

$$s = \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{3 \cdot N_A}{8 \cdot R \cdot T} \cdot \left(\frac{\mu_A^2}{r_A^3} + \frac{\mu_B^2}{r_B^3} - \frac{\mu_{\neq}^2}{r_{\neq}^3} \right) \tag{5-89}$$

If a reaction between neutral, dipolar molecules occurs with the formation of an activated complex with a dipole moment $\mu_{\neq}$ greater than either μ_A or μ_B, there will be an increase in the rate constant with increasing ε_r according to Eq. (5-88). This is because a medium with higher ε_r favours the production of any highly dipolar species as, in this case, the activated complex. In applying Eqs. (5-87) and (5-88) to experimental data, a model for the activated complex has to be constructed in order to evaluate reasonable values for $\mu_{\neq}$ and $r_{\neq}$. This has been done, for example, for the acid and base hydrolysis of carboxylic esters [11, 242].

Another Coulombic energy approach for the calculation of electrostatic solvent effects on reactions between dipolar molecules was made by Amis [12, 21, 244]. He related the rate constant to the energy of activation by the well-known Arrhenius equation $k = A \cdot \exp(-E_a/RT)$. It is assumed that the effect of the relative permittivity on the rate is given by Eq. (5-90):

$$\ln k = \ln k_{\infty} - \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{2 \cdot \mu_A \cdot \mu_B \cdot N_A}{R \cdot T \cdot \varepsilon_r \cdot r^3} \tag{5-90}$$

where k is the rate constant in any medium of relative permittivity ε_r, k_{∞} is the rate constant in a medium of relative permittivity with a value of infinity, and μ_A and μ_B are the dipole moments of the two dipolar reactants A and B in vacuum. Eq. (5-90) predicts that a plot of $\ln k$ versus $1/\varepsilon_r$ will give a straight line, the slope of which should give a reasonable value of $r = r_A + r_B$, the distance of approach for the two dipolar molecules to react.

It should be mentioned that there is no inconsistency in the fact that $\ln(k/k_0)$ is a function of $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ according to Eq. (5-87), and of $1/\varepsilon_r$ according to Eqs. (5-88) and (5-90). It can readily be seen, by carrying out explicitly the division of $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ according to Eq. (5-91), that $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ is linear in $1/\varepsilon_r$ to a good approximation [40].

$$\frac{\varepsilon_r - 1}{2\varepsilon_r + 1} = \frac{1}{2} - \frac{3}{4\varepsilon_r} + \frac{3}{8\varepsilon_r^2} - \frac{3}{16\varepsilon_r^3} + \cdots \tag{5-91}$$

For $\varepsilon_r = 8$, the error involved in neglecting terms in ε_r^2 and higher powers is less than one percent [40].

Laidler has pointed out [11, 242] that Eq. (5-88) is best considered as a semi-quantitative formulation, which gives only a rough prediction of the effect of a change in ε_r on the rate of dipole–dipole reactions. This also applies to Eqs. (5-87) and (5-90). Nevertheless, in many cases a satisfactory correlation between rate constants and function of solvent relative permittivity has been obtained, as, for instance, in the Menschutkin reaction between trialkylamines and haloalkanes forming quaternary tetraalkylammonium salts [2, 56, 58, 60, 61, 64, 65, 245–247].

Glasstone, Laidler, and Eyring [2] were the first to correlate rate data for some Menschutkin reactions according to Eq. (5-87), and they found, in particular, that a linear correlation between $\lg(k/k_0)$ and $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ is observed in the binary solvent mixture benzene/alcohol, while in benzene/nitrobenzene a monotonous deviation from the linear dependence is observed.

Fig. 5-10 shows typical dependences of $\lg(k/k_0)$ on the Kirkwood parameter $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ obtained for the Menschutkin reaction between triethylamine ($\mu = 2.9 \cdot 10^{-30}$ Cm = 0.9 D) and iodoethane ($\mu = 6.3 \cdot 10^{-30}$ Cm = 1.9 D) in binary solvent mixtures [56].



Fig. 5-10. Correlation between $\lg(k/k_0)$ [56] and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for the Menschutkin reaction between triethylamine and iodoethane at 40 °C in binary acetone/benzene and acetone/1,4-dioxane mixtures (rate constants relative to acetone as common standard solvent).

Binary solvent mixtures have the advantage that changing their composition will predominately change the electrostatic solute/solvent interactions, whereas non-electrostatic and specific interactions remain the same within the whole solvent series and will be cancelled out. Fig. 5-10 shows that Kirkwood's equation holds for such solvent systems, but the slopes of the straight lines are markedly different. One might, however, expect them to be the same if Eq. (5-87) were applicable.

If one compares the rate constants for the same Menschutkin reaction with Kirkwood's parameter in thirty-two pure aprotic and dipolar non-HBD solvents [59, 64], one still finds a rough correlation, but the points are widely scattered as shown in Fig. 5-11.



Fig. 5-11. Correlation between $\lg(k/k_0)$ [59, 64] and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for the Menschutkin reaction between triethylamine and iodoethane in 32 aprotic and dipolar aprotic solvents at 25 °C (rate constants relative to *n*-hexane as 'slowest' solvent):
1) *n*-hexane, 2) cyclohexane, 3) tetrachloromethane, 4) 1,4-dioxane, 5) benzene, 6) toluene, 7) diethyl ether, 8) iodobenzene, 9) trichloromethane, 10) bromobenzene, 11) chlorobenzene, 12) ethyl benzoate, 13) ethyl acetate, 14) 1,1,1-trichloroethane, 15) chlorocyclohexane, 16) bromocyclohexane, 17) tetrahydrofuran, 18) 1,1-dichloroethane, 19) 1,1,2,2-tetrachloroethane, 20) dichloromethane, 21) 1,2-dichloroethane, 22) acetophenone, 23) 2-butanone, 24) acetone, 25) propionitrile, 26) benzonitrile, 27) nitrobenzene, 28) *N*,*N*-dimethylformamide, 29) acetonitrile, 30) nitromethane, 31) dimethyl sulfoxide, and 32) propylene carbonate.

The values of the second-order rate constants are taken from the compilation made by M. H. Abraham and P. L. Grellier, J. Chem. Soc., Perkin Trans. 2 *1976*, 1735.



Fig. 5-12. Correlation between $\lg(k/k_0)$ [65] and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for the Menschutkin reaction between 1,4-diazabicyclo[2.2.2]octane and (2-bromoethyl)benzene in non-HBD (○) and HBD (●) solvents at 54.5 °C (rate constants relative to tetrachloromethane as 'slowest' standard solvent).

Extending the media used for the Menschutkin reaction to protic solvents such as alcohols leads to an even worse correlation, as shown in Fig. 5-12 for the quaternization of 1,4-diazabicyclo[2.2.2]octane with (2-bromoethyl)benzene studied in a total of thirty-six solvents [65]. The group of protic solvents is separated from the assembly of non-HBD solvents, each group showing a very rough but distinct correlation with the function of relative permittivity. Such behaviour has also been observed for several other Menschutkin reactions [60, 61].

Although Eq. (5-87) is often qualitatively obeyed, as has been frequently mentioned, there is no exact linear correlation between the rate of Menschutkin reactions and the functions of relative permittivity as in the case of Fig. 5-12 [246, 247]. A complete absence of a regular effect of changes in the dielectric properties of the solvent on the reaction rate has also been observed [248, 249]. Sometimes a satisfactory correlation has been obtained because the reaction under consideration was studied in only a limited

number of solvents. It can readily be seen from Fig. 5-11 that studying this Menschutkin reaction by chance in the six solvents nos. 3, 11, 12, 26, 27, and 28 would simulate a very good correlation.

The fact that the curves of Figs. 5-10 to 5-12 (and others) have positive slopes is direct evidence for a partially ionized, dipolar activated complex. In cases for which Eq. (5-87) is reasonably well obeyed (especially for binary solvent mixtures), a reasonable value for $\mu_{\neq}$ may be calculated from the slope according to Eq. (5-89) [2, 8] (for a compilation of such calculations, see references [23, 64]). The calculated dipole moments $\mu_{\neq}$ range from *ca.* $17 \cdot 10^{-30}$ Cm (*ca.* 5 D) to *ca.* $30 \cdot 10^{-30}$ Cm (*ca.* 9 D) and show a considerable separation of charges for the activated complex of Menschutkin reactions, in agreement with the observed rate accelerations with increasing solvent polarity. However, it seems probable that if Eq. (5-87) is to hold over a substantial range of solvent compositions, rather large-scale cancellations of non-electrostatic effects must take place [23].

As the data for the Menschutkin reactions indicate, the character of the solute–solvent interactions is more complex than described by Eq. (5-87). It is evident that functions of relative permittivity alone, as given in Eq. (5-87), are not useful for describing the solvent effect on reactions between dipolar reactants, except in certain special cases, such as when a mixture of two solvents is used. In addition to electrostatic forces, non-electrostatic interactions, such as dispersion forces and hydrogen-bonding, must also be involved in Menschutkin reactions.

As seen from Fig. 5-11, although the three halobenzenes (points nos. 8, 10, and 11) have similar values of ε_r, when used as solvents they lead to different reaction rates. Iodobenzene (no. 8), with the lowest ε_r value, gives the largest rate. This observation strongly suggests that the polarizability of the solvent is an important factor in stabilizing the dipolar activated complex of this reaction. This was confirmed by Reinheimer *et al.* [57], who studied some Menschutkin reactions in benzene and its chloro, bromo, and iodo derivatives. They showed that the rate of the reaction increases with increasing polarizability of the solvent.

In case of alcoholic, protic solvents, it should be particularly noted from Fig. 5-12 that the values of $\lg(k/k_0)$ are much lower than expected on the basis of their relative permittivities. It has been pointed out [64] that hydrogen bonding between the alcohols and the trialkylamine – thus reducing the Gibbs energy of the reactants – is not the only cause of this low accelerating power. This anomalous effect of aliphatic alcohols on the rates of Menschutkin reactions is due to the fact that the activated complex for these reactions resembles a relatively nonpolar solute. Its structure lies somewhere between reactants and ion pair, probably nearer to the reactants. Compared to the situation in dipolar non-HBD solvents, which may effect stabilization through nonspecific interaction [64], such an activated complex will be destabilized by aliphatic alcohols. Within a series of aliphatic alcohols and in spite of the fact that the values of ε_r fall from 32.7 to 3.4, the reaction between bromoethane and *N*,*N*-dimethylaniline is accelerated by a factor of almost ten on going from methanol to octanol [245]. A detailed discussion of all aspects of the Menschutkin reaction, including solvent effects, can be found in an excellent review by Abboud *et al.* [786].

Eq. (5-87), which is a general rate equation predicting the effect of solvent on bimolecular reactions between dipolar molecules, can readily be modified to treat



Fig. 5-13. Correlation between lg k_1 [40] and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for the solvolysis of 2-chloro-2-methylpropane in binary 1,4-dioxane/water (A), ethanol/water (B), acetone/water (C), and methanol/water (D) mixtures at 25 °C; *cf.* Eq. (5-13) in Section 5.3.1.

unimolecular reactions of dipolar molecules A by neglecting the term for the second reactant B. A classical example of this reaction type is the solvolysis of 2-chloro-2-methylpropane ($\mu = 7.1 \cdot 10^{-30}$ Cm = 2.1 D). This is an example of an S_N1 reaction, the solvent dependence of which has been studied in many solvents and binary solvent mixtures (*cf.* Eq. (5-13) in Section 5.3.1) [40, 47, 250].

Assuming that the influence of non-electrostatic terms and specific solute/solvent interactions are negligible, a linear relationship is expected between $\lg(k/k_0)$ and the Kirkwood parameter $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$. Such plots are shown in Figs. 5-13 and 5-14 for binary solvent mixtures and pure solvents. It can be seen that there is no simple relationship between $\lg(k/k_0)$ and $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for any of the solvents and solvent mixtures examined. As seen from Fig. 5-13, there is a large dispersion of the plot into separate lines for each solvent pair and curvature of the separate lines. Fig. 5-14 shows that there is a very poor correlation for pure solvents. In protic solvents, as a result of considerable specific solvation of the leaving anion through hydrogen bonding, the rate of solvolysis is higher than that expected from the relative permittivities of the solvents.

A somewhat better correlation using values of lg k_1 at 120 °C was obtained by Koppel and Pal'm [250], but again protic and some non-HBD solvents do not conform to the expected linear pattern. From the slope of the estimated straight line, a value of $\mu_{\neq} = 31 \cdot 10^{-30}$ Cm = 9.2 D for the activated complex was calculated, which although reasonable must treated with caution, since the slope of the line depends largely on only a few values of lg k_1 in non-HBD solvents [250]. Nevertheless, it indicates that in the activated complex the charges are considerably separated, in agreement with the qualitative considerations made in Section 5.2.1.

The breakdown of the simple linear relationship between $\lg(k/k_0)$ and $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, required by Eq. (5-87), is obviously due to the failure of the approximations involved in deriving this equation, neglection of non-electrostatic and specific solute–solvent interactions, and, in the case of binary solvent mixtures, due also to the selective solvation of the reactants and activated complex by one component of the mixture (*cf.*



Fig. 5-14. Correlation between $\lg(k/k_0)$ [40, 47] and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ for the solvolysis of 2-chloro-2-methylpropane in 17 pure non-HBD (○) and HBD (●) solvents at 25 °C (rates relative to diethyl ether as 'slowest' standard solvent). Values of lg k are taken from the compilation in reference [47].
1) 1,4-dioxane, 2) benzene, 3) diethyl ether, 4) chlorobenzene, 5) acetone, 6) nitrobenzene, 7) 1-methyl-2-pyrrolidinone, 8) acetonitrile, 9) *N*,*N*-dimethylformamide, 10) nitromethane, 11) acetic acid, 12) 1-butanol, 13) ethanol, 14) methanol, 15) formamide, 16) formic acid, and 17) water.

Section 2.4). It can be unequivocally concluded from a large amount of literature data, that the number of instances of deviation for reactions between dipolar molecules from the simple electrostatic relationship, given in Eqs. (5-87), (5-88), and (5-90), greatly exceeds the number of cases where the relationships hold. Thus, allowances must also be made for electrostatic, nonelectrostatic, and specific solvation effects in the general effect of the medium.

It would appear from these observations that the solvation capability might be better characterized using a linear Gibbs energy relationship approach than functions of relative permittivity. There are now numerous examples known, for which the correlation between the rates of different reactions and the solvation capability of the solvent can be satisfactorily described with the help of semiempirical parameters of solvent polarity (*cf.* Chapter 7).

5.4.4 Reactions between Neutral Molecules and Ions

Numerous organic reactions are of the ion-dipole type, as, for example, the S_N2 reactions given in Eqs. (5-17) and (5-18) in Section 5.3.1. Considering the reaction between an ion A of charge $z_A \cdot e$ and a neutral, dipolar molecule B of dipole moment μ_B according to $A^{z_A \cdot e} + B \rightleftharpoons (AB)^{\neq z_A \cdot e} \rightarrow C + D$, Laidler and Eyring [2, 251] obtained Eq. (5-92) for the rate constant in a medium of zero ionic strength:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \left[\frac{z_A^2 \cdot e^2 \cdot N_A}{2 \cdot R \cdot T} \cdot \left(\frac{1}{\varepsilon_r} - 1 \right) \cdot \left(\frac{1}{r_A} - \frac{1}{r_{\neq}} \right) - \frac{N_A}{R \cdot T} \cdot \frac{\mu_B^2}{r_B^3} \cdot \left(\frac{\varepsilon_r - 1}{2\varepsilon_r + 1} \right) \right] \tag{5-92}$$

where k and k_0 are the rate constants in media of relative permittivity ε_r and unity, respectively, and N_A is Avogadro's number. This equation predicts that $\ln k$ plotted against $1/\varepsilon_r$ for a molecule of zero dipole moment reacting with an ion of charge $z_A \cdot e$ should give a straight line, the slope of which would be $z_A^2 e^2 N_A / 2RT(1/r_A - 1/r_{\neq})$. This relationship should be especially true if the rate of the reaction is studied in mixtures of two solvents so that the relative permittivity can be varied by changing the proportions of each solvent. Since $r_{\neq}$ will be larger than r_A, the rate should be somewhat greater in a medium of lower values of ε_r. A quantitative test of Eq. (5-92) would require reasonable values of r_A and $r_{\neq}$. There seems to be no published data, however, that confirm Eq. (5-92).

A somewhat different equation relating the rate constant to the relative permittivity for a reaction $A^{z_A \cdot e} + B^{z_B \cdot e} \rightleftharpoons (AB)^{\neq (z_A + z_B)e} \rightarrow C + D$, in which the electrostatic interactions are more important than nonelectrostatic ones, was derived by Laidler and Landskroener [11, 242]:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \left[\frac{e^2 \cdot N_A}{2 \cdot R \cdot T} \cdot \left(\frac{1}{\varepsilon_r} - 1 \right) \cdot \left(\frac{z_A^2}{r_A} + \frac{z_B^2}{r_B} - \frac{(z_A + z_B)^2}{r_{\neq}} \right) + \frac{3 \cdot N_A}{8 \cdot R \cdot T} \cdot \left(\frac{2}{\varepsilon_r} - 1 \right) \cdot \left(\frac{\mu_A^2}{r_A^3} + \frac{\mu_B^2}{r_B^3} - \frac{\mu_{\neq}^2}{r_{\neq}^3} \right) \right] \tag{5-93}$$

For a reaction between two dipoles having no net charge, the second term disappears, and the solvent effect is given entirely by the last term; Eq. (5-93) then equals Eq. (5-88) in Section 5.4.3. For a reaction between an ion and a dipole (or between two charged dipoles) both terms must be included. The simplest case is for the reaction of a monovalent, structureless ion A of charge $z_A \cdot e (\mu_A = 0)$ with a neutral molecule B $(z_B \cdot e = 0)$ of dipole moment μ_B. Eq. (5-93) then has the form:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \left[\frac{e^2 \cdot N_A}{2 \cdot R \cdot T} \cdot \left(\frac{1}{\varepsilon_r} - 1 \right) \cdot \left(\frac{1}{r_A} - \frac{1}{r_{\neq}} \right) + \frac{3 \cdot N_A}{8 \cdot R \cdot T} \cdot \left(\frac{2}{\varepsilon_r} - 1 \right) \cdot \left(\frac{\mu_B^2}{r_B^3} - \frac{\mu_{\neq}^2}{r_{\neq}^3} \right) \right] \tag{5-94}$$

This equation predicts that $\ln k$ will vary linearly with the reciprocal of the relative permittivity, and gives an explicit expression for the slope in terms of radii and dipole moments. If data are available for a series of mixed solvents, one procedure for the application of this equation is to plot $\ln k$ against $1/\varepsilon_r$ and to determine the slope. Then it is possible to see whether its sign and magnitude may be predicted, using reasonable values for the radii and the dipole terms, in terms of Eq. (5-94). Of course, this procedure will depend on devising a suitable model for the activated complex. For Eq. (5-94), the same restrictions are valid as for Eq. (5-88). At best, Eq. (5-94) gives a semiquantitative formulation, which allows only very rough predictions.

Another derivation of the relative permittivity dependence of $\ln k$ has been made by Amis [12, 21, 244] using a Coulomb energy approach for the ion–dipole interaction. Considering the mutual potential energy between an ion A of charge $z_A \cdot e$ and a dipole B of dipole moment μ_B at a distance r_{AB} leads eventually to Eq. (5-95):

$$\ln k = \ln k_\infty + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{z_A \cdot e \cdot \mu_B \cdot N_A}{R \cdot T \cdot \varepsilon_r \cdot r_{AB}^2} \tag{5-95}$$

where k_∞ is the rate constant in a medium with relative permittivity of infinite magnitude. Taking the charge of the ion into account, Eq. (5-95) predicts that a plot of $\ln k$ versus $1/\varepsilon_r$ should be a straight line of positive slope if $z_A \cdot e$ is positive, and of negative slope if $z_A \cdot e$ is negative. This equation has been applied to both positive and negative ionic reactants reacting with dipole molecules [12, 21, 57, 252]. Illustrative examples of the applicability of Eqs. (5-94) or (5-95) are the alkaline hydrolysis of methyl propionate ($\mu = 6 \cdot 10^{-30}$ Cm = 1.8 D) in acetone/water mixtures [252], and the S_N2 reaction between the azide anion and 1-bromobutane ($\mu = 7 \cdot 10^{-30}$ Cm = 2.1 D) in pure dipolar non-HBD solvents [67]. In the case of alkaline hydrolysis, the rate-determining step would be that between the negatively charged hydroxide ion and the dipolar ester. As suggested by the above mentioned equations, plots of $\lg(k/k_\infty)$ versus $1/\varepsilon_r$ give the requisite straight lines as shown for both reactions in Fig. 5-15. The positive slope of the line for the S_N2 reaction indicates that in non-HBD solvents $r_{\neq} > r_A$, according to Eq. (5-94), whereas the negative slope of the line for the ester hydrolysis in protic solvents requires that $r_A > r_{\neq}$, which can be explained in terms of specific solvation of the hydroxide ion by hydrogen bonding [67, 252]. Further examples of the application of Eq. (5-95) are given in references [12, 21].

5.4.5 Reactions between Ions

The combination of simple inorganic ions usually proceeds very rapidly and the rate of reaction is governed by the diffusion of the ions towards each other. There are, however,



Fig. 5-15. Correlation between $\lg(k/k_\infty)$ and $1/\varepsilon_r$ for the alkaline hydrolysis of methyl propionate in eight acetone/water mixtures at 25 °C (○) [252], and for the S_N2 reaction between the azide anion and 1-bromobutane in six pure dipolar non-HBD solvents at 25 °C (●) [67] (rate constants relative to the solvent with the largest dielectric constant).

many reactions between ions which involve the making and breaking of covalent bonds or electron transfer, which may be as slow as reactions between neutral molecules (*cf.*, for example, reaction (5-19) in Section 5.3.1). Only reactions of this type will be considered in this Section.

Again, electrostatic theory will be useful in delineating the general effect of the changes on the ionic reactants and the relative permittivity of the solvent. The reaction between ions was the first reaction to which the simple electrostatic model was applied to calculate the effect of the relative permittivity and the ionic strength of the medium on the reaction rate. The change in Gibbs energy during the formation of an ion pair from the ions A and B in a standard medium with relative permittivity ε_r is equal to the electrostatic energy of the approach of two point charges $z_A \cdot e$ and $z_B \cdot e$ to a distance r_{AB} according to the following equation (N_A is Avogadro's number):

$$\Delta G_{\text{electrostatic}} = \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{z_A \cdot z_B \cdot e^2 \cdot N_A}{\varepsilon_r \cdot r_{AB}} \tag{5-96}$$

From this equation, which applies to infinitely dilute solutions only, it is possible to write an expression for the dependence of the rate constant on the relative permittivity at zero ionic strength:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{z_A \cdot z_B \cdot e^2 \cdot N_A}{R \cdot T \cdot r_{AB}} \cdot \left(1 - \frac{1}{\varepsilon_r}\right) \tag{5-97}$$

where k and k_0 are the rate constant in media of relative permittivity ε_r and unity (*i.e.* in the gas phase), respectively. r_{AB} is the distance to which the two ionic reactants must approach in order to react and can be assumed to be equal to $r_A + r_B$. Eq. (5-97), first proposed by Scatchard [253], predicts a linear plot of $\ln k$ versus $1/\varepsilon_r$ with a negative slope if the charges of the ions have the same sign, and a positive slope if the charges are of opposite sign.

For the reaction $A^{z_A \cdot e} + B^{z_B \cdot e} \rightleftharpoons (AB)^{\neq (z_A + z_B)e} \rightarrow C + D$, Laidler and Eyring [2, 251] derived another expression for the solvent dependence of rate constants using a slightly different model for the distribution of charges in the activated complex:

$$\ln k = \ln k_0 + \frac{1}{4\pi \cdot \varepsilon_0} \cdot \frac{e^2 \cdot N_A}{2 \cdot R \cdot T} \cdot \left(\frac{1}{\varepsilon_r} - 1\right) \cdot \left(\frac{z_A^2}{r_A} + \frac{z_B^2}{r_B} - \frac{(z_A + z_B)^2}{r_{\neq}}\right) \tag{5-98}$$

This equation is equivalent to the second term of Eq. (5-93). For reactions between two simple ions, the final term of Eq. (5-93) is less important and can generally be neglected. Eq. (5-98) is valid for reactions carried out in solutions of infinite dilution only, and reduces to Eq. (5-97) if $r_A = r_B = r_{\neq}$.

Some tests of Eqs. (5-97) and (5-98) have been made, particularly by carrying out a reaction between ions in a series of mixed solvents of varying values of ε_r. Examples are the reaction between bromoacetate and thiosulfate ions [11, 251], the reaction between negative divalent tetrabromophenolsulfophthalein ions and hydroxide ions [254], and the ammonium cation/cyanate anion reaction yielding urea [255]. On the whole, the relationships (5-97) and (5-98) are a good approximation, although there are usually serious deviations especially at low values of ε_r (obviously due to association between unlike ions). Deviations from linearity may be explained as being due to the failure of the simple approximations made in deriving these equations, due to selective solvation in binary solvent mixtures, and due to the neglection of such phenomena as the mutual polarization of ions or dipoles and the resulting inductive interactions, as well as specific solvation.

The entropy changes of reactions between ions deserve special mention. For reactions between ions of unlike sign there is an entropy increase in going from reactants to activated complex. This is because the activated complex will have less charge than the reactants and will become partially desolvated. For ions of the same sign, however, the activated complex will be more charged than the reactants. This one would expect to be strongly solvated. There is a loss of entropy, therefore, when the activated complex is formed. For examples which confirm these relationships, see references [5, 11, 20].

Since the activity coefficients of the reactants vary with the ionic strength I of the solution, the value of which is determined by the molar concentrations c_i of the ions i of charge z_i according to $I = 1/2 \cdot \sum_i c_i \cdot z_i^2$, the rate of reactions in solution must also depend on the ionic strength. This effect of the ionic strength of the solution on the rate constant described by Eq. (5-99) is termed the *Primary Salt Effect.*

$$\ln k = \ln k_0 + 2 \cdot z_A \cdot z_B \cdot A \cdot \sqrt{I} \tag{5-99}$$

A is a constant for a given solvent and temperature ($A = 0.51 \cdot 2.303\,L^{1/2} \cdot mol^{-1/2}$ for aqueous solutions at 25 °C), and k_0 is the rate coefficient extrapolated to infinite dilution. Eq. (5-99), which was first derived by Brønsted [256], Bjerrum [257], and Christiansen [258] applying the Debye–Hückel theory [259] to the influence of neutral salts upon reaction rates in solution, predicts a linear relationship if ln k is plotted against the square root of the ionic strength. Since the simple Debye–Hückel relationship does not apply at those concentrations where ion association becomes important, Eq. (5-99) is only valid for solutions below *ca.* 10^{-2} mol/L for 1-1 electrolytes. Eq. (5-99) is perfectly accurate in dilute solutions.

Its conceptual significance is closely related to the Hughes–Ingold rules as given in Table 5-4 in Section 5.3.1. An increase in the ionic strength of a solution leads to an increase in its polarity. Thus, reactions between ions of like charge ($z_A \cdot z_B$ positive) are accelerated by an increase in the ionic strength, *i.e.* the addition of electrolytes, whereas reactions between ions of opposite charge ($z_A \cdot z_B$ negative) are retarded. If one of the reactants is a neutral molecule ($z_A \cdot z_B$ zero), the rate is expected to be independent of the ionic strength to a first approximation.

However, even in reactions involving neutral molecules ($z_A \cdot z_B$ zero), electrolyte effects can be found. Such reactions between neutral, nonpolar molecules can yield ionic products *via* a dipolar activated complex. This dipolar activated complex can be stabilized by an ionic medium, and a large positive salt effect is found. In reactions such as the solvolyses of 2-halo-2-methylpropanes, ions are produced and the ionic strength increases automatically during the reaction. This can lead to a kind of autocatalytic acceleration of the solvolysis. Small negative salt effects are frequently found in reactions between ions and neutral molecules because there is a greater charge dispersion during the activation process.

Eq. (5-99) was checked in detail in a large number of studies, but an extensive treatment of primary and secondary salt effects on reaction rates is beyond the scope of this Section. The reader is referred to references [2–5, 11, 12, 19–21, 28], particularly to the excellent comprehensive reviews of Davies [260], Blandamer *et al.* [828], and Loupy *et al.* [829].

In conclusion, it can be said that the electrostatic theory of solvent effects is a most useful tool for explaining and predicting many reaction patterns in solution. However, in spite of some improvements, it still does not take into account a whole series of other solute/solvent interactions such as the mutual polarization of ions or dipoles, the specific solvation *etc.*, and the fact that the microscopic relative permittivity around the reactants may be different to the macroscopic relative permittivity of the bulk solvent. The deviations between observations and theory, and the fact that the relative permittivity cannot be considered as the only parameter responsible for the changes in reaction rates in solution, has led to the creation of different semiempirical correlation equations, which correlate the kinetic parameters to empirical parameters of solvent polarity (see Chapter 7).

5.5 Specific Solvation Effects on Reaction Rates

The intermolecular forces responsible for the association of dissolved ions or molecules with solvent molecules, consist of other forces beside the non-specific directional, induc-

tive, and dispersion forces which formed the essential basis for the quantitative relationships discussed in the preceding chapter. This other class of intermolecular forces consists of specific interactions such as hydrogen bonding and EPD/EPA forces (*cf.* Section 2.2). In the following discussion it will be shown, with the aid of some representative examples, how a consideration of the specific solvation of the reactants as well as the activated complexes can contribute to a better understanding of the rates, mechanisms, and steric outcome of organic reactions. In this discussion, not even an approximately complete treatment of the multitude of specific solvent effects can be attempted; for further reading, see references [6, 10, 21, 26, 261–263, 452–454].

5.5.1 Influence of Specific Anion Solvation on the Rates of S_N and other Reactions

In aliphatic nucleophilic substitution reactions, the solvation of the departing anions is particularly important. In protic solvents, this takes place mainly through hydrogen bonding, thus the activated complexes may be described as in Eqs. (5-100)*) and (5-101).

$$\mathrm{R{-}X + H{-}S \overset{S_N1}{\rightleftharpoons} [R^{\delta\oplus}\cdots X^{\delta\ominus}\cdots H{-}S]^{\ddagger} \longrightarrow R^{\oplus} + X^{\ominus}\cdots H{-}S}$$

$$\xrightarrow{+Y^{\ominus}} \mathrm{R{-}Y + X^{\ominus}\cdots H{-}S} \qquad (5\text{-}100)$$

$$\mathrm{Y^{\ominus} + R{-}X + H{-}S \overset{S_N2}{\rightleftharpoons} [Y^{\delta\ominus}\cdots R\cdots X^{\delta\ominus}\cdots H{-}S]^{\ddagger}}$$

$$\longrightarrow \mathrm{Y{-}R + X^{\delta\ominus}\cdots H{-}S} \qquad (5\text{-}101)$$

In these equations, the energy necessary for the ionization of the R–X bond will be lowered by that of the X···H interaction.

Therefore, protic solvents usually have an accelerating effect on S_N reactions. This is one of the main reasons why S_N reactions of haloalkanes and sulfonate esters are carried out in solvents which consist wholly or partially of water, alcohols, or carboxylic acids. The relative strengths of the hydrogen bonding in the initial and transition states often overwhelms the electrostatic effect of solvents on the Gibbs energy of activation. On the other hand, the attacking nucleophile $Y^{\ominus}$ in the S_N2 reaction (5-101) can also be specifically solvated by protic solvents, so that its reactivity, and therefore the rate of the S_N2 reaction, will be diminished. Examples of specific (electrophilic) solvation of anionic nucleophiles and leaving groups in S_N reactions can be found in references [264–269; 581–585]; see references [581, 582] for reviews.

Since the protic solvent is usually in large excess, its participation in the reaction cannot generally be established by means of kinetic measurements. However, if the reaction is carried out in a non-HBD solvent (*e.g.* C_6H_6, CCl_4), the effect of addition of small amounts of a protic solvent is easily observable. Thus, the S_N2 reaction between

* It should be pointed out that the formation of a carbenium ion in the course of an S_N1 reaction does not in fact occur as simply as described in Eq. (5-100), but takes place *via* intermediate contact- and solvent-separated ion pairs before the free solvated ions are formed; *cf.* Eqs. (2-19) and (2-20) in Section 2.6.

bromomethane and pyridine in benzene is accelerated by the addition of small amounts of alcohols or phenols. The more acidic the added solvent, the greater its ability to form hydrogen bonds, and the greater the rate acceleration [264]. The electron-withdrawing effect of a protic solvent on the departing group X can be so pronounced that reactions that are second-order in less strongly solvating media (*e.g.* ethanol), become first-order in acidic media (*e.g.* formic acid). The protic solvents exert an electrophilic pull on the departing anions in much the same way that heavy metal ions ($Ag^{\oplus}$, $Hg^{\oplus\oplus}$) catalyze nucleophilic substitution reactions of haloalkanes.

These results are supported by the observation that dipolar non-HBD solvents such as *N*,*N*-dimethylformamide or dimethyl sulfoxide, in spite of their high relative permittivities (36.7 and 46.5) and their high dipole moments ($12.3 \cdot 10^{-30}$ and $13.0 \cdot 10^{-30}$ Cm), favour neither the ionization of haloalkanes nor S_N1 reactions (*cf.* Section 2.6). Dipolar non-HBD solvents cannot act as hydrogen-bond donors and are therefore poor at solvating the departing anions. Thus, the anchimerically assisted ionization of 4-methoxyneophyl tosylate, shown in Eq. (5-102), is nine times faster in acetic acid than in dimethyl sulfoxide. This is in spite of the fact that the relative permittivity of acetic acid is eight times smaller than that of dimethyl sulfoxide; also the dipole moment of acetic acid is smaller than that of dimethyl sulfoxide [265].



(5-102)

Solvent	$(C_2H_5)_2O$	$(CH_2)_4O$	CH_3COCH_3	$HCON(CH_3)_2$	CH_3SOCH_3
k_1^{rel}	1	17	169	980	3600

Solvent	C_2H_5OH	CH_3OH	CH_3CO_2H	H_2O	HCO_2H
k_1^{rel}	$1.2 \cdot 10^4$	$3.2 \cdot 10^4$	$3.3 \cdot 10^4$	$1.3 \cdot 10^6$	$5.1 \cdot 10^6$

It is furthermore remarkable that an approximately linear relationship between $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ and lg k_1 values for reaction (5-102), measured in 19 solvents, is found only for non-HBD solvents (*cf.* Eq. (5-87) in Section 5.4.3), whereas protic solvents are much better ionizing media than their relative permittivity would suggest [265]. For example, acetic acid and tetrahydrofuran have very similar relative permittivities (6.2 and 7.6, respectively), and yet ionization in acetic acid exceeds that in tetrahydrofuran by a factor of $2 \cdot 10^3$! The reason for this extraordinary rate acceleration is again that the departing tosylate is better solvated due to hydrogen bonding in the protic solvent. The ability of the protic solvent to form hydrogen bonds is not reflected in its relative permittivity or in the dipole moment [265].

Another instructive example of electrophilic or H-bonding assistance of protic solvents (or co-solvents) in S_N1 reactions is the accelerated acetolysis rate of 2-bromo-2-methylpropane upon the addition of phenols to a tetrachloromethane/acetic acid solution of the reactant [582]; see reference [582] for further examples. The usefulness of phenol as a solvent for S_N1 solvolysis reactions, in particular phenolysis of 1-halo-1-phenylethanes, has been stressed by Okamoto [582]. In spite of its low relative permittivity ($\varepsilon_r = 9.78$ at 60 °C), its low dipolarity ($\mu = 4.8 \cdot 10^{-30}$ Cm = 1.45 D), and its low nucleophilicity, it represents a solvent of high ionizing power due to its electrophilic driving force.

A further instructive study on the influence of protic and dipolar aprotic solvents on the rate of S_N1 heterolysis of tertiary $R_3C–X$ (with X = Cl, Br, I, 2,4-dinitrophenolate) shows that the rate-accelerating anion solvation due to H-bonding by protic solvents decreases dramatically on increasing the radius of the halide ions. Therefore, the differential solvation transferred from anion-solvating methanol to cation-solvating dimethyl sulfoxide as solvent is reversed on going from the chloro- to the iodoalkane: at 60 °C $k_1(\text{DMSO})/k_1(\text{MeOH}) = 0.05$ (*t*-BuCl) < 0.57 (*t*-BuBr) < 6.9 (*t*-BuI); the 2,4-dinitrophenolate ion behaves like the iodide [830].

While increasing anion solvation by protic solvents has an accelerating effect on S_N1 reactions as described above, it is often a decelerating factor in S_N2 reactions. Thus, reaction (5-103) between (iodomethyl)benzene and radioactively labeled sodium iodide in acetone is clearly decelerated by the addition of protic solvents such as water, ethanol or phenol, as demonstrated in Fig. 5-16 [266].

$$\overset{*}{I}^{\ominus} + \underset{\displaystyle CH_2\text{-}I}{\overset{\displaystyle C_6H_5}{|}} \xrightarrow[0\,^\circ C]{k_2} \overset{*}{I}\text{-}\underset{\displaystyle CH_2}{\overset{\displaystyle C_6H_5}{|}} + I^{\ominus} \qquad (5\text{-}103)$$

The decrease in reaction rate is greatest for the addition of phenol, whereas the addition of the aprotic solvent CCl_4 leaves the reaction rate practically unchanged.



Fig. 5-16. The variation of k_2 for S_N2 reaction (5-103) with the ratio [solvent]/[acetone] [266].

The inhibitory effect of the protic solvents H–S (water < ethanol < phenol) correlates with their increasing tendency to form hydrogen bonds. This shows that the decrease in reaction rate is caused by specific solvation of the iodide ion, and depends on the position of the following equilibrium [266]:

$$(I^{\ominus})_{Acetone} + H\text{-}S \rightleftharpoons (I^{\ominus}\cdots H\text{-}S) + Acetone \qquad (5\text{-}104)$$

Similar conclusions can be drawn from specific solvent effects on the following S_N2 reactions: (a) substitution of 1-halobutanes by iodide ion in acetone in the presence of small amounts of water [267]; (b) substitution by $N_3^{\ominus}$ and $SCN^{\ominus}$ in a series of *n*-octyl derivatives (*n*-$C_8H_{17}X$ with X = Cl, Br, I, OTs, OMs as leaving groups) in different solvents (CH_3OH in contrast to CH_3SOCH_3) [583]; (c) Menschutkin reaction between 1,4-diazabicyclo [2.2.2]octane and (2-haloethyl)benzenes in binary solvent mixtures, *i.e.* a main non-HBD solvent with low concentrations of protic co-solvents [584]. (d) S_N2 reaction of *n*-octyl methanesulfonate with fluoride ions (associated with tetrahexylammonium ions) in chlorobenzene, the rate of which is dramatically reduced (by a factor of 822) on addition of small amounts of water through specific hydration of F^- to give $F^- \cdots (H_2O)_n$ (with $n = 0 \ldots 8.5$) [831].

The nucleofugacity scales observed for the S_N2 reaction (b) in nonpolar solvents (*e.g.* cyclohexane) are different from those obtained in protic solvents (*e.g.* methanol) and dipolar non-HBD solvents (*e.g.* dimethyl sulfoxide) [583]. For the Menschutkin reaction (c), specific rate inhibition has been observed on the addition of protic solvents which form hydrogen-bonds between the amine and the main solvent (*e.g.* C_6H_5OH, CH_3OH, $CHCl_3$). The retardation is proportional to the Brønsted acidity of the protic solvent added. Conversely, hydrogen-bond association between the halide nucleofuge and the protic solvent causes a specific catalytic effect, which increases as the charge density on the leaving group increases. These two effects, deactivation of the amine and activation of the halide leaving group, are always in competition and the outcome depends on the chemical nature of the reactants [584].

In this connection, it is noteworthy that the retarding effect of protic solvents on a given S_N2 reaction can also depend on the nature of the non-HBD solvent used [268, 269, 584]. Addition of a basic solvent may lead to a rate acceleration since the base can now compete with the nucleophile for hydrogen-bond formation with the protic solvent. Thus, the S_N2 reaction (5-105) is accelerated by addition of 1,4-dioxane to a protic solvent such as methanol as reaction medium [268].

$$CH_3O^{\ominus} + CH_3\text{-}I \xrightarrow[25\,^{\circ}C]{k_2} CH_3O\text{-}CH_3 + I^{\ominus} \qquad (5\text{-}105)$$

Solvent (1,4-Dioxane in cL/L)	Methanol/1,4-Dioxane			
	0	20	40	60
k_2^{rel}	1	1.9	3.4	5.0

Most probably this rate increase is due to the fact that the specific solvation of the methoxide ion is decreased because of an association between 1,4-dioxane and methanol:



(5-106)

(5-107)

In other words, by the addition of 1,4-dioxane, the reaction medium will become richer in free, not specifically solvated methoxide ions. Accordingly, the rate of the S_N2 reaction (5-105) increases [10, 268]. This interpretation is supported by the fact that reaction (5-105) is accelerated further by addition of the more highly basic solvent dimethyl sulfoxide, although admittedly this addition also leads to an increase in the relative permittivity of the reaction medium [269].

Similar results are found for the above-mentioned Menschutkin reaction (c). The added protic solvent can also combine with the main solvent. Such solvent/solvent association leads to a diminution of the specific inhibitory and catalytic effect of protic solvents on the S_N2 reaction (c) [584]. The basicity of the main solvent determines the extent of deactivation of the protic solvent through H-bond association; this is analogous to Eq. (5-107).

There is also direct IR spectroscopic evidence that addition of a basic co-solvent to aqueous solutions reduces the percentage of non-associated "free" OH groups which are necessary for hydrogen-bonding to S_N reactants [585]. When basic co-solvents are added, they scavenge the free OH groups in water, thus lessening the specific protic solvent effect [585].

Nucleophiles with a heteroatom adjacent to the reaction center (*e.g.* $HOO^{\ominus}$, $ClO^{\ominus}$, $R_2C{=}N{-}O^{\ominus}$, $HONH_2$, NH_2NH_2) are found to be more reactive toward carbon electrophiles in S_N2 reactions in solution than would be expected from their basicities. This corresponds to a positive deviation from a Brønsted-type plot (*i.e.* $\lg k_2$ against pK_B). For instance, the hydroperoxide ion ($HOO^{\ominus}$) is 10^4 times less basic than the hydroxide ion ($HO^{\ominus}$), but it reacts *ca.* 50 times faster with (bromomethyl)benzene [459]. This behaviour is known as the *α-effect* [586] and various explanations for it have been given; *cf.* [587, 588] and references therein. If the α-effect is an intrinsic property of the anion, then it should manifest itself in the gas phase. However, it has been reported that in the gas phase $HOO^{\ominus}$ shows no evidence of enhanced nucleophilicity, compared to $HO^{\ominus}$, in its behaviour toward saturated (fluoromethane) and unsaturated carbon (methyl formate) [587]. If this is true, then the α-effect should arise because of a specific differential solvation of the α- and normal nucleophiles ($HOO^{\ominus}/HO^{\ominus}$, NH_2NH_2/R-NH_2, *etc.*), which would lead to reduced solvation of the α-nucleophile. It is, however, not easy to understand why α-nucleophiles should be less solvated than normal nucleophiles; see the comments given in references [588]. A thorough study of the rate of reaction of an α-nucleophile (butane-2,3-dione monooximate, Ox^-) and a normal nucleophile (4-chlorophenolate, 4-$ClPhO^-$) with a phosphorus ester (4-nitrophenyl diphenylphosphinate) in binary water/DMSO mixtures has shown that the magnitude of the α-effect [$=k(Ox^-)/k(4\text{-}ClPhO^-)$] may be due to advanced desolvation of the oximate nucleophile at the transition state relative to the O–P bond formation, corresponding to an imbalanced or asynchronous effect [832].

Our final reaction is an instructive example of specific anion solvation in a non-S_N reaction. The decarboxylation rate of 6-nitrobenzisoxazole-3-carboxylate as the tetra-

methylguanidinium salt exhibits a dramatic rate increase on going from protic to non-HBD solvents, as shown in Eq. (5-108) [589].



(5-108)

Solvent	H_2O	CH_3OH	C_2H_5OH	CH_2Cl_2	Et_2O	CH_3CN	CH_3SOCH_3	HMPT
k_1^{rel}	1	34	135	$6.4 \cdot 10^3$	$1.2 \cdot 10^4$	$3.9 \cdot 10^5$	$1.4 \cdot 10^6$	$9.5 \cdot 10^7$

The rate is slowest in aqueous solution and is enhanced in non-HBD solvents by a factor of up to 10^7! Obviously, specific hydrogen-bonding between protic solvent molecules and the carboxylate group suppresses the decarboxylation. This is consistent with the fact that the decarboxylation rate of *4-hydroxy*benzisoxazole-3-carboxylate (R = OH), a compound with its own intramolecular protic environment, is very slow and does not show a substantial solvent effect. Furthermore, not only hydrogen-bonding but also ion-pairing seems to stabilize the carboxylate ion, at least in less polar solvents such as C_6H_6, CH_2Cl_2, and Et_2O [590]. Thus, the most reactive compound is the free carboxylate ion, which lacks any specific stabilization.

Quantum-chemical calculations on the thermolysis of unsubstituted benzisoxazole-3-carboxylate, a reaction showing "an awesome solvent effect" [833], are in agreement with a concerted decarboxylation and ring-opening step. Hydration of the carboxylate increases the activation barrier (E_A) from 57 kJ/mol in the uncomplexed system to 130 kJ/mol for a model carboxylate with two bifurcated water molecules (including four hydrogen bonds to the carboxylate group), in good agreement with the experimental value of 134 kJ/mol found in water [589, 833]. The extreme solvent-dependence of this decarboxylation reaction suggests its use as a probe for many kinds of biochemical environments (*e.g.* micelles, bilayers, binding pockets of catalytic antibodies) [834–836].

In conclusion, it can be stated, according to a general rule proposed by Palit in 1947 [270], that solvents which impede the active centre of a reactant through hydrogen bonding or by some other means, will suppress the reactivity of that reactant. Conversely, solvents capable of promoting a favourable electron shift, necessary to the reaction, by specific solute/solvent interactions, will enhance the reaction rate.

It should be mentioned that specific anion solvation is not only possible through hydrogen-bonding with protic solvents and ion-pairing, it is also possible through coordination with macrocyclic organic ligands, particularly with protonated cryptands [591, 592]. For example, well-defined anion complexes, chloride ion cryptates, have been identified in aqueous solution by means of ^{35}Cl NMR spectroscopy [591]. The anion is located inside the intramolecular cryptand cavity and is held by an array of hydrogen bonds. A review on the supramolecular chemistry of anions can be found in reference [837].

5.5.2 Protic and Dipolar Aprotic Solvent Effects on the Rates of S_N Reactions

In the course of a nucleophilic substitution reaction, a new bond is formed between the attacking nucleophile and the substrate. Since the nucleophile "attacks" with a free

electron pair, a reagent should be all the more nucleophilic, the more readily the electron pair can be engaged in chemical bonding. Thus, the nucleophilicity of an anion is determined – among other things – by its basicity*) and its polarizability. One or the other property may dominate. While polarizability is hardly influenced by solvent, the basicity of an anion is strongly solvent dependent (*cf.* Section 4.2).

In the course of the activation process, the solvent shell of the nucleophile must be removed at the place of attack, while a new solvent shell around the activated complex is simultaneously forming. Thus, the Gibbs energy of activation will be higher, and the rate lower, the more strongly the molecules of the solvent shell are bound to the nucleophile. The less solvated a nucleophile, the more reactive it will be. Conversely, a nucleophile is stabilized by increasing solvation, its chemical potential falls, and its reactivity decreases. Since, according to Bunnett [271], nucleophilic reactivity depends on no less than seventeen factors (solvation effects included), it is very difficult to establish a general nucleophilicity order for all common nucleophiles in different solvents. Numerous attempts to establish such orders of nucleophilicities, which should be independent of any particular reaction electrophile, can be found in the literature [272–275].

When considering the solvation of anions, according to Parker [6] it is reasonable to distinguish between two classes of solvents: protic and dipolar aprotic solvents**) (*cf.* Section 3.4 and Fig. 3-4). The main difference between these two classes of solvents lies in their ability to solvate anions. Small anions with a high charge density (*i.e.* the ratio of charge to volume), which are strong hydrogen-bond acceptors, are more strongly solvated in protic solvents. The hydrogen-bonding interaction is greatest for small anions (*e.g.* $F^{\ominus}$, $Cl^{\ominus}$, $HO^{\ominus}$), and least for large anions where the charge is dispersed (*e.g.* $SCN^{\ominus}$, $I^{\ominus}$, picrate). Dipolar non-HBD solvents, which are polarizable, have mutual polarizability interactions with polarizable anions. This mutual polarizability interaction should be greatest for large, polarizable anions, and least for small weakly polarizable anions. Of course, mutual polarizability is not an exclusive prerogative of dipolar non-HBD solvents; some protic solvents may also be very polarizable. From the point of view of the "principle of hard and soft acids and bases" [275], hard anions are better solvated by hard (hydrogen-bonding) solvents, and soft anions have strong interactions with soft (dipolar non-HBD) solvents (*cf.* also Section 3.3.2). In other words, small ions lose in solvation and large, polarizable anions gain in solvation on transfer from protic to dipolar non-HBD solvents [6]. These considerations are essentially confirmed by the nucleophilic reactivities observed for some S_N reactions in different media, as presented in Table 5-15 [276–287].

---

* Basicity is measured in terms of a thermodynamic equilibrium involving coordination with H^+. Nucleophilicity is measured in terms of the rates of reaction with the most varied electrophiles. Hence, although a correlation between basicity and nucleophilicity is often found (generally S_N reactions are faster with the stronger bases), it is by no means *a priori* necessary. An example of a direct simple relationship between nucleophilicity and basicity in S_N2 reactions is the reaction of 9-substituted fluorenide ions with (chloromethyl)benzene in dimethyl sulfoxide solution [595].

** Solvents referred to as *dipolar aprotic* are not in fact aprotic. In reactions where strong bases are employed their protic character can be easily recognized (*e.g.* $H_3C{-}SO{-}CH_3 + NH_2^{\ominus} \rightleftharpoons H_3C{-}SO{-}CH_2^{\ominus} + NH_3$). The term dipolar *aprotic* solvent is in fact rather misleading. Therefore, it has been recommended by Bordwell *et al.* [594] that the designation *dipolar aprotic* should be replaced by *dipolar nonhydroxylic* or better still by *dipolar non-HBD* solvents. The abbreviation *HBD* stands for *hy*drogen-*b*ond *d*onor. See also Chapter 3.4 and the footnote on page 82.

Table 5-15. Relative nucleophilic reactivities of free[a] anions for S_N2 reactions in various protic and dipolar non-HBD solvents (nos. 1 . . . 5) as well as in molten salts (no. 6 and no. 7) and in the gas phase (no. 8).

No.	Reaction	Solvent (temperature)	Sequence of decreasing reactivity (relative reaction rates)	References
(1)	$CH_3–I + Me^{\oplus}X^{\ominus}$	CH_3OH (25 °C)	$C_6H_5S^{\ominus} > S_2O_3^{2\ominus} \gg I^{\ominus} > SCN^{\ominus} \approx CN^{\ominus} > CH_3O^{\ominus} > Br^{\ominus} > N_3^{\ominus} > C_6H_5O^{\ominus} > Cl^{\ominus} > CH_3CO_2^{\ominus} > F^{\ominus}$	[276]
(2)	$CH_3–OTos + Me^{\oplus}X^{\ominus}$	CH_3OH (25 °C)	$C_6H_5S^{\ominus} > CH_3O^{\ominus} > I^{\ominus} > SCN^{\ominus} > Br^{\ominus} > Cl^{\ominus}$	[277]
		$HCON(CH_3)_2/H_2O$[b]	$I^{\ominus} > Br^{\ominus} > Cl^{\ominus}$	[278]
		$HCON(CH_3)_2$ (0 °C)	$Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ (9:3:1)	[278]
(3)	$n\text{-}C_4H_9–OBros + (n\text{-}C_4H_9)_4N^{\oplus}X^{\ominus}$	CH_3COCH_3 (25 °C)	$Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ (18:5:1)	[279]
(4)	$n\text{-}C_3H_7–OTos + (n\text{-}C_4H_9)_4N^{\oplus}X^{\ominus}$	CH_3SOCH_3 (25 °C)	$S_2O_3^{2\ominus} > HO^{\ominus} \approx CH_3O^{\ominus} > F^{\ominus} > C_6H_5O^{\ominus} > N_3^{\ominus} > Cl^{\ominus} > Br^{\ominus} > I^{\ominus} > SCN^{\ominus}$	[280]
		CH_3SOCH_3 (50 °C)	$Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ (5.2:3.2:1)	[281]
		CH_3SOCH_3/H_2O[c]	$I^{\ominus} > Br^{\ominus} > Cl^{\ominus}$ (6.7:1.3:1)	[281]
(5)	$C_6H_5CH_2–OTos + K^{\oplus}X^{\ominus}$[d]	CH_3CN (30 °C)	$N_3^{\ominus} > CH_3CO_2^{\ominus} > CN^{\ominus} > F^{\ominus} > Cl^{\ominus} \approx Br^{\ominus} > I^{\ominus} > SCN^{\ominus}$ (10:9.6:2.4:1.4:1.3:1.3:1:0.3)	[282]
(6)	$CH_3–OTos + (C_2H_5)_4N^{\oplus} + X^{\ominus}$	$R_4N^{\oplus}R_4B^{\ominus}$[e] (35 °C)	$Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ (2.1:1.2:1)	[283]
(7)	$(n\text{-}C_5H_{11})_4N^{\oplus} + X^{\ominus}$	$(n\text{-}C_5H_{11})_4N^{\oplus}X^{\ominus}$[f] (180 °C)	$Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ (620:7.7:1)	[284]
(8)	$CH_3–Br + X^{\ominus}$	Gas phase	$HO^{\ominus} > CH_3O^{\ominus} \approx F^{\ominus} > CH_3S^{\ominus} \gg CN^{\ominus} > Cl^{\ominus} > Br^{\ominus}$	[285, 290]

[a] "Free" means without association with the corresponding cation, no ion pairing.
[b] 91:9 cL/L (0 °C).
[c] 70:30 cL/L (50 °C).
[d] The potassium salts are solubilized in acetonitrile with [18]crown-6, producing extremely reactive "naked" anions because of the weak anion solvation forces in acetonitrile solutions and the complete dissociation of the potassium salts [282].
[e] Molten triethyl-*n*-hexylammonium triethyl-*n*-hexyl-boride as solvent [283].
[f] Molten tetra-*n*-pentylammonium halides as solvent [284].

It is apparent that the order of anion nucleophilicity is almost completely reversed on transfer from protic to dipolar non-HBD solvents. Especially for halide ions, the relative reactivity is completely reversed in the two classes of solvents: whereas the order of reactivity is $I^{\ominus} > Br^{\ominus} > Cl^{\ominus} > F^{\ominus}$ in the protic solvent methanol (reactions no. 1 and no. 2 in Table 5-15), in dipolar non-HBD solvents such as *N*,*N*-dimethylformamide (no. 2), acetone (no. 3), dimethyl sulfoxide (no. 4), and acetonitrile (no. 5) the sequence of nucleophilicity is reversed. The traditional order of halide nucleophilicities, $I^{\ominus} > Br^{\ominus} > Cl^{\ominus}$ [261], applies only when the nucleophile is deactivated through solvation by

protic solvents ($X^{\ominus} \cdots H\text{–}S$), whereas the natural order, $Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$, is observed in dipolar non-HBD solvents. The sequences observed in dipolar non-HBD solvents give a much better picture of the true nucleophilicity of the weakly solvated anions, which also correlates better with their basicity.

With acetonitrile as solvent, there appears to be a general levelling of the nucleophilicities of anions, as indicated by the relatively small variation of rate constants obtained for reaction no. 5 in Table 5-15 [282]. Quite surprisingly, one of the best nucleophiles in acetonitrile is $CH_3CO_2^{\ominus}$, which is normally considered a very poor nucleophile in protic solvents, whereas $SCN^{\ominus}$, one of the more potent nucleophiles in aqueous solution, is approximately 30 times slower than $CH_3CO_2^{\ominus}$. Obviously, the so-called "naked", and therefore very reactive, anions in acetonitrile are solvated by much weaker forces than in protic solvents, and the variations in anion solvation in acetonitrile do not appreciably effect the relative nucleophilic reactivities of the anions [282].

Reactions no. 6 and no. 7 in Table 5-15 demonstrate that with molten quaternary ammonium salts as solvents, where deactivation by anion solvation is absent, the halide ions show the same nucleophilic order as in dipolar non-HBD solvents [283, 284]. This is in accordance with the theory of protic/dipolar non-HBD medium effects on $X^{\ominus}$ nucleophilicity [6]. It has been suggested that fused-salt experiments should provide a good model for the determination of intrinsic relative nucleophilicities of anions towards saturated carbon atoms [284].

The results for reaction no. 8 in Table 5-15 indicate that nucleophilic reactivities of anions obtained in the gas phase are essentially in the same order as in molten salts and in dipolar non-HBD solvents [285, 290]. This again suggests that specific solvation of the anions is responsible for the reversed order obtained in protic solvents relative to dipolar non-HBD solvents. Whereas the relative nucleophilicities in acetonitrile are similar to those found in the gas phase [282, 285, 290], the absolute gas-phase rates are some orders of magnitude greater than those in acetonitrile. The specific rates of displacement reactions of anions with halomethanes exceed those in solution by factors of up to $\geq 10^{10}$ [285, 290]. These large differences in absolute rates demonstrate the moderating influence of the solvent on all the reactivities [282]. See also Chapter 5.2.

In spite of evidence for increased stabilization of increasingly large anions in increasingly polarizable solvents due to dispersion interactions, there is some question as to whether the decreased reactivity of halide ions in dimethyl sulfoxide ($Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$; *cf.* reaction no. 4 in Table 5-15) is due to increased solvation in the series $Cl^{\ominus} \rightarrow Br^{\ominus} \rightarrow I^{\ominus}$. Because the solvation order of these halide ions in dimethyl sulfoxide is $I^{\ominus} < Br^{\ominus} < Cl^{\ominus}$ (the same as in water!) as shown by solvation enthalpies, ΔH_{solv}, of the corresponding alkali halides in dimethyl sulfoxide and in water [281], Parker [6] suggested that the observed reactivity order $Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$ in dipolar non-HBD solvents might be the order of increasing solvation in these solvents. Alternatively, this could also be the intrinsic order of nucleophilicity, which is only partially offset by decreasing solvation in the same order [281]. If the unsolvated $Cl^{\ominus}$ is a sufficiently stronger nucleophile than unsolvated $I^{\ominus}$, the more strongly solvated $Cl^{\ominus}$ in dimethyl sulfoxide may remain more reactive than the comparatively less well solvated $I^{\ominus}$. In protic solvents, however, the difference in solvation is much larger. Now the extremely strong solvation of $Cl^{\ominus}$ in protic solvents diminishes the reactivity of this anion to less

than that of the more weakly solvated $I^{\ominus}$. Therefore, the reversed nucleophilic order of the halide ions in dimethyl sulfoxide relative to water must be attributed to the smaller differences in halide solvation in dimethyl sulfoxide than in water [281]. This [281] and other estimates [288, 289] are all in agreement with the observation that solvent stabilization of halide ions, although small compared to water, goes in the same direction, $Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$, in dipolar non-HBD solvents.

Accordingly, gas-phase results for the clustering of $Cl^{\ominus}$, $Br^{\ominus}$, and $I^{\ominus}$ with Me_2SO and H_2O show that the bonding to dimethyl sulfoxide also decreases in the order $Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$, *i.e.* with increasing ion radius [596]. Thus, anion solvation decreases with an increase in the ion radius for both protic and non-HBD solvents; however, the decrease is appreciably less in dipolar non-HBD solvents. This is the principle reason for the much higher rates of anion-molecule S_N2 reactions in dipolar non-HBD solvents [596].

An important contribution to a quantitative estimation of nucleophilic reactivity towards cations in different solvents was made by Ritchie *et al.* [274, 597]. According to Eq. (5-109),

$$\lg k = \lg k_0 + N_+ \tag{5-109}$$

where k is the rate constant for the reaction of a cation (particularly triarylmethyl cations, aryldiazonium ions, and aryltropylium cations) with a given nucleophile/solvent system, and k_0 is the rate constant for reaction of the same cation with a standard system (*e.g.* $HO^{\ominus}$ in H_2O), a parameter N_+ was defined, which represents the difference in nucleophilicity of the two systems. The N_+ values are measures of an inherent property of the (nucleophile *and* solvent) system, which is associated with the differential solvation of the nucleophile in different solvents. A selection of N_+ values is listed in Table 5-16 [597]. The higher the N_+ value, the less strongly solvated and the more reactive is the nucleophile in a given solvent. The large differences in N_+ values for the same nucleophile in protic and non-HBD solvents deserve special mention in this context; *cf.* for example entries (3), (13), and (17) for the cyanide ion, entries (15) and (19) for the monothioglycolate ion, and entries (6), (7), and (11) for hydrazine in Table 5-16.

The N_+ scale, which is derived from nucleophile/electrophile combination reactions, differs from the nucleophilicity scales generated from rates of S_N2 reactions (*cf.* Table 5-15). The N_+ equation, an example of a so-called "constant selectivity relationship", has been applied to many other electrophile/nucleophile reactions in solution, although often with only limited success.

A considerable improvement in the construction of reference scales for the quantitative characterization of nucleophiles and electrophiles in solution has been made by Mayr *et al.* [598] by means of Eq. (5-109a),

$$\lg k(20\ ^{\circ}C) = s(N + E) \tag{5-109a}$$

where N and E are substrate nucleophilicity and electrophilicity parameters, respectively, and s is the nucleophile-dependent slope, measuring the susceptibility of $\lg k$ to changes in nucleophilicity of π-, n-, and σ-nucleophiles in reactions with carefully selected reference electrophiles. For example, 23 differently substituted electrophilic diarylcarbenium (benzhydryl) ions and 38 neutral nucleophilic π-systems (*e.g.* arenes, alkenes, allyl silanes, silyl enol ethers, enamines) have been used as basis sets for estab-

Table 5-16. Nucleophilic parameters N_+ for various (nucleophile + solvent) systems, based on reactions of malachite green or the tris-4-anisylmethyl cation [597].

No.	Nucleophile (solvent)	N_+
(1)	$H_2O(H_2O)$	0.73
(2)	$NH_3(H_2O)$	3.89
(3)	$CN^{\ominus}(H_2O)$	4.12
(4)	$HO^{\ominus}(H_2O)$	*4.75*[a)]
(5)	$HO–NH_2(H_2O)$	5.05
(6)	$H_2N–NH_2(H_2O)$	6.01
(7)	$H_2N–NH_2(CH_3OH)$	6.89
(8)	$BH_4^{\ominus}(H_2O)$	6.95
(9)	$CH_3O^{\ominus}(CH_3OH)$	7.51
(10)	$N_3^{\ominus}(H_2O)$	7.54
(11)	$H_2N–NH_2(CH_3SOCH_3)$	8.17
(12)	$HOO^{\ominus}(H_2O)$	8.52
(13)	$CN^{\ominus}(CH_3SOCH_3)$	8.64
(14)	$N_3^{\ominus}(CH_3OH)$	8.78
(15)	$HOCH_2–CH_2S^{\ominus}(H_2O)$	8.87
(16)	$C_6H_5S^{\ominus}(H_2O)$	9.10
(17)	$CN^{\ominus}[HCON(CH_3)_2]$	9.44
(18)	$C_6H_5S^{\ominus}(CH_3OH)$	10.41
(19)	$HOCH_2–CH_2S^{\ominus}(CH_3SOCH_3)$	12.71

[a)] All N_+ values are relative to $N_+ = 4.75$ for hydroxide ion in water.

lishing general reactivity scales for electrophiles and nucleophiles, which cover more than sixteen orders of magnitude [598b]. It has been shown that the reactivity parameters derived in this way from the reaction of diarylcarbenium ions with π-nucleophiles are also applicable for characterizing the reactivities of other nucleophiles such as alkynes, transition metal π-complexes, *n*-nucleophiles (*e.g.* amines, anions), and σ-nucleophiles (*e.g.* hydride donors such as Bu_3SnH), not belonging to the basis set [598b,c]. Most of the electrophile/nucleophile reactions were carried out in dichloromethane as reference solvent. Using the rate constants of a total of 209 combinations of diarylcarbenium ions and π-nucleophiles from the basis set, Eq. (5-109a) represents a well-founded new and helpful linear Gibbs energy relationship; for other LGE relationships, see Section 7.1.

Since anions are solvated to a much lesser extent in dipolar non-HBD solvents than in protic solvents, they exist in these solvents as more or less "naked" and therefore extremely reactive ions. For S_N2 and S_NAr reactions involving anionic nucleophiles, a change from protic to dipolar non-HBD solvent often causes a very dramatic acceleration of these reactions [6]*). Some typical examples are reactions (5-110) to (5-113), for

* To speak of dipolar non-HBD solvents having an accelerating effect on the rates of bimolecular substitution reactions involving anionic nucleophiles seems to be looking at things in reverse order. The view that protic solvents have a retarding effect on such reactions seems to be much more consistent with the experimental data. However, the above mentioned description of the protic/dipolar non-HBD solvent effect on reaction rates has been widely used in the literature.

Table 5-17. Relative rates of the S_N2 anion-molecule reactions (5-110) [291] and (5-111) [67], and of the S_NAr reactions (5-112) [291, 292] and (5-113) [293] in protic and dipolar non-HBD solvents at 25 °C.

Solvents	$\lg(k_2^{Solvent}/k_2^{MeOH})$ [a] for reaction			
	(5-110) [b]	(5-111) [c]	(5-112) [d]	(5-113) [e]
Protic Solvents:				
CH_3OH	0	0	0	0
H_2O	0.05	0.8	–	–
$CH_3CONHCH_3$	–	0.9	–	–
$HCONH_2$	1.2	1.1	0.8	–
$HCONHCH_3$	1.7	–	1.1	–
Dipolar non-HBD Solvents:				
$(CH_2)_4SO_2$ (Sulfolane)	–	2.6	4.5	–
CH_3NO_2	4.2	–	3.5	0.8
CH_3CN	4.6	3.7	3.9	0.9
CH_3SOCH_3	–	3.1	3.9	2.3
$HCON(CH_3)_2$	5.9	3.4	4.5	1.8
CH_3COCH_3	6.2	3.6	4.9	0.4
$CH_3CON(CH_3)_2$	6.4	3.9	5.0	1.7
	6.9	–	5.3	–
$[(CH_3)_2N]_3PO$	–	5.3	7.3	–

[a] $k_2^{MeOH}/(L \cdot mol^{-1} \cdot s^{-1})$ is the second-order rate constant for the reaction in methanol; $k_2^{Solvent}$ is for reaction in the given solvent. $\lg k_2^{Solvent} = \lg(k_2^{Solvent}/k_2^{MeOH}) + \lg k_2^{MeOH}$.
[b] $\lg k_2^{MeOH} = -5.5$.
[c] $\lg k_2^{MeOH} = -5.1$.
[d] $\lg k_2^{MeOH} = -7.2$.
[e] $\lg k_2^{MeOH} = -3.85$ (50 °C).

which rate data in several solvents relative to solvent methanol are reported in Table 5-17 as the logarithms of rate ratios.

(5-110)

(5-111)


(5-112)


(5-113)

The rate increases (relative to methanol) involve factors in the range of 10^2 to 10^7 going from a protic to a dipolar non-HBD solvent. Even larger factors, as in the reaction 2,4-$(NO_2)_2C_6H_3I$ with $Cl^{\ominus}$ in hexamethylphosphoric triamide (ca. 10^9!) are not uncommon [291]. For reaction (5-112), the solvent change from methanol to hexame-

thylphosphoric triamide corresponds to a decrease in the Gibbs energy of activation, $\Delta\Delta G^{\neq}$, of 36.4 kJ/mol (8.7 kcal/mol) [292]. Hexamethylphosphoric triamide is usually the "fastest" of dipolar non-HBD solvents [291]. That these protic/dipolar non-HBD solvent effects are mainly anion-solvation phenomena is confirmed by the observation that similarly large rate ratios are not observed for reaction (5-113), which involves a neutral nucleophile (piperidine). Menschutkin reactions between tertiary amines – neutral nucleophiles – and haloalkanes are also rather insensitive to protic/dipolar non-HBD solvent transfer [6, 23]. As expected for such an effect, attributed mainly to hydrogen-bonding interactions, protic/dipolar non-HBD solvent effects on the chemistry of anions are strongly temperature dependent; differences between protic and dipolar non-HBD solvents are greater at low temperatures than at high temperatures [291].

In addition, it should be mentioned that for protic/dipolar non-HBD solvent effects on S_N2 reactions $Y^{\ominus} + RX \rightleftharpoons [YRX^{\ominus}]^{\neq} \rightarrow YR + X^{\ominus}$ not only the solvation of the anion must be taken into account. An important factor in determining the rate of this reaction in different solvents may be the solvation of the reactant molecule RX and the activated complex $[YRX^{\ominus}]^{\neq}$. Although ions have much greater solvation enthalpies than nonelectrolytes, the solvation of nonelectrolytes should not be neglected, because important criteria are the differences in solvation. Many anionic activated complexes, especially those of S_NAr reactions, are expected to be highly polarizable and will have strong dispersion interactions in highly polarizable solvents such as dimethyl sulfoxide or *N*,*N*-dimethylformamide. On the other hand, due to the delocalization of the negative charge, they are weak hydrogen-bond acceptors and are not particularly well solvated by protic solvents. For example, solvation of large polarizable S_NAr transition state anions decreases in the solvent order $[(CH_3)_2N]_3PO$, $CH_3SOCH_3 > HCON(CH_3)_2 > CH_3OH > CH_3NO_2$, CH_3CN [291]. These differences in solvation of activated complexes may also contribute to the large rate accelerations obtained on protic/dipolar non-HBD solvent transfer for bimolecular substitution reactions*). However, in most cases S_N reactions are faster in dipolar non-HBD solvents because the reactant anion $Y^{\ominus}$ is much more solvated by protic than by dipolar non-HBD solvents and this outweighs any effects due to solvation of the transition state anion or reactant molecule [6].

Nucleophilic reactions with anions for which hydrogen bonding is less important are not so greatly influenced by changes from protic solvents to dipolar aprotic ones. For example, the rate constant for the degenerate thiolate/disulfane interchange reaction shown in Eq. (5-113a) is larger in dimethyl sulfoxide than in water

$$R^1\text{–}S^- + \overset{R}{\underset{|}{S}}\text{–}S\text{–}R^2 \underset{24\ ^\circ C}{\overset{k_2}{\rightleftharpoons}} [R^1\text{–}S^{\delta-}\cdots\overset{R}{\underset{|}{S}}\cdots{}^{\delta-}S\text{–}R^2]^{\neq} \rightarrow R^1\text{–}S\text{–}\overset{R}{\underset{|}{S}} + {}^-S\text{–}R^2 \qquad (5\text{-}113a)$$

as expected, but only by a factor of ca. 2700 (with $R^1 = R = R^2 = -CH_2CH_2OH$) [838]. In D_2O/DMSO mixtures, $\lg k_2$ is directly proportional to the mole fraction of D_2O, even at small fractions of D_2O. Evidently, thiolate ions are not specifically solvated by water, in contrast to alcoholate ions. The rate acceleration in DMSO seems to

* For an extensive discussion concerning the solvation of activated complexes of bimolecular substitution reactions for numerous variations of RX and $Y^{\ominus}$, as well as different mechanisms, see references [6, 23, 291].

be due to better solvation of the charge-delocalized, polarizable activated complex of this exchange reaction by DMSO, which reduces the activation barrier.

Since small anions such as $F^{\ominus}$, $HO^{\ominus}$, and $CN^{\ominus}$ are often so poorly solvated by pure dipolar non-HBD solvents, it is difficult to find a soluble salt to act as the anion source. For example, electrolytes such as KF, KOH, and KCN are only slightly soluble in pure dimethyl sulfoxide. There are two possibilities for overcoming this experimental difficulty. First, the solubility of electrolytes in dipolar non-HBD solvents can be increased by using the corresponding tetraalkylammonium salts. Among these, tetraalkylammonium fluorides are the most prominent. In solutions of tetraalkylammonium fluorides in dipolar non-HBD solvents, the fluoride ion is virtually unsolvated. These "naked" fluoride ions are both strong nucleophiles [599] and very effective bases [600]. They are more nucleophilic and basic than the fluoride ions of lower alkali metal fluorides (LiF, NaF, KF) in protic solvents; see reference [839] for reviews on the solvation of fluoride ions in nonaqueous solvents. Secondly, protic/dipolar non-HBD solvent mixtures can be used as media for substitution reactions. Mixtures of dimethyl sulfoxide with water or alcohols have proven especially useful. This is because salts such as KOH, KF, KCN, CH_3ONa, and CH_3CO_2Na are reasonably soluble in solvent mixtures such as CH_3SOCH_3/H_2O (90:10) or CH_3SOCH_3/CH_3OH (90:10). Water or alcohols in combination with pure dipolar non-HBD solvents (*e.g.* 15 cL/L water in HMPT) provide a source of nucleophilic oxygen for the conversion of haloalkanes or sulfonate esters into the corresponding alcohols or ethers [601].

All the features discussed for protic/dipolar non-HBD solvent transfer are also observed for transfer from protic to dipolar non-HBD/protic mixtures, but to a slightly lesser extent. In general, all S_N2 anion-molecule reactions are faster in mixtures than in pure protic solvents. They show a continuous, but not necessarily linear, rate increase with an increase in the dipolar non-HBD component of the protic/dipolar non-HBD mixture [6]. Even small amounts of dipolar non-HBD component may cause a considerable acceleration in reaction rate.

The use of dipolar non-HBD instead of protic solvents as reaction media often has considerable practical synthetic advantages, which have been summarized by Parker [6], Madaule-Aubry [294], Liebig [295], and Schmid [26]. A selection of common and less common dipolar non-HBD solvents is given in Table 5-18, together with some physical constants useful for their practical application. Reviews on particular dipolar non-HBD solvents have appeared; these are included in Table 5-18 (*cf.* also references [75–91] in Chapter 3).

Finally, some important examples emphasizing the versatility and synthetic utility of dipolar non-HBD solvents as reaction media will be given.

$RCO_2^{\ominus}$, an indifferent nucleophile in protic solvents, enjoys a large rate enhancement, permitting rapid alkylation with haloalkanes in hexamethylphosphoric triamide [301, 302]. When the Williamson ether synthesis is carried out in dimethyl sulfoxide [303], the yields are raised and the reaction time shortened. Displacements on unreactive haloarenes become possible [304] (conversion of bromobenzene to *tert*-butoxybenzene with *tert*-$C_4H_9O^{\ominus}$ in dimethyl sulfoxide in 86% yield at room temperature). The fluoride ion, a notoriously poor nucleophile or base in protic solvents, reveals its hidden capabilities in dipolar non-HBD solvents and is a powerful nucleophile in substitution reactions on carbon [305].

Table 5-18. A selection of twenty-one organic dipolar non-HBD solvents in order of increasing dipole moment (*cf.* also Appendix, Table A-1).

Structure (↔)[a)]	Name	$\mu/(10^{-30}$ Cm)	t_{mp}/°C	t_{bp}/°C	References
	Acetone	9.0	−94.7	56.1	
	Tetramethylurea (Temur)	11.6	−1.2	175.3	[602, 603]
	Nitromethane	12.0	−28.6	101.2	[604]
	1,3,2-Dioxathiolane-2-oxide (Ethylene sulfite)	12.3		170–171 (dec.)	[605]
	*N*,*N*-Dimethylacetamide (DMAC)	12.4	−20.1	166.1	[606]; *N*,*N*-Diethylacetamide: [633]
	*N*,*N*-Dimethylformamide (DMF)	12.7	−60.4	153.1	[606–608]
	1-Methylpyrrolidin-2-one	13.6	−24.4	202	[609]; 1-Ethyl-pyrrolidin-2-one: [633]
	1,3-Dimethylimidazolidin-2-one (*N*,*N*′-Dimethyl-ethylene urea, DMEU)	13.6	8.2	225.5	[603]
	3-Methyloxazolidin-2-one	13.7	15.9	87–90 (1.3 mbar)	[610, 611]
	Acetonitrile	13.0	−43.8	81.6	[296, 608, 612]
	Dimethyl sulfoxide	13.5	18.5	189.0	[608, 613–616]
	3,4,5,6-Tetrahydro-1,3-dimethyl-2(1H)-pyrimidinone (*N*,*N*′-Dimethylpropylene urea, DMPU)	14.1	−23.8	246.5	[603, 617–618]
	1-Methyl-hexahydro-azepin-2-one, N-Methyl-ε-capro-lactam	14.1		12 (25 mbar)	
	Dimethylsulfone	14.2	110	238	[619]

Table 5-18. (Continued)

Structure (↔)[a]	Name	$\mu/(10^{-30}$ Cm)	t_{mp}/°C	t_{bp}/°C	References
	*S*,*S*-Dimethylsulfoximine (DMSOI)		52–53	100 (6.5 mbar)	[620]
	Tetraethylsulfamide (TES)	14.6[b]		249–251	[621]
	Tetrahydrothiophene-1,1-dioxide (Sulfolane)	16.0	28.4	287.3	[608, 622–624]
	1,3-Dioxolan-2-one (Ethylene carbonate)	16.0	36	156	[625, 626]
	Methylphosphonic acid bis(dimethylamide)	16.0	−3.2	62–63 (4 mbar)	[627, 628]
	4-Methyl-1,3-dioxolan-2-one (Propylene carbonate)	16.5	−54.5	241.7	[608, 625, 629, 630]
	Hexamethylphosphoric acid triamide (HMPT)	18.5	7.3	233	[608, 617, 618, 631]

[a] All formulae are oriented in such a way that the dipole moment of the solvent molecule is approximately in a parallel direction with this arrow.
[b] Value for tetramethylsulfamide; H. Nöth, G. Mikulaschek, and W. Rambeck, Z. Anorg. Allg. Chem. *344*, 316 (1966).

The observation that bimolecular reactions of anions are often much faster in dipolar non-HBD solvents than in protic solvents of comparable relative permittivity is of great practical significance, not only for substitution reactions but also for elimination, proton abstraction, and addition reactions [6].

Reduced solvation of commonly used E2 bases ($HO^{\ominus}$, $RO^{\ominus}$) in dipolar non-HBD solvents may elevate their reactivities to such an extent that E2 reactions of quite inert substrates occur [306]. Halide ions in dipolar non-HBD solvents are sufficiently strong bases to promote dehydrohalogenations of haloalkanes [73, 74]. Even the fluoride ion is the most efficient in this reaction [307, 308, 600]; the elimination rates decrease in the order $F^{\ominus} > Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$.

A shift from $CH_3O^{\ominus}/CH_3OH$ to *tert*-$C_4H_9O^{\ominus}/CH_3SOCH_3$ can produce dramatic increases in rates (up to a factor of 10^9) of reactions which depend on proton abstraction from a C–H bond [31, 304]. The subsequent reaction of the carbanion may involve isomerization, elimination, oxidation, and condensation, hence demonstrating the importance of the potassium *tert*-butoxide/dimethyl sulfoxide system in organic syntheses. The base-catalyzed alkene isomerization reaction (double bond migration) has been used to investigate the efficacy of a wide variety of base/dipolar non-HBD solvent systems [309, 310].

Addition of anionic nucleophiles to alkenes and to heteronuclear double bond systems (C═O, C═S) also lies within the scope of this Section. Chloride and cyanide ions are efficient initiators of the polymerization and copolymerization of acrylonitrile in dipolar non-HBD solvents, as reported by Parker [6]. Even some 1,3-dipolar cycloaddition reactions leading to heterocyclic compounds are often better carried out in dipolar non-HBD solvents in order to increase rates and yields [311]. The rate of alkaline hydrolysis of ethyl and 4-nitrophenyl acetate in dimethyl sulfoxide/water mixtures increases with increasing dimethyl sulfoxide concentration due to the increased activity of the hydroxide ion. This is presumably caused by its reduced solvation in the dipolar non-HBD solvent [312, 313]. Dimethyl sulfoxide greatly accelerates the formation of oximes from carbonyl compounds and hydroxylamine, as shown for substituted 9-oxofluorenes [314]. Nucleophilic attack on carbon disulfide by cyanide ion is possible only in *N*,*N*-dimethylformamide [315]. The fluoride ion, dissolved as tetraalkylammonium fluoride in dipolar difluoromethane, even reacts with carbon dioxide to yield the fluorocarbonate ion, $F–CO_2^-$ [840].

The superoxide ion, $O_2^{\ominus}$, produced by the electron-transfer reduction of dioxygen ($O_2 + e^{\ominus} \rightleftharpoons O_2^{\ominus}\cdot$), is a strong Brønsted base and an effective nucleophile. Because of rapid hydrolysis and disproportionation, the lifetime of $O_2^{\ominus}\cdot$ in aqueous solution is limited. This has led to investigations of its reaction chemistry in dipolar non-HBD solvents [632]. Under these conditions, the superoxide ion attacks haloalkanes by S_N2 displacement of the halides to eventually give dialkyl peroxides in a multi-step reaction [632].

5.5.3 Quantitative Separation of Protic and Dipolar Aprotic Solvent Effects for Reaction Rates by Means of Solvent-Transfer Activity Coefficients

By considering the changes in *standard molar Gibbs energy of solvation*, $\Delta G^{\circ}_{\mathrm{solv}}$, of reactants and activated complexes, when a reaction is transferred from one solvent to another, it is possible to approach the problem from a quantitative point of view. The observed changes in Gibbs energy of solvation when an electrolyte $M^{\oplus}X^{\ominus}$ is transferred from a reference solvent O to another solvent S is commonly reported as the *solvent-transfer activity coefficient* ${}^{\mathrm{O}}y^{\mathrm{S}}$*) [6, 291, 316–319, 454, 634, 635]. This coefficient makes it possible to relate the difference in chemical potential, μ, of a solute MX at infinite dilution on transferring it from an arbitrarily chosen reference solvent O to another solvent S at temperature T according to Eq. (5-114)**):

---

* Also called the *medium effect*, *solvent activity coefficient*, or *transfer activity coefficient*, and also written as $y_t^{\infty}(\mathrm{MX}, \mathrm{O} \rightarrow \mathrm{S})$. It is a constant characteristic of the solute MX (or the solute ions $M^{\oplus}$ and $X^{\ominus}$) and the two solvents O and S.

** The change in standard molar Gibbs energy on transferring an electrolyte MX from one solvent to another may also be expressed as the *standard molar Gibbs energy of transfer*, ΔG_t° [6, 454, 634, 635]. The two quantities, solvent-transfer activity coefficient, ${}^{\mathrm{O}}y^{\mathrm{S}}$, and Gibbs energy of transfer, ΔG_t°, are related by the following simple equation:

$$\Delta G_t^{\circ}(\mathrm{MX}, \mathrm{O} \rightarrow \mathrm{S}) = \mu^{\infty}_{\mathrm{MX}}(\text{in S}) - \mu^{\infty}_{\mathrm{MX}}(\text{in O}) = RT \cdot \ln({}^{\mathrm{O}}y^{\mathrm{S}}_{\mathrm{M}^{\oplus}} \cdot {}^{\mathrm{O}}y^{\mathrm{S}}_{\mathrm{X}^{\ominus}}),$$

where $\mu^{\infty}_{\mathrm{MX}}$(in S) and $\mu^{\infty}_{\mathrm{MX}}$(in O) are the standard chemical potentials of MX referred to at infinite dilution in solvents S and O, respectively. In this Section, results are quoted as solvent-transfer activity coefficients using the molar concentration scale; *cf.* Eq. (2-12a) in Chapter 2.3.

$$\mu^{\infty}_{MX}(\text{in S}) = \mu^{\infty}_{MX}(\text{in O}) + RT \cdot \ln {}^{O}y^{S}_{MX} \tag{5-114}$$

A positive value of $\ln {}^{O}y^{S}_{MX}$ implies that the solute MX is better solvated by solvent O than by solvent S. A negative value indicates that solvation by solvent S is greater. Values of $\ln {}^{O}y^{S}_{MX}$ are obtained from various measurements, *e.g.* solubility, distribution, vapour pressure, and electrochemical measurements. Although solvent-transfer activity coefficients can be defined for individual ions $M^{\oplus}$ and $X^{\ominus}$ they cannot be independently measured. In order to divide the solvent-transfer activity coefficient ${}^{O}y^{S}_{MX}$ into separate contributions ${}^{O}y^{S}_{M^{\oplus}}$ and ${}^{O}y^{S}_{X^{\ominus}}$, an extrathermodynamic assumption must be considered; *cf.* Chapter 2.3 and [454, 634, 635].

The most convenient and generally accepted extrathermodynamic assumption is that ${}^{O}y^{S}_{Ph_4As^{\oplus}} = {}^{O}y^{S}_{Ph_4B^{\ominus}}$, using $Ph_4As^{\oplus}Ph_4B^{\ominus}$ as the reference electrolyte ($Ph = C_6H_5$). The tetrahedral ions of tetraphenylarsonium tetraphenylboride are of comparable structure and size. It is proposed, therefore, that the anion and the cation are similarly influenced on transfer from one solvent to another. This assumption makes it possible to calculate reasonable values for single-ion solvent-transfer activity coefficients (and single-ion Gibbs energies of transfer; *cf.* Table 2-9 in Chapter 2.3) between solvent pairs. Table 5-19 shows some selected values of $\lg {}^{O}y^{S}$ for anions and cations. These have been taken from Marcus's extensive critical compilation [634, 635]. All $\lg {}^{O}y^{S}$ values are expressed relative to water as the reference solvent O at 25 °C.

The striking increase in $\lg {}^{W}y^{S}_{ion}$ values for small anions with high charge density, on going from water to any of the dipolar non-HBD solvents, is consistent with the large rate increase observed in reactions of these anions and gives quantitative significance to the qualitative discussion in Section 5.5.2. The rather large increase in the solvent-transfer activity coefficient of $CH_3CO_2^{\ominus}$, compared with $Cl^{\ominus}$, for the isodielectric solvent change from CH_3OH to CH_3CN and $HCON(CH_3)_2$, reflects the considerable stabilization of this anion through hydrogen-bonding in both methanol and water. As expected, the larger and more polarizable anions show considerably smaller increases in $\lg {}^{W}y^{S}_{ion}$ values on transfer from water to dipolar non-HBD solvents (*e.g.* $I^{\ominus}$, $SCN^{\ominus}$). An extreme example of the difference between small, non-polarizable and large, polarizable anions is seen in the $\lg {}^{W}y^{S}$ values of $F^{\ominus}$ and $(C_6H_5)_4B^{\ominus}$ for the solvent change $CH_3OH \rightarrow HCON(CH_3)_2$: $\lg {}^{W}y^{DMF}_{F^{\ominus}}$ and $\lg {}^{W}y^{DMF}_{Ph_4B^{\ominus}}$ differ by 14.7 units, which corresponds to a difference in Gibbs energy of transfer of 84 kJ/mol.

In addition, it is worth emphasizing the contrasting behaviour of anions and cations on going from water to protic and to dipolar non-HBD solvents. Cations are generally smaller and less polarizable than anions and they are not hydrogen-bond acceptors. Some cations, however, are pronounced Lewis acids and are substantially better solvated in solvents which contain basic oxygen atoms and behave like EPD solvents (*cf.* Section 3.3.2). Therefore, small closed-shell cations such as $Li^{\oplus}$, $Na^{\oplus}$, and $K^{\oplus}$ are much more extensively solvated by solvents such as $[(CH_3)_2N]_3PO$, $HCON(CH_3)_2$, and CH_3SOCH_3, as well as by water, than by methanol. The silver cation has a strong, apparently specific interaction with solvents such as $[(CH_3)_2N]_3PO$, $HCON(CH_3)_2$, CH_3SOCH_3, and CH_3CN. The interaction of $Ag^{\oplus}$ with CH_3COCH_3 and CH_3OH is much weaker. Large organic cations [*e.g.* $(CH_3)_4N^{\oplus}$, $(C_6H_5)_4As^{\oplus}$] are very poorly solvated by water but are much better solvated by dipolar non-HBD solvents.

When solvent-transfer activity coefficients are applied to reaction rates in terms

Table 5-19. Some representative values of solvent-transfer activity coefficients for anions and cations at 25 °C with water (W) as reference solvent; molar concentration scale[a]. For the corresponding Gibbs energies of transfer, see Table 2-9 in Section 2.3.

Ions	lg ${}^{W}y^{S}$ for solvent S[b]						
	S = CH_3OH	$HCONH_2$	CH_3CN	CH_3COCH_3	CH_3SOCH_3	$HCON(CH_3)_2$	$[(CH_3)_2N]_3PO$
$F^{\ominus}$	2.8	4.4	12.4	13.9	12.8	14.9	–
$Cl^{\ominus}$	2.3	2.4	7.4	10.0	7.0_5	8.5	10.2
$Br^{\ominus}$	1.9_5	1.9	5.5	7.4	4.8	6.3	8.0_5
$I^{\ominus}$	1.3	1.3	2.9	4.4	1.8	3.6	5.2_5
$N_3^{\ominus}$	1.6	1.9	6.5	7.5	4.5	6.3	8.6
$CN^{\ominus}$	1.5	2.3	6.1	8.4	6.1	7.0	–
$SCN^{\ominus}$	0.98	1.2	2.5	–	1.7	3.2	3.5
$ClO_4^{\ominus}$	1.1	−2.1	0.35	–	–	0.70	−1.2
$CH_3CO_2^{\ominus}$	2.8	3.5	10.7	–	–	11.6	–
$(C_6H_5)_4B^{\ominus}$	−4.2	−4.2	-5.7_5	−5.6	-6.5_5	-6.7_5	−6.8
$Li^{\oplus}$	0.77	-1.7_5	4.4	–	−2.6	-1.7_5	–
$Na^{\oplus}$	1.4	−1.4	2.6_5	–	-2.3_5	−1.7	–
$K^{\oplus}$	1.7	−0.75	1.4	0.70	−2.3	−1.8	−2.8
$Ag^{\oplus}$	1.1_5	−2.7	-4.0_5	1.6	−6.1	-3.6_5	−7.7
$(CH_3)_4N^{\oplus}$	1.0_5	–	0.53	0.53	−0.35	−0.93	–
$(C_6H_5)_4As^{\oplus}$	−4.2	−4.2	-5.7_5	−5.6	−6.6	-6.7_5	−6.8

[a] Values recalculated from Table 2.4 in Y. Marcus, Pure Appl. Chem. *55*, 977 (1983) [634]. Some values for F^- were taken from G. T. Hefter, Pure Appl. Chem. *63*, 1749 (1991). See also G. Hefter, Y. Marcus, and W. E. Waghorne, Chem. Rev. *102*, 2773 (2002).

[b] A positive value of lg ${}^{W}y^{S}$ indicates that the ion is better solvated by water than by solvent S; a negative value means that the ion is more strongly solvated by transferring it to the solvent S. A difference in lg ${}^{W}y^{S}$ of unity corresponds to a difference in Gibbs energy of solvation of $\lg 10 \cdot RT \cdot \lg {}^{W}y^{S}$, *i.e.* 5.7 kJ/mol at 25 °C.

of the absolute rate theory, where reactions are considered to involve an equilibrium between the reactants and the activated complex, Eq. (5-115) is obtained for a bimolecular reaction such as $Y^{\ominus} + RX \rightleftharpoons [YRX^{\ominus}]^{\neq} \rightarrow YR + X^{\ominus}$ [6, 291, 292].

$$\lg(k^S/k^O) = \lg {}^{O}y^{S}_{Y^{\ominus}} + \lg {}^{O}y^{S}_{RX} - \lg {}^{O}y^{S}_{[YRX^{\ominus}]^{\neq}} \qquad (5\text{-}115)^{*)}$$

The specific rate constant for this bimolecular reaction in a solvent S is related to the rate constant in the reference solvent O through the appropriate solvent-transfer activity coefficients. Eq. (5-115) shows to what extent solvent effects on the reaction rate are due to changes in the solvation of reactant anions, $Y^{\ominus}$, of reactant nonelectrolytes, RX, and of anionic activated complexes, $[YRX^{\ominus}]^{\neq}$. Anionic and uncharged activated complexes will behave in exactly the same way upon solvent transfer as "real" anions and non-electrolytes of comparable structure. Anionic activated complexes such as $[YRX^{\ominus}]^{\neq}$ should behave like large polarizable anions, and, therefore, should be better solvated in polarizable, dipolar solvents than in protic solvents.

The rate constants k in the reference solvent O and in solvent S can be readily measured. Values of $\lg {}^{O}y^{S}$ for RX and $Y^{\ominus}$ are obtained as described above. Thus, by applying Eq. (5-115), $\lg {}^{O}y^{S}$ values for the activated complex can be calculated, providing information about the structure and charge distribution of activated complexes. Values of $\lg {}^{O}y^{S}_{[YRX^{\ominus}]^{\neq}}$ from Eq. (5-115) are compared with $\lg {}^{O}y^{S}$ values for relevant solutes, or for other well-established activated complexes, whose structure and charge distribution are such that they might act as models for the activated complex under consideration. The model with a $\lg {}^{O}y^{S}$ value close to $\lg {}^{O}y^{S}_{[YRX^{\ominus}]^{\neq}}$ may be used as a guide to the structure of $[YRX^{\ominus}]^{\neq}$. Parker and coworkers have estimated the $\lg {}^{O}y^{S}$ values of activated complexes for numerous variations of RX, $Y^{\ominus}$ and mechanisms (*e.g.* S_N2, S_NAr) for substitution reactions of various charge types [6, 291, 292]. Using the Gibbs energy of transfer approach*), similar calculations of solvent effects on the Gibbs energy of reactants and activated complexes have been collected by Abraham [23], especially for Menschutkin reactions [64].

Examples of the use of this procedure are shown in Table 5-20. These data for the S_N2 reaction (5-116) are taken from Parker *et al.* [6, 291], whose extensive compilation compares relative rates and solvent-transfer coefficients on transfer from the reference solvent methanol to the isodielectric solvent *N*,*N*-dimethylformamide at 25 °C.



(5-116)

On transfer from methanol to *N*,*N*-dimethylformamide, the increase in the rate of these four analogous S_N2 reactions varies from $10^{2.0}$ to $10^{4.6}$, the rate being faster in

---

* An alternative way to analyze the solvent effect on this S_N2 reaction in terms of Gibbs energy of transfer, ΔG_t°, of reactants and activated complex from reference solvent O to another solvent S is given by the following equation [6, 23, 64, 292]:

$$\Delta G_t^{\neq} = \Delta G_t^{\circ}(Y^{\ominus}) + \Delta G_t^{\circ}(RX) - \Delta G_t^{\circ}([YRX^{\ominus}]^{\neq}).$$

For an attempt to correlate solvent transfer Gibbs energies of activation, ΔG_t°, of some S_N1 and S_N2 reactions with the donor and acceptor numbers of the solvent, see reference [292a].

Table 5-20. Relative reaction rates and solvent-transfer activity coefficients[a] for reactants and activated complex of S_N2 reaction (5-116) upon solvent transfer from methanol (abbreviated to M) to *N*,*N*-dimethylformamide (DMF) at 25 °C [6, 291].

Leaving Group X	$\lg\left(\frac{k_2^{DMF}}{k_2^{M}}\right)$	$\lg {}^{M}y^{DMF}_{N_3^{\ominus}}$	$\lg {}^{M}y^{DMF}_{CH_3X}$	$\lg {}^{M}y^{DMF}_{[N_3CH_3X^{\ominus}]^{\neq}}$	$\lg {}^{M}y^{DMF}_{X^{\ominus}}$
$Cl^{\ominus}$	3.3	4.9	−0.4	1.2	6.5
$Br^{\ominus}$	3.9	4.9	−0.3	0.7	4.9
$I^{\ominus}$	4.6	4.9	−0.5	−0.2	2.6
$TsO^{\ominus}$	2.0	4.9	−0.6	2.3	3.5

[a] The more positive the value of $\lg {}^{M}y^{DMF}$, the greater the solvation by methanol relative to *N*,*N*-dimethylformamide.

the dipolar non-HBD solvent. Whereas the medium has only a small effect on the CH_3X Gibbs energy (these molecules being slightly better solvated in *N*,*N*-dimethylformamide almost independently of X), the anion $N_3^{\ominus}$ is substantially destabilized on transfer from CH_3OH to $HCON(CH_3)_2$. The increase in rate, superimposed by a relative increase in solvation of the activated complex in *N*,*N*-dimethylformamide in the order $[N_3CH_3OTs^{\ominus}]^{\neq} < [N_3CH_3Cl^{\ominus}]^{\neq} < [N_3CH_3Br^{\ominus}]^{\neq} < [N_3CH_3I^{\ominus}]^{\neq}$, clearly reflects this destabilization. It is, however, modified by a small destabilization of the activated complex when X = $TsO^{\ominus}$, $Cl^{\ominus}$, and $Br^{\ominus}$, thus showing the anionic nature of the activated complex. For X = halide, this increase in stability correlates with the increasing polarizability of the activated complexes and corresponds to the increase in polarizability from $Cl^{\ominus} < Br^{\ominus} < I^{\ominus}$. Differences in solvation of activated complexes do not appear to contribute greatly to $\lg(k_2^{DMF}/k_2^{M})$ when halide ions are the leaving groups. However, the comparatively low rate increase obtained for methyl tosylate must be due to the more positive value of $\lg {}^{M}y^{DMF}_{[N_3CH_3OTs^{\ominus}]^{\neq}}$, because the tosylate ion itself has a less positive $\lg {}^{M}y^{DMF}$ value than does $Br^{\ominus}$ or $Cl^{\ominus}$, yet the activated complex containing tosylate has $\lg {}^{M}y^{DMF}_{\neq}$ more positive than the activated complexes containing halide. This effect has been explained by assuming that a looser activated complex with a greater localization of negative charge is present in the case of tosylate. Transition state anions with increased localization of negative charge on the leaving and incoming groups become increasingly more solvated, for the reasons already given. Departing tosylate is a better hydrogen-bond acceptor relative to free tosylate ion than is departing halide relative to free halide ion. Among other observations this leads to the generalization that, other things being equal, those $Y^{\ominus} + RX$ reactions possessing the "tightest" activated complex (*i.e.* most covalently bound Y and X) give the greatest accelerations on transfer from protic to dipolar non-HBD solvents. For a more detailed description of "tight and loose S_N2 activated complexes", see the reviews of Parker [6, 291].

Assuming that $\lg {}^{O}y^{S}_{RX} - \lg {}^{O}y^{S}_{[YRX^{\ominus}]^{\neq}}$ is roughly constant for closely related reactions, Parker obtained, from Eq. (5-115), the simple linear Gibbs energy relationship (5-117),

$$\lg(k^S/k^O) = \lg {}^{O}y^{S}_{Y^{\ominus}} + C \qquad (5\text{-}117)$$

This involves a limited number of constants C, varying only with the nature of the substrates [6, 291]. It is possible to use Eq. (5-117) to estimate the rate constant for a given reaction in a new solvent S from its value in another reference solvent O, to within a factor of two. This is very good in view of the fact that rates can vary by a factor of up to 10^{10} on switching solvents. Assuming the validity of Eq. (5-117), the easiest way of estimating lg ${}^{O}y^{S}_{Y^{\ominus}}$ is to measure a reaction rate for $Y^{\ominus}$ in two different solvents. A test of Eq. (5-117) for the S_N2 reaction between bromide ion and methyl tosylate in twelve solvents is given in reference [68].

Further examples of the dissection of initial state and transition state medium effects for reactions in protic and dipolar non-HBD solvents have been given by Buncel [467, 636], Abraham [23, 64, 637], Haberfield [638], and Blandamer *et al.* [639].

5.5.4 Acceleration of Base-Catalysed Reactions in Dipolar Aprotic Solvents

The specific solvation of anions in protic solvents, mediated mainly by hydrogen-bonding, diminishes not only their nucleophilic reactivity but also their basicity. All anions, therefore, are much stronger bases in dipolar non-HBD solvents than in protic solvents. This substantial increase in kinetic and thermodynamic basicity of anions in dipolar non-HBD solvents is very profitable in preparative organic chemistry in view of the activation of weakly acidic C—H, O—H, and N—H bonds. Especially useful for this purpose are solutions of potassium *tert*-butoxide in dimethyl sulfoxide, which is among the most basic media available for the organic chemist. The basicity of the *tert*-$C_4H_9O^{\ominus}/CH_3SOCH_3$ system is comparable to that of the $NH_2^{\ominus}$/liquid NH_3 system. Extremely basic solutions are also generated by the addition of dimethyl sulfoxide to aqueous or alcoholic solutions of alkali metal hydroxides or tetraalkylammonium hydroxides*). Substitution of dimethyl sulfoxide for alcohols or water, not only replaces the proton donor physically, but deactivates the remaining protic solvent by virtue of its own great H-bond acceptor capability according to Eq. (5-118) [841].

$$RO^{\ominus}\cdots H\text{-}OR + (CH_3)_2\overset{\delta\oplus}{S}=\overset{\delta\ominus}{O} \rightleftharpoons RO^{\ominus} + (CH_3)_2\overset{\delta\oplus}{S}=\overset{\delta\ominus}{O}\cdots H\text{-}OR \qquad (5\text{-}118)$$

Many reactions become possible only in such superbasic solutions, while others can be carried out under much milder conditions. Only some examples of preparative interest (which depend on the ionization of a C—H or N—H bond) will be mentioned here. The subsequent reaction of the resulting carbanion may involve electrophilic substitution, isomerization, elimination, or condensation [321, 322]. Systematic studies of solvent effects on intrinsic rate constants of proton-transfer reactions between carbon acids and carboxylate ions as well as amines as bases in various dimethyl sulfoxide/water mixtures have been carried out by Bernasconi *et al.* [769].

---

* The use of pure dimethyl sulfoxide with alkoxide ($CH_3O^{\ominus}$, *tert*-$C_4H_9O^{\ominus}$) or hydride ($H^{\ominus}$) ions leads to even more strongly basic systems which contain, in part, the very strongly basic dimethylsulfinyl ("dimsyl") anion, formed in the following equilibrium reaction:

$$RO^{\ominus} + CH_3\text{—}SO\text{—}CH_3 \rightleftharpoons RO\text{—}H + {}^{\ominus}|CH_2\text{—}SO\text{—}CH_3$$

Rates of isotopic H/D exchange and racemization of optically active 2-methyl-3-propionitrile in the presence of potassium methoxide can be increased by a factor of $5 \cdot 10^7$ on going from pure methanol ($\varepsilon_r = 32.7$) to dimethyl sulfoxide ($\varepsilon_r = 46.5$) [31, 231, 304].



(5-119)

Solvent CH_3SOCH_3 in cg/g	CH_3OH/CH_3SOCH_3						
	0	25	50	76	90	98.5	100
k_2^{rel} (25 °C)	1	32	160	4900	$1.3 \cdot 10^5$	$5.0 \cdot 10^7$	ca. 10^{9} [a]

[a] extrapolated

Much of the progress made in the area of base-catalysed alkene isomerization resulted from the use of dipolar aprotic solvents. Under homogeneous conditions, this brings about high reactions rates at low temperatures [309, 310, 323, 324]. For example, the base-catalysed rearrangement of allyl to propenyl ethers [325], and of allyl to propenyl amines [326] is best carried out using *tert*-C_4H_9OK/CH_3SOCH_3 as the reaction medium [323]. The alkenylation and aralkylation of a variety of alkylaromatic compounds in a homogeneous *tert*-C_4H_9OK/dipolar aprotic solvent system is greatly influenced by the solvent. In the case of the reaction between 4-isopropylpyridine (Ar = 4-pyridyl) and isoprene according to Eq. (5-120), the rate is faster in dimethyl sulfoxide ($t_{1/2} = 1.15$ min!) [324].



(5-120)

The Wittig reaction proceeds more rapidly in dimethyl sulfoxide than in other customary solvents due to the enhanced deprotonation of the phosphonium salts to alkylidenephosphoranes [327]. Even camphor, a rather unreactive ketone, undergoes the Wittig reaction easily in base/CH_3SOCH_3 systems [327].

Similarly spectacular increases in rate were found for the transformation of ketone hydrazones into hydrocarbons using the Wolff–Kishner reduction. This is mainly due to the enhanced ionization of the N—H bond. This reaction occurs even at room temperature in *tert*-C_4H_9OK/CH_3SOCH_3 and proceeds by solvent-mediated transfer of a proton from nitrogen to carbon [328, 329].



(5-121)

Since the hydrazone anion is stable in the absence of a proton donor, the overall reaction must depend on the presence of an alcohol molecule in the vicinity of the hydrazone anion. This is in agreement with the observation that rate constants obtained in ROH/CH_3SOCH_3 mixtures, as a function of dimethyl sulfoxide concentration, exhibit a maximum at low alcohol concentration [330]. The rate-determining step of this reaction appears to be a concerted proton-transfer from the protic solvent to the hydrazone anion and a proton abstraction from the hydrazone anion by the basic solvent [329].

Since the Cope elimination reaction of amine oxides proceeds with intramolecular proton abstraction by the *N*-oxide group according to Eq. (5-122), it is not surprising that with dimethyl sulfoxide as solvent the reaction occurs at room temperature [328, 331]. Protic solvents require temperatures of 120 to 150 °C because hydrogen bonding of the oxygen terminal makes the reaction more difficult.



(5-122)

Further examples of strongly basic solvent systems are solutions of ionic fluorides in dipolar non-HBD solvents such as acetonitrile, *N*,*N*-dimethylformamide, dimethyl sulfoxide, and tetrahydrofuran. Tetraalkylammonium fluorides $R_4N^{\oplus}F^{\ominus}$ (R_4 = tetraethyl, tetra-*n*-butyl, benzyltrimethyl), which are soluble in dipolar non-HBD solvents, have received the most attention. Since they are highly hygroscopic, they are usually associated with varying amounts of protic solvents (mainly water). The amount of these protic solvents influences the effective basicity of the fluoride ion by hydrogen bonding ($RO—H\cdot\cdot F^{\ominus}$). In solutions of anhydrous tetramethylammonium fluoride in acetonitrile, dimethyl sulfoxide, trichloromethane, and dichloromethane, the fluoride ion is basic enough to abstract a proton from CH_3CN and CH_3SOCH_3, and nucleophilic enough to undergo halogen exchange with $CHCl_3$ and CH_2Cl_2 at room temperature [640].

Among the fluoride ion promoted reactions which occur in dipolar non-HBD solvents are alkylations of alcohols and ketones, esterifications, Michael additions, aldol and Knoevenagel condensations as well as eliminations; for a review, see reference [600]. In particular, ionic fluorides are useful in the dehydrohalogenation of haloalkanes and haloalkenes to give alkenes and alkynes (order of reactivity $R_4N^{\oplus}F^{\ominus} > K^{\oplus}$ ([18]crown-6) $F^{\ominus} > Cs^{\oplus}F^{\ominus} \approx K^{\oplus}F^{\ominus}$). For example, tetra-*n*-butylammonium fluoride in *N*,*N*-dimethylformamide is an effective base for the dehydrohalogenation of 2-bromo- and 2-iodobutane under mild conditions [641]; *cf.* Eq. (5-123).



(5-123)

Wittig reactions of phosphonium fluorides in dipolar non-HBD solvents can be carried out without an additional base: *e.g.* 4-nitrobenzaldehyde reacts slowly with (4-nitrobenzyl)triphenylphosphonium fluoride in refluxing acetonitrile to give 4,4′-nitro-stilbene in good yield (84 cmol/mol) [642]. In this reaction, the fluoride ion attacks its own cation to give the corresponding Wittig ylide as intermediate.

Further examples of solvent effects on base-catalysed reactions can be found in references [297–300, 321, 322, 600].

5.5.5 Influence of Specific Cation Solvation on the Rates of S_N Reactions

The nucleophilic reactivity of an anion depends not only on the extent of its specific solvation, but also on the degree of association with the corresponding cation. An ion-pair associated anion (or cation) is much less reactive than a free, non-associated ion*). As early as 1912, Acree postulated that the reactivity of an anionic nucleophile should be depressed when its salt is incompletely dissociated [332]. Due to incomplete dissociation of the ionophore, the reaction rate constant will fall as its concentration increases. The simple model given in Eq. (5-124) is consistent with the observation that in all cases ion association deactivates the nucleophile [289].



(5-124a)





(5-124b)

Formation of the activated complex from an ion pair differs from formation from a free anion in that in reaction (5-124a) the activation process suffers from a loss in coulombic interaction energy between the centres of opposite charges. Therefore, in the presence of $M^{\oplus}$, the activation energy is increased. This model also suggests that the ratio k_2^f/k_2^p should increase with increasing intensity of cation–anion interaction [289]. Ion pairs are more stable the higher the charge densities of the component ions, and hence the stronger the electrostatic attraction. Accordingly, the nucleophilicity of anions in weakly dissociating solvents (*i.e.* solvents with low relative permittivities; *cf.* Sections 2.6 and 3.2) depends on the nature of the binary salt, and therefore on the cation. The relative reactivities of free and associated ions has been examined in detail, not only for

* A covalent compound dissociates into free ions in stages involving the formation of at least two types of ion pairs, contact and solvent-separated ion pairs; *cf.* Section 2.6 and Eqs. (2-19) to (2-21) [289, 333].

nucleophilic aliphatic substitution [333–335], but also for electrophilic aliphatic substitution reactions. In the latter, carbanionic ion pairs or compounds are involved [321, 322, 336–339]. The reactivity of carbanions is affected in the same way by the degree of interaction between the carbanion and the counter ion. For a comprehensive review on all kinds of salt effects in organic and organometallic chemistry, see reference [829].

According to Coulomb's law [*cf.* Eq. (2-17) in Section 2.6], the interaction between cation and anion can be minimized by increasing either the relative permittivity of the medium or the interionic distance. There are several ways of separating anions from cations and thus increasing the anionic reactivity [643, 842]:

(i) enlarge the size of the opposite cation by substituting bulky 'onium cations such as quaternary tetraalkylammonium ions for the small alkali metal cations;

(ii) use EPD solvents of high Lewis basicity as reaction media or additives because they strongly solvate small alkali metal cations, thus increasing the interionic distance; they also weakly solvate anions;

(iii) encage cations by means of macro(poly)cyclic ligands such as crown ethers or cryptands. Cation complexation leads to ion-pair dissociation through formation of *ligand-separated ion pairs.*

Examples of each of these three types of anion activation*) are given.

An example of the S_N2 rate dependence on the nature of the counterion is given by the reaction of *n*-butyl 4-bromobenzene sulfonate with lithium- and tetra-*n*-butylammonium halides in the weakly dissociating solvent acetone ($\varepsilon_r = 20.6$) [279].

$$CH_3CH_2CH_2CH_2\text{-}OBros + X^{\ominus} \xrightarrow[25\,^{\circ}C]{k_2} CH_3CH_2CH_2CH_2\text{-}X + {}^{\ominus}OBros \qquad (5\text{-}125)$$

Anion $X^{\ominus}$		$I^{\ominus}$	$Br^{\ominus}$	$Cl^{\ominus}$
k_2^{rel}	$Li^{\oplus}$ salt[a)]	6.2	> 5.7	> 1.0
		∨	∧	∧
	$(n\text{-}C_4H_9)_4N^{\oplus}$ salt[a)]	3.7	< 18	< 68

a) Salt concentration 0.04 mol/L

The nucleophilic reactivity of the lithium salts changes in the same order as in protic solvents ($I^{\ominus} > Br^{\ominus} > Cl^{\ominus}$; *cf.* Table 5-15). However, the order is completely reversed for the ammonium salts ($Cl^{\ominus} > Br^{\ominus} > I^{\ominus}$), and this latter order is the same as that found in dipolar non-HBD solvents such as *N*,*N*-dimethylformamide [278]. The small lithium cation, with its high charge density, has a strong tendency to form ion pairs with anions, whereas the electrostatic interaction between the large tetraalkylammonium ion and anions is comparatively weak. Quaternary ammonium salts, therefore, should be practically fully dissociated in acetone solution. Thus, the reactivity order obtained with these salts corresponds to that of the free, non-associated halide ions. On the other hand, the sequence obtained with the lithium salts also reflects the dissociation equilibria of these salts in acetone solution [279].

* The term anion activation suggests that methods (i) ... (iii) cause an increase in reaction rate. This does not hold for reactions in which the ion pair is more reactive than the free anion as a result of a cation-assisted reaction pathway. Methods (i) ... (iii) exhibit cation *de*activation in conjunction with the anion activation!

Similarly, the reactivity of phenolate ions as the tetra-*n*-butylammonium salt has been shown to be $3 \cdot 10^4$ times higher than that of the corresponding potassium salt in the S_N2 alkylation reaction with 1-halobutanes, carried out in 1,4-dioxane [340]. Whereas the rate of alkylation of the potassium salt increases by a factor of ca. 10^5 on going from 1,4-dioxane ($\varepsilon_r = 2.2$) to *N*,*N*-dimethylformamide ($\varepsilon_r = 36.7$), the alkylation rate of the quaternary phenolate is essentially insensitive to the same solvent change. Obviously, the phenolate ion combined with the larger ammonium ion is already very reactive because of the relatively weak cation–anion interaction in the ion pair. In such cases, dissociation to a truly free anion does not seem to be required in order to explain the high reactivity [340].

Examination of the S_N2 reaction between ethyl tosylate and halide ions in hexamethylphosphoric triamide ($\varepsilon_r = 29.3$) with a variety of counter ions [$Li^{\oplus}$, $(n\text{-}C_4H_9)_4N^{\oplus}$] has shown that the rates obtained with lithium salts are always higher than those with the corresponding tetra-*n*-butylammonium salts [341]. This is in contrast to the situation observed in acetone [279]. This means that, in this particular solvent, lithium salts are more dissociated than tetraalkylammonium salts. This has indeed been confirmed by conductivity measurements [341, 342]. The lithium cation apparently has specific interactions with strong EPD solvents such as $[(CH_3)_2N]_3PO$ (*cf.* Section 3.3.2).

Another effect, called the "cesium effect" [843, 844], is also connected with the observation that salts of anions with large counterions are highly dissociated in dipolar aprotic solvents, and consequently more anion-reactive. For example, the higher solubility of cesium carbonate in dipolar aprotic solvents and the fact that this salt is far more dissociated than the corresponding Li^+, Na^+, or K^+ salts, makes this carbonate a superior base in organic synthesis. Amongst the alkali metal cations, the ionic radius of Cs^+ ($r = 334$ pm) is more than twice that of Li^+ ($r = 152$ pm).

Ion-pair dissociation can be achieved not only by using large cations with correspondingly low charge densities, but also by employing EPD solvents as either reaction media or as additives to salt solutions in other solvents. EPD solvents of high Lewis basicity are particularly good specific cation solvators and weak anion solvators, thus giving rise to highly reactive anions. The cation-solvating capacity of EPD solvents can be quantitatively described in terms of their high donor numbers *DN* (*cf.* Table 2-3 in Section 2.2.6) as well as their large negative solvent-transfer activity coefficients for cations (*cf.* Table 5-19 in Section 5.5.3). Good cation-solvating EPD solvents include most of the common dipolar non-HBD solvents (*cf.* Table 5-18 in Section 5.5.2), as well as open-chain polyethers such as oligoethylene glycol dialkyl ethers ("glymes"), which contain the repeating unit $(—CH_2—CH_2—O—)_n$ ($n \geq 2$) [345].

Alkali metal derivatives of CH-acidic compounds are usually highly aggregated in solvents of low relative permittivity such as benzene ($\varepsilon_r = 2.3$). The accelerating influences of a variety of EPD solvents used as additives in the alkylation of diethyl sodio-*n*-butylmalonate with 1-bromobutane in benzene according to Eq. (5-126) was discovered by Zaugg *et al.* [350]; *cf.* Table 5-21.



(5-126)

Table 5-21. Relative rate constants of the alkylation of diethyl sodio-*n*-butylmalonate with 1-bromobutane in benzene at 25 °C with different additives [350].

Added Cation Solvator (0.648 mol/L)	$k_2^{rel.}$
No additive (benzene alone)	1
Tetrahydrofuran	1.1
CH_3COCH_3	1.3
C_2H_5OH	4.4
$CH_3O—CH_2CH_2—OCH_3$	6.4[a]
1,4-Dioxane	18
$HCON(CH_3)_2$	19
1-Methylpyrrolidin-2-one	30
$[(CH_3)_2N]_3PO$[b]	54

[a] In pure 1,2-dimethoxyethane (8.7 mol/L) the reaction is 80 times faster than in benzene.
[b] Only 0.324 mol/L added.

Table 5-21 shows that the addition of even small proportions of EPD solvents affects the reaction rate markedly. The rate acceleration thus obtained is produced by a specific solvation of sodium ion, which tends to dissociate the high-molecular mass ion-pair aggregate of the sodio-malonic ester that exists in benzene solution (degree of aggregation n is equal to 40 . . . 50 in benzene). This indicates that the kinetically active species is a lower aggregate of the free carbanion. Further evidence for a specific cation solvation is derived from the six-fold rate difference observed in tetrahydrofuran ($\varepsilon_r = 7.6$) and 1,2-dimethoxyethane ($\varepsilon_r = 7.2$), despite the fact that these two solvents possess nearly equal relative permittivities. The latter solvent is able to solvate sodium ions in the manner shown in Eq. (5-127). Especially noteworthy is the high reactivity exhibited on the addition of dicyclohexyl[18]crown-6. In benzene solution containing only 0.036 mol/L of this crown ether, the alkylation rate is already equal to that observed in neat 1,2-dimethoxyethane [351].

The reactivity of sodio-butyrophenone toward 1-bromo-2-methylpropane is greatly enhanced in monoglyme (2560-fold) and diglyme (11400-fold) with respect to the rate in diethyl ether according to Zook *et al.* [352]. The ethylation of sodio-butyrophenone (0.13 mol/L) by bromoethane (1.6 mol/L) in diglyme at 30 °C is 75% complete in 152 s, whereas the corresponding reaction time for a comparable ethylation in diethyl ether is 234 h [353]. Again, specific solvation of the cation in the sodium enolate aggregates may explain these results. The reactivity toward bromoethane increases in the order $Li^{\oplus} < Na^{\oplus} < K^{\oplus}$-enolate, corresponding to the decreasing charge density of the cations [353].

Finally, anion activation can also be achieved by the complexation of the cation using suitable macro(poly)cyclic ligand molecules [643]. Organic ligands that contain enforced cavities of dimensions at least equal to those of the smaller cations (and anions) have been called *cavitands* [644]. Cation complexation by such macro(poly)cyclic ligands leads to dissociation of the ion pairs as well as to salt solubilization. Complex

formation with lipophilic organic ligands transforms small cations into voluminous lipophilic cationic species, which are much more soluble in organic solvents of low polarity*). Therefore, the increase in reaction rates induced by specific cation-complexing ligands usually results from an increase in the anion reactivity *as well as* an increase in the reagent concentration.

Some general examples of anion activation by different types of cation complexation are given in Eqs. (5-127)...(5-131). Among the specific multidentate complexing agents for cations are:

(a) open-chain *podands*, such as the oligoethylene glycol dialkyl ethers ("glymes"), investigated by Vögtle *et al.*; *cf.* Eq. (5-127) [345].

(b) cyclic *coronands*, such as the crown ethers, first introduced by Pedersen in 1967 [343]; *cf.* Eq. (5-128) [348, 645–647];

(c) macrobi-/tricyclic spherical *cryptands*, first introduced by Lehn *et al.* in 1969 [344, 648]; *cf.* Eqs. (5-129) and (5-130). Eq. (5-130) illustrates that in the case of the ammonium ion the hydrogen-bond acceptor ability of the macrotricyclic ligand can also be important [349, 648];

(d) cyclic *spherands*, first introduced by Cram *et al.* in 1979 [644, 649]; *cf.* Eq. (5-131).

Whereas the outside of these cationic complexes is lipophilic, the cations are held inside the hydrophilic cavity of the organic ligands. In the case of Eq. (5-127), the



(5-127)

Pentaglyme, an open-chain *podand*



(5-128)

[18]Crown-6, a cyclic *coronand*

* Using water/organic solvent two-phase systems, the distribution of a salt in the organic layer is enhanced by lipophilic cation-complexing ligands. This is the basis of *liquid-liquid phase-transfer catalysis*. Phase-transfer catalysts help to transfer a water-soluble reactant salt across the interface into the organic phase where a homogeneous reaction can occur, thus enhancing the rate of reaction. For further explanations of phase-transfer catalyzed reactions, see Section 5.5.13, particularly Fig. 5-18, and reviews [656–658, 879–883].



(5-129)

[2.2.2]Cryptand, a bicyclic spherical *cryptand*



(5-130)

A tricyclic spherical *cryptand*



(5-131)

A cyclic *spherand*

podands cavity is built up during complexation by a template effect. The stability constants K_s increase from Eq. (5-127) to Eq. (5-131). Typical orders of magnitudes for K_s values in methanol are $10^2 \ldots 10^4$ for podates, $10^4 \ldots 10^5$ for coronates, and $10^6 \ldots 10^8$ for cryptates*). The appropriate choice of ligand allows a selective complexation of a particular cation. For economic reasons, the simple and cheap open-chain glymes have found greater application in anion activation. Only a few representative examples of enhanced anion reactivity caused by cavitands will be mentioned here.

The effect of both podands and coronands on the S_N2 alkylation of potassium phenolate with 1-bromobutane in 1,4-dioxane ($\varepsilon_r = 2.2$) has been investigated by Ugel-

* It should be mentioned that cation complexation by crown-type ligands can itself be solvent-dependent. For example, the dissociation rate of potassium [2.2.2]cryptate in EPD solvents increases with the donor number of the solvent [650]. Moreover, coronands themselves can interact with organic solvent molecules [651]. Such cation-solvent and ligand-solvent interactions can influence the formation of cation-ligand complexes.

stad *et al.* [354]. The addition of tetraethylene glycol dimethyl ether in concentrations equivalent to the phenolate concentration leads to an alkylation rate 11 times faster than that in pure 1,4-dioxane, whereas the same amount of dicyclohexyl[18]crown-6 causes an 8700-fold rate increase.

The base-catalysed hydrolysis of the sterically hindered ester methyl 2,4,6-trimethylbenzoate can be carried out in KOH/DMSO at room temperature in 100% yield using a [2.2.2]cryptand as the potassium-complexing ligand [652].

Liotta *et al.* [282, 355, 356, 359] investigated the chemistry of so-called "naked anions"*), *i.e.* poorly solvated fluoride, cyanide, and acetate ions, solubilized as potassium salts in acetonitrile or benzene containing [18]crown-6 as an additive. The reactivity of these naked anions in reactions with haloalkanes has been examined. According to the haloalkane structure, the reaction leads to substitution and/or elimination products. Solubilised fluoride ions are both potent nucleophiles and strong bases, whereas solubilised cyanide and acetate ions are good nucleophiles but rather weak bases. A variety of fluoroalkanes [282, 355] and nitriles [356, 361] can be prepared in good yields using crown ethers as cation solvators. The carboxylate ion is generally considered a bad nucleophile, but [18]crown-6 complexed potassium acetate (also called "bare acetate") reacts readily with haloalkanes in acetonitrile to yield the corresponding esters [359, 360]. The nitrite ion also displays a remarkable enhanced nucleophilicity in the presence of [18]crown-6. Under these conditions, nitro compounds are formed from haloalkanes in good yields [361].

Other anions ($X^{\ominus} = Br^{\ominus}$, $I^{\ominus}$, $HO^{\ominus}$, $CH_3O^{\ominus}$) also exhibit enhanced nucleophilic reactivity in crown-ether mediated reactions of $K^{\oplus}X^{\ominus}$ [357, 358]. Montanari *et al.* [653] compared the effects of phosphonium salts, coronands, and cryptands on the S_N2 reaction of *n*-octylmethanesulfonate with various nucleophiles in chlorobenzene.

Apart from these examples of nucleophilic substitution reactions, there are many more applications of reactions which are enhanced by the addition of macro(poly)cyclic ligands; examples can be found in some excellent reviews [346–348, 362, 643, 645–648].

Cation-assisted reactions, on the other hand, are hindered by complexation of the cation [648]. Reduction of carbonyl groups by metal hydrides [654] and the addition of organolithium compounds [655] are examples. This indicates that the coordination between carbonyl group and cation is an important step in these reactions. For applications of concentrated $LiClO_4/Et_2O$ (LPDE) solutions in cation-catalysed reactions, see Section 5.3.3 and references [803–806].

In this context, it should be mentioned that free cations in suitable solvents can be used as templates for the synthesis of macro(poly)cyclic compounds (*e.g.* catenanes). Such template-based syntheses are mediated by slightly solvated metal cations, which assemble the reaction partners as ligands, allowing them to easily react with each other; the desired product is obtained after removal of the template; for a review, see reference [845].

---

* Completely unsolvated "naked anions" cannot be prepared in solution with coronands or even with cryptands as cation solvators. Even in this case ion-pairing still occurs leading to complexed ion pairs [646]. Totally unsolvated "naked anions" can exist only in the gas phase (*cf.* Section 5.2.).

5.5.6 Solvent Influence on the Reactivity of Ambident Anions

Ambident anions are mesomeric, nucleophilic anions which have at least two reactive centers with a substantial fraction of the negative charge distributed over these centers*)**). Such ambident anions are capable of forming two types of products in nucleophilic substitution reactions with electrophilic reactants***). Examples of this kind of anion are the enolates of 1,3-dicarbonyl compounds, phenolate, cyanide, thiocyanide, and nitrite ions, the anions of nitro compounds, oximes, amides, the anions of heterocyclic aromatic compounds (*e.g.* pyrrole, hydroxypyridines, hydroxypyrimidines) and others; *cf.* Fig. 5-17.



Fig. 5-17. Some ambident or ambifunctional anions. The arrows indicate the sites of dual reactivity towards electrophiles.

If the substitution reaction is kinetically controlled, then the composition of the products is determined by the relative nucleophilicity of each of the donor atoms in the ambident anion in relation to the given electrophilic reactant. Among the factors influencing the mode of reaction (counterion, additives, concentration, temperature, pressure, leaving group, structure of the alkylating agent), the solvent plays a major role in the orientation of the electrophile, and this has been examined in several reviews [364–367, 367a, 367b]. As early as 1923, Claisen noticed that the O/C alkylation ratio observed in the reaction of phenol with 3-bromopropene in the presence of potassium carbonate depends strongly on the solvent used as reaction medium [369]. In acetone, the major product formed was allyl phenyl ether, whereas in solvents such as benzene or toluene, *ortho*-allyl phenol was obtained as the main product [369].

---

* The term *ambident anion* was proposed by Kornblum [363], whereas Gompper suggested that such anions should be called *ambifunctional* [364, 367].

** In principle, *ambident cations* may be defined in the same way, but the solvent dependence of their dual reactivity with nucleophiles has not yet been investigated thoroughly [368].

*** When a reaction can potentially give rise to two (or more) constitutional isomers but actually produces only one, the reaction is said to be *regioselective.* For example, the ambident nucleophile $NCO^{\ominus}$ usually gives only isocyanates R—NCO and not the isomeric cyanates, R—OCN.

Table 5-22. Oxygen versus carbon alkylation in the reaction of two sodium arenolates with haloalkanes at room temperature [370, 371].

Solvents	% *O*-Alkylation	% *C*-Alkylation[a)]
a) *Reaction of sodium phenolate with 3-chloropropene according to equation (5-132)*		
1,4-dioxane	100	0
$(CH_3)_3COH$	100	0
C_2H_5OH	100	0
CH_3OH	100	0
$HCON(CH_3)_2$	100	0
$(CH_2)_4O$	96	0
H_2O	49	41
CF_3CH_2OH[b)]	37	42
C_6H_5OH[c)]	22	78
b) *Reaction of sodium β-naphtholate with (bromomethyl)benzene*		
$HCON(CH_3)_2$	97	0
CH_3SOCH_3	95	0
$CH_3O—CH_2CH_2—OCH_3$	70	22
$(CH_2)_4O$	60	36
CH_3OH	57	34
C_2H_5OH	52	28
H_2O	10	84
CF_3CH_2OH	7	85

[a)] Dialkylated products included.
[b)] Reaction with 3-bromopropene.
[c)] At 43 °C.

Some representative data, taken from the extensive work of Kornblum *et al.* concerning the alkylation of phenolates [*cf.* Eq. (5-132)] and β-naphtholates, are collected in Table 5-22 [370, 371].



(5-132)

In protic solvents, the solvent effect on orientation has been interpreted in terms of preferential deactivation of the more electronegative atom by specific solvation through H-bonding*). As shown in Table 5-22, carbon alkylation competes more sig-

* Kornblum termed this shielding of the center with maximal electron density in ambident anions by protic solvents, which prevents the reaction at this center, "selective solvation" [370]. In order to avoid confusion with selective solvation of ions or dipolar molecules in binary solvent mixtures (*cf.* Section 2.4), the designation "specific solvation" is preferred. An IR and NMR spectroscopic study of the specific solvation of OCN^-, SCN^-, and $SeCN^-$ ions has shown that, in neat protic solvents (*e.g.* MeOH, $HCONH_2$), these anions form intimate hydrogen bonds, albeit to different extents. In methanol solution, a significant proportion of the cyanate ion is hydrogen-bonded at the oxygen end, while the other two anions are much less solvated than OCN^-, forming hydrogen bonds to the nitrogen end and non-axially to the C=N π-bond to only a limited extent [846].

nificantly with oxygen alkylation as the protic solvents become stronger proton donors (*cf.* $CH_3CH_2OH \rightarrow H_2O \rightarrow CF_3CH_2OH \rightarrow C_6H_5OH$). In 2,2,2-trifluoroethanol, a very strong proton donor, the yield of *C*-alkylated isomers approaches 42% and 85%, respectively. Owing to H-bonding, protic solvents selectively and effectively solvate the center with the maximum electron density. As a result of this, the accessibility of this center is reduced and substitution at the other donor atom (in this case a carbon atom) may successfully compete with that at the oxygen atom. This protic solvent effect is also observed for other ambident anions. For example, in the alkylation of the enolate salt of acetoacetic ester, the yield of the *O*-alkylated isomer falls sharply on going from dipolar non-HBD solvents to protic solvents such as alcohols [372]. Even relatively small amounts of added protic solvents influence the mode of reaction. Thus, in the alkylation of the enolate salt of acetoacetic ester with ethyl tosylate in hexamethylphosphoric triamide, the addition of equimolar amounts of protic solvents such as water or *tert*-butanol significantly decreases the degree of *O*-alkylation [373]. High pressure promotes specific solvation of the more electronegative site in protic solvents, and increased alkylation occurs at the alternative position [378].

The tendency towards reaction at the center with the maximum electron density increases when dipolar non-HBD solvents are employed owing to the lack of specific solvation (*cf.* solvents $HCON(CH_3)_2$ and CH_3SOCH_3 in Table 5-22). Thus, in the alkylation of the enolate ions of 1,3-dicarbonyl compounds, the greatest yields of the *O*-alkylated isomers are obtained in hexamethylphosphoric triamide, followed by dipolar non-HBD solvents of the amide type [372–375].

In poorly cation-solvating solvents, there is also a marked dependence of reaction site on association with the corresponding cation. In non-dissociating solvents, it is likely that the cation will be preferentially coordinated to the atom in the ambident anion with the maximal electron density and this hinders reaction at this site. As shown in Table 5-22, by the results observed for the alkylation of sodium β-naphtholate, screening of the electronegative oxygen atom due to association with the accompanying sodium cation leads to an increase of *C*- *versus* *O*-alkylation in aprotic, non-dissociating solvents such as 1,2-dimethoxyethane ($\varepsilon_r = 7.2$) and tetrahydrofuran ($\varepsilon_r = 7.6$), compared with dipolar non-HBD EPD solvents. Alkylation of the less electronegative site can also be favoured by the addition of excess $M^{\oplus}$ to suppress dissociation of ambident ion pairs. The lithium enolates of 1,3-dicarbonyl compounds are an exception. They display a pronounced tendency towards *O*-alkylation (especially when compared to sodium enolates), which is not very sensitive to changes in the solvent (*e.g.* THF $\rightarrow$ DMF) [659]. It has been suggested that the tendency of lithium enolates towards *O*-alkylation is partly the result of an $Li^{\oplus}$ leaving group interaction and partly an intrinsic property of the enolate lithium ion pair [659].

The ion-pair dissociation of ambident alkali enolates, which results in increasing O/C alkylation ratios, can be promoted not only by dissociating solvents but also by specific cation solvation. In the latter case, EPD solvents (*cf.* DMF and DMSO in Table 5-22b) or macro(poly)cyclic ligands such as coronands ("crown ethers") or cryptands are used [376, 377, 660]. For example, the alkylation of sodium β-naphtholate with (bromomethyl)benzene or iodomethane in the presence of benzo[18]crown-6 gives high O/C alkylation ratios when tetrahydrofuran or benzene are the solvents [660]. In dissociating solvents such as *N*,*N*-dimethylformamide or acetonitrile, however, so far no

influence of added crown ether on the O/C product ratio has been observed [660]. The influence of ion-pairing on the alkylation of preformed alkali metal 2,4-pentanedionates in dimethyl sulfoxide solution has been systematically studied [661]. High pressure results in a shift from paired to free ambident anions and increased *O*-alkylation in non-HBD solvents [378]. The molar O/C methylation ratio for the reaction of lithium, sodium, and potassium acetophenone enolates with dimethyl sulfate has been studied in fourteen aprotic solvents and correlated with solvent polarity and Lewis basicity, as well as with the ionic radii of the alkali metal cations [847].

To summarize, it can be stated that the freer the ambident anion in every respect, the larger the O/C-alkylation ratio in the case of 1,3-dicarbonyl compounds [365]. Thus, if *O*-alkylation products are desired in the alkylation of enolates, dipolar non-HBD and dissociating solvents such as *N*,*N*-dimethylformamide, dimethyl sulfoxide, or, especially, hexamethylphosphoric triamide should be used. If *C*-alkylation is desired, protic solvents like water, fluorinated alcohols, or, in the case of phenols, the parent phenol will be the best choice [365].

Taking into account the fact that the solvation of ambident anions in the activated complex may differ considerably from that of the free anion, another explanation for the solvent effect on orientation, based on the concept of hard and soft acids and bases (HSAB) [275] (see also Section 3.3.2), seems preferable [366]. In ambident anions, the less electronegative and more polarizable donor atom is usually the softer base, whereas the more electronegative atom is a hard Lewis base. Thus, in enolate ions, the oxygen atom is hard and the carbon atom is soft, in the thiocyanate ion the nitrogen atom is hard and the sulfur atom is soft, *etc.* The mode of reaction can be predicted from the hardness or softness of the electrophile. In protic solvents, the two nucleophilic sites in the ambident anion must interact with two electrophiles, the protic solvent and the substrate RX, of which the protic solvent is a hard*) and RX a soft acid. Therefore, in protic solvents it is to be expected that the softer of the two nucleophilic atoms (C *versus* O, N *versus* O, S *versus* N) should react with the softer acid RX.

In non-HBD, non-dissociating solvents, a corresponding proposal can be made: hard counterions (alkali metal cations) should associate preferably with the hard site, and the substrate RX with the soft site in the activated complex composed of RX and the ambident ion pair [366]. With increasing hardness of the counterion (increasing charge density), the fraction of *C*-alkylation should increase in non-HBD solvents and decrease on solvent insertion into the ion pair. Indeed, the *C*-ethylation of $M^{\oplus}$ (ethyl acetoacetate)$^{\ominus}$ in dimethyl sulfoxide or hexamethylphosphoric triamide increases in the order $M^{\oplus} = R_4N^{\oplus} < Cs^{\oplus} < K^{\oplus} < Na^{\oplus} < Li^{\oplus}$ [373].

In the gas phase, the reaction of ethyl cations, $C_2H_5^{\oplus}$, with the ambident 2,4-pentanedione (which is 92% enolized at 25 °C in the gas phase) leads predominantly (>95%) to alkylation at the hard oxygen site and not at the soft carbon atom, as predicted by the HSAB concept [662]. Accordingly, the gas-phase alkylation of the enolate ion of cyclohexanone gives only the *O*- and no *C*-alkylation product [848], and the gas-phase acylation of acetophenone enolate with trifluoroacetylchloride leads predominantly to the *O*-acylation product (O/C ratio = 6.0) [849].

* Hydrogen atoms behave as hard Lewis acids in the formation of a hydrogen bond.

A theoretical explanation of the HSAB concept using quantum-chemical perturbation theory was given by Klopman [379]. He showed that the softness of a particular Lewis acid (acceptor atom) is determined by the energy of the lowest unoccupied molecular orbital and the softness of a Lewis base (donor atom) by the energy of the highest occupied molecular orbital, taking into account that the softness or hardness is influenced by the solvent. Limitations of the application of the HSAB concept to the reactivity of ambident anions have been discussed by Gompper *et al.* [367]. An alternative quantitative approach for the prediction of the dual reactivity of ambifunctional anions towards electrophiles has been proposed ("principle of allopolarization"), using so-called selectivity factors $S_f = Q_X/Q_Y$ (Q_X and Q_Y are the yields observed in reactions of ambident anions with two reaction sites X and Y) and polarity indices $P = l_X/l_Y$ (l_X and l_Y are the charge densities at reaction sites X and Y, taken from HMO calculations) as the defining parameters.

In analogy with ambident anions, mesomeric *ambident cations* do exist, but the solvent influence on their dual reactivity with nucleophiles has not been thoroughly investigated; see reference [368] for a review.

A nice example of the solvent-dependent dual reactivity of an electrophilic cryptocationic species has been given by Hünig *et al.* [663]. Ambident electrophilic α-enones react with nucleophiles such as the anion of the benzaldehyde O-(trimethylsilyl)-cyanohydrin ($Nu|^{\ominus}$) in diethyl ether exclusively by 1,4-addition. In tetrahydrofuran (THF) or 1,2-dimethoxyethane (DME), the 1,2-adduct is formed predominantly; on the addition of HMPT or [12]crown-4 it is formed exclusively; *cf.* Eq. (5-133).



(5-133)

This dramatic solvent effect is a result of the difference in the extent of ion-pairing of $Li^{\oplus}Nu^{\ominus}$ in diethyl ether (→ contact ion pair) and THF or DME (→ solvent-separated ion pair), modified by the α-enone-$Li^{\oplus}$ complex [663].

5.5.7 Solvent Effects on Mechanisms and Stereochemistry of Organic Reactions

Elaborating on the preceding dicussion on the dual reactivity of ambident anions, this Section contains a selection of reactions of various types which have in common the fact that a change in solvent drastically changes the mechanism as well as the stereochemistry usually associated therewith. The examples are intended to demonstrate how, out of two or more alternatives, a single pathway for a reaction can be favoured through a

proper choice of solvent. This, of course, is of considerable preparative interest. In this connection, it should be emphasized that alternative routes of a reaction may be separated by very small differences in Gibbs activation energy. It requires less than 12 kJ/mol (3 kcal/mol) of activation energy to change a 10:90 ratio to a 90:10 product mixture, and the sensitivity of this ratio can frequently be exploited by a proper choice of the reaction medium.

The stereochemical outcome of a nucleophilic substitution reaction at a saturated carbon is a function of the reaction mechanism (S_N1, S_N2, or S_Ni), which, in turn, can depend on the nature of the solvent, as already mentioned (*cf.* Sections 5.3.1 and 5.5.1). The fact that, in contrast to dipolar non-HBD solvents, the protic solvents diminish the nucleophilicity of anions and simultaneously favour the ionization of polarized bonds, makes it possible to displace a given reaction toward either the S_N1 or S_N2 type. A remarkable example is found in reaction (5-134) [380]. The nucleophilic substitution of cholesteryl tosylate in dipolar non-HBD solvents (*e.g.* $HCON(CH_3)_2$, CH_3SOCH_3) gives preferably the 3α-derivative with inversion of configuration. In contrast, in protic solvents (*e.g.* CH_3OH, $HCONHCH_3$), a mixture of the 3*β*- and 3,5-cyclo-6*β* derivatives is formed, corresponding to an S_N1 mechanism with initial formation of a homoallyl carbenium ion. However, a change from a protic to a dipolar non-HBD solvent never results in complete suppression of the S_N1 reaction [380].



(5-134)

An "S_N2 (intermediate)" mechanism has been proposed as a result of the consideration of the dependence of S_N1/S_N2 solvolysis rates on electrophilic (EPA) and nucleophilic (EPD) solvent assistance [664, 665].

S_N1 reactions of secondary and tertiary substrates $R_3C—X$ (*e.g.* 2-adamantyl tosylate [665], *t*-butyl heptafluorobutyrate [666]) proceed via activated complexes with high carbenium ion character to give ion-pair intermediates. They exhibit electrophilic solvent assistance in protic solvents SOH (H-bonding to the leaving group), but they are practically insensitive to changes in solvent nucleophilicity; *cf.* Eq. (5-135).



(5-135)

S_N2 reactions of primary and secondary substrates $R_3C—X$ are accelerated relative to S_N1 reactions by additional rearside nucleophilic attack on the substrate, which reduces the carbenium ion character in the activated complex. S_N2 reactions involving a concerted mechanism are also called "S_N2 (one-stage)" reactions [665]. They span a huge spectrum of nucleophilic solvent assistance – varying from classical S_N2 processes involving strong nucleophilic solvent assistance (*e.g.* methyl tosylate) to weakly nucleophilic solvent-assisted processes with relatively high carbenium ion character in the activated complex (*e.g.* 2-propyl tosylate). This variability of S_N2 solvolysis reactions can be regarded as a solvent-dependent spectrum of intermediate mechanisms rather than a varying mixture of only two distinct processes S_N1 and S_N2. To account for this gradation in mechanism, an "S_N2 (intermediate)" mechanism has been proposed [664, 665]. This mechanism involves a pentacoordinated intermediate, which is a nucleophilically solvated ion pair similar to the activated complex in the S_N2 (one-stage) reaction. The solvent/carbocation interaction is covalent in character; *cf.* Eq. (5-136).



(5-136)

The heterolysis of the bond between carbon and the leaving group in $R_3C—X$ is supported by nucleophilic solvent assistance, the extent of which depends on the solvent and increases in the order $(CF_3)_2CHOH < CF_3CO_2H < CF_3CH_2OH < HCO_2H < H_2O < CH_3CO_2H < CH_3OH < CH_3CH_2OH$ [665]. If there is evidence for an intermediate as well as for nucleophilic solvent assistance, then the mechanism can be considered "S_N2 (intermediate)". Examples showing a gradual change of mechanism from S_N2 (one-stage) through S_N2 (intermediate) to S_N1 have been given [665].

The "S_N2 (intermediate)" mechanism has not been without criticism; see, for example, [667, 668, 784] and references cited therein. An alternative mechanism can be formulated, involving ion pairs formed by heterolysis of the $R_3C—X$ bond *before* rate-limiting nucleophilic solvent attack occurs. Internal return to the educt would then occur more rapidly than attack by the nucleophile to give the products. For example, the solvolyses of secondary 1-arylethyl tosylates, $ArCH(OTs)CH_3$, can be also explained in terms of an ion-pair mechanism in which nucleophilic solvent attack on the ion pair plays a major role [667]. In less nucleophilic solvents, this attack is rate-limiting, whereas attack of more nucleophilic solvents is fast, resulting in a rate-limiting initial ionization of $R_3C—X$. The ion pair generally interacts with the solvent, although not by a specific interaction (*i.e.* covalent bonding with a single nucleophilic solvent molecule) as in the "S_N2 (intermediate)" mechanism [667].

It is difficult to objectively prove or disprove either the S_N2 (intermediate) or the ion-pair mechanism of S_N2 solvolysis reactions. According to Olah *et al.* [669], "the S_N2 intermediate can indeed be best characterized as a trivalent carbocationic centre solvated from both sides by the negatively charged nucleophile and the leaving group". According to Tidwell *et al.* [667], "this view deemphasizes the covalent character of the

solvent-carbocation interaction, and is operationally equivalent to an ion pair". A recent cluster model study of contact ion-pair formation of $Me_3C–Cl$ in aqueous solution has shown that its S_N1 hydrolysis presumably proceeds by an S_N2 (intermediate) mechanism, involving an electrophilically *and* nucleophilically solvated activated complex, subsequent formation of the electrophilically *and* nucleophilically solvated contact ion pair, and eventual formation of the products [850]; *cf.* Eq. (5-136).

The *ortho*/*para* alkylation ratio of the intramolecular nucleophilic substitution reaction of sodio-4-(3-hydroxyphenyl)butyl tosylate, according to Eq. (5-137), was shown to be solvent-dependent [381]. In general, an increase in the *ortho*/*para* alkylation ratio resulted when the polarity of the solvent was decreased. The orientation of the reaction was explained in terms of the nature of the association between the metal and phenolate ions in different solvents [381].



	para-product	*ortho*-product
in $(CH_2)_4O$:	13 cmol/mol	87 cmol/mol
in CH_3OH:	49 cmol/mol	51 cmol/mol

(5-137)

The α/γ alkylation ratio of the intramolecular nucleophilic substitution reaction of an α,β-unsaturated cyclohexenone tosylate can also be controlled by the right choice of solvent. The desired γ-alkylation – the final step of the total synthesis of the sesquiterpene β-vetivone – is favoured by NaOH in CH_3SOCH_3/H_2O, in contrast to $(CH_3)_3COK/(CH_3)_3COH$, which promotes α-alkylation [670].

That a single solvent molecule clustered to a nucleophile can drastically change the reaction pathway has been demonstrated by studying the reaction of phenyl acetate with methoxide ion in the gas phase [671, 672]. Alkaline hydrolysis of esters in solution is known to proceed by attack of the nucleophile at the carbonyl carbon atom to form a tetrahedral intermediate, followed by cleavage of the acyl–oxygen bond ($B_{AC}2$ mechanism); *cf.* Eq. (5-138a).

In the gas phase, however, nucleophilic aromatic substitution according to Eq. (5-138b) is the preferred pathway! Partially solvated nucleophiles such as $CH_3O^{\ominus}\ldots HOCH_3$, formed in the gas phase, again react with phenyl acetate by the $B_{AC}2$ mechanism. In the gas phase, charge-dispersed activated complexes such as those formed in the S_NAr reaction obviously have lower activation barriers than charge-localized activated complexes leading to the tetrahedral intermediate of the $B_{AC}2$ mechanism. The latter can only be stabilized by dispersing the localized negative charge over the carbonyl oxygen atom by means of H-bonding with protic solvent molecules. The attachment of even one HBD solvent molecule to the nucleophile is enough to change the reaction mechanism from S_NAr to $B_{AC}2$ [671].

The mechanism and stereochemical course of electrophilic substitution reactions at saturated carbon atoms (S_E1, S_E2, or S_Ei) may also be affected by the medium [337, 382, 383]. This is especially true for S_E reactions of organomercury compounds, as shown by Reutov *et al.* [384], Petrosyan [673], and Hughes and Ingold [385]. The isotopic exchange reaction of (α-ethoxycarbonyl)benzylmercuric bromide with radiomercuric



bromide in anhydrous dimethyl sulfoxide, shown in Eq. (5-139), is first-order in substrate but zeroth-order in Hg^*Br_2 ($Hg^* = {}^{203}Hg$) and proceeds with complete racemization of the optically active substrate [384, 385]. These observations are in agreement with an S_E1 mechanism. This was the first time that such a mechanism was established for organomercury compounds. The ionization of the C—Hg bond is facilitated by specific solvation of the leaving group, *i.e.* $(CH_3)_2S{=}O \rightarrow HgBr^{\oplus}$. However, when the



same reaction is carried out in aqueous acetone, aqueous ethanol, or in pyridine, second-order kinetics are followed and the reaction occurs with retention of configuration. The reaction pathway followed in these solvents is presumably of the S_E2 type. Subsequent to the work of Reutov [384] and Hughes and Ingold [385], solvent-induced alterations of mechanisms have been observed for other reactions of organometallic compounds, par-

Table 5-23. Stereoselectivity of electrophilic bromine addition to *cis*- and *trans*-stilbene, carried out at 0 °C in the dark [79, 386]; *cf.* Eqs. (5-29) and (5-140).

Solvents	cmol/mol *racemic* 1,2-dibromo-1,2-diphenylethane[a)]	
	from *cis*-stilbene	from *trans*-stilbene
CS_2	81.4	5.5
CCl_4	77.0	11.0
	67.4	–
C_2H_5OH	52.0	–
$CCl_3CO_2CH_3$	51.0	21.0
CCl_3CN	34.0	18.5
$C_6H_5NO_2$	29.5	16.5

[a)] The difference to 100 cmol/mol consists of *meso*-1,2-dibromo-1,2-diphenylethane.

ticularly other organomercury compounds [337, 338, 382, 383] as well as organotin compounds [674].

For electrophilic additions of halogens to alkenes, not only is the reaction rate strongly solvent-dependent [79–81] (*cf.* Eq. (5-29) in Section 5.3.2), but the stereochemical course may also be affected by the polarity of the medium [79, 386–388]. For example, the stereoselectivity of bromine addition to *cis*- and *trans*-stilbene according to Eq. (5-140) has been found to be solvent-dependent, as shown in Table 5-23 [79, 386].



(5-140)

Whereas in nonpolar solvents such as CS_2 and CCl_4, bromine adds to *cis*-stilbene in a highly stereoselective manner to give 81.4 cmol/mol *racemic* stilbene dibromide, and to *trans*-stilbene to give 94.5 cmol/mol *meso*-stilbene dibromide, in polar solvents, the degree of stereoselectivity is considerably reduced in both cases. Similar results have

been observed for bromine addition to 1-phenylpropene [387]. These results can be interpreted in terms of reaction scheme (5-140), which involves both ethene bromonium ions (symmetrical or unsymmetrical) and α-halogeno carbenium ions as intermediates. The observed solvent dependence of this polar bromination reaction may be the result of differential solvation requirements of the bromonium ion and the carbenium ion in their equilibria, leading to more carbenium ion character, as opposed to bromonium ion character, as the solvent polarity increases. In nonpolar solvents, the increased relative bromonium ion stability may be attributed to "internal solvation" of the carbenium ion by the neighbouring bromine atom. In polar solvents, however, the carbenium ion is more stabilized, enabling it to rotate about the C—C single bond (A ⇌ B), particularly at higher temperatures. In other words, the cyclic halonium ion becomes relatively more stable than the carbenium ion as the solvent becomes less capable of nucleophilic solvation. The observation that there is an increase in stereoselectivity for electrophilic alkene halogenation in nonpolar solvents at low temperatures may serve as a qualitative example of solvent and temperature effects on halonium ⇌ carbenium ion equilibria. These effects have been quantitatively determined by McManus and Peterson [388]. The fact that in polar solvents *racemic* and *meso*forms are not obtained in equal amounts, although the reaction seems to proceed through an α-halogeno carbenium ion in these solvents, is simply explained by the different conformational energies of A and B. The conformer B in scheme (5-140), leading to *meso*form, is sterically more favoured than A. Generally, solvent polarity plays a significant role in determining the stereochemistry of addition reactions only in borderline cases such as stilbene, where there is stabilization of the carbenium ion by the neighbouring phenyl group. Halogen addition to other alkenes, such as diethyl fumarate and diethyl maleate, also exhibits high stereoselectivity in polar solvents [79].

Examples of the solvent-dependent competition between nucleophilic substitution and β-elimination reactions (*i.e.* S_N1 versus E_1 and S_N2 versus E_2) have already been given in Section 5.3.1 (*cf.* Table 5-7). A nice example of a dichotomic β-elimination reaction, which can proceed via an E_1 or E_2 mechanism depending on the solvent used, is shown in Eq. (5-140a); *cf.* also Eqs. (5-20) and (5-21) in Section 5.3.1. The thermolysis of the potassium salt of racemic 2,3-dibromo-1-phenylpropanoic acid (**A**), prepared by bromine addition to (*E*)-cinnamic acid, yields, in polar solvents (*e.g.* water), apart from carbon dioxide and potassium bromide, the (*E*)-isomer of 1-bromo-2-phenylethene, while in solvents with low or intermediate polarity (*e.g.* butanone) it yields the (*Z*)-isomer [851].



(5-140a)

In water, ionization of the C–Br bond occurs first (E_1 mechanism) to give the intermediate resonance-stabilized benzylic zwitterion **C**. After fast rotation about the C–C bond, carbon dioxide leaves conformer **D** perpendicularly to the plane of the carbenium ion, to give mainly the most stable (*E*)-isomer of *β*-bromostyrene. In butanone, after fast rotation about the C–C bond, elimination of CO_2 and Br^- occurs in a concerted single-step (E_2 mechanism) for stereoelectronic reasons (Br and CO_2^- must be *anti* to one another) to give conformer **B**, which decomposes exclusively to the thermodynamically less stable (*Z*)-isomer. In more polar solvents, the partly zwitterionic activated complex, leading to zwitterion **C** in the rate-determining step, will clearly be more stabilized by solvation than the less dipolar activated complex leading directly to the (*Z*)-isomer of *β*-bromostyrene from conformer **B** [851].

In cycloaddition and cycloreversion reactions, the mechanism and product distribution may also be strongly influenced by the reaction medium. Two examples have already been briefly mentioned in Sections 5.3.2 [102] and 5.3.3 [124]. In the reaction of dimethylketene with enamines such as *N*-isobutenylpyrrolidine, a two-step process *via* a zwitterionic intermediate, leading preferably to a *δ*-methylene *δ*-lactone as 2:1 adduct, competes with a concerted mechanism to give a cyclobutanone derivative according to Eq. (5-141) [102]. In cyclohexane, 92% of the enamine reacts concertedly to give the cyclobutanone derivative, whereas 8% goes through the zwitterionic intermediate. The path *via* the zwitterion increases in importance with increasing solvent polarity and reaches 57% in acetonitrile [102].

An interesting example of the solvent-dependent cycloreversion of a substituted $3H$-Δ^1-pyrazoline (formed by the 1,3-dipolar cycloaddition of diazomethane to an allylic bromide) has been given by Kolsaker *et al.* [675].

The stereoselectivity of the Diels–Alder reaction between methyl acrylate and cyclopentadiene depends to some extent on solvent polarity [124]; *cf.* Eq. (5-43) in Section 5.3.3. The *endo*/*exo* ratio for the cycloaddition product has been found to vary from 74:26 cmol/mol in triethylamine to 88:12 cmol/mol in methanol at about 30 °C. It has



(5-141)

Solvent	*c*-C_6H_{12}	C_6H_6	C_6H_5Cl	$CHCl_3$	CH_3COCH_3	CH_3CN
%concerted reaction	92.0	79.4	72.8	62.3	52.1	43.0

been observed that for the same reaction not only the *exo*/*endo* stereoisomer ratio is affected by solvent polarity, but also the product ratio of *exo* (or *endo*) adduct to *endo*-dicyclopentadiene [389]. The latter is formed as a by-product in the Diels–Alder reaction. As expected, the *exo* (or *endo*) adduct/*endo*-dicyclopentadiene ratio increases with increasing solvent polarity [389]. The rate and regioselectivity of Diels–Alder reactions with inverse electron demand between the ketene aminal $CH_2{=}C(NMe_2)_2$ and 3-phenyl-1,2,4,5-tetrazine as well as substituted 1,2,4-triazines is strongly solvent-dependent [676]. The "*ortho*"/"*meta*" product ratio of regioisomers decreases with increasing solvent polarity in non-HBD solvents [676].

Examples of the solvent-influenced competition between concerted [4 + 2]Diels–Alder type cycloaddition reactions and 1,4-dipolar reaction pathways with zwitterionic intermediates can be found in references [677–679]. For example, in solvents of low polarity ($CHCl_3$, CH_2Cl_2), homofuran reacts with tetracyanoethene to form the seven-membered [4 + 2]cycloadduct **A** in quantitative yield. In solvents of high polarity (CH_3CN), however, the [2 + 2]cycloadduct **B** predominates, formed *via* a 1,4-dipolar activated complex and a zwitterionic intermediate [679]; *cf.* Eq. (5-142).



(5-142)

Appreciable solvent effects are also obtained in photochemically induced [2 + 2]-cycloadditions and cycloreversions [390–392]. Examples are given in Eqs. (5-143) [390] and (5-144) [680].

The [2 + 2]photodimerization of 2-cyclopenten-1-one leads to the regioisomeric cyclobutane derivatives shown in Eq. (5-143) [390]. The *anti*/*syn* product ratio decreases as the solvent polarity increases. This is due to the more dipolar, and hence more strongly solvated activated complex being favoured, which leads to the dipolar *syn*-cycloadduct.

When 1,2-bis(methoxycarbonyl)-3,4-bis(2-naphthyl)cyclobutane is irradiated in solution (cyclohexane/tetrahydrofuran and tetrahydrofuran/acetonitrile mixtures) in the presence of triethylamine, it undergoes [2 + 2]cycloreversion *via* an exciplex with triethylamine [680]. Interestingly, as shown in Eq. (5-144), the mode of cycloreversion changes with the solvent polarity, reflecting the different dipolar electronic structure of the exciplex. In nonpolar solvent mixtures, a 'horizontal' cleavage of the cyclobutane



(5-143)

Solvent	C_6H_6	C_6H_5Cl	CH_2Cl_2	CH_3COCH_3	CH_3CN
$\lg \frac{[anti]}{[syn]}$	0.61	0.39	0.28	0.16	−0.14

ring takes place, in highly polar solvent mixtures a 'vertical' one. It has been assumed that in the latter case the intermediate exciplex exhibits increased charge-transfer character [680].



(5-144)

Photo sensitized oxygenation of alkenes with singlet oxygen can, in principle, proceed *via* three competitive reaction pathways: [4 + 2]cycloaddition to *endo*-peroxides, ene reaction of allylic hydroperoxides, and [2 + 2]cycloaddition to 1,2-dioxetanes (see reference [681] for a review). With suitable olefinic substrates, the chemical outcome of such photo-oxygenation reactions can be strongly influenced by the solvent. This is shown in the somewhat simplified Eq. (5-145).

Since [4 + 2]cycloaddition and ene reactions are generally assumed to proceed in a concerted manner *via* isopolar activated complexes, they should exhibit virtually the same small, often negligible, response to changes in solvent polarity. This is what, in fact, has been found; *cf.* for example [138, 682, 683]. However, two-step [2 + 2]-cycloaddition reactions of singlet oxygen to suitably substituted electron-rich alkenes proceed *via* dipolar activated complexes to zwitterionic intermediates (1,4-dipoles or perepoxides). In this case, the relative amounts of 1,2-dioxetane and allylic hydroperoxides or *endo*-peroxides should vary markedly with solvent polarity if two or even all three of the reaction pathways shown in Eq. (5-145) are operative [681, 683, 684].

In the photo-oxygenation of enol ethers, where the ene reaction and [2 + 2]-cycloaddition compete, polar solvents favour cycloaddition whereas nonpolar solvents favour ene product formation [681, 683–685]. For example, 2,3-dihydro-4-methyl-4*H*-pyran reacts with singlet oxygen to yield both a 1,2-dioxetane and an allylic hydro-



(5-145)

peroxide as primary products in nonpolar cyclohexane and dipolar acetonitrile, with product ratios of 13:87 and 84:16 cmol/mol, respectively, as shown in Eq. (5-146) [685].



(5-146)

Further evidence in support of zwitterionic intermediates in the [2 + 2]-cycloaddition of singlet oxygen to electron-rich alkenes has been obtained by Jefford *et al.* [684]. The photo-oxygenation of 2-(methoxymethylidene)adamantane creates a zwitterionic intermediate (peroxide or perepoxide), which can be captured by acetaldehyde to give 1,2,4-trioxanes in addition to 1,2-dioxetanes; *cf.* Eq. (5-147).

The ease of capture itself is also solvent-dependent: polar protic solvents (methanol, ethanol) stabilize the zwitterion by H-bonding, thus rendering it unreactive towards external electrophiles such as acetaldehyde, and facilitating its closure to a 1,2-dioxetane. On the other hand, nonpolar solvents (toluene, tetrahydrofuran, ethyl acetate) prevent the dispersion of charge, thus enhancing the reactivity towards aldehydes and favouring 1,2,4-trioxane formation [684]. The zwitterionic intermediate in Eq. (5-147) can be formulated as either peroxide or perepoxide (peroxirane). A distinction between these two species is difficult to make. In both cases, the positive charge is stabilized as an oxonium ion while the negative charge remains localized on the terminal oxygen atom. According to calculations, a peroxide would have a permanent dipole moment of *ca.* $34 \cdot 10^{-30}$ Cm (10.1 D) and a perepoxide a dipole moment of *ca.* $19 \cdot 10^{-30}$ Cm (5.6 D) [686].



(5-147)

Further examples of solvent effects on photo-oxygenation reactions with singlet oxygen can be found in references [687–689].

Similar solvent effects can be found in fragmentation reactions [109]. Such reactions may proceed heterolytically or homolytically depending on the solvent used as reaction medium. Thus, the observed solvent effect for the thermolysis of 4-CH_3–C_6H_4–N=N–S–C_6H_4–4-$C(CH_3)_3$ indicates that this benzenediazoaryl sulfane decomposes homolytically in apolar non-HBD solvents according to Eq. (5-148a) [393].

The marked increase in rate observed in protic solvents such as acetic acid and aqueous ethanol, however, may be due to heterolytic decomposition according to Eq. (5-148b). Suitably substituted benzenediazoaryl sulfanes such as 4-CH_3–C_6H_4–N=N–S–C_6H_4–4-NO_2 should be more inclined to decompose heterolytically because of mesomeric stabilization of the diazonium and thiolate ions formed. Indeed, the thermolysis of this compound is extremely sensitive to solvent polarity, heterolysis presumably being predominant in polar solvents [393]. It was subsequently found that thermolysis of (*Z*)-benzenediazoalkyl sulfanes leads to isomerization to the (*E*)-form as well as to competing decomposition [394]. Only the decomposition is strongly accelerated with increasing solvent polarity, whereas the rate of (*Z*)/(*E*) isomerization is almost independent of the solvent.



(5-148a)

(5-148b)

Solvent	*i*-C_8H_{18}	C_6H_6	C_5H_5N	CH_3SOCH_3	CH_3CO_2H	C_2H_5OH/H_2O (96:4 cL/L)
$k_1^{rel\ a)}$	0.32	1	0.47	0.91	47	44
$k_1^{rel\ b)}$	0.60	1	25	100	45	190

a) 4-CH_3–C_6H_4–N=N–S–C_6H_4–4-$C(CH_3)_3$
b) 4-CH_3O–C_6H_4–N=N–S–C_6H_4–4-NO_2

According to Zollinger *et al.* [690], depending on the solvent used, the thermal dediazoniation of arenediazonium ions in solution can proceed *via* competitive heterolytic and homolytic reaction pathways; *cf.* the somewhat simplified Eq. (5-149).

(5-149)

In solvents of low nucleophilicity (*e.g.* $(CF_3)_2CHOH$, CF_3CH_2OH, CH_3CO_2H, H_2O with $pH < 1$), the familiar heterolytic dediazoniation takes place to give an aryl cation and products derived therefrom. The solvent effect on heterolytic dediazoniation rates is small, the slowest (in 1,4-dioxane) and fastest rate (in 2-propanol) differing only by a factor of 9 [690].

However, in solvents of high nucleophilicity (*e.g.* DMSO, HMPT, pyridine), the formation of products derived from aryl radicals is favoured. Therefore, in these solvents, a homolytic dediazoniation must be taking place. This mechanism involves addition of a nucleophilic solvent molecule to the electrophilic β-nitrogen atom of the diazonio group, followed by homolysis to give a radical pair and nitrogen. The homolytic dediazoniation is not only enhanced by sufficiently nucleophilic solvent molecules but also by the addition of other nucleophiles, which form relatively stable radicals through electron transfer. Under comparable reaction conditions, the rates of homolytic dediazoniation are higher than the corresponding rates of heterolysis [690].

Dichotomous homolytic and heterolytic decomposition reactions can also be found in the thermolysis of peroxycarboxylic esters [195, 207–209]; *cf.* Eq. (5-62) and

The dissociation equilibrium of associated $Li^{\oplus}AlH_4^{\ominus}$ ion pairs in a variety of ethereal solvents has been investigated by ^7Li and ^{27}Al NMR spectroscopies [768]. $\delta(^{27}Al)$ is nearly independent of the solvent as is $^1J_{Al-H}$, whereas $\delta(^7Li)$ depends on the $LiAlH_4$ concentration and the EPD character of the ethereal solvent. According to this result, diglyme seems to be the solvent of choice for reactions in which the $AlH_4^{\ominus}$ should be least affected by ion-pair formation [768].

Another interesting example of solvent-dependent control of chemoselectivity is the reaction of a bromo α-enone with an organocopper reagent, as shown in Eq. (5-154) [698]. The lithium dimethylcuprate displacement reaction of haloalkanes and the conjugate addition of cuprate reagents to α-enones exhibit opposite responses on the addition of EPD solvents. An appropriate choice of substrate and reaction medium should therefore make it possible to select either the $LiCuMe_2$ coupling reaction with a haloalkane or its conjugate addition.



(5-154)

Indeed, in diethyl ether, lithium dimethylcuprate usually reacts with the α-enone group to give a methyl-substituted bromo ketone. Addition of hexamethylphosphoric triamide (HMPT), however, slows down this reaction to such an extent that displacement of the bromo substituent takes place [698]. Another remarkable example of the influence of HMPT on chemoselectivity is the reaction of an arsonium ylide, $Ph_3As{=}CH{-}CH{=}CH{-}Ph$, with benzaldehyde in tetrahydrofuran solution, yielding either an epoxide (in THF) or an alkene (in THF/HMPT) [699].

Not only chemoselectivity but also *enantioselectivity* and *diastereoselectivity* of organic reactions can be controlled by choice of solvent; for reviews on asymmetric induction, see references [700, 701].

Some examples of enantioselective syntheses, carried out in chiral media, have already been given at the end of Section 3.2, which deals with chiral solvents; *cf.* also Table A-2 (Appendix). In general, asymmetric inductions as a result of chiral solvents or chiral cosolvents are disappointingly low [700]. The reason for this is that the differential solvation by the chiral solvent of the two enantiomorphic activated complexes which lead to either the (*R*)- or (*S*)-product is not sufficient. That is, the difference in Gibbs energy of activation, $\Delta\Delta G^{\neq} = \Delta G^{\neq}_{(R)} - \Delta G^{\neq}_{(S)}$, is not large enough to favour only one of the two enantiomeric products. It should be remembered that a difference of $\Delta\Delta G^{\neq} = 10.8$ kJ/mol (2.6 kcal/mol) at 20 °C would be sufficient to get an (*S*)/(*R*) product ratio of 99:1 (*i.e.* an enantiomeric excess $ee = 98\%$).

Therefore, a more direct covalent linking of the chiral information with the reactants is necessary in order to obtain diastereomorphic activated complexes with greater differential solvation, thus leading to better diastereoselectivity.

The Grignard reaction of benzaldehyde with ethylmagnesium bromide in diethyl ether in the presence of molar amounts of (*R*, *R*)-TADDOL as a chiral additive leads predominantly to (*S*)-1-phenyl-1-propanol, while the same reaction carried out in tetrahydrofuran gives mostly the corresponding (*R*)-alcohol according to Eq. (5-154a) [852]. TADDOL is the acronym for *t*etra*a*ryl-1,3-*d*ioxolane-4,5-*d*imethan*ol*, which reacts with



(5-154a)

the excess Grignard reagent to presumably generate a chiral magnesium alcoholate, Et–Mg–OR*. This then adds to the carbonyl group of benzaldehyde either at the *re*- or the *si*-side, depending on the ethereal EPD solvent used, which is also part of the solvation shell surrounding the magnesium ion. Otherwise, for a variety of other carbonyl compounds and Grignard reagents, this TADDOL-mediated addition reaction invariably leads to the (*R*)-alcohol with high enantioselectivity in tetrahydrofuran as solvent [852].

Further detailed investigations of the solvation of carbonyl compounds as the predominant factor influencing the diastereoselectivity of nucleophilic additions of various organometallics (*n*-BuLi, *t*-BuLi, *n*-BuMgBr) to α-R_3SiO-substituted aldehydes can be found in reference [853].

The Grignard reaction of (±)-3-phenylbutanone with phenylmagnesium bromide to give stereoisomeric 2,3-diphenylbutan-2-ols, shown in Eq. (5-155), represents another example [702].



(5-155)

Solvent	Et_3N	Et_2O	1,4-dioxane	$(CH_2)_4O$	diglyme	$(MeOCH_2)_2$
%[(*S*, *R*) + (*R*, *S*)]	26	36	49	61	66	73

In this particular case, the observed diastereoselectivity, $ds = \%[(S,R) + (R,S)]$, increases with increasing solvent polarity. The (SR,RS)-carbinol is preferentially formed in more polar solvents, whereas the (RR,SS)-carbinol dominates in less polar solvents. This result can be understood by a careful analysis of the two diastereomorphic activated complexes in terms of steric and polar effects. The activated complex which leads to the (SR,RS)-carbinol appears to be more dipolar and hence more strongly solvated; see reference [702] for details.

Another nice example is the addition of lithium dimethylcuprate to a chiral oxazolidine (prepared from (E)-cinnamaldehyde and $(-)$-ephedrine), followed by hydrolysis, to give 3-phenylbutanal; *cf.* Eq. (5-156) [703].



(5-156)

Diastereoselective addition in Et_2O/HMPT (1:1) leads to the (S)-aldehyde with an enantiomeric excess of 40%, whereas in *n*-hexane the (R)-aldehyde is formed with 80% (!) enantiomeric excess. The (R)-configured aldehyde is also obtained in benzene and in dichloromethane, but with lower *ee* values of 50% and 25%, respectively. Inverse results were obtained with a chiral oxazolidine prepared from (E)-cinnamaldehyde and $(+)$-ephedrine. Here, the (S)-aldehyde with $ee = 79\%$ is formed in *n*-hexane, and the (R)-aldehyde in Et_2O/HMPT (1:1) with $ee = 50\%$ [703]. This result may be due to different structures of the organocopper reagent, and hence of the diastereomorphic activated complexes, in nonpolar solvents (*n*-hexane, benzene, dichloromethane) and in EPD solvents (Et_2O/HMPT) [703].

A striking solvent effect was observed in the reduction of a chiral α-keto amide, C_6H_5–CO–CO–NR_2 ($NR_2 = (S)$-proline methyl ester), with sodium tetrahydridoborate, leading to mandelic acid after hydrolysis [704]. When the α-keto amide was reduced in pure tetrahydrofuran or methanol, the resulting enantiomeric excess of (S)-mandelic acid produced was 36% and 4%, respectively. However, when a tetrahydrofuran/methanol (99:1 cL/L) solvent mixture was used, the enantiomeric excess increased to 64% (!). In other solvent mixtures, a catalytic amount of a protic solvent (CH_3OH or H_2O) was found to be necessary for good asymmetric induction [704].

Extremely diastereoselective solvent-influenced alkylations were obtained with propionates of chiral alcohols derived from (+)-camphor, as shown in Eq. (5-157) [705].

Metallation of the propionate with lithium cyclohexylisopropylamide (LiCA) in a solvent-controlled reaction gives either the (Z)- (in THF) or (E)-lithium enolate (in THF/HMPT, 4:1). Rearside shielding of both diastereomeric enolates, provided by the 3,5-dimethylphenyl group attached to the sulfonamide moiety at C-2, prevents alkylation of these enolates from the backface. Hence, frontface attack with iodo-*n*-tetradecane leads to diastereomeric esters with either (R)- or (S)-configuration at the α-carbon atom of the propionate group. Using LiCA/HMPT mixtures with increasing

HMPT molar ratio in the metallation step leads to a smooth changeover from diastereoselective formation of the (*R*)-ester to diastereoselective formation of the (*S*)-ester. The turning point is at an LiCA/HMPT molar ratio of 1:1. Obviously, this configurational conversion is caused by the kinetically controlled formation of the diastereomeric enolates by deprotonation with either LiCA or the LiCA/HMPT complex. The EPD solvent HMPT acts as a lithium-solvating agent in this reaction, causing preferential formation of the (*E*)-lithium enolate. An explanation for this dramatic solvent effect has been given by analysis of the steric requirements for the enolization of esters [706]. The better solvation of $Li^{\oplus}$ in the presence of HMPT and the enhanced reactivity of the amide base favours the activated complex leading to the (*E*)-enolate. In the less coordinating solvent THF, the interaction between $Li^{\oplus}$ and the ester carbonyl oxygen atom must be strong, thus favouring the activated complex leading to the (*Z*)-enolate; see reference [706] for details. Addition of cation-solvating HMPT *after* the deprotonation step in reaction (5-157) also leads to an increase in diastereoselectivity, *i.e.* preferential formation of the (*R*)-ester. This solvent effect may be caused by disruption of conformationally unfavourable interactions between $Li^{\oplus}$ and the 3,5-dimethylphenyl group of the sulfonamide moiety; *cf.* Eq. (5-157) [705].



(5-157)

Solvent-controlled diastereoselectivities have also been observed in Diels–Alder cycloaddition reactions of cyclopentadiene with bis-(−)-menthyl fumarate [707] and with the acrylate of (*S*)-ethyl lactate, $CH_2{=}CH{-}CO{-}OCH(CH_3){-}CO_2Et$ [708]. In the latter reaction, giving four diastereomeric cycloadducts, diastereoselectivities of up to 85:15 have been obtained in *n*-hexane [708]. The diastereoselectivities decrease with increasing solvent polarity, while the *endo*/*exo* selectivity increases. This is in agreement with the pattern found for simple achiral acrylates [124]; *cf.* Eq. (5-43) in Section 5.3.3.

We shall conclude this section with reference to the (*E*)/(*Z*) isomerization of imino compounds $R_2C{=}N{-}R'$, which, in principle, may proceed by a rotation mech-

anism or an inversion mechanism. Determination of the solvent dependence of this isomerization has been used in deciding between the two possible reaction mechanisms [397, 398]. The small solvent effects usually observed are in agreement with an inversion mechanism [398]. An inversion mechanism was also postulated for the $(E)/(Z)$ isomerization of push-pull substituted azobenzenes [529, 561]; *cf.* Eq. (5-40) in Section 5.3.2.

Only a few representative, particularly interesting examples of solvent effects on the mechanisms and stereochemistry of organic reactions have been mentioned in this Section. The significance of these, often very specific, solvent effects is well-recognized by organic chemists, but a detailed understanding of them is frequently lacking and hence they are the subject of ongoing research.

5.5.8 Influence of Micellar and Solvophobic Interactions on Reaction Rates and Mechanisms

The modification of chemical reactions through the incorporation of the reactant molecules into aqueous micelles or other organized assemblies has received considerable attention in recent years. Reactions are known for which rates, mechanisms, and even the stereochemistry have been significantly affected by the addition of so-called *amphiphiles* to the reaction medium.

Solvophobic and especially hydrophobic interactions provide the driving force for the aggregation of organic ions known as amphiphiles in dilute aqueous solutions (*cf.* Sections 2.2.7 and 2.5) [399]. Amphiphilic ions possessing long unbranched hydrocarbon chains have both pronounced hydrophobic and hydrophilic properties and exhibit, therefore, the important property of forming, over a narrow concentration range termed the *critical micelle concentration* (*cmc*), molecular aggregates in solution, called *micelles* (*cf.* Fig. 2-12 in Section 2.5)*). These micelles, rather than individual amphiphilic ions, may cause alterations of rates and mechanisms of organic reactions in aqueous solutions of surfactants. A suitable choice of surfactant can lead to rate increases of 5- to 1000-fold compared to the same reaction in the absence of surfactant. The catalysis of organic reactions by ionic micelles can be explained in terms of electrostatic and hydrophobic interactions of the reactants and activated complexes with the micelle. The reaction substrate is partitioned between micellar and bulk aqueous phases by hydrophobic binding of the substrate to a micelle. Then, by simple electrostatics, this complex either attracts (rate acceleration) or repels (rate retardation) an incoming ionic reactant. The field of micellar catalysis has been the subject of comprehensive reviews [289, 400–403, 711–713]. Therefore in this section, only a few typical examples shall demonstrate the influence of hydrophobic interactions on organic reactions.

According to Eq. (5-158), the reaction of 1-fluoro-2,4-dinitrobenzene with phenolate or thiophenolate ions is accelerated by micelles of cetyl-trimethylammonium bromide (CTABr) in aqueous solution by factors of 230 and 1100, respectively [404].

---

* Depending on their chemical structure, surfactants capable of forming micelles are usually classified into cationic (*e.g.* ammonium salts), anionic (*e.g.* sulfates, carboxylates), ampholytic (*e.g.* zwitterionic salts), and non-ionic surfactants (usually containing polyoxyethene chains); *cf.* Table 2-10 in Section 2.5.

This enhancement of the reaction of bulk-phase anions $C_6H_5Y^{\ominus}$ with the organic nonelectrolyte 2,4-$(NO_2)_2C_6H_3F$, the latter partitioned between the bulk and micellar phases, is expected in the presence of cationic amphiphiles such as CTABr from purely



(5-158)

Surfactant	no CTABr added	with cationic CTABr
$k_2^{rel}(Y = O)$	1	230
$k_2^{rel}(Y = S)$	1	1100

electrostatic considerations. Amphiphiles with opposite charge to that of the reactant ion accelerate, whereas amphiphiles of like charge inhibit the reactions of neutral substrates. The catalytic efficiency increases with increasing diffusion of neutral substrate into the micelles, that is, with increasing lipophilicity of the reactant molecules. A compilation of micellar catalytic effects expected for organic reactions of different charge type in the presence of cationic or anionic micelles can be found in reference [289].

In contrast to reaction (5-158), addition of micelle-forming surfactants can also slow down chemical reactions. For example, the spontaneous hydrolysis of phenyl chloroformate at 25 °C according to Eq. (5-158a) is retarded by a factor of *ca.* 16 on addition of sodium dodecylsulfate (SDS) above the *cmc*. Below the *cmc*, the SDS

(5-158a)

monomers do not have a significant effect on the observed rate constant [855]. The hydrolysis of phenyl chloroformate follows an addition/elimination pathway, the addition of water being rate-determining. The observed micellar effect can be attributed to the lower polarity of the micellar interface at which the reaction takes place and to the electrostatic destabilization of the zwitterionic activated complex by the negatively charged head groups ($-SO_3^-$) of the surfactant.

Even at reactant concentrations well below critical micelle concentrations, hydrophobic interactions may result in marked rate accelerations. Thus, the rates of bimolecular aminolysis of 4-nitrophenyl decanoate and acetate by *n*-decylamine and ethylamine have been determined in aqueous solution and a distinct rate enhancement in the aminolysis of the long chain ester by the long chain amine was reported [405, 406]. As shown in Table 5-24, the ratio $k_2^{decylamine}/k_2^{ethylamine}$ for 4-nitrophenyl decanoate is 317*). That is, *n*-decylamine attacks the long-chain ester 47 times faster than expected on the

* Even larger rate increases (up to 10^7-fold) were observed for the corresponding amine-catalyzed termolecular reaction of 4-nitrophenyl ester with alkylamines, carried out in water or aqueous ethanol at 25 °C [406].

Table 5-24. Second-order rate constants for the aminolysis of 4-nitrophenyl esters in water/acetone (99:1) at 35 °C according to $CH_3(CH_2)_nCO–OC_6H_4–4\text{-}NO_2 + C_2H_5(CH_2)_nNH_2 \rightarrow CH_3(CH_2)_nCO–NH(CH_2)_nC_2H_5 + HOC_6H_4–4\text{-}NO_2$, with $n = 0$ and 8 [405].

Ester	$k_2^{\text{ethylamine}}$	$k_2^{\text{decylamine}}$	$k_2^{\text{decylamine}}/k_2^{\text{ethylamine}}$
4-Nitrophenyl acetate	5.74	39.1	6.8
4-Nitrophenyl decanoate	0.42	133	317
$k_2^{\text{decanoate}}/k_2^{\text{acetate}}$	0.073	3.4	47

basis of its reactivity toward 4-nitrophenyl acetate. This rise in the observed rate constant is best explained by the association of the reactant molecules to form a kind of "micro-micelle" prior to reaction. This is due to hydrophobic interactions between the long hydrocarbon chains of both reactants in aqueous solution. In accordance with this interpretation, the ratios $k_2^{\text{decylamine}}/k_2^{\text{ethylamine}}$ fall to values near unity for both esters when the reactions are carried out in aqueous 1,4-dioxane (50:50), a medium in which hydrophobic interactions are seriously disrupted [405].

Another remarkable example is the hydrolysis of the amide ($H_{25}C_{12}–NMe_2^+–CH_2–CO–NH–C_6H_4–4\text{-}NO_2)Br^-$ at 25 °C, which is accelerated about 10^8-fold on addition of sodium hexadecanoate as compared to the addition of sodium acetate [856]. This dramatic rate increase under very mild reaction conditions can again be attributed to hydrophobic association between the long-chain amide and the long-chain catalyst, as well as to the electrostatic attraction between the cationic ammonium centre of the reactant and the anionic carboxylate group of the catalyst.

Diels–Alder cycloaddition reactions have undergone impressive improvements, taking advantage of hydrophobic interactions existing between the essentially nonpolar reactants in the aqueous medium. The use of water as a solvent in Diels–Alder reactions leads to greatly enhanced reaction rates and selectivities. This remarkable result has been pioneered by Breslow *et al.* [801] and further explored by Grieco *et al.* [714]; for reviews, see references [715–718].

For example, the rate of the Diels–Alder cycloaddition reaction between 9-(hydroxymethyl)anthracene and *N*-ethylmaleimide, as shown in Eq. (5-159), is only slightly altered on changing the solvent from dipolar acetonitrile to nonpolar isooctane, as expected for an isopolar transition state reaction; *cf.* Section 5.3.3. In water, however,



(5-159)

Solvent	*i*-C_8H_{18}	*n*-C_4H_9OH	CH_3OH	CH_3CN	H_2O	H_2O + 4.86 M LiCl
k_2^{rel}	7.4	6.2	3.2	1	211	528

the reaction is 211 times faster than in acetonitrile! The addition of lithium chloride, known as a solute that increases hydrophobic effects, increases the rate in water by a

further 2.5-fold. This exceptional behaviour of the solvent water is best explained in terms of hydrophobic interactions, which promote the association of the diene and the dienophile during the activation process [801]. For some Diels–Alder reactions of cyclopentadiene, which can give both *endo* and *exo* addition of dienophiles, the *endo*/*exo* product ratio increases when water is used as solvent [801]. This is related to the well known influence of polar solvents on such ratios [124] as well as to the need to minimize the activated complex surface area in aqueous solution.

The Diels–Alder cycloaddition reaction of 2,6-dimethyl-1,4-benzoquinone with methyl (*E*)-3,5-hexadienoate, carried out in toluene as solvent, gives only traces of the cycloadduct shown in Eq. (5-160), even after seven days. However, when the solvent is changed to water and sodium (*E*)-3,5-hexadienoate is used as the diene, 77 cmol/mol of the desired cycloadduct is obtained after one hour and esterification with diazomethane [714]*). Again, hydrophobic interactions between diene and dienophile in the aqueous medium seem to be responsible for this remarkable and synthetically useful rate acceleration.



(5-160)

A sugar-assisted solubilization of the diene in aqueous Diels–Alder reactions has been proposed, using glucose as the hydrophilic moiety [744]. Based on the aqueous Diels–Alder cycloaddition of cyclopentadiene with diethyl fumarate, an increase in reaction rate with the increasing solvophobicity of the medium, characterized by Abraham's solvophobicity parameter *Sp* (see Section 7.3), has been found [745]. Addition of β-cyclodextrin to the reaction medium further promotes this cycloaddition. Increasing medium solvophobicity and the addition of β-cyclodextrin also influences the diastereoselectivity (*i.e.* the *exo*/*endo* product ratio) of this Diels–Alder reaction: the amount of *endo* product increases on addition of β-cyclodextrin and with increasing medium solvophibicity, due to the more compact activated complex leading to the *endo* adduct [745].

An interesting phenomenon related to the acceleration of Diels–Alder reactions in aqueous media is found when clays such as montmorillonite are suspended in an organic solvent (*e.g.* ethanol or dichloromethane) [719]. The layered structure of the clay enables it to trap pools of internal water, which are obviously capable of exerting hydrophobic effects.

It should also be mentioned in this context that the pressure dependence of the rate of Diels–Alder cycloaddition reactions is much less in water than in nonaqueous

* Under the conditions of the aqueous Diels–Alder reaction, the initially formed *cis*-cycloadduct equilibrates to the more stable *trans*-fused ring system.

solvents, mainly because of hydrophobic interactions of the apolar reactants. A careful study of the cycloaddition of butanone to isoprene has shown that in aqueous solution the volume of activation, $\Delta V^{\neq}$, is only about -34 cm^3/mol, whereas in dichloromethane as solvent $\Delta V^{\neq}$ is close to the expected value of ca. -40 cm^3/mol [857]; see also Section 5.5.11.

Not only Diels–Alder cycloadditions but also 1,3-dipolar cycloaddition reactions can be subject to hydrophobic rate enhancements. For example, the reaction of *C,N*-diphenylnitrone with di-*n*-butyl fumarate at 65 °C to yield an isoxazolidine is about 126 times faster in water than in ethanol, while in nonaqueous solvents there is a small 10-fold rate decrease on going from *n*-hexane to ethanol as solvent – in agreement with an isopolar transition-state reaction [*cf.* Eq. (5-44) in Section 5.3.3] [858]. Because water and ethanol have comparable polarities, the rate increase in water cannot be due to a change in solvent polarity. During the activation process, the unfavourable water contacts with the two apolar reactants are reduced, resulting in the observed rate enhancement in aqueous media. Upon addition of LiCl, NaCl, and KCl (5 M) to the aqueous reaction mixture the reaction rate increases further, whereas addition of urea (2 M) leads to a rate decrease, as expected for the structure-making and structure-breaking effects of these additives on water [858].

The hydrophobicity-driven association of reactant molecules in aqueous solution has even been found in aldol reactions. The trimethylsilyl ether of cyclohexanone adds to benzaldehyde in aqueous solution at 20 °C in the absence of a catalyst to give aldol addition products with a *syn/anti* stereoselectivity opposite to that of the acid-catalyzed reaction carried out in dichloromethane [746].

The useful ability of micelles to produce high local concentrations of bound organic reactants at low bulk concentrations has been used in photodimerization reactions; see references [712, 713] for reviews. The increase in reactant concentration per unit volume of the micelle promotes the probability of encounters between two molecules resulting in up to 1000-fold rate enhancements. The photodimerization of acenaphthylene illustrates this effect nicely; *cf.* Eq. (5-161) [720, 721].



(5-161)

At acenaphthylene concentrations as low as $2 \cdot 10^{-3}$ M, facile photodimerization takes place in benzene in the presence of nonionic or anionic surfactants, whereas in pure benzene at these concentrations no detectable amount of the two dimers is found [720]. Furthermore, the *cis/trans* product ratio is slightly dependent on the type of surfactant used [720, 721]. The product ratio during the course of photodimerization of 1-substituted acenaphthylenes is influenced by both solvent polarity and the addition of

surfactants. The photodimerization reaction in micelles yields stereoisomer ratios similar to those obtained in polar solvents [721].

Most chemical studies in this field have concentrated on the effect of micellar surfactants on reaction rates and only a few attempts have been made to investigate the effect that micelles might have in altering the relative extent of competing reactions. For example, in studying the competitive hydrolysis and aminolysis of aryl sulfates in aqueous solution, Fendler *et al.* have found [407] that cationic micelles such as cetyltrimethylammonium bromide (CTABr) are able to alter the balance between S—O bond fission and C—O bond fission as shown in Eq. (5-162).



(5-162)

The reaction of amines with 2,4-dinitrophenyl sulfate can result in the formation of phenol and sulfate ion (by S—O bond fission), or alternatively in the production of *N*-substituted anilines and hydrogen sulfate ions (by C—O bond fission). Under non-micellar conditions, C—O bond cleavage is the dominating reaction, while cationic micelles are able to induce complete suppression of aniline formation. This dramatic effect has been explained in terms of a change in the micro-environment of both the reactants and activated complexes through contributions from hydrophobic and electrostatic interactions [407].

To summarize, solubilization of reactants in micelles can lead to the following effects: local concentration, cage, pre-orientational, microviscosity, and polarity effects [713]. *Local concentration effects* are attributable to the increase in substrate concentration per unit volume of micelle due to the tendency of hydrophobic organic substrates to be solubilized in micelles. The ability of micelles to hold reactive intermediates together long enough for intramicellar reactions between them to occur is called the *cage effect*; *cf.* Section 5.5.10. The *pre-orientational effect* is the capability of micelles to solubilize substrates in a specific orientation, thus determining the regioselectivity of their reactions. Since the viscosity inside the micelles is generally much higher compared to the surrounding aqueous solution, a substrate molecule incorporated into a micelle has less translational and rotational freedom. This could be reflected in its chemical reactivity and is called the *microviscosity effect.* Finally, the micropolarity of the hydrophobic interior of a micelle and of the micelle/solution interface is different from the polarity of the bulk aqueous solution. A change in substrate reactivity due to the variation in micropolarity is called a *polarity effect.* Solvatochromic dyes can be used as molecular probes (*e.g.* pyridinium *N*-phenolate betaine dyes) to determine directly the actual micropolarity experienced by a substrate in the region of the micelle; *cf.* [722–724].

Space limits us to only a few representative examples of this type of hydrophobic interaction in this Section.

Oil and water do not mix, but on addition of a suitable surfactant a microemulsion can be formed depending on the relative concentrations of the three components. Microemulsions (*i.e.* surfactant/water/oil mixtures) can also be used as reaction media; see references [859–862] for reviews. Microemulsions are isotropic and optically clear, thermodynamically stable, macroscopically homogeneous, but microscopically heterogeneous dispersions of oil-in-water (O/W) or water-in-oil (W/O), where "oil" is usually a hydrocarbon. The name *microemulsion*, introduced by Schulman *et al.* in 1959 [863], derives from the fact that oil droplets in O/W systems or water droplets in W/O systems are very small (ca. 10 . . . 100 nm: *nanodroplets*). Unlike conventional emulsions, microemulsion domains fluctuate in size and shape with spontaneous coalescence and breakup. The oil/water interface is covered with surfactant molecules and this area can amount to as much as 10^5 m^2 per litre (!) of microemulsion.

A great variety of chemical reactions can be advantageously carried out in microemulsions [860–862]. In one of the first papers in this field, Menger *et al.* described the imidazole-catalyzed hydrolysis of 4-nitrophenyl acetate in water/octane microemulsions with AOT as an anionic surfactant [=sodium bis(2-ethyl-1-hexyl)-sulfosuccinate] [864]. The solubilized water, containing the imidazole catalyst, is confined in spherical pools encased by surfactant molecules, which have only their anionic head groups ($-SO_3^-$) immersed in the aqueous droplets. When the ester, dissolved in water-insoluble organic solvents, is added to this water/octane/AOT/imidazole system, it readily undergoes the catalysed hydrolysis under mild reaction conditions (25 °C).

More recently, microemulsions have been developed for the oxidative/hydrolytic destruction of sulfanes (→ sulfoxides) and phosphoric acid derivatives using sodium hypochlorite and cetyl-trimethylammonium chloride (CTAC). The compounds studied serve as model compounds for chemical warfare agents such as 'mustard gas' [bis(2-chloroethyl)sulfane] and sarin (GB: *i*-propyl-methylphosphonofluoridate). Big stocks of these must now be destroyed after implementation of the Chemical Weapons Convention (CWC), which came into force in April 1997 [865].

5.5.9 Liquid Crystals as Reaction Media

Liquid crystals possess physical properties which lie somewhere between those of solids and liquids; *cf.* Section 3.1 and [725]. The rigidity which is present in a solid matrix is absent in liquid crystals, thus permitting molecular motion as well as conformational flexibility of the dissolved solute molecules. At the same time, due to the order in the liquid-crystalline phase, the randomness in motion and conformational flexibility of the dissolved solute molecules is to some extent restricted. If the structures of the solute and solvent molecules are compatible, then solute molecules can be incorporated into the liquid-crystalline phase without disrupting its order. Thus, the reactivity of substrate molecules incorporated into liquid crystals without destroying their order should be different from that in isotropic solvents. Apart from the first report on the influence of liquid crystals on chemical reactions by Svedberg in 1916 [726], the use of liquid crystals as

solvents for chemical reactions has become a subject of further research only in recent decades; see references [713, 866, 867] for reviews.

For example, reactions which would lead to a product with steric demands that do not fit in the liquid-crystalline order can be retarded. That is, the rigidity of the solvent molecules can prevent uni- or bimolecular reactions which would be feasible in isotropic solvents. Alternatively, the ability of liquid crystals to orient dissolved solute molecules acts as a driving force in bimolecular reactions, particularly in entropy-controlled reactions, which are subject to severe orientational constraints in the activation process. Two factors seem to be primarily responsible in influencing the reactivity of solutes ordered in liquid crystals: (*a*) the solvation ability of the liquid-crystalline solvent, *i.e.* the efficiency with which the solute molecules are solvated, and (*b*) the degree of distortion which the reacting system must undergo during the activation process. Only a few examples showing the influence of solvent order on chemical reactions can be presented here; for reviews of organic photochemical reactions in organized media, see [713].

The rates and activation parameters for the thermal *cis* → *trans* isomerization of *N*,*N*′-distearoylindigo have been determined in both isotropic and liquid-crystalline solvents [727].



(5-163)

cis-*N*,*N*′-Diacylindigo R = *n*-$C_{17}H_{35}$ *trans*-*N*,*N*′-Diacylindigo

Solvent	C_6H_6 (48 °C)	$C_6H_5CH_3$ (48 °C)	isotropic BS (46 °C)	smectic BS*) (24 °C)
k_1^{rel}	68	53	31	1 (!)
$\Delta H^{\neq}/(kJ \cdot mol^{-1})$	93		91	130
$\Delta S^{\neq}/(J \cdot K^{-1} \cdot mol^{-1})$	−25		−33	+88

In isotropic nonpolar solvents such as benzene, toluene, and *n*-butyl stearate (BS; $t > 27$ °C), the long alkyl chains have no influence on the rate of the *cis* → *trans* isomerization reaction. However, in the smectic phase of the liquid-crystalline solvent *n*-butyl stearate, the isomerization rate is considerably slower than in isotropic solvents. The correspondingly higher activation enthalpy and the increased positive activation entropy are obviously caused by the migration of the two long stearoyl chains involved in the *cis* → *trans* isomerization. The solute alkyl chains are intertwined with ordered solvent molecules and their translocation will be resisted by nearby *n*-butyl stearate molecules. This anchoring effect, and not the shape changes of the isomerizing indigoid

* *n*-Butyl stearate has an enantiotropic smectic B phase from 14 . . . 27 °C [727].

moiety, is what is responsible for the observed rate decrease in smectic *n*-butyl stearate [727].

Similar rate decelerations in liquid-crystalline solvents have been observed for the thermal *cis* → *trans* isomerization of a bulky tetrasubstituted ethene in cholesteric phases [728]. On the other hand, the activation parameters for the thermal *cis* → *trans* isomerization of less-dipolar substituted azobenzenes show no dependence on the solvent order. This indicates that the *cis* isomers and their corresponding activated complexes present a similar steric appearance to the solvent environment [729]. This result is more consistent with an isomerization mechanism which proceeds by inversion rather than by rotation; *cf.* Eq. (5-40) in Section 5.3.2 and [527–529, 561]. The latter reaction represents a nice example of the use of liquid-crystalline solvents as mechanistic probes [729].

As a consequence of the alignment of solute molecules in liquid-crystalline solvents, the ratio of products formed in competitive reaction pathways can be different from that observed in isotropic liquids. This is illustrated by the Norrish type II photolysis of alkyl phenyl ketones with varying alkyl chain length in the isotropic, smectic, and



(5-164)

Solvent	*n*-C_7H_{16}	*n*-$C_{17}H_{36}$*)	CH_3CO_2—*n*-C_4H_9	*n*-$C_{17}H_{35}CO_2$—*n*-C_4H_9**)
$\frac{ketone}{cyclobutanol}$	1.1	1.2	2.2	21 (!)

solid phase of *n*-butyl stearate (BS) [730]. For example, the ratio of elimination to cyclization products formed in the photolysis of *n*-heptadecyl phenyl ketone was shown to exhibit a strong phase dependence with a 10-fold increase in the smectic *n*-butyl stearate phase relative to the isotropic solvent *n*-butyl acetate; *cf.* Eq. (5-164).

The small increase in product ratio observed on changing the solvent from *n*-heptane to *n*-butyl acetate is caused by the increase in solvent polarity. When the photolysis was carried out in solid *n*-heptadecane, no specific change was observed in the ketone/cyclobutanol product ratio.

These results have been explained as follows [730]: the cyclization of the intermediate 1,4-biradical, formed after γ-H-abstraction, requires rotation of the phenyl-substituted radical centre away from its equilibrium position. That is, the phenyl group

* Solid at 20 °C; $t_{mp} = 22.5$ °C [730].
** *n*-Butyl stearate has an enantiotropic smectic B phase from 14 . . . 27 °C [727].

must be placed in a direction perpendicular to the long axis of the alkyl phenyl ketone and the surrounding *n*-butyl stearate molecules. This should cause a disruption of the smectic solvent order and, as a consequence, cyclobutanol formation will be unfavourable in *n*-butyl stearate. For the competitive elimination reaction, the intramolecular motions required in the activation process should cause only a small distortion of the liquid-crystalline solvent order. Thus, the formation of elimination products is enhanced in the ordered solvent.

For alkyl phenyl ketones with shorter alkyl chains (*e.g.* *n*-butyl and *n*-decyl phenyl ketone), the elimination-to-cyclization ratio is virtually unaltered in the smectic solvent [730]. These ketones are not as rigidly incorporated into the liquid-crystalline lattice as *n*-heptadecyl phenyl ketone and thus are able to disrupt their local smectic environment to a greater extent. Hence, the corresponding conformations of the 1,4-biradical which lead to the elimination products are not more favourable than those which lead to cyclization products in the liquid-crystalline solvent.

In general, the greater the similarity in size and shape between the solute and liquid-crystalline solvent molecules, the easier it is for the solute to incorporate itself into the liquid-crystalline phase. If the result of a chemical reaction depends on the solvent order, then the largest effects can be expected for those solute molecules which fit best into the liquid-crystal structure. Since the structure of *n*-heptadecyl phenyl ketone is identical to *n*-butyl stearate except for the terminal phenyl and butoxy groups, it exhibits the largest solvent effect of the ketones studied [730].

Another illustrative example is the photodimerization of *n*-octadecyl *trans*-cinnamate, which has been studied in the isotropic, smectic, and crystalline phases of *n*-butyl stearate, BS [731]; *cf.* Eq. (5-165).

Irradiation of cinnamate esters leads to *trans*/*cis*-isomerization, dimerization, and ester cleavage reactions. The latter represents less than 5% of the total reaction pathway in this particular case. As expected, dimerization is favoured over isomerization with increasing initial solute concentrations. The *trans*/*cis*-isomerization involves only a relatively small perturbation of the surrounding solvent molecules. Therefore, it occurs with almost equal facility in the isotropic, smectic, and solid phases of *n*-butyl stearate. However, the regioselectivity of the photochemical [2 + 2]cycloaddition reaction is strongly phase-dependent. Of the possible dimers, only the two stereoisomers shown in Eq. (5-165) were detected in the irradiated samples. A strong preference for *head-to-tail* dimerization has been found in the smectic and solid phases. This result has been explained by assuming that not only solvent-mediated solute alignments but also dipole-dipole induced interactions between dipolar cinnamate molecules must exist in the mesophase. The dipole-dipole interaction leads to pairwise, antiparallel associations of the cinnamate molecules with interdigitation among neighbouring solute molecules. Irradiation of these antiparallel oriented solute associates preferentially yields the *head-to-tail* dimers. In summary, the regioselectivity of the photodimerization of *n*-octadecyl *trans*-cinnamate is controlled by the combination of two factors: solute alignment by ordered solvents and dipole-dipole interactions between solute molecules [731].

Similar results have been observed in the photodimerization reaction of acenaphthylene [732, 733]; *cf.* Eq. (5-161) in Section 5.5.8. A considerable increase in the production of the *trans*-adduct was reported in cholesteric liquid-crystalline media compared to the isotropic solvent benzene, in which the *cis*-adduct is the dominant product



(5-165)

Solvent	BS at 32 °C (isotropic)	BS at 18 °C*) (smectic)	BS at 8 °C (solid)
$\frac{\textit{head-to-tail}^{**)}}{\textit{head-to-head}}$	3.3	14.2	15.1

[733]. That is, the ordered structure of the cholesteric mesophase affects the formation of the *trans*-adduct advantageously. Furthermore, the *trans*/*cis* product ratio depends significantly on the initial acenaphthylene concentration. In isotropic solutions, the dimerization of singlet-excited acenaphthylene molecules is known to yield exclusively the *cis*-adduct, whereas a mixture of *cis*- and *trans*-adducts results from triplet-excited solute molecules. The lowering of *cis*-adduct production in the mesophase has been attributed to the enhanced efficiency of the triplet reaction in comparison with the singlet reaction, as shown by quantum yield measurements [732]. The increase in triplet reaction efficiencies has been ascribed to the increase in the fraction of acenaphthylene–acenaphthylene collisions which have coplanar or parallel-plane orientations with respect to the surrounding solvent molecules, and not to the increase in the total number of collisions per unit time [732]. See references [713, 732, 733] for a more detailed discussion of this photodimerization reaction.

Surprisingly, some Diels–Alder cycloaddition reactions show no variation in *endo*/*exo* product ratio with changes in solvent phase. Ordered liquid-crystalline solvents are not able to differentiate between *endo*- and *exo*-activated complexes in the Diels–Alder reaction of 2,5-dimethyl-3,4-diphenylcyclopentadienone with dienophiles of varying size (cyclopentene, cycloheptene, indene, and acenaphthylene), when it is carried out in isotropic (benzene), cholesteric (cholesteryl propionate), and smectic liquid-crystalline solvents at 105 °C [734].

Cholesteric liquid crystals are optically active nematic phases as a result of their gradual twist in orientational alignment. Therefore, cholesteric liquid-crystalline solvents are expected to induce enantioselectivity in chemical reactions; see reference [713] for a review on photoasymmetric induction by chiral mesophases. The existing results are not very promising. So far, the maximum photoasymmetric induction reported has

* *n*-Butyl stearate exists as an enantiotropic smectic B phase from 14 . . . 27 °C [727].
** Dimer ratios.

been for hexahelicene formation in cholesteric phases, with an enantiomeric excess of the right-handed helix of about 1% [735]. The decarboxylation of ethyl-phenylmalonic acid in cholesteryl benzoate as a liquid-crystalline solvent gave 2-phenylbutyric acid (overall yield 80 cmol/mol), which was shown to be optically active with an enantiomeric excess of 18% of the (*R*)-(−) enantiomer; *cf.* Eq. (5-166) [765]. In contrast to the



(5-166)

18% *ee* found in the cholesteric solvent, decarboxylation of ethyl-phenylmalonic acid in bornyl acetate, an isotropic chiral solvent, yielded 2-phenylbutyric acid which was essentially racemic (*ee* = 0%) [765]. Using cholesteryl 4-nitrobenzoate as a liquid-crystalline solvent, the *ortho*-Claisen rearrangement of γ-methylallyl 4-tolyl ether yielded optically active 2-(α-methylallyl)-4-methylphenol, but the absolute configuration and optical purity of the rearrangement product were unknown [766]. Various explanations for the failure of asymmetric induction in thermal reactions in liquid-crystalline media have been given in the literature [736, 737].

Not only reaction rates, but also the positions of chemical equilibria can be influenced by liquid crystals as reaction media. A nice example is the ionization equilibrium of chloro-tris(4-methoxyphenyl)methane according to $Ar_3C{-}Cl \rightleftharpoons Ar_3C^+ + Cl^-$, which is more shifted in favour of the nearly planar triarylcarbenium ion in nematic liquid crystals as compared to in an isotropic reaction medium [868]. Obviously, the disc-shaped carbenium ion fits better into the rod-like nematic phase than the tetrahedral covalent ionogen, which distorts the internal structure of the nematic liquid crystal.

All in all, liquid-crystalline media are not generally useful solvents for controlling the rates and stereochemistries of chemical reactions. In each case, careful consideration of the fine details regarding the structure of educts and activated complex, their preferred orientations in a liquid-crystalline solvent matrix, and the disruptive effects that each solute has on the solvent order has to be made. A mesophase effect can only be expected when substantial changes in the overall shape of the reactant molecule(s) occur during the activation process [734].

5.5.10 Solvent Cage Effects

When two chemical species react together at every encounter*), chemical change can take place only as fast as the reactants can diffuse together. In contrast to the gas phase, there may be many collisions per encounter in solution. Considering a pair of molecules that have just encountered each other or, what in this regard is equivalent, a pair of molecules that have just arisen from the decay of a parent molecule, these molecules can

* The whole process of two species coming together and remaining together for a number of subsequent collisions in solution has been called an "encounter".

become separated from each other only as a result of their diffusional motion through the inert solvent. If this diffusion is comparatively slow in solution, diffusion may be answerable for the observable rates of many reactions, since the slowest step in a sequence determines the net rate of change. Such reactions are called "encounter-" or "diffusion-controlled" [18, 408, 409, 409a].

Reactions that occur with an activation energy of less than about 20 kJ/mol (5 kcal/mol) depend upon the rate at which the reactants move together through the solution. Since the movement of a molecule through an inert solvent has itself an activation energy of diffusion of about 20 kJ/mol, this movement becomes the slowest step for diffusion-controlled reactions. Almost all radical–radical recombination reactions require so little activation energy that they are usually diffusion-controlled in solvents of normal viscosity. Thus, the recombination rates for simple alkyl radicals in inert solvents lie in the range $10^8 \ldots 10^{10}$ L · mol^{-1} s^{-1} at 25 °C [410]. Ion combination reactions such as the acid-base reaction between the solvated proton and solvated hydroxide ion are also known to be diffusion-controlled. This reaction has one of the largest rate constants known for liquid-phase reactions, $1.4 \cdot 10^{11}$ L · mol^{-1} s^{-1} in water at 25 °C [411].

When a pair of radicals (or ions) have reached adjacent positions through diffusion, or have been generated from a single parent molecule as in the decomposition of radical initiators, they are hemmed in by a cage of solvent molecules – an effect that has been called the "solvent cage effect"*). The two species must diffuse to become statistically distributed in the solvent, but, because of the activation energy of diffusion, they collide with each other several times before they separate, and recombination may occur during these collisions**). This type of reaction is called a "cage reaction" [413].

Rate constants for diffusion-controlled reactions can be calculated from the laws of diffusion [18, 869]. For a simple cage reaction $A\cdot + \cdot B \rightarrow AB$, in which $A\cdot$ reacts with $\cdot B$ every time the two approach one another to within a distance R, the following equation can be derived,

$$k_2 = 4\pi \cdot R \cdot N_A \cdot (D_A + D_B)$$

where k_2 represents the overall second-order rate constant for uncharged reactants, R is the average radius of the solvent cage, N_A is Avogadro's number, and D_A and D_B are

* The concept of a *solvent cage* restricting the separation of two reactants was first introduced by Frank and Rabinowitsch [412] in predicting a reduction in the quantum yield for photodissociation processes in solution compared with the gas phase. Thus, when iodine in solution is dissociated by a flash, the quantum yield is much less than unity indicating that most of the iodine atoms recombine before escaping from the solvent cage. The *solvent cage* is an aggregate of solvent molecules that surrounds the fragments formed by thermal or photochemical bond cleavage of a precursor species. A more imaginative description of the cage effect has been given by Houston [18]: "The difference between a reaction in the gas phase and one in solution is much like the difference between a romantic encounter on an empty beach and one on a crowded dance floor. The romance of closeness is not appreciably changed by the surrounding solvent of dancers, but it is more difficult to find one another in a crowd, and correspondingly difficult to separate once the dance has ended. The solvent tends to slow the rate of approach of the reactants, so that they must diffuse toward one another through the solution, but it also keeps them together for many 'collisions' once they come in contact. This latter phenomenon is often referred to as the *cage effect*."

** Other radical reactions which can compete with recombination inside the solvent cage are disproportionation [414] and reaction with the adjacent solvent molecules.

the diffusion coefficients of reactants A· and ·B through the solvent under consideration. For a cage reaction between a pair of reactants whose average diameter is $R = 2.0$ nm and diffusion coefficients of $D_A \approx D_B \approx 10^{-9}\ m^2 \cdot s^{-1}$, typical for many species in aqueous solution, a rate constant of $k_2 = 3.0 \cdot 10^{10}\ L \cdot mol^{-1} \cdot s^{-1}$ can be calculated according to this equation [18]. For oppositely charged reactants A^+ and B^-, an additional multiplicative factor due to the mutual electrostatic attraction has to be added.

With azoalkanes, peroxides, and other initiators, varying percentages of the radicals formed react together as a geminate pair in the solvent cage, as seen, for instance, in the photolysis of azomethane according to Eq. (5-167) [415–417]. The efficiency of free-radical production, F, in the decomposition of azomethane can be described in terms of an effective rate constant, k_d, which approximately describes the diffusional separation of caged radical pairs to give free radicals, a process competing with radical–radical reactions within the solvent cage, the rate constant of which is designated as k_c. The combination reaction must obviously be very fast, in order to compete successfully with diffusion. It is usually assumed that k_c is nearly independent of the nature of the solvent, and that all the variation in cage products may be attributed to k_d. Very clear evidence for the existence of a cage effect in reaction (5-167) was found from the results of cross-



(5-167)

over experiments [416, 417]. When a mixture of azomethane and perdeuterioazomethane is photolyzed in the gas phase, the $CH_3 \cdot$ and $CD_3 \cdot$ radicals formed recombine rapidly to yield CH_3CH_3, CD_3CD_3, and CH_3CD_3 in such proportions (1:1:2) that it is clear that $CH_3 \cdot$ and $CD_3 \cdot$ are randomly mixed before recombination. When, however, the same experiment is repeated in the inert solvent *i*-octane, no CH_3CD_3 is detected when CH_3CH_3 and CD_3CD_3 are formed along with N_2. Evidently, solvent molecules keep methyl radicals formed from the same parent molecule together until they recombine. The cage effect can occur in the gas phase at high pressures, as has been demonstrated for the photolysis of azomethane in propane at 49 bar. Here, propane and intact reactant molecules act as a quasi-solvent and prevent formation of CH_3CD_3 [416]. For further gas-phase analogous cage effects, see reference [870].

Cage effects also account for the fact that not all the radicals produced from the decomposition of initiators such as azobisisobutyronitrile (AIBN) are effective in initiating radical polymerizations. In the somewhat simplified reaction Scheme (5-168) depicting the thermolysis of AIBN, two types of cyanopropyl radicals are shown, one still within the solvent cage, whereas the others have reached their statistical separation

in the solution. Adding certain reactive compounds called scavengers to the solution should divert the free radicals away from dimer formation. In other words, radical



(5-168)

scavengers should reduce the yield of tetramethylsuccinonitrile derived from the free radicals (but not that formed from the caged ones) to a limiting value. This then represents the amount of product formed within the solvent cage. Indeed, the yield of tetramethylsuccinonitrile from the thermolysis of AIBN in tetrachloromethane at 80 °C falls from about 96 cmol/mol when no scavenger is present to a constant value of about 19 cmol/mol using *n*-butanethiol, a good hydrogen donor, as a scavenger [418]. This means that 19 cmol/mol of the initiator radicals are lost in solvent cage recombinations and only 81 cmol/mol of AIBN molecules are efficient in free-radical production under these reaction conditions. The initiator efficiency of AIBN is, in fact, lower under most circumstances, since only few substrates approach the reactivity of *n*-butanethiol as a scavenger. The fraction of AIBN molecules leading to initiation of styrene polymerization increases as the concentration of the monomer increases, reaching a limiting value of 68 cmol/mol above 2 mol/L styrene in an inert solvent [419]. The remaining 32 cmol/mol may be attributed to the reactions of the geminate radical pairs.

Solvent cage effects have been observed not only in the decomposition of azoalkanes, but also in the thermolysis of other compounds, such as peroxides (*e.g.* diacyl peroxides [420], peresters [421]), which generate two radicals simultaneously. For example, diacetyl peroxide initially labeled with ^{18}O in the carbonyl oxygen was partially thermolysed in *i*-octane at 80 °C and it was found that *ca.* 38 cmol/mol of all geminate radical pairs formed in the decomposition recombine to give diacetyl peroxide with scrambling of the label [420]. Solvent cage effects have even been found for ion-producing reactions, albeit only at high external pressures. The solvolysis of 2-bromopropane in a 4:1 methanol/ethanol mixture at 46 °C becomes diffusion-controlled at pressures higher than 40 kbar [738]. Liquid-crystalline solvents and micellar systems can also provide solvent cages. Liquid crystals and micelles have the capacity to hold two reactive species together for a longer period of time compared to isotropic, homogeneous solutions. Micellar aggregates are particularly effective in imposing cage constraints on chemical reactions. Photochemical examples of the consequences of restrict-

ing solute diffusion in liquid crystals and in micelles can be found in references [713, 871]; *cf.* also Sections 5.5.8 and 5.5.9.

The prediction of solvent effects on k_d, as reflected by the efficiency of free-radical production, F, has been attempted by considering the correlation between F and macroscopic solvent parameters such as solvent viscosity [413]. That the proportion of cage reactions depends on the viscosity of the solvent used was shown by Kochi [422], who decomposed a series of diacyl peroxides in *n*-pentane and decalin. In decalin, the more viscous solvent, photolysis of two different diacyl peroxides gave high yields of the symmetrical dimers and only a minor yield of cross-dimer, thus indicating the strong solvent cage effect. In the more fluid solvent *n*-pentane, the cage effect is less important and a higher proportion of the cross-products was formed. Usually, however, no simple relationship with solvent viscosity was found. Studying the photolysis of azomethane in a wide range of different media, Martin *et al.* [423] were able to correlate the observed amount of cage product ethane to macroscopic solvent parameters other than viscosity, namely solvent internal pressure and cohesive energy density (*cf.* Section 5.4.2).

We shall conclude this Section with an example of solvent cage effects of ion-molecule recombination reactions as found in the ozonolysis of alkenes in non-polar solvents [739, 740]. According to the Criegee mechanism [424], unsymmetrically



(5-169)

substituted alkenes ought to give two zwitterions and two carbonyl compounds after decomposition of the unstable primary ozonide, as shown in reaction scheme (5-169). If both scission and recombination are statistical, the three possible final ozonides *A*, *B*, and *C* should be formed in a 1:2:1 ratio, provided there is no preferred breakdown of the initial ozonide and no solvent cage. The experimentally determined values of this molar ratio differ from the statistical value, the yield of the symmetrical cross-ozonides always being lower, as observed for 2-pentene [425] and 2-hexene [426]. This means that part of the recombination reaction must occur in a solvent cage. With an increase in the initial concentration of alkene, the relative proportions of "normal" and "cross" ozonides approach the statistical values. When he failed to find cross-ozonides in the ozonolysis of 3-heptene, Criegee postulated that the zwitterion and carbonyl fragments were formed and recombined in a solvent cage, thus preventing cross-recombination [427]. The earlier failure of Criegee to obtain any cross-ozonides in the case of 3-heptene can

now probably be attributed to the alkene concentration used [428]. At the concentration of alkene normally used in ozonolysis the predominant product is the normal parent ozonide of the unsymmetrically substituted alkene. A plot of the ratio of normal ozonides to cross-ozonides *versus* alkene concentration shows that cross-ozonide formation decreases with dilution [428]; *cf.* also [428a].

Reduced solvent cage effects have, however, been observed for ozonations in polar solvents such as dichloromethane and ethyl acetate, in contrast to nonpolar hydrocarbons; *cf.* [739, 740]. Increasing the polarity of the ozonation solvent leads to an increase in the ratio of cross to normal ozonides formed from unsymmetrical alkenes. Obviously, the increased solvent polarity increases the separation of the carbonyl oxide and the carbonyl compound, allowing them to react more independently rather than in a solvent cage. Accordingly, in contrast to nonpolar solvents, an increase in reaction temperature has no effect on the ozonide ratio. This is due to the fact that in polar solvents the solvent cage effect is minimal. For the ozonolysis of an equimolar mixture of ethene and tetradeuterioethene, leading to two normal and one cross-ozonide, it has been estimated that in polar solvents such as CH_2F_2 at least 90% of the final ozonide formation occurs outside the solvent cage in which the primary ozonide decomposition took place. This percentage decreases to about 10% out-of-cage recombination in a nonpolar solvent such as *i*-butane [741].

In conclusion, the relative yields of the normal and cross-ozonides are a function of initial alkene concentration, solvent polarity, and temperature. High alkene concentration, polar solvents, and high temperatures maximize the cross-ozonide yields, with proportions close to statistical amounts in favourable cases [739, 740].

For recent comprehensive reviews on the ozonation reaction, dealing not only with solvent effects on the relative yields of normal and cross-ozonides, but also with solvent effects on the total ozonide yield and the *cis*/*trans* ozonide ratio from *cis*- and *trans*-alkenes, see references [739, 740]. Supplementary solvent effects have been reported for the ozonolysis of 3-aryl-1-methylindenes yielding solvent-dependent mixtures of final *exo*- and *endo*-ozonides [742].

5.5.11 External Pressure and Solvent Effects on Reaction Rates

Not only the internal pressure of a solvent can affect chemical reactions (see Section 5.4.2 [231, 232]), but also the application of external pressure can exert large effects on reaction rates and equilibrium constants [239, 429–433, 747–750]. According to Le Chatelier's principle of least restraint, the rate of a reaction should be increased by an increase in external pressure if the volume of the activated complex is less than the sum of the volumes of the reactant molecules, whereas the rate of reaction should be decreased by an increase in external pressure if the reverse is true. The fundamental equation for the effect of external pressure on a reaction rate constant k was deduced by Evans and Polanyi on the basis of transition-state theory [434]:

$$\left(\frac{\partial \ln k}{\partial p}\right)_T = -\frac{\Delta V^{\neq}}{RT} \tag{5-170}$$

provided that k is expressed in pressure-independent concentration units (mole fraction or molality scale) at a fixed temperature and pressure*). The volume of activation, $\Delta V^{\neq}$, is interpreted as the difference between the partial molar volume of the activated complex including its molecules of solvation, $V^{\neq}$, and the sum of the partial molar volumes of the reactants with their associated solvent molecules, $\sum r \cdot V^{R}$, at the same temperature and pressure: $\Delta V^{\neq} = V^{\neq} - \sum r \cdot V^{R}$, where r is the stoichiometric number of the reactant R and V^{R} its partial molar volume.

To produce significant changes in $\ln k$, pressures of several hundred bars are commonly used. While a change in reaction rate by a factor of 2 to 4 can be anticipated for a change in temperature of 10 °C, a pressure change of about 700 bar is required to bring about the same effect even for reactions with the relatively large activation volume of ± 40 cm$^3 \cdot$ mol^{-1} [435]. Normally, the magnitude of $\Delta V^{\neq}$ lies somewhere between $+25$ and -25 cm$^3 \cdot$ mol^{-1}, which requires pressures of *ca.* 1200 bar to produce rate changes of a factor of 3. The minus sign in Eq. (5-170) means that pressure accelerates reactions which are characterized by a volume shrinkage in passing from the reactants through the activated complex (negative $\Delta V^{\neq}$) and retards those with a volume expansion (positive $\Delta V^{\neq}$). In many cases, a plot of $\ln k$ against pressure is linear, indicating that for these reactions the quantity $\Delta V^{\neq}$ is independent of pressure. However, above *ca.* 10 kbar, reactions do not obey the ideal rate equation (5-170) since activation volumes are pressure-dependent; the values of $\Delta V^{\neq}$ generally decrease as pressure increases. Extensive compilations of activation and reaction volumes in solution for a great variety of pressure-dependent chemical reactions can be found in references [430, 749].

The activation volume changes arise from two sources: (a) making and breaking of chemical bonds, and (b) interaction of reactants and activated complex with the surrounding solvent molecules. Therefore, $\Delta V^{\neq}$ can be considered as being composed of $\Delta V_i^{\neq}$, the *intrinsic* change in molar volume of the reactant molecules themselves in forming the activated complex, as well as $\Delta V_e^{\neq}$, the change in molar volume of the solvating solvent molecules during the activation process, according to Eq. (5-171)**):

$$\Delta V^{\neq} = \Delta V_i^{\neq} + \Delta V_e^{\neq} \tag{5-171}$$

$\Delta V_e^{\neq}$ is related to the well-known volume contraction normally observed on dissolution of electrolytes and called *electrostriction*. The contraction of the solvent surrounding an ion is best expressed by the Drude–Nernst equation (5-172):

---

* If k is expressed in the pressure-dependent molarity scale (*i.e.* mol $\cdot$ L^{-1}), its pressure-dependence is given by

$$\left(\frac{\partial \ln k}{\partial p}\right)_T = -\frac{\Delta V^{\neq}}{RT} - \kappa(n-1)$$

where κ is the compressibility of the solution (usually that of the solvent) and n is the kinetic order of the reaction. For first-order reactions ($n = 1$), the equations for the pressure-dependence of k are identical. For reactions of higher order, however, the additional compressibility term must be taken into account.

** For modifications of the somewhat simplified Eq. (5-171) and corresponding discussions, see references [872, 873].

$$\Delta V_e = -\frac{N_A \cdot z^2 \cdot e^2}{2r \cdot \varepsilon_r^2} \frac{\partial \varepsilon_r}{\partial p} \tag{5-172}$$

The solvent is assumed to be a continuum of relative permittivity ε_r, and the ion to be a hard sphere of radius r with charge $z \cdot e$; N_A is the Avogadro number. Eq. (5-172) represents the electrostatic contraction change in volume for a mole of ions. According to the Drude–Nernst equation, electrostriction should be proportional to the square of the charge on the ions, inversely proportional to the ionic radii, and should increase in proportion to the value of $(1/\varepsilon_r^2)/(\partial \varepsilon_r/\partial p)$, which depends on the nature of the solvent. It follows that ΔV_e will vary strongly in reactions where charges are created or neutralized, which, in turn, should show up as a solvent dependence of $\Delta V^{\neq}$.

$\Delta V_i^{\neq}$ represents the change in volume due to changes in bond lengths and angles. It is this contribution to $\Delta V^{\neq}$ that is connected to the reaction mechanism in terms of the relative positions of the atoms in reactants and the activated complex. The absolute size of $\Delta V_i^{\neq}$ has been concluded to be approximately $+10\ \mathrm{cm^3 \cdot mol^{-1}}$ for bond cleavage and approximately $-10\ \mathrm{cm^3 \cdot mol^{-1}}$ for bond formation in reactions of organic molecules [430].

In a reaction, $A \rightleftharpoons [X]^{\neq} \rightarrow C + D$, bonds will be stretched to form $[X]^{\neq}$, so that $\Delta V_i^{\neq}$ would be expected to be positive for unimolecular dissociative reactions and negative for the reverse bimolecular associative reaction of C and D. The $\Delta V_e^{\neq}$ contribution is caused by the rearrangement of the solvent molecules due to steric requirements of the reaction and to the change in charge density on activation. The latter effect can be predicted on the basis of the qualitative theory of solvent effects introduced by Hughes and Ingold [16, 44] (*cf.* Section 5.3.1), representing the reacting ions or dipolar molecules as spheres and the solvent as a dielectric. Creation or concentration of charge on the reacting species, on passing from the initial to the transition state, will increase the intermolecular electrostatic forces between the solute and the permanent or induced dipoles in the solvating molecules. This leads to a reduction in volume, called *electrostriction*, of the solvate complex. Since the extent of electrostriction varies as the square of the charge on the sphere according to Eq. (5-172), the association of ions with like charges will increase the electrostriction in the solvent and $\Delta V_e^{\neq}$ will be large and negative, whereas association of two oppositely charged reactants will be accompanied by a large positive $\Delta V_e^{\neq}$. Because the degree of solvation of ionic or strongly dipolar species may be extensive, the $\Delta V_e^{\neq}$ term often predominates over the $\Delta V_i^{\neq}$ term. The activation volume, therefore, reflects not only the intrinsic differences in molecular dimensions of reactants and activated complex, but also the difference in their degree of solvation. Pressure is therefore a probe uniquely suited for the study of solvation changes during a reaction.

The importance of the solvent in determining the effects of pressure on reaction rates has been recognized in general terms for a long time, but the first satisfactory discussion was given by Buchanan and Hamann in 1953 [436]. A schematic compilation of pressure and solvent effects on reactions of different charge type, established by Dack [27, 239], is given in Table 5-25. The entire basis of the effect of solvent polarity on $\Delta V^{\neq}$, as shown in Table 5-25, is that less polar solvents have higher compressibilities and are therefore more constricted by ionic or dipolar solutes than more polar solvents, which exhibit smaller compressibilities owing to the strong intermolecular interactions already present in the absence of a solute. This consideration would also suggest a cor-

Table 5-25. Effect of external pressure and solvent polarity on the rate of reactions of different charge type and on their volume of activation, $\Delta V^{\neq}$ [27, 239].

No.	Reactants	Activated Complex	Effect of increased pressure on reaction rate	Effect of increased solvent polarity on $\Delta V^{\neq}$
1			Decrease	None
2			Increase	None
3			Increase	More negative
4			Decrease	More positive
5			Increase	More negative
6			Decrease	More positive
7			Decrease	More positive
8			Decrease	More positive

relation between $\Delta V^{\neq}$ and the entropy of activation, $\Delta S^{\neq}$, because an increase in electrostriction due to an intensification of the electric field around the solute corresponds to a decrease in both volume and entropy owing to reduced freedom of motion within the solvent complex. That is, a large negative $\Delta V^{\neq}$ value should, in principle, correspond to a large negative $\Delta S^{\neq}$ value for reactions whose temperature and pressure dependence has been investigated in solvents of different polarity. Indeed, a linear relationship of $\Delta V^{\neq}$ with $\Delta S^{\neq}$ has been reported for many reactions.

Two typical examples shall illustrate the predictions made by Table 5-25. Clear-cut examples of reaction type 2 are Diels–Alder cycloaddition reactions*). The solvent

* The first Diels–Alder reaction studied under high pressure was the dimerization of cyclopentadiene [751]. For recent, more detailed studies of the pressure-dependence of Diels–Alder cycloaddition reactions in solvents of different polarity, as well as discussions of the corresponding mechanistic aspects, see references [857, 874].

Table 5-26. Effect of external pressure and solvent polarity on reaction rate and activation volume of the Diels–Alder reaction between isoprene and maleic anhydride at 35 °C [437]; *cf.* Eq. (5-42) in Section 5.3.3.

Solvents	$k_2 \cdot 10^4/s^{-1}$ [a]		$k_2^{rel.}$ [b]	$\Delta V^{\neq}/(cm^3 \cdot mol^{-1})$ at 1 bar [c]
	at 1 bar	at 1336 bar		
Dichloromethane	5.28	26.1	4.9	−39.8
Dimethyl carbonate	1.82	–	–	−39.3
Acetone	2.18	11.8	5.4	−39.0
Diisopropyl ether	0.597	3.56	6.0	−38.5
1-Chlorobutane	1.59	9.31	5.9	−38.0
Acetonitrile	6.25	33.9	5.4	−37.5
Ethyl acetate	1.22	6.33	5.2	−37.4
1,2-Dichloroethane	5.50	32.2	5.9	−37.0
Nitromethane	9.86	44.4	4.5	−32.5

[a] Rate constants based on mol fraction scale. $\Delta\Delta V^{\neq} = 7.3\ cm^3 \cdot mol^{-1}$
[b] $k_2^{rel.} = k_2(1336\ bar)/k_2(1\ bar)$.
[c] Limit of error $\pm 0.8\ cm^3 \cdot mol^{-1}$.

effect on the activation volumes for such a cycloaddition reaction is reported in Table 5-26 [437]. In agreement with an isopolar cyclic activated complex being intrinsically smaller than the reactants, large negative values of $\Delta V^{\neq}$ have been found for this reaction. As expected, solvent polarity has comparatively little influence on $\Delta V^{\neq}$ ($\Delta\Delta V^{\neq} = 7.3\ cm^3 \cdot mol^{-1}$ between nitromethane and dichloromethane). This suggests that the solute–solvent interactions of the activated complex are small and similar to those of the reactants. Nevertheless, the small solvent effects obtained might be explained in terms of solvent internal pressure [27, 438], this acting on the rates of non-polar reactions in the same direction as external pressures*).

Similar pressure effects have been observed in the 1,3-dipolar cycloaddition of diazo-diphenylmethane to various alkenes. This is in agreement with a concerted mechanism involving an isopolar activated complex [752, 753]; *cf.* Section 5.3.3. For the 1,3-dipolar cycloaddition of diazo-diphenylmethane to dimethyl acetylenedicarboxylate at 25 °C in *n*-hexane ($\Delta V^{\neq} = -24\ cm^3 \cdot mol^{-1}$) and in acetonitrile ($\Delta V^{\neq} = -15\ cm^3 \cdot mol^{-1}$), the solvent-induced difference $\Delta\Delta V^{\neq}$ is only 9 $cm^3 \cdot mol^{-1}$; this corresponds to a very small rate constant solvent effect, $k_2(CH_3CN)/k_2(n\text{-}C_6H_{14}) = 3.4$ [752].

Contrary to reactions going through isopolar transition states, reactions of types 3 to 8 in Table 5-25, which involve formation, dispersal or destruction of charge, should exhibit large solvent effects on their activation volumes. This is shown in Table 5-27 for the S_N2 substitution reaction between triethylamine and iodoethane [441], an example of the well-known Menschutkin reaction, the pressure dependence of which has been investigated thoroughly [439–445, 755].

Corresponding to reaction type 5 in Table 5-25, increasing pressure leads to an increase in reaction rate, which is more pronounced in less polar solvents. The $\Delta V^{\neq}$

* For common organic solvents, the internal pressures range from 1800 to 5000 bar at 25 °C [438].

Table 5-27. Effect of external pressure and solvent polarity on reaction rate and activation volume of the Menschutkin reaction between triethylamine and iodoethane at 50 °C [441]; *cf.* also Table 5-5 in Section 5.3.1 [59].

Solvents	$k_2 \cdot 10^6/(\mathrm{L \cdot mol^{-1} \cdot s^{-1}})$		$k_2^{\mathrm{rel.}}$ [a]	$\Delta V^{\neq}/(\mathrm{cm^3 \cdot mol^{-1}})$ at 1 bar [b]
	at 1 bar	at 1961 bar		
n-Hexane	0.123	1.35	11.0	−51.5
Acetone	318	2600	8.2	−48.5
Benzene	27.8	–	–	−46.2
Chlorobenzene	92.0	627	6.8	−44.6
Methanol	50.6	208	4.1	−40.6
Nitrobenzene	934	4450	4.8	−37.0

[a] $k_2^{\mathrm{rel.}} = k_2(1961\ \mathrm{bar})/k_2(1\ \mathrm{bar})$. $\Delta\Delta V^{\neq} = 15\ \mathrm{cm^3 \cdot mol^{-1}}$

[b] The limit of error for $\Delta V^{\neq}$ lies between ± 1.6 and $\pm 2.6\ \mathrm{cm^3 \cdot mol^{-1}}$.

values observed in solvents of different polarity demonstrate clearly that nonpolar solvents undergo more electrostriction than polar media ($\Delta\Delta V^{\neq} = 15\ \mathrm{cm^3 \cdot mol^{-1}}$ between *n*-hexane and nitrobenzene). Somewhat more positive activation volumes have been reported for the Menschutkin-type reaction of triphenylphosphane with iodomethane at 30 °C: $\Delta V^{\neq} = -28.0\ \mathrm{cm^3 \cdot mol^{-1}}$ in acetonitrile and $\Delta V^{\neq} = -17.6\ \mathrm{cm^3 \cdot mol^{-1}}$ in the slightly more polar propylene carbonate; the reaction is 245 times faster in propylene carbonate than it is in nonpolar diisopropyl ether [755].

Reaction type 3 in Table 5-25 is best represented by the S_N1 solvolysis of 2-chloro-2-methylpropane; *cf.* Eq. (5-13) in Section 5.3.1. Considering the heterolysis of the C—Cl bond, one would expect the activation volume to be positive because of the C—Cl stretching during the activation process. However, a negative activation volume of $\Delta V^{\neq} = -22.2\ \mathrm{cm^3 \cdot mol^{-1}}$ has been found for this solvolysis at 30 °C in ethanol/water (80:20 cL/L), indicating a strong volume contraction due to solvation of the dipolar activated complex (electrostriction) [756]. Typical activation volumes for haloalkane solvolyses in protic solvents are in the range of $-15 \ldots -30\ \mathrm{cm^3 \cdot mol^{-1}}$.

Both S_N1 and S_N2 solvolyses of haloalkanes have negative activation volumes at low external pressures because the activation volume is mainly determined by the electrostriction of the surrounding solvent molecules [*cf.* reaction types 3 and 7 in Table 5-25 and Eq. (5-171)]. The electrostriction of solvents is usually high at zero or low external pressures because common solvents are quite compressible [875]. However, at very high external pressures, the compressibility of solvents decreases greatly. As a consequence, the electrostriction around a solute ion or dipole, which is related to the macroscopic compressibility, drops significantly at high pressures (*i.e.* at several tens of kilobars). Under such drastic experimental conditions, an S_N1 solvolysis should now have a positive and an S_N2 solvolysis a negative activation volume, which allows a clear distinction between the two mechanisms. This has indeed been found for the solvolysis of 2-chloro-2-methylpropane (S_N1) and iodomethane (S_N2) in glycerol as solvent and nucleophile at high external pressures [738]. The S_N1 reaction now has (above 16 kbar) a positive $\Delta V^{\neq}$ because the C–Cl bond length is increased in the activated complex, while $\Delta V^{\neq}$ remains negative (up to 70 kbar) for the S_N2 reaction because a solvent molecule forms a partial

O···C bond in the activation process. An analogous high-pressure experiment (up to 82 kbar) showed that the solvolysis of 2-bromopropane in methanol/ethanol (4:1) follows an S_N1 mechanism, with a negative $\Delta V^{\neq}$ at low pressures ($p < 40$ kbar), and a positive $\Delta V^{\neq}$ at higher pressures ($p > 40$ kbar) [738].

Many other examples of pressure-dependent organic reactions of the types collected in Table 5-25 are given in references [239, 429–433, 747–750]. In general, the influence of external pressure and solvent polarity can be used to draw conclusions about whether or not the activated complex is more dipolar than the initial reactants, and helps to clarify reaction mechanisms in the framework of the transition-state theory (TST).

High-pressure effects can be used to study dynamic solvent effects on reactions in highly viscous liquid phases, that is, in such phases in which reactions are shifted from the equilibrium TST-valid region to the nonequilibrium TST-invalid region at high pressures; *cf.* Section 5.3.5 and a recent review by Asano [754]. Commonly, chemical transformations and the corresponding rearrangement of the solvent molecules in the solvation shell are concerted processes and reactions can be described by one reaction coordinate. If the thermal fluctuations of the solvent are not fast enough to maintain the thermal solvation equilibrium between reactants and activated complex, then the solvent rearrangement during the activation process can be rate-determining. The *cis* → *trans* isomerization reactions at N=N and C=N double bonds of azobenzenes and azomethines are effected by inversion at nitrogen [*cf.* Eq. (5-40) in Section 5.3.2] and the activation volumes are close to zero as expected. However, when azobenzene is 4,4′-donor/acceptor-substituted, the reaction proceeds by rotation through a dipolar activated complex in polar solvents, and is accompanied by the expected large negative activation volume. The rates of both reactions are independent of the viscosity of the solvent. These unimolecular isomerization reactions have activation energies of 50–70 kJ/mol and they are slow enough to proceed in the TST-valid solvation equilibrium region at normal pressure. However, in viscous solvents (*e.g.* 2-methylpentane-2,4-diol, glycerol triacetate), increasing the external pressure leads first to a normal rate behaviour, and then, at higher pressures, this becomes a strong pressure-dependent rate retardation. This is best explained by a shift of the reaction from the TST-valid region at lower pressure to the TST-invalid region at higher pressure. Under these circumstances, the reactant molecule has to wait for the formation of the solvation shell that stabilizes the activated complex before it undergoes the isomerization. The solvent rearrangement is then the rate-determining step and this rate constant was found to correlate with solvent viscosity. Thus, for the reaction systems studied, the chemical transformation and the solvent rearrangement have to be described by two independent reaction coordinates [754].

Finally, it should be mentioned that external pressure has found more and more application in organic synthesis, not least because of the increasing commercial availability of high-pressure devices. For the following types of organic reactions, a rate enhancement with increasing pressure is expected [749]: (a) associative reactions in which the number of molecules decreases in forming the products; *e.g.* cycloaddition and condensation reactions; (b) reactions which proceed via cyclic isopolar activated complexes; *e.g.* Cope and Claisen rearrangements; (c) reactions which proceed *via* dipolar activated complexes; *e.g.* Menschutkin-type S_N2 reactions, aromatic electrophilic

substitution reactions; and (d) reactions with steric hindrance. Comprehensive reviews on organic syntheses under high external pressure can be found in references [430, 749].

Whilst dealing with pressure-dependent chemical reactions, the use of supercritical fluids (sc-fluids) as reaction media should be briefly mentioned here; see Table 3-4 and Figure 3-2 in Section 3.2, as well as references [757, 758]. An important property of supercritical fluids as solvents is the possibility of manipulating the physicochemical properties of these solvents through small changes in *pressure* and temperature. This can influence solubilities, mass transfer (diffusivity), and rate constants of the reacting systems which are dissolved in supercritical solvents. The effect of external pressure on reaction kinetics measured in supercritical solvents can occur through the pressure-dependence of concentrations of the reactants or through the pressure-dependence of the rate constants.

The Diels–Alder cycloaddition reaction of maleic anhydride with isoprene has been studied in supercritical-fluid CO_2 under conditions near the critical point of CO_2 [759]. The rate constants obtained for supercritical-fluid CO_2 as solvent at 35 °C and *high* pressures (>200 bar) are similar to those obtained using normal liquid ethyl acetate as the solvent. However, at 35 °C and pressures approaching the critical pressure of CO_2 (7.4 MPa), the effect of pressure on the rate constant becomes substantial. Obviously, $\Delta V^{\neq}$ takes on large negative values at temperatures and pressures near the critical point of CO_2. Thus, pressure can be used to manipulate reaction rates in supercritical solvents under near-critical conditions. This effect of pressure on reacting systems in sc-fluids appears to be unique. A discussion of fundamental aspects of reaction kinetics under near-critical reaction conditions within the framework of transition-state theory can be found in reference [759].

5.5.12 Solvent Isotope Effects

Solvent isotope effect (SIE) is a term frequently used to describe changes in kinetic and equilibrium processes produced by replacing a normal solvent by its isotopically substituted counterpart. Since replacement of hydrogen by deuterium gives the largest relative mass change and easily measurable results from a perturbation of those molecular properties which are sensitive to mass, the ratio of a measurement X in light water (H_2O) to the corresponding value in heavy water (D_2O), X_{H_2O}/X_{D_2O}, is usually what is meant by the term solvent isotope effect of that property [446–449, 760, 761]. Comparatively little is known of solvent isotope effects in solvents other than water (*e.g.* CH_3OH/CH_3OD, CH_3CO_2H/CH_3CO_2D) [447]. Kinetic solvent isotope effects (KSIE), k_{H_2O}/k_{D_2O}, range from 0.5 to about 6, with the most common values falling between 1.5 and 2.8 [447]. The isotope effect for reactions carried out in an isotopically substituted solvent can be used to indicate direct or indirect solvent participation in the reaction. Unfortunately, the observed effects are a combination of three factors: (a) the solvent can be a reactant; for instance, if an O—H or O—D bond of the solvent is broken in the rate-determining step, there will be a *primary isotope effect*; (b) the reactant molecules may become labelled with deuterium by a fast H/D exchange reaction, and then the newly labeled molecule can cleave in the rate-determining step; (c) the intermolecular solute/solvent interactions (*i.e.* the solute solvation) may be different in the labelled and

Table 5-28. Some important physical properties of light and heavy water [446, 451, 635].

Property	Unit	Value for H_2O	Value for D_2O
Relative molecular mass, M_r	$g \cdot mol^{-1}$	18.015	20.031
Melting point (at 1013 hPa), t_m	°C	0.00	3.82
Boiling point (at 1013 hPa), t_{bp}	°C	100.00	101.42
Temperature of maximal density, t	°C	4.0	11.2
Density, ρ[a)]	$g \cdot cm^{-3}$	0.997047	1.10448
Molar volume, V_m[a)]	$cm^3 \cdot mol^{-1}$	18.069	18.133
Viscosity, η[a)]	mPa · s	0.912	1.121
Vapour pressure, p[a)]	kPa	3.170	2.740
Molar heat of vaporization, ΔH_v[a)]	$kJ \cdot mol^{-1}$	43.869	46.375
Refractive index, n_D[a)]	1	1.33250	1.32841
Relative permittivity, ε_r[a)]	1	78.46	78.06
Dipole moment, μ	C · m	$6.12 \cdot 10^{-30}$	$6.14 \cdot 10^{-30}$
Ionization constant, K_w[a)]	$mol^2 \cdot L^{-2}$	$1.008 \cdot 10^{-14}$	$0.112 \cdot 10^{-14}$
Solubility parameter, δ[a)]	$MPa^{1/2}$	47.9[b)]	48.7
Solubility of NaCl in moles of salt per 55.5 moles of solvent[a)]	$mol \cdot kg^{-1}$	6.1	5.8
Critical temperature, t_C	°C	374.0	370.7
Critical pressure, p_C	MPa	22.055	21.941
Critical volume, V_C	$cm^3 \cdot mol^{-1}$	55.9	56.3
Critical density, ρ_C	$g \cdot cm^{-3}$	0.322	0.356

[a)] At 25 °C.
[b)] *Cf.* Table 3-3 in Section 3.2.

nonlabelled solvent. This can change the differential solvation of educts and activated complex and hence the Gibbs energy of activation of the reaction. This is called a *secondary isotope effect*. In many cases, not only the first and third factors, but often the second as well, are simultaneously operative and it is difficult to separate them [762].

In Table 5-28 a number of the physical properties of light and heavy water are compared [446, 451, 635].

Whereas light and heavy water have nearly identical relative permittivities and dipole moments, it can be concluded from the greater boiling point, heat of vaporization, density, and viscosity of heavy water, that liquid D_2O is more structured than the already highly structured H_2O at room temperature (*cf.* Fig. 2-1 in Section 2.1). This is also consistent with the fact that salts acting as structure-breakers are generally less soluble in D_2O than H_2O [446].

Bearing this in mind, and assuming that the direction and magnitude of medium-influenced kinetic solvent isotope effects of type (c) are determined by the different solvation of reactants and activated complex, Swain and Bader draw the following conclusion [450]*): A reaction which destroys the structure of water during formation of the activated complex (by creation of charge) will proceed more rapidly in light water. A reaction which returns structure to the solvent when the activated complex is reached (by neutralization or dispersal of charge) will exhibit an enhanced rate in heavy water.

* Another view of the origin of solvent isotope effects has been given by Bunton and Shiner [763]; see also [764].

In addition to these non-specific effects, there also can exist differences in specific interactions such as hydrogen bonding between reactants or activated complex and the solvent. The rather scarce experimental results available confirm these simple predictions only in part. Most of the systems studied involve acid or base catalysis and consequently the participation of water as a reactant. Thus, the observed solvent isotope effects often include a composite of primary and secondary isotope effects, because any protons which are exchangeable with solvent protons or deuterons give rise to primary and secondary isotope effects. For example, in the autoprotolysis of water according to $2\ L_2O \rightleftharpoons L_3O^+ + LO^-$ (L = H, D) with $K_w(H_2O)/K_w(D_2O) = 9.0$ (*cf.* Table 5-28), the bond from oxygen to one of the isotopically substituted atoms L is broken (=primary isotope effect) and the bond from oxygen to the other hydrogen remains intact so that substitution here contributes to a secondary isotope effect. As the whole reaction takes place in water ($H_2O \rightarrow D_2O$), there is a solvent isotope effect as well.

Since proton-transfer is an integral part of kinetic solvent isotope effects in acid- and base-catalyzed reactions, and the solvent is involved in the reaction as a reaction partner, further treatment of this topic lies beyond the scope of this book. The reader is therefore referred to some excellent and comprehensive reviews relating to solvent isotope effects [20, 446–449, 760, 761].

5.5.13 Reactions in Biphasic Solvent Systems and in Neoteric Solvents

The increasing awareness about environmental protection and safety at work has led to an increasing search for safer procedures and safer solvents during the last decade. The efforts in this respect are often summarized by the key phrase *green chemistry*. Green chemistry is dedicated to environmentally benign chemical syntheses and industrial processing, and includes sustainable development as a desirable goal for achieving economic and societal objectives [876]. A few illustrative examples of these developments, that is, reactions in biphasic solvent systems and in neoteric solvents such as perfluorohydrocarbons, supercritical carbon dioxide, and room temperature liquid salts, are given in this section. Neoteric (=recent, new, modern) indicates a class of novel reaction media with remarkable new properties; see reference [167] to Chapter 3. Some chemical and physical properties of perfluorohydrocarbons, ionic liquids, and supercritical fluids have already been described in Sections 3.1 (Table 3-1) and 3.2 (Table 3-4), and recommendations for the substitution of hazardous solvents by safer ones are given in Section A.10 (Table A-14). Recent reviews on the substitution of solvents by safer media and processes can be found in reference [877].

Conventionally, organic syntheses are carried out in homogeneous media. Often, however, is it difficult to find solvents in which covalent organic compounds and ionic inorganic reagents (bases, nucleophiles, oxidizing agents, *etc.*) as well as catalysts are sufficiently soluble to obtain a homogeneous reaction mixture. Furthermore, isolation of the product(s) can entail a fairly lengthy work-up procedure, involving quenching with a large amount of water, distillative removal of the reaction solvent, followed either by isolation of the product by filtration and washing, or else multiple extraction with a water-immiscible solvent, washing the extracts with water, drying, and evaporation of the extraction solvent – to mention only a few possibilities.

One of the oldest techniques for overcoming these problems is the use of biphasic water/organic solvent systems using phase-transfer methods. In 1951, Jarrouse found that the reaction of water-soluble sodium cyanide with water-insoluble, but organic solvent-soluble 1-chlorooctane is dramatically enhanced by adding a catalytic amount of tetra-*n*-butylammonium chloride [878]. This technique was further developed by Makosza *et al.* [879], Starks *et al.* [880], and others, and has become known as liquid-liquid *phase-transfer catalysis (PTC)*; for reviews, see references [656–658, 879–882]. The mechanism of this method is shown in Fig. 5-18 for the nucleophilic displacement reaction of a haloalkane with sodium cyanide in the presence of a quaternary ammonium chloride as PT catalyst.

Sodium cyanide and the haloalkane are soluble in water and in the organic phase, respectively. In the aqueous phase, Na^+CN^- reacts with water-soluble Q^+Cl^- to produce Q^+CN^-, which is also soluble in the water-saturated organic phase. Then, this Q^+CN^- reacts in the organic phase with R–Cl to yield R–CN and Q^+Cl^-, which transfers back to the aqueous phase, ready for further reaction cycles. This procedure profits from the fact that quaternary ammonium salts (mostly chlorides, bromides, or hydrogen sulfates) with longer *N*-alkyl chains (usually with eight or more carbon atoms) have affinity for both the polar aqueous and the nonpolar organic phase. In order to minimize hydration of the nucleophiles in the organic phase, thereby achieving the best results, nonpolar solvents such as dichloromethane, 1,2-dichloroethane, toluene, or benzene are used, which are practically immiscible with water. According to Fig. 5-18, this PTC reaction involves two reactions (one in the aqueous and one in the organic phase), transfer of Q^+CN^- from the aqueous to the organic phase, and transfer of Q^+Cl^- from the organic to the aqueous phase, as well as equilibrium partitioning of Q^+CN^- and Q^+Cl^- between the aqueous and organic phases. Thus, the overall reaction rate depends on the intrinsic rate constants in the aqueous and organic phases, the mass-transfer rate of Q^+CN^- and Q^+Cl^-, and the equilibrium partition coefficients of Q^+CN^- and Q^+Cl^-, which are all affected by the interaction of the components and their environments. The organic solvent provides the medium for interaction of the reactants and plays, therefore, the dominant role in influencing the PTC reaction rate



Fig. 5-18. Phase-transfer catalyzed nucleophilic displacement reaction of a haloalkane, R–Cl, with sodium cyanide, Na^+CN^-, to yield a nitrile, R–CN, in the presence of sub-stoichiometric amounts of a quaternary ammonium chloride, Q^+Cl^-, as PT catalyst.

and the degree of conversion of the reactants. The polarity of the organic phase, in conjunction with the structure of the anion and the catalyst cation, affects the selectivity of the PT catalyst partitioning into the organic phase. For example, increasing the polarity and hydrogen-bonding ability of the organic phase has a favourable effect on the extraction of anions with high charge density (=high charge/volume ratio). An excellent review on the influence of organic solvents on a great variety of PTC reactions has recently been given by Wang [882].

The advantages of PTC reactions are moderate reaction conditions, practically no formation of by-products, a simple work-up procedure (the organic product is exclusively found in the organic phase), and the use of inexpensive solvents without a need for anhydrous reaction conditions. PTC reactions have been widely adopted, including in industrial processes, for substitution, displacement, condensation, oxidation and reduction, as well as polymerization reactions. The application of chiral ammonium salts such as *N*-(9-anthracenylmethyl)cinchonium and -cinchonidinium salts as PT catalysts even allows enantioselective alkylation reactions with *ee* values up to 80–90%; see reference [883] for a review. Crown ethers, cryptands, and polyethylene glycol (PEG) dialkyl ethers have also been used as PT catalysts, particularly for solid–liquid PTC reactions; *cf.* Eqs. (5-127) to (5-130) in Section 5.5.4.

Another particularly interesting biphasic solvent system is the combination of nonpolar perfluoro-substituted alkanes, dialkyl ethers, or trialkylamines (so-called *fluorous solvents*) with more polar, conventional organic solvents. The term *fluorous* was introduced in analogy to the term *aqueous* [885]. Fluorous solvents are nonpolar, hydrophobic, chemically inert, and non-toxic, with a higher density than the corresponding non-fluorinated solvents ($\rho \approx 1.7$–$1.9\ g \cdot cm^{-3}$). They usually have a limited, temperature-dependent miscibility with conventional organic solvents, forming biphasic solvent systems at ambient temperature. With different solubilities for educt, reagent, catalyst, and product, biphasic solvent systems with a fluorous phase can facilitate the separation of the product from the reaction mixture. The fluorous biphase system (FBS) consists of a fluorous phase preferentially containing fluorous-soluble nonpolar educts, reagent, and catalyst, and a second phase preferentially containing polar educts and the product, which is usually an organic solvent with limited solubility in the fluorous phase. Reagents and catalysts can be made fluorous phase-soluble by the introduction of perfluoroalkyl groups of appropriate length and number, mostly of the type F_3C–$(CF_2)_n$–$(CH_2)_2$– (so-called *fluorous ponytails* or *fluorous tags*). The insulating ethylene spacer –$(CH_2)_2$– is necessary to attenuate the strong ($-I$)-effect of the strongly electronegative fluorine substituents, which can otherwise change the chemical properties of the reagent and catalyst in an undesirable manner. Such fluorous biphase systems are well-suited for converting nonpolar educts to products of higher polarity, as the partition coefficients of educts and products will be higher and lower, respectively, in the fluorous phase. As a consequence, there is little or no solubility limitation on the educts and easy separation of the products. A fluorous biphase reaction can proceed either in the fluorous phase or at the interface between the two phases, depending on the solubility of the educts in the fluorous phase.

Furthermore, some fluorous biphase systems can become a single phase with increasing temperature. For example, *n*-heptane and perfluoro(methylcyclohexane) are immiscible at room temperature (*ca.* 20 °C) and form a two-phase system. At higher



Fig. 5-19. Catalyzed chemical reaction between polar educt A and nonpolar educt B and a reagent in a biphasic solvent system with temperature-dependent mutual miscibility of the polar and nonpolar (fluorous) solvents. A more detailed illustration of the experimental possibilities for catalysis in fluorous solvents is given in reference [890].

temperatures (*ca.* 40 °C), these solvents become miscible and form a monophasic binary solvent mixture, which can serve as a homogeneous reaction medium for educts, reagent, and catalyst, and the catalytic reaction can take place. After completion of the reaction, cooling down the reaction mixture leads to formation of two separate phases once more. In the ideal case, the product is now dissolved in one phase, the remaining educts, reagent, and catalyst in the other, making the isolation of the product very easy. The isolation and re-use of the often rather expensive catalyst is also made easier by this method. For a hypothetical catalyzed reaction, this procedure is illustrated in Fig. 5-19.

This new experimental technique, using *fluorous solvents* or *fluorous biphasic systems* (FBS) with *fluorous biphase catalysis* (FBC), was developed by Vogt and Kaim [884] and by Horváth and Rábai [885] in 1991 and 1994, respectively. Since then, this method has found many applications in synthetic organic chemistry and has already been reviewed repeatedly [886–893]. Incidentally, temperature-dependent two-phase → one-phase transitions are not limited to combinations of fluorous solvents with organic solvents. For example, certain mixtures of water and 1-cyclohexylpyrrolidin-2-one form one phase at ambient temperature and a two-phase system at higher temperatures (>*ca.* 50 °C), also allowing interesting separation possibilities.

The reason why fluorous alkanes are immiscible with normal alkanes possibly stems from their different conformations: *n*-alkanes exist in well-known zig-zag conformations, whereas perfluoro-*n*-alkanes adopt more helical conformations because of the larger van der Waals radius of fluorine ($r = 135$ pm) as compared to that of hydrogen ($r = 120$ pm). Molecules of fluorous solvents are also subject to very weak van der Waals interactions due to the low polarizability of the electrons of the CF_2 groups. As a

consequence of these particular conformations, which have many cavities to accommodate solutes, as well as the low degree of solvent/solvent interactions, perfluorohydrocarbons exhibit good solubilities for gases such as O_2, N_2, H_2, and CO_2 [*e.g.* up to 57 mL of O_2 in 100 mL of perfluoro(methylcyclohexane)]. Therefore, fluorous solvents are good media for aerobic oxidation reactions.

For example, the aerobic catalytic oxidation of alcohols ($R–CH_2OH$ or $R_2CHOH \rightarrow R–CHO$ or $R_2C{=}O$), aldehydes ($R–CH{=}O \rightarrow R–CO_2H$), sulfanes ($R_2S \rightarrow R_2SO$ or R_2SO_2), and cyclic alkenes ($R–CH{=}CH–R \rightarrow$ epoxides), using fluorous biphasic systems and specially designed perfluoroalkyl-substituted transition metal complexes (Ni, Ru, Cu) as catalysts, has been studied by Knochel *et al.* [894, 895]. The oxidation of 4-chlorobenzaldehyde with oxygen (at *ca.* 1 bar) in the presence of an Ni catalyst (3 cmol/mol) in monophasic toluene/perfluorodecalin (1:1) at 64 °C yielded, after 12 hours of stirring, 87 cmol/mol of 4-chlorobenzoic acid. After cooling to room temperature and the formation of two phases, the acid deposited as a precipitate could be isolated by filtration, and the catalyst-containing fluorous phase could be separated and re-used for further oxidations. After six reaction cycles, the yield of the acid was still 70 cmol/mol [894]. The aerobic oxidation of 4-nitrobenzyl alcohol in the presence of catalytic amounts of a perfluoroalkyl-substituted copper(I) complex and TEMPO (2,2,6,6-tetramethylpiperidine-1-oxide) has been carried out in biphasic chlorobenzene/perfluorooctane at 90 °C under a gentle stream of oxygen, to give 4-nitrobenzaldehyde in 93 cmol/mol yield. After cooling, the two phases were separated, the product was isolated from the chlorobenzene layer, and the catalyst-containing fluorous phase was directly re-used for further reaction runs. Even after eight cycles, 86 cmol/mol of analytically pure 4-nitrobenzaldehyde could still be isolated [895].

These two simple examples demonstrate the usefulness of the FBC procedure, combining the advantages of both homogeneous and heterogeneous catalysis: homogeneous reaction conditions due to the miscibility of fluorous solvents with conventional organic solvents only at higher temperatures, and the facile separation of the usually expensive catalyst from the biphasic system formed at lower temperatures, allowing its re-use for further reaction cycles. This method also avoids the contamination of the product phase with traces of metal stemming from the catalyst. In contrast to aqueous biphase catalysis, fluorous biphasic systems are compatible with moisture-sensitive compounds such as organometallics. Further examples can be found in references [885–893], particularly in [891]; a *user's guide* to practical applications of fluorous catalysts and reagents is given in reference [890].

Although perfluorinated solvents are still expensive, the advantages of FBS and FBC also make this method of interest for industrial applications. Of particular importance in this respect is the conversion of terminal alkenes into aldehydes containing one more carbon than the starting alkene, by reacting the alkene with synthesis gas (CO/H_2) in the presence of cobalt or rhodium catalysts in a homogeneous phase (hydroformylation, oxo, or Roelen reaction). According to Horváth *et al.* [885], this hydroformylation reaction can be advantageously carried out in the fluorous biphasic system toluene/perfluoro(methylcyclohexane), using conventional rhodium catalysts but with tris-(perfluoroalkyl)-substituted phosphanes as ligands. The 1-alkene substrate is soluble in cold toluene and the rhodium catalyst in cold perfluoro(methylcyclohexane). By heating this reaction mixture to 100 °C in the presence of CO/H_2 in an autoclave, a homoge-

neous phase is formed in which the transition metal catalyzed reaction takes place. Cooling to room temperature results in separation of the aldehyde product (in the organic phase) from the fluoro-ponytailed catalyst (in the fluorous phase). This FBS-based method has some advantages over the modern aqueous biphasic oxo reaction, called the Ruhrchemie/Rhône-Poulenc process, in which water-soluble rhodium catalysts with triply-sulfonated triarylphosphane ligands are used; for a detailed discussion, see reference [896].

Another remarkable class of solvents are *ionic liquids* (molten salts), which can be used as monophasic solvents or in biphasic solvent systems. Some of their chemical and physical properties have already been mentioned in Section 3.1; see Fig. 3-1 and Table 3-1 in Section 3.1, as well as references [3, 24–30, 112–114, 167, 187–191] to Chapter 3. Ionic liquids have been known since 1914, when Walden discovered ethylammonium nitrate ($EtNH_3^+NO_3^-$), which has a melting point of just 12 °C; see reference [156] to Chapter 3. Subsequently, high-temperature melting inorganic salts and eutectic inorganic salt mixtures were used as reaction media for many inorganic and organic reactions. Three examples of the application of such inorganic eutectics as solvents for organic reactions have already been given in Section 3.1. However, for the most part, this approach has not been very successful, often because of corrosion problems with the reaction vessels. For a long time, ionic liquids were only of interest to electrochemists as electrolytes. The rediscovery of room temperature liquid salts, this time consisting of large organic cations such as 1,3-dialkylimidazolium, 1,3- or 1,4-dialkylpyridinium, tetraalkylammonium, and tetraalkylphosphonium, in combination with mostly inorganic anions such as Cl^-, Br^-, BF_4^-, PF_6^-, $F_3C–CO_2^-$, $F_3C–SO_3^-$, $(F_3C–SO_2)_2N^-$, and $AlCl_4^-$, with melting points under 100 °C (often below room temperature) has, during the last decade, led to a renaissance in the application of these ionic liquids as reaction media. Ambient temperature ionic liquids based on 1-alkyl-3-methylimidazolium salts were first reported by Wilkes *et al.* in 1982 as tetrachloroaluminates; see reference [162a] to Chapter 3. Replacement of the moisture-sensitive $AlCl_4^-$ ion by the BF_4^- ion and other stable anions led, in 1992, to air- and water-stable, room temperature ionic liquids (see ref. [162b] to Chapter 3), which are now finding increasing application as reaction media for a great variety of chemical reactions, mainly due to the work of Seddon, Hussey, Chauvin, and others; see references [167, 190] and reference [187] to Chapter 3, as well as [905, 906].

Ionic liquids have very good dissolution properties for most inorganic and organic compounds, high thermal stabilities, densities greater than unity ($\rho > 1\ g \cdot cm^{-3}$), no measurable vapour pressure, and they are non-flammable. The two last-mentioned properties in particular make ionic liquids promising as "green solvents for the future" [897, 898]. Further advantages are the optimization of physical solvent characteristics and coordinating abilities through a broad choice of cation/anion combinations, reaction rate enhancements because of missing reactant solvation shells, higher reaction selectivities, and often better yields. Because of the manifold cation/anion combinations, these solvents can be constructed to possess a particular set of desired properties and are therefore also called *tailor-made* or *designer* solvents [898]. For example, the miscibility of ionic liquids with water changes with their structure: 1-alkyl-3-methylimidazolium tetrafluoroborates with alkyl chain lengths of less than six carbon atoms are miscible with water at 25 °C, but at or above six carbon atoms they form a separate phase when

mixed with water. The ionic liquid 1-(*n*-butyl)-3-methylimidazolium hexafluorophosphate, $[BMIM]^+PF_6^-$, even forms triphasic mixtures with water and alkanes. This multiphasic behaviour of some ionic liquids is analogous to that of fluorous biphasic systems and has important implications for clean synthetic processes. Using biphasic solvent systems consisting of an immiscible covalent organic solvent and ionic liquid, organic products can easily be separated from the educts, reagent, and catalysts, which remain dissolved in the ionic phase after the reaction, by decantation of the organic phase or by extraction of the ionic phase.

Ionic liquids can be used as replacements for many volatile conventional solvents in chemical processes; see Table A-14 in the Appendix. Because of their extraordinary properties, room temperature ionic liquids have already found application as solvents for many synthetic and catalytic reactions, for example nucleophilic substitution reactions [899], Diels–Alder cycloaddition reactions [900, 901], Friedel–Crafts alkylation and acylation reactions [902, 903], as well as palladium-catalyzed Heck vinylations of haloarenes [904]. They are also solvents of choice for homogeneous transition metal complex catalyzed hydrogenation, isomerization, and hydroformylation [905], as well as dimerization and oligomerization reactions of alkenes [906, 907]. The ions of liquid salts are often poorly coordinating, which prevents deactivation of the catalysts.

Only two examples will be mentioned here in more detail. The S_N2 alkylation reaction of ambident potassium 2-naphtholate (2-naphthol + KOH) with 1-bromobutane in $[BMIM]^+PF_6^-$ at room temperature proceeds at a similar rate as observed in dipolar aprotic solvents such as DMF or DMSO, to regioselectively afford 1-butyl 2-naphthyl ether in 98 cmol/mol isolated yield (*O*/*C* alkylation ratio > 99:1) [899]. In contrast to the reaction in dipolar aprotic solvents, which are difficult to remove from the product, the ether product can be simply extracted into an organic solvent such as toluene, leaving the ionic liquid behind. The by-product (potassium bromide) of the reaction can be extracted with water, and the ionic liquid can be used again.

For Friedel–Crafts acylations of aromatic compounds, a Lewis acid catalyst (*e.g.* $AlCl_3$) is necessary, which can be delivered by the chloroaluminate(III) ion of the ionic liquid used as the reaction medium [902, 903]. Mixing 1-ethyl-3-methylimidazolium (EMIM) chloride with aluminium trichloride results in the equilibrium formation of the ambient-temperature ionic liquid $[EMIM]^+AlCl_4^-$. The Lewis acid/base properties of this system are described by the equilibrium $2\ AlCl_4^- \rightleftharpoons Al_2Cl_7^- + Cl^-$, where $Al_2Cl_7^-$ is the Lewis acid and Cl^- the Lewis base. The composition of such tetrachloroaluminate(III) ionic liquids is best described by the apparent mole fraction x of added $AlCl_3$: ionic liquids with $x(AlCl_3) < 0.5$ contain an excess of Cl^- ions over $Al_2Cl_7^-$ ions and are called 'basic', those with $x(AlCl_3) > 0.5$ containing an excess of $Al_2Cl_7^-$ over Cl^- are called 'acidic', and salt melts with $x(AlCl_3) = 0.5$ are called 'neutral'. In Lewis acidic mixtures of $[EMIM]^+Cl^-$ and $AlCl_3$ with $x(AlCl_3) > 0.5$, benzene reacts quantitatively with acetyl chloride at room temperature to give acetylbenzene as the only product in less than five minutes [902]. The proposed reaction mechanism includes the formation of the intermediate electrophile according to $H_3C–CO–Cl + Al_2Cl_7^- \rightleftharpoons H_3C–CO^+ + 2\ AlCl_4^-$. Acylation of naphthalene with acetyl chloride in $[EMIM]^+Cl^-/AlCl_3$ with $x(AlCl_3) = 0.67$ within 5 min at 0 °C regioselectively affords 89 cmol/mol of 1-acetylnaphthalene and only 2 cmol/mol of the corresponding 2-isomer [903].

Further illustrative examples of the application of ionic liquids as reaction media can be found in the recent reviews of Welton (ref. [188] to Chapter 3), Wasserscheid and Keim (ref. [189] to Chapter 3), Seddon [897], and Freemantle [898].

In addition to fluorous solvents and ionic liquids, *supercritical fluids* (*sc-fluids*, *scf's*), such as supercritical carbon dioxide (sc-CO_2), constitute a third class of neoteric solvents that can be used as reaction media. Although sc-fluids have been known for a long time and have been advantageously used as eluants in extraction and chromatography processes (see Sections A.6 and A.7 in the Appendix), their application as reaction media for chemical processes has become more popular only during the last decade. Some of their physical properties and the supercritical conditions necessary for their existence have already been described in Section 3.2 (see Figure 3-2 and Table 3-4); see also references [209, 211–220, 224–230] to Chapter 3 for reviews on sc-fluids and their applications (particularly for sc-CO_2 and sc-H_2O).

When carbon dioxide is heated beyond its critical point, with a critical temperature of $t_C = 31.0\ ^\circ C$, a critical pressure of $p_C = 7.38$ MPa, and a critical density of $\rho_C = 0.47\ g \cdot cm^{-3}$, the gaseous and the liquid phase merge into a single supercritical phase (sc-CO_2) with particular new physical properties: very low surface tension, low viscosity, high diffusion rates, pressure-dependent adjustable density and solvation capability (''solvation power''), and miscibility with many reaction gases (H_2, O_2, *etc.*). It can dissolve solids and liquids. The relative permittivity of an sc-fluid varies linearly with density, *e.g.* for sc-CO_2 at 40 °C, $\varepsilon_r = 1.4 \rightarrow 1.6$ on going from 108 to 300 bar. This small change in the already low relative permittivity, together with the empirical determination of solvent polarity by solvatochromic dyes, indicates that pure sc-CO_2 is always a fluid of very low polarity, comparable with that of hydrocarbons such as cyclohexane (see refs. [221, 222] to Chapter 3). The polarity of sc-CO_2 can be substantially increased by the addition of even small amounts of polar co-solvents. Carbon dioxide has no permanent dipole moment, but a significant quadrupole moment, which contributes to solute dipole/solvent quadrupole interactions. Quadrupolar solvent effects on solvation and reactivity of solutes dissolved in sc-CO_2 have been studied in detail [908].

It should be noted that the local density of an sc-fluid around solute molecules is subject to large fluctuations and can differ considerably from its bulk density. The region near a solute molecule is quite dynamic, with solvent molecules rapidly entering and leaving this area, but nonetheless enhancing the local density. As a result, the solubility of reactants in sc-fluids can be varied by adjusting the pressure and temperature, whereby higher density generally corresponds to higher solubility. Appreciable solubilities in pure sc-fluids usually require densities of $\rho > \rho_C$, and supercritical fluids are only suitable as reaction media if they fulfil this criterion. For the solubility of reactants in sc-fluids, three factors are important. The solubility increases with increasing vapour pressure of the solute. Solutes of low polarity are more soluble than polar or ionic compounds. The solubility in sc-CO_2 can be enhanced by the introduction of ''CO_2-philic'' perfluoroalkyl or polysiloxane substituents into the reactants, reagents, and/or catalysts [909]. Many polyfluoro-substituted polymers are highly soluble in both liquid and sc-CO_2. Addition of reactants, reagents, and catalysts to an sc-fluid can change its phase behaviour. Therefore, visual control of the reaction mixture in a window-equipped reactor is always necessary, in order to make sure that a single homogeneous phase is still present after addition of the substrates.

The high compressibility of sc-fluids allows continuous variation of their densities and related properties from gaseous to liquid-like values with comparatively small variations in temperature and/or pressure. In this way, the positions of equilibria and the rates of chemical reactions can be continuously changed ("reaction tuning"), as shown by the following two examples.

The azo/hydrazone tautomeric equilibrium 4-phenylazo-1-naphthol $\rightleftharpoons$ 1,4-naphthoquinone-1-phenylhydrazone [*cf. (16a)* $\rightleftharpoons$ *(16b)* in Section 4.3.2] has been studied in various liquids and in liquid and supercritical carbon dioxide [910]. The less dipolar azo tautomer ($\mu \approx 6 \cdot 10^{-30}$ Cm) is dominant in the dilute gas phase, in compressed ethane, and in liquid *n*-hexane. However, in liquid and in sc-CO_2 (at 35 °C and 27 MPa), the tautomeric equilibrium is shifted towards the more dipolar hydrazone form ($\mu \approx 17 \cdot 10^{-30}$ Cm), to give similar amounts of the two tautomers. This shift can be attributed to the quadrupole moment of CO_2 and its Lewis acidity. If only dipole/induced dipole forces were operative, similar results would be expected for sc-CO_2 and ethane [910]. Incidentally, the electron-accepting, Lewis acidic property of CO_2 is nicely demonstrated by its chemical reaction with dimethylamine to yield the ionic liquid dimethylammonium dimethylcarbamate (Dimcarb; *cf.* Section 3.1, Table 3-1, entry 28).

The rate constants of the unimolecular thermolysis of α-chlorobenzyl methyl ether to yield benzaldehyde and chloromethane [*cf.* Eq. (5-41) in Section 5.3.2] have been determined in supercritical 1,1-difluoroethane ($t_C = 113$ °C; $p_C = 4.5$ MPa). Near the critical point at 130 °C, the rate constant increases by about two orders of magnitude on increasing the pressure from 4.48 to 345 MPa [911]. At 130 °C and 30 MPa, the rate constant is comparable to that obtained in conventional tetrachloromethane. The activation volume found for this reaction is $\Delta V^{\neq} = -6$ L/mol, which suggests that the sc-solvent clusters are more ordered around the activated complex than around the reactant. This is in agreement with the formation of a highly dipolar activated complex, which is stabilized by solvation through dipolar $H_3C–CHF_2$. The rate constant of this reaction can be adjusted over a continuum with a single sc-fluid instead of a series of common liquid solvents. Thus, this example nicely demonstrates how supercritical solvents can be used to bridge, over a continuum, the gap between gas-phase and liquid-phase kinetics [911]. For reviews on solvation in supercritical fluids and on homogeneous organic reactions used as mechanistic probes in sc-fluids, see references [912, 913].

Supercritical fluids as solvents for *synthetic* organic reactions have found increasing interest only during the last decades. The earliest reported sc-fluid organic reaction seems to be the hydrolysis of aniline in sc-H_2O to give phenol and ammonia [914]. Meanwhile, a great variety of other organic reactions have been studied in sc-fluids, predominately in sc-CO_2 and sc-H_2O. Supercritical carbon dioxide and water are obviously the most attractive solvents for organic syntheses in both the laboratory and on a technical scale from the point of view of a "green chemist" [876]. They are environmentally and toxicologically benign, non-flammable, comparatively inert, inexpensive, and easy to recycle. However, the use of sc-H_2O demands rather harsh reaction conditions because of its high critical temperature and pressure ($t_C = 374$ °C; $p_C = 22.1$ MPa). Therefore, mainly sc-CO_2 has been used as an sc-fluid for organic syntheses, in particular for homogeneously and heterogeneously catalyzed hydrogenation reactions, hydroformylation of 1-alkenes, Pd-catalyzed C–C coupling reactions between halo-

arenes and vinylic substrates (including the Heck reaction), alkene metathesis reactions, oxidation reactions with molecular oxygen or peroxides, Diels–Alder cycloaddition reactions, homogeneous and heterogeneous polymerization reactions, free-radical chain reactions, enzymatically catalyzed reactions, and reactions with simultaneous use of CO_2 as an sc-fluid and a C_1 reaction partner. More recent compilations of synthetic organic reactions carried out in sc-fluids can be found in references [909, 915, 916]; see also references [211–220, 224–230] to Chapter 3. Only a few illustrative examples can be mentioned here.

The laboratory-scale hydrogenation of unsaturated organic compounds is usually slow because H_2 is only sparingly soluble in common solvents. In contrast, the high miscibility of sc-CO_2 with H_2 makes this sc-fluid very useful for such hydrogenation reactions using heterogeneous noble metal catalysts. Thus, the hydrogenation of cyclohexene in sc-CO_2 with a polysiloxane-supported Pd catalyst (5% Pd) in a continuous flow reactor is extremely rapid and gives cyclohexane in practically quantitative yield [917]. In addition to the high miscibility of sc-CO_2 and H_2, the reduced viscosity of sc-fluids and the increased diffusion rates therein enhances the transport to and from the catalyst, which is no longer a limiting factor. The ability to control the reaction conditions with great precision can be used to manipulate the selectivity of reactions carried out in sc-fluids. For example, the hydrogenation of acetylbenzene in sc-CO_2 (at 12 MPa) at 90 °C and a H_2/substrate ratio of 2:1 predominantly affords 1-phenylethanol (yield 90 cmol/mol) as the product. However, at 300 °C and a H_2/substrate ratio of 6:1, the more hydrogenated product ethylcyclohexane is formed (yield 90 cmol/mol) [917]. An excellent review on sc-fluids in heterogeneous catalysis can be found in reference [220] to Chapter 3.

A nice example of a homogeneous catalytic hydrogenation is the reduction of carbon dioxide to formic acid, carried out in sc-CO_2 in the presence of a soluble ruthenium(II)-trimethylphosphane catalyst and triethylamine at 50 °C and 21 MPa [918]. The turnover frequency (TOF = mole of product per mole of catalyst per hour) of this rapid reaction is rather high, with values up to 1400. This reaction, carried out at 50 °C in sc-CO_2, is 18 times faster than in conventional tetrahydrofuran under otherwise identical reaction conditions. This formic acid synthesis can be coupled with subsequent reactions: by addition of methanol or dimethylamine, this supercritical reaction system provides a highly efficient one-pot route to methyl formate and *N*,*N*-dimethylformamide, respectively [918]. Another example of a reaction in which carbon dioxide acts as both reactant and reaction medium is the formation of tetraethyl-2-pyranone from hex-3-yne and CO_2 in the presence of an Ni(II)-diphosphane catalyst at 102 °C under supercritical reaction conditions [919].

A further useful application of sc-CO_2 as a reaction medium is the free-radical side-chain bromination of alkylaromatics, replacing conventional solvents such as tetrachloromethane or chlorofluorohydrocarbons having no abstractable hydrogen atoms [920]. For example, bromination of ethylbenzene in sc-CO_2 at 40 °C and 22.9 MPa yields 95 cmol/mol (1-bromoethyl)benzene, with practically the same regioselectivity as obtained in conventional tetrachloromethane as the solvent. Even the classical Wohl–Ziegler bromination of benzylic or allylic substrates using *N*-bromosuccinimide (NBS) can be conducted in sc-CO_2 [920]. Irradiation of a solution of toluene, NBS, and AIBN (as initiator) in sc-CO_2 at 40 °C and 17.0 MPa for 4 hours gave (bromomethyl)-

benzene in quantitative yield, with no detectable amount of the potential side product 4-bromotoluene. In neither case is there any change in reaction mechanism on going from tetrachloromethane to sc-CO_2 as solvent [920].

The first Diels–Alder reaction conducted in sc-CO_2 was reported by Hyatt in 1984 [921]: cycloaddition of cyclopentadiene to methyl acrylate gave a mixture of methyl *endo*- and *exo*-5-norbornene 2-carboxylate in high yield at ambient temperature [*cf.* Eq. (5-43) in Section 5.3.3]. The solvent-dependent *endo*/*exo* product ratio found in sc-CO_2 (at 40 °C and 8.5 MPa) was of the same order as that found for liquid CO_2 and for hydrocarbons such as cyclohexane. According to Eq. (7-24) in Section 7.3, the logarithm of the *endo*/*exo* product ratio (called the Ω value) was introduced by Berson as an empirical measure of solvent polarity. A comparison of the Ω values from the *endo*/*exo* ratios obtained in sc-CO_2 places this sc-fluid close to hydrocarbons with respect to its polarity, in agreement with solvatochromic measurements (*cf.* references [221, 222] to Chapter 3). Numerous other Diels–Alder cycloaddition reactions have since been successfully studied in sc-CO_2. The reaction between cyclopentadiene and ethyl acrylate has been explored in dichloromethane, tetrahydrofuran, and *n*-hexane, as well as in sc-CO_2. The reaction rates found in the three nonpolar conventional solvents were slightly higher, but within a factor of two of those in sc-CO_2 [922]. For a more recent reinvestigation of the well-studied Diels–Alder reaction between isoprene and maleic anhydride in sc-CO_2, see reference [923].

As an example of the use of sc-CO_2 in an enzymatic reaction, the lipase-catalyzed esterification of oleic acid with racemic (±)-citronellol should be mentioned. At 31 °C and 8.4 MPa, the (−)-(*S*)-ester is formed enantioselectively in sc-CO_2 with an optical purity of nearly 100% [924]. The reaction rate is enhanced by increasing pressure, *i.e.* by increasing the solvation capability or solvent polarity of sc-CO_2. A linear correlation has been found between reaction rate and the solvatochromic solvent polarity parameter $E_T(30)$; see Section 7.4 for the definition of $E_T(30)$.

A great variety of chemical reactions have also been conducted advantageously in sc-H_2O, *e.g.* the hydrolysis of esters, amides, and nitriles, dehydration of alcohols (cyclohexanol → cyclohexene, glycerol → propenal, butane-1,4-diol → THF), partial oxidation with molecular oxygen (cyclohexane → cyclohexanone), to mention only a few possibilities. Excellent reviews on the chemistry in supercritical water can be found in references [224, 225, 228–230] to Chapter 3, as well as in a special report [925]. Some physical properties of sc-H_2O have already been described in Section 3.2. A gradual decrease of the density and relative permittivity of water with increasing temperature is paralleled by increasing water solubility of organic compounds. At 300 °C, the polarity and density of water roughly approach those of acetone at room temperature. The enhanced self-ionization of water at high temperatures makes it a stronger acid and a stronger base, thus promoting acid- and base-catalyzed reactions at higher temperatures. Thus, sc-H_2O frequently participates not only as a solvent and reactant, but also as a catalyst. The properties of supercritical ionophore (electrolyte) solutions have been studied by geochemists in order to understand mineral-forming processes deep in the Earth, where conditions can be supercritical. Because oxygen, carbon dioxide, methane, and other alkanes are completely miscible with sc-H_2O, even combustion (flameless or with flames!) can occur in this sc-fluid. Thus, oxidation in sc-H_2O can be used to detoxify hazardous organic materials (all kinds of waste, chemical warfare agents, *etc.*) nearly

quantitatively in short times. As the main oxidation products are water, carbon dioxide, and (for halo-organics) simple acids, the final aqueous mixture can mostly be disposed of without further treatment. Because of the rather harsh reaction conditions in sc-H_2O ($t_C = 374\ ^\circ C$, $p_C = 22.1$ MPa $= 218$ atm), the most promising application of sc-H_2O as a reaction medium seems to lie not in synthesis, but in the destruction of hazardous organic compounds [925].

In addition to supercritical water (sc-H_2O), *superheated water* (SHW) or *high-temperature water* (HTW) has recently attracted attention as a new medium for organic reactions. High-temperature water can be broadly defined as liquid water above 200 °C under autogenic pressure. The physical properties of superheated water, which are quite different from those of ambient liquid water, have already been described at the end of Section 3.2. Superheated water has a lower relative permittivity, fewer and weaker intermolecular hydrogen bonds, and a higher isothermal compressibility than normal liquid water. The solvent polarity of water at 300 °C is roughly equivalent to that of acetone at 25 °C ($\varepsilon_r = 20.6$), as measured by its relative permittivity. Whereas the solubility of most gases in ambient liquid water initially decreases with increasing temperature, at higher temperature a minimum is soon reached, and then the gas solubility increases. The physical and chemical properties of high-temperature water vary with temperature and pressure over wide ranges, approaching those of polar organic solvents at ambient temperature. Therefore, high-temperature water can support dipolar, isopolar, and free-radical transition state reactions, depending on the reaction conditions. In addition, high-temperature water can be used as medium for chemical syntheses, waste destruction, plastics recycling, coal liquefaction, and biomass processing. For recent reviews on the application of superheated or high-temperature water for scientific and engineering purposes see references [228–230] to Chapter 3.

etc.

)

chromophore

cf.

9 b

p ! p

l ¼ b

b

l ¼ n ¼

b

Solvents and Solvent Effects in Organic Chemistry, Third Edition Christian Reichardt
Copyright 8 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

solvatochromism

negative solvatochromism
positive solvatochromism)

s ! s
n ! s p ! p n ! p
p ! p n ! p
p
p
aromatic compounds polyenes polyynes polymethines cf.



p

e.g

peri-

chromism peri ¼ chroma¼

enzymichromism

...ðn þ Þp... n ¼ ; ; ; ;...
a a 0 ¼ H
; 0¼ N; O; P; S
¼ 0 ¼ 0¼ N ¼ 0¼ O
0 0 ¼ N 0¼ O

meropolymethine dyes

merocyanines

0

p
aa aa $ l aa aa m
n ¼ ; ; ;...

m

m

p

aromatic compounds

e.g. polyenes e.g.

polyynes e.g. poly-

methine dyes aromatic

p
l ¼ Dn¼ n

polyenicp b

p ! p l ¼ Dn¼

Dl ¼ !

polymethinicp



---

l

Dl

!



Dn¼ Dl ¼ p ! p

N

cf.

Dn¼ Dl ¼

p ! p

E cf.

N



a) i a)

I

!

a)

ca. ca.

N

N

tert

o o^{0}

e.g.

!

m < m . . .
m > m
. . .

m m

cf.

p ! p

0 0

Dn¼ Dl ¼ n

Dn¼ Dl ¼ t

cf.

Positive solvatochromic compounds						
				iso		
	c)	c)				
				n		
						
				n		
 Phenol blue						
				iso		
				n		

m a	m a	n	n	Dn	b)







Negative solvatochromic compounds



d) d)









m a	m a ñ	ñ	Dñ	b)
				
				
				
				
				
				

a) ¼ : m

b) Dñ ¼ ñ ñ
c)
d) t



cis



m > m m ? m

m ¼

m ¼

Z cf.

m < m

p

aa bb aa bb aa

0





i

l

Dl

l

Dl þ

0

d ! p

p ! d

cf.

Nonpolar solute in a nonpolar solvent.

n

n n þ

n Dn

e.g. e.g.

b e.g.

Nonpolar solute in a polar solvent.

n

e.g

Dipolar solute in a nonpolar solvent.

)

n

Dipolar solute in a polar solvent.

m < m

cf.

m 0 m

m ! m^{0}

ca. ca.

cf. 21

32



m < m i.e.

m > m

) m

)

(a)

(b)



electrochromism.

J bb

aa bb aa bb $ dl — — p — — dm m ¼ a)
(a) (b)

d

d b,c)
J =
J =
n b b) d)

a) 1958
b)
c) 261
d) c

(b)
(a) (c)
(a)

(b)

(a) p
(b) p

J
bb

$

(a) (b)

(c)

solvent polarity

(b)

e.g.

(b)

(c)

p ! p cf.



(a) (b) (c)

c

l

Dl þ

!

(a) (c)

(b)

(b)

m < m m > m

p ! p

p

ca.

trans ! cis

e.g.

a a

p ! p

!

p ! p n

et al.

g e

ħ ¼ħ ½ðm m j m Þ =a ½ðe Þ Lðn Þ

½ðm m Þ =a ½ðn Þ ð Þ

ħ ħ ħ a
m m
j LðeÞ
LðnÞ LðxÞ¼ðx þ ÞxÞ¼e n
e n
LðeÞ LðnÞ LðnÞ
m; m; j ; a

g e)

hcaDħ ¼ hcaðħ ħ Þ

¼ ðW W ÞþðW W Þ

–ðħm ħmÞf0ð1 f0aÞ ðm mÞ

ð ħm ħmÞfð1 faÞ m

ħmð1 f0aÞ ð1 faÞ fða a Þ

h i
–ð1 f0aÞfm þ ð1 faÞf0ðm mÞ ð Þ

ħ ħ
h c
a ð¼ ħ

f f0

f f

f ¼f 1 ¼½ðe Þ¼ pe a ð e þ Þ1 ð Þ

f0¼f 01 ¼½ðn Þ¼ pe a ð n þ Þ1 ð Þ

e e n
ħ! a

W W ¼

W W ¼ hcaD f ^{0} D

n

ða a Þ)

f^{0} cf.

a ¼

m ¼ ðm mÞ ¼

e ¼ ca.
e ¼ ca.
m m m

et al.

p

p cf.

p ! p

n ! p n ! s N

bb aa bb ¼

etc n ! p)

n

p

m aa

n ! p

mðn ! p Þ ¼

m ¼

hypsochromic n ! p

n p

bathochromic

n ! p

n

p

hypsochromic n ! p

n ! p

! !

!

n

n ! p

cf

n ! p

ñ ¼ l ¼



n ! p

n

c ¼ :

Dn ¼ Dl ¼ þ

Dn ¼ Dl ¼ þ

Dn ¼ þ Dl ¼ Dn ¼

þ Dl ¼

n ! p p p

n ! p n ¼

l ¼ n Dn ¼ þ Dl ¼

n

Dn ¼ þ Dl ¼ !

n ! p

b

n ! p p ! p

mðp ! p Þ m

m e.g. m ¼ mðp ! p Þ ¼

p ! p

n ! p

n ! p p ! p

	ñðnÞ ! p Þ	ñðpÞ ! p Þ
n		
N N		

ca.

	Dñ¼þ Dl ¼	Dñ¼ Dl ¼þ

n ! p p ! p
n ! p p ! p

n ! p

0

n ! p p ! p

ñ ¼ l ¼ n



Dn¼ Dl ¼ þ

n

Dn¼ Dl ¼ þ

n ! p

bb aa

N aa bb

n ! p p ! p



n

l n! p

Dl

n ! p Dl ¼

n

E

cf

N Dn¼ þ Dl ¼

n !

n ! p

n ! s

n ! s

solvatochromism
”uoro-
solvatochromism solvato”uorochromism

!

Stokes shift Dn¼n n

i.e m > m

t

t

t x t)

t w t cf.)

t
t
t w t t
t
t A t t x t

...



S^0 S^0

S S t w t

”uorophores

a p

m m

a

m > m

0

Dn¼

Dl ¼ þ i.e.

ca.

m ¼ m ¼



(a) (b)

c	Dl	Dn
l :	þ	
l :	þ	

i c t

(a)

(b)

m A

m A

t i c t

et al.

et al.

¼ ¼


hνA
ICT
S0,np
S1,np
S1,ct
R = SO3⊖Na⊕, SO2N(CH3)2
modestly solvent-dependent
strongly solvent-dependent
hνF,np
hνF,ct

; ; ;

; !

; !

; ;

et al.

cf.

N N

Dn¼ Dl ¼ þ n !

m ¼

p



$\mu_g \leq 5 \cdot 10^{-30}$ C m (1.6 D)

$\mu_e \approx 83 \cdot 10^{-30}$ C m (25 D)

n Dl Dn

l : þ

< >



2,5-Bis[3-(*N*-methyl-*N*-phenylamino)propen-1-yliden]cyclopentanone
$\Delta\tilde{\nu} = -3970\ cm^{-1}$ ($\Delta\lambda = +119$ nm) for toluene → water [346, 347]



2-(2-Naphthoxy)ethyl 4-cyanobenzoate
$\Delta\tilde{\nu} = -4000\ cm^{-1}$ ($\Delta\lambda = +88$ nm)
for cyclohexane → *t*-butanol [348]



2-[4-(Dimethylamino)styryl]benzoxazole
$\Delta\tilde{\nu} = -4050\ cm^{-1}$ ($\Delta\lambda = +92$ nm)
for *n*-hexane → water [349]



4-Aminophthalimide
$\Delta\tilde{\nu} = -5010\ cm^{-1}$ ($\Delta\lambda = +115$ nm)
for diethyl ether → water [350–353]



2-(Dimethylamino)-6-propanoylnaphthalene (PRODAN)
$\Delta\tilde{\nu} = -6100\ cm^{-1}$ ($\Delta\lambda = +130$ nm)
for cyclohexane → water [354, 355]



2-(Dimethylamino)-7-nitrofluorene
$\Delta\tilde{\nu} = -7020\ cm^{-1}$ ($\Delta\lambda = +260$ nm)
for triethylamine → dimethyl sulfoxide [429]



A representative molecular fluorescence marker
$\Delta\tilde{\nu} = +3600\ cm^{-1}$ ($\Delta\lambda = -105$ nm)
for *n*-heptane → methanol [433]

selective preferential solvation
p ! p

cf.

et al.
Py cf.
p ! p
. . .
! etc. !
!

Py ¼ I I

i.e.

cf.

Dm¼ m m

Dħ¼ ħ ħ ¼
e n

ħ ðaÞ g ! e ħ ðfÞ
e ! g t w t

hcaDn¼hcaðn ðaÞ n ðf ÞÞ

¼hcaðn ðaÞ n ðf ÞÞ

þ $\frac{\text{pe a}}{\text{e þ}}$ $\frac{\text{e}}{\text{n þ}}$ $\frac{\text{n}}{\text{jm mj}}$ ð Þ

n ðaÞ n ðf Þ

a Dn

a

a

e

Dn¼n n ¼$\frac{\text{p e}}{\text{h c a}_{w}}$ ðm mÞ Df þ : ð Þ

Df ¼f ðeÞ f ðn Þ ¼½ð Þ=ð e þ Þ ½ð Þ=ð n þ Þ

f ðeÞ f ðn Þ

Df

Df ¼ Df ¼

e n

Df ¼

½ð Þ=ðe þ Þ ½ð Þ=ðn þ Þ

Dm Dn Df

Cotton e¤ect

n ! p

)

R

y De a

n ! p

cf.

n ! p l ¼

n

n ! p

n

n ! p

De

b

a

a

R

n ! p

l ¼ and

Dn ¼ þ Dl ¼

n ! p

!

e.g

e.g

S

½F ¼ þ : ½F ¼ :

b



½F a) þ þ þ þ

a) c ¼ : ½F ¼ ðb½ M Þ= M ¼

b

b cf.

n ! p

cf.

þ trans

n cf.

32a Ð 32b

a

½F

l ¼ þ

þ

n ! p

R; R

induced optical activity

þ

n ! p

R

n ! p

y l A

þ þ



Contact ion pair
in non-dissociating
solvents

½a ½F2

S R; R þ

½a

e.g

e.g.

solvent

aa

ñ g



bb

(a) ñ
A
ñ (b)
ñ Dñ =

g

Dñ

ñ ñ

e.g. ñ b

e.g.

g a

dl bb dm ¼ dm aa dl ¼

dl aa dm ¼

bb aa aa

ñ b e.g. ñ a e.g.

bb

aa aa

	ñ b bb a)	ñ a aa b)	ñ a n a,c)

ðñ Þ

n

N N

d)

e)

	Dñ ¼ ñ ñ ¼ :	Dñ ¼ ñ ñ ¼	Dñ ¼ ñ ñ ¼
a)			
b)			
c) trans			
d)			
e)			

ħ b

N N

N N

ħ b

ħ b

ħ b ħ a

n ħc a ħ a

ħ c ħ a ħ a N

N ħ a tert

ħ b ħ c

c ¼

b

c

etc.

e.g. aa

ca.

bb

cf.

bb



ħ b



ħ b

n

n cf.

ħ b

cf.

bb

n

. . .

cf.

ħ b ¼ :

ħ b ¼ :

ħ b ¼ :

ħ b

ħ b

n ! p

bb

dl bb dm dm aa dl aa

ñ a ¼ ñ a

cf. ñ a

ñ a ñ a cf.

bb aa a

Dñ=ñ

e

$$\frac{Dn}{n} \frac{1}{4} \frac{n \quad n}{n} \frac{1}{4} C \frac{e}{e\ þ} \qquad ð \quad Þ$$

n n

e C

ca.

$$\frac{Dn}{n} \frac{1}{4} \frac{n \quad n}{n} \frac{1}{4} C \frac{n}{n\ þ} \qquad ð \quad Þ$$

n

Dn=n

$$\frac{Dn}{n} \frac{1}{4} C \; \beta \; C \quad \frac{e}{e \; \beta} \beta \; C \quad \frac{n}{n \; \beta} \quad \delta \qquad \Phi \qquad \delta \quad \Phi$$

$$\frac{Dn}{n} \frac{1}{4} C \; \beta \; - \delta C \; \beta \; C \; \Phi \; \frac{e}{e \; \beta} \quad \delta \qquad \Phi \qquad \delta \quad \Phi$$

C C C

ðe Þ=ð e þ Þ
C ðC þ C Þ C C

f ðeÞ ¼ ðe Þ=ð e þ Þ
f ðn Þ ¼ ðn Þ=ð n þ Þ

et al. f ðeÞ f ðn Þ

$$\frac{Dn}{n} \frac{1}{4} C \; \beta \; C \quad f \delta e \Phi \beta C \quad f \delta n \Phi \beta C \quad f \delta e \Phi \; f \delta n \Phi \qquad \delta \quad \Phi$$

f ðeÞ f ðn Þ

Dn b ¼
n !
f ðeÞ
f ðe; n Þ

et al.

Dn=n
Dn=n

G

et al. cf.

A

cf.
A Dn=n
bb
A aa

g a

g

p p

a a

g





N N

)

g

t

g

g

p _aa_j^m $ ^mj_aa_p

g a
a

	g	a	a)	a	a)

N N

a) ¼ Dg ¼ : Dað Þ ¼ : Dað Þ ¼ þ :
k : k k



g
a

a a a

(a) (b)



ð Þ

a a p
(b)
(a) a
a
(a) (b)

(b)

(b)

g

g g_e ¼
: cf.

t t

a t

a

cf.

a ħ aa

ħ n ! p t

bb aa

a t

ħ n ! p

a a a

a

g

g ¼ :

g ¼ :

a



(a) (b)

ð Þ

N

a

a

(b)

a x ¼ : . . . :
a

ca. a

cf.

p
et al.

N N

et al.
et al. et al.

t et al.

t
cf.

g t

a



ð Þ

a
a
a

a

(a) (b) etc. (c) (d) etc.

c ¼

a

!

l m

m

I ¼ =

cf.

cf.

N N

n

w

Dw

d ¼ d þ $\frac{p}{}$ Dw

¼ d þ $\frac{p}{}$ ðw w Þ ð Þ

etc.

Ddca. : ... :

Ddca. ... Dd>

cf.

cf.

a) l

	d l	d m	Dd b)	d c)	d d)	d e)	f)

n

N N

	Dd¼	Dd¼ :	DDd¼ :	Dd¼ :	Dd¼ :

a)
b)
c) n
n
d)

e)
f)

bb $ l aa m

et al.

cf.

ñ bb bb
b ¼



ð Þ

Ddð	b	Þ	a,b)		þ	þ	þ	þ	þ
Ddð	b	Þ	a,c)	þ					

a)

b)
c)

bb

N N O

N

cf.

p

(a) (b)

cf.



ð Þ

---

d a)

a) c ¼ :

DdA

DG^0

aa

p aa

DG^0

cf.

et al.

: . . . þ :

s

s s

s ¼ s þ s ð Þ

s

s ¼ ðH H Þ=H ð Þ

H

H

s

s ¼s þ s þ s þ s þ s ð Þ

s

s

cf.

s

s

s s s

e.g.

e.g.

s

i.e.

ðe Þ=ð e þ Þ

s

vice versa s

s

s s

s

s

s d

Dd¼d d ¼d þ d þ d þ d þ d ð Þ



Dd
ðd Þ ðd Þ
I ¼ -
DdA
d
i.e.
d d
! n ! !
! m
: ! : ! : ! ! þ :
all

a priori

e.g.

DdA

d c a Dd a)

N N

a) Dd ¼ dð Þ Þ dð Þ Þ

cc aa

cf.

aa

aa j j

j

p

cf.

p

j

cf.

cf.

n l m

aa cc aa

ca.

solute/solvent

solute/solute

cf.



ð Þ

---

d a)

a) c¼ :

ca.

i.e.

(a)

(b)

(b)

cf.

e.g.

cf.

s

aromatic solvent-induced shift)

Ddð Þ ¼d d ð Þ

d e.g.

d e.g.

Dd

Dd G

Dd

: ... :

: ... :

aaa aa a0

Dd

Dd

cf.



Δδ{1H-ASIS} of pulegone
in ppm



Δδ {1H-ASIS} of camphor
in ppm

Dd

t

Dd

cf.

Dd

Ddð Þ ¼ ð Þ ð Þ Þ ð Þ

et al.

cf.

aa

Dd

cf.

Dd

cf.

cf. et al.

R S

n

n

þ þ

þ þ

þ

ðI þ Þ

vice versa.

cf.

J a J ¼ :

J ¼ : J a

J ¼ : J ¼ : J a

aa

J a

s

J a

s

aa

J a

J a

J a

J a

J a

J a

J

J a

J a

J a

J a

J a

cf.

aa aa aa

J a

aa

J a

n

J a

J

a

J a

7 Empirical Parameters of Solvent Polarity

7.1 Linear Gibbs Energy Relationships

Solvent e¤ects on organic reactivity (cf. Chapters 4 and 5) and on absorption spectra (cf. Chapter 6) have been studied for more than a century (cf. Chapter 1). Organic chemists have usually attempted to understand these solvent e¤ects in terms of the polarity of the solvent. Solvent polarity is a commonly used term related to the capacity of a solvent for solvating dissolved charged or neutral, apolar or dipolar, species. This concept of solvent polarity is easily grasped qualitatively, but it is di‹cult to de“ne precisely and even more di‹cult to express quantitatively. Attempts to express it quantitatively have mainly involved physical solvent properties such as relative permittivity, dipole moment, or refractive index (cf. Section 3.2). From idealized theories, the solvent dielectric constant (i.e. the relative permittivity ε_r) is often predicted to serve as a quantitative measure of solvent polarity. However, this approach is often inadequate since these theories regard solvents as a non-structured isotropic continuum, not composed of individual solvent molecules with their own solvent/solvent interactions, and they do not take into account speci“c solute/solvent interactions such as hydrogen-bonding and EPD/EPA interactions, which often play a dominant role in solute/solvent interactions. Similarly, solvent dipole moments are inadequate measures of solvent polarity since the charge distribution of a solvent molecule may not only be given by its dipole moment but also by its quadrupole or higher multipole moments, leading to dipolar, quadrupolar, octupolar, etc. solvent molecules [121]. Therefore, a more general de“nition of the commonly used term *solvent polarity* would be useful.

The author stated in 1965 [1, 3] that the polarity of a solvent is determined by its solvation capability (or solvation power) for reactants and activated complexes as well as for molecules in their ground and excited states. This, in turn, depends on the action of all possible, speci“c and nonspeci“c, intermolecular forces between solvent and solute molecules. These intermolecular forces include Coulomb interactions between ions, directional interactions between dipoles, inductive, dispersion, hydrogen-bonding, and charge-transfer forces, as well as solvophobic interactions (see Chapter 2). Only those interactions leading to de“nite chemical alterations of the solute molecules through protonation, oxidation, reduction, complex formation, or other chemical processes are excluded. This more pragmatic de“nition of solvent polarity has meanwhile been included in the •IUPAC Recommendations 1994• for terms used in physical organic chemistry [291].

It is obvious that such a de“nition of solvent polarity cannot be measured by an individual physical quantity such as the relative permittivity. Indeed, very often it has been found that there is no correlation between the relative permittivity (or its di¤erent functions such as $1/\varepsilon_r$, $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, etc.) and the logarithms of rate or equilibrium constants of solvent-dependent chemical reactions. No single macroscopic physical parameter could possibly account for the multitude of solute/solvent interactions on the molecular-microscopic level. Until now the complexity of solute/solvent interactions has also prevented the derivation of generally applicable mathematical expressions that would allow the calculation of reaction rates or equilibrium constants of reactions carried out in solvents of di¤erent polarity.

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

In such a situation other indices of solvent polarity are sought. The lack of simple theoretical expressions for calculating solvent e¤ects and the inadequacy of de“ning ••solvent polarity•• in terms of single physical constants has stimulated attempts to introduce empirical scales of solvent polarity, based on convenient, well-known, solvent-sensitive reference processes. A common approach is to assume that some particular reaction rate, equilibrium, or spectral absorption is a suitable model for a large class of other solvent-dependent processes. If one carefully selects an appropriate, su‹ciently solvent-sensitive reference process, one can assume that this process re”ects all possible solute/solvent interactions that are also present in related solvent-in”uenced processes. It should therefore give an empirical measure of solvent polarity … or, more precisely, an empirical measure of the solvation capability of a particular solvent for the given reference process. This reference process can be considered as a probe of the solvation shell of the standard solute … a probe that sums up a wide variety of possible intermolecular interactions such as ion-dipole, dipole-dipole, dipole-induced dipole, hydrogen-bonding, interactions, etc. Naturally, the most useful model processes should be those best understood on a molecular basis. Model processes used to establish empirical scales of solvent polarity have been reviewed [1…9, 122…124, 289]. More recent reviews on empirical solvent polarity scales can be found in references [292…296].

At best, this approach provides a quantitative index to solvent polarity, from which absolute or relative values of rate or equilibrium constants for many reactions, as well as absorption maxima in various solvents, can be derived. Since they re”ect the complete picture of all the intermolecular forces acting in solution, these empirical parameters constitute a more comprehensive measure of the polarity of a solvent than any other single physical constant. In applying these solvent polarity parameters, however, it is tacitly assumed that the contribution of intermolecular forces in the interaction between the solvent and the standard substrate is the same as in the interaction between the solvent and the substrate of interest. This is obviously true only for closely related solvent-sensitive processes. Therefore, an empirical solvent scale based on a particular reference process is not expected to be universal and useful for all kinds of reactions and absorptions. Any comparison of the e¤ect of solvent on a process of interest with a solvent polarity parameter is, in fact, a comparison with a reference process.

This kind of procedure,i.e. empirical estimation of solvent polarity with the aid of actual chemical or physical reference processes, is very common in chemistry. The well-known Hammett equation for the calculation of substituent e¤ects on reaction rates and chemical equilibria, was introduced in 1937 by Hammett using the ionization of meta- or para-substituted benzoic acids in water at 25°C as a reference process in much the same way [10]. Usually, the functional relationships between substituent or solvent parameters and various substituent- or solvent-dependent processes take the form of a linear Gibbs energy relationship, frequently still referred to as a linear free-energy (LFE) relationship [11…15, 125…127].

Let us consider a certain reaction series, one in which only small changes are involved in going from one reaction to another. These changes may be structural, such as a series of di¤erently substituted compounds, or may involve a single reaction carried out in a series of di¤erent solvents or solvent mixtures. It has been found that the changes in rate and in equilibrium constant occurring in one reaction series can be often related to those in another, closely related series. Thus, plotting the logarithms of rate

or equilibrium constants for one reaction series against the corresponding constants for a second, related series frequently gives a straight line, which can be expressed by Eq. (7-1).

$$\lg k_i^B = m \cdot \lg k_i^A + c \tag{7-1}$$

k_i^A and k_i^B are rate or equilibrium constants of two reaction series A and B, which are subject to the same changes in the structure or the surrounding medium.

Since the relationship between the equilibrium constant K, for a reaction and the difference between the standard molar Gibbs energies of the products and reactants, ΔG°, is given by Eq. (7-2),

$$\lg K = -\frac{\Delta G^\circ}{\ln 10 \cdot R \cdot T} \tag{7-2}$$

and a similar expression using the standard molar Gibbs energy of activation, $\Delta G^{\neq}$, for the rate constant k of a reaction can be written as in Eq. (7-3),

$$\lg\left(\frac{k}{R \cdot T / N_A \cdot h}\right) = -\frac{\Delta G^{\neq}}{\ln 10 \cdot R \cdot T} \tag{7-3}$$

Eq. (7-1) essentially describes a relationship between standard molar Gibbs energies.* It is often convenient to express linear Gibbs energy relationships in terms of ratios of constants by referring all members of a reaction series to a reference member of the series; thus, the correlation in Eq. (7-1) can also be expressed by Eq. (7-4).

$$\lg(k_i^B/k_0^B) = m \cdot \lg(k_i^A/k_0^A) \tag{7-4}$$

where k_0^A and k_0^B are the constants for the reference substituent or the reference solvent.

Such relationships are useful in two ways. The first application is in the study of reaction mechanisms. The correlation of data for a new reaction series by means of a linear Gibbs energy relationship establishes a similarity between the new series and the reference series. The second use of linear Gibbs energy equations is in the prediction of reaction rates or equilibrium constants dependent on substituent or solvent changes. Let us consider a reaction between a substrate and a reagent in a medium M, which leads, via an activated complex, to the products**:

---

* Linear Gibbs energy relationships are manifestations of so-called extrathermodynamic relationships. Extrathermodynamic approaches are combinations of detailed models with the concepts of thermodynamics. Since it involves model building, this kind of approach lacks the rigour of thermodynamics, but it can provide information not otherwise accessible. Although linear Gibbs energy relationships are not a necessary consequence of thermodynamics, their occurrence suggests the presence of a real connection between the correlated quantities, and the nature of this connection can be explored.

** The designation of one reactant as the *substrate* and another as the *reagent* is arbitrary but useful in considering chemical reactivity. The substrate always undergoes some change in the reaction series, the reagent does not. A catalyst is always considered to be a reagent.

$$(\text{Substrate})_M + (\text{Reagent})_M \rightleftharpoons [S\cdots R]^{\neq}_M \rightarrow (\text{Products})_M \qquad (7\text{-}5)$$

There are three ways of introducing small changes in order to establish a reaction series:

(a) First, we can change the substrate by introducing different substituents. In the case of meta- and para-substituted benzene derivatives, this leads to the well-known Hammett equation (7-6), where k_X is a rate or equilibrium constant for meta- or para-

$$\lg k_X - \lg k_0 = \lg(k_X/k_0) = s \cdot \varrho \qquad (7\text{-}6)$$

substituted substrates, s is the substituent constant, and ϱ is the reaction constant ($\varrho = 1$ for the standard reaction by definition) [10]. A typical Hammett correlation is shown in Fig. 7-1 for the S_N2 alkylation reaction of substituted pyridinium-N-phenolate betaines with iodomethane [16]. As expected for this S_N2 reaction (negative ϱ-value), electron-releasing substituents increase the reaction rate, whereas electron-withdrawing groups have a decelerating effect.



(b) Second, we can change the reagent to give, for example, a Gibbs energy relationship called the Brønsted…Pedersen catalysis equation, developed as a result of studies on the base-catalyzed decomposition of nitroamide [17]. This equation establishes a quantitative relationship between acid and base strengths and their effectiveness as catalysts in reactions subject to general acid or base catalysis: the stronger the acid (or base), the better it is as a catalyst. This Brønsted catalysis law, introduced in 1924, was the first linear Gibbs energy relationship.



Fig. 7-1. Hammett correlation between s-values and the logarithms of the relative rate constants of the S_N2 alkylation reaction of substituted pyridinium N-phenolate betaines with iodomethane in trichloromethane at 25°C [16].

(c) Finally, we can change the surrounding medium M while leaving the other reaction parameters unchanged. Provided the selected reaction is sufficiently solvent-sensitive, this gives us the desired empirical solvent parameters [1…9, 122…124, 292…296]. Thermodynamically, solvation may be considered in the same general terms as the modification of the properties of the substrate molecule by substituent changes, the solvating molecules being equivalent to loosely attached substituents [18]. One important difference between substituent and solvent effects on chemical reactivity is that substituents may change the chemical reactivity of a given substrate only in a discontinuous manner, whereas solvents, especially solvent mixtures, allow a continuous modification of the substrate reactivity. Empirical parameters of solvent polarity based on solvent effects on chemical equilibria and reaction rates will be described in Sections 7.2 and 7.3.

In principle, the same considerations as in Eq. (7-5) can be made for the spectral excitation of a substrate, dissolved in a medium M, with photons $h \cdot n$. Although linear Gibbs energy relationships usually deal only with relative reactivities, in the form of reaction-rate and equilibrium data, this approach can be extended to various physical

$$\underset{\text{Ground state}}{(\text{Substrate})_M} + h \cdot n \rightarrow \underset{\text{Excited state}}{(\text{Substrate})^*_M} \qquad (7\text{-}7)$$

measurements such as spectroscopic investigations of the members of a reaction series in various spectral ranges (UV/Vis [19], IR [19], NMR [20], etc.). Spectroscopic measurements can very often be obtained under conditions of greater precision and variety than available in reactivity measurements.

In order to establish a reaction series, the substrate in Eq. (7-7) can be altered in two ways:

(a) Variation by substituents leading to spectroscopic Hammett equations in the form presented in Eq. (7-8),

$$\frac{E_{T,X} - E_{T,O}}{\ln 10 \cdot R \cdot T} = s \cdot \varrho_A \qquad (7\text{-}8)$$

and first introduced by Kosower, Hofmann, and Wallenfels [21]. $E_{T,X}$ and $E_{T,O}$ are the transition energies in kJ/mol (or kcal/mol) of the substituted and unsubstituted substrate, respectively; s is the Hammett substituent constant, and ϱ_A has been denoted as an absorption constant [22, 23]. The transition energies are divided by $(\ln 10 \cdot RT)$ in order to convert them into an appropriate form for Hammett substituent constants, which are commonly derived from equilibrium or rate constants. A representative example of this kind of Hammett correlation for the long-wavelength $p \rightarrow p^*$ absorption of substituted pyridinium N-phenolate betaines is shown in Fig. 7-2 [23].

As expected, electron-withdrawing substituents cause a bathochromic shift of this absorption band, which exhibits a strong intramolecular charge-transfer character (cf. Section 6.2.1), whereas electron-releasing substituents give rise to a corresponding hypsochromic shift.



Fig. 7-2. Hammett correlation between σ-values and the modified transition energies of the long-wave-length charge-transfer $p \rightarrow p^*$ absorption of substituted pyridinium N-phenolate betaines in methanol at 25 °C [23]; cf. Fig. 6-2 in Chapter 6.

(b) Provided the position of the absorption band is sufficiently solvent-sensitive, changing the medium M, in which the substrate is to be dissolved, permits the introduction of spectroscopic solvent polarity scales. Empirical parameters of solvent polarity based on spectroscopic measurements will be described in Section 7.4. The most comprehensive solvent polarity scale is based on solvatochromic pyridinium N-phenolate betaines, already mentioned in Section 6.2.1 and in Figs. 7-1 and 7-2 [293].

In the early use of linear Gibbs energy relationships, simple single-term equations such as the Hammett equation were considered sufficient to fit given sets of experimental data from reaction series. Later on, more complicated multi-term equations with more than one product term were formulated in order to model the simultaneous influence of several effects on chemical reactions or optical excitations; one product term per effect [15]. The connection between such multiparameter relationships and solvent effects will be described in Section 7.7.

Simple and multiple linear Gibbs energy relationships can be generally interpreted in two distinct ways [126, 127]:

(a) Traditionally, as relationships expressing combinations of fundamental chemical or physical effects, universally present in chemical reactions. Deviations from such relationships have to be explained by new effects in addition to the old, known effects. If all such effects are discovered and quantified, retrospective rationalization and prediction of

chemical processes (i.e. chemical reactions and equilibria) is fully possible and it is conceivable that at some time in the future basic research will no longer be needed. The main problems are to "nd reaction series which are best suited to discovering all these fundamental e¤ects, and to establish how large a deviation can be tolerated before another fundamental e¤ect has to be postulated.

(b) Chemometrically*), as locally valid linearizations of very complicated unknown functional relationships using empirical models of similarity. That is, linear Gibbs energy relationships are considered as approximate models with local validity for similar reaction series only, e.g. one model for reactions of substituted benzene derivatives, another model for reactions of substituted aliphatic compounds. In the case of solvent e¤ects, one model would apply to solvatochromic non-HBD, another to solvatochromic HBD reference dyes. In contrast to the traditional interpretation (a), this means that a given linear Gibbs energy relationship is not necessarily universally valid. In any new chemical reaction series there will be regularities in the observed data that cannot be predicted from the behaviour of previously investigated chemical model processes. In other words, the data observed in a chemical reaction series can be divided into two parts: one part which will coincide with other known reaction series, and another part which will be system-speci"c. The latter part is often not negligible and can sometimes be substantial. The larger the common part, the closer the similarity between a new reaction series and the reference reaction series previously investigated. The mathematics of "tting PC and FA models to a matrix of chemical data permits one to judge objectively the applicability of a given linear Gibbs energy relationship, without making assumptions about a single term model "tting part of the data matrix.

A fascinating discussion on the problem of whether linear Gibbs energy relationships are (a) fundamental laws of chemistry, or (b) only local empirical rules, can be found in references [126, 127]. Attempts at discriminating between these two kinds of interpretation of linear Gibbs energy relationships based on an evaluation of the practicality of models derived from either view are described. It seems that from a more rigorous mathematical point of view, interpretation (b), stressed by Wold and Sjöström [126], appears to be the correct one, whereas from a more practical, descriptive point of view, interpretation (a), stressed by Taft and Kamlet [127], will be the one preferred by experimental chemists. The "nal answer is ambiguous.

Another question now seems to be more pertinent. Are empirical parameters of solvent polarity still necessary in view of the great progress that has been made during the last decades in the development of theoretical methods and computational strategies for describing and understanding the complex phenomenon of solvation of reactants, activated complexes, and products, as well as of molecules in their ground and excited states? Essentially, three general approaches have been used for the theoretical description of solute/solvent interactions: quantum-chemical methods, supramolecular methods, and semicontinuum quantum-chemical methods (see the end of Section 2.3 and

* Chemometrics stands in this context for analysis of multivariate chemical data by means of statistical methods such as principal component analysis (PCA) or factor analysis (FA) (cf. Section 3.5).

references [27d, 355…361] to Chapter 2 for recent reviews). The results obtained from all these theoretical calculations are very impressive. However, when applied to actual chemical or physical processes in a solution under study, all methods still have their limitations and ”aws, particularly concerning the inclusion of nonspeci“c solute/solvent interactions. Therefore, for obtaining comparatively simple and quick information about how a solvent change may in”uence a particular reaction under study, both qualitatively and quantitatively, the use of linear Gibbs energy relationships still seems to be a good choice. A chemist in his daily laboratory work is often faced with simple questions such as: would the reaction under study pro“t from a change from solvent A to solvent B, would the reaction run faster in B than in A, and if yes to what extent, 10-fold or 100-fold? For these decisions, the comparison with other analogous, well-studied solvent-dependent processes and the empirical parameters derived therefrom will certainly continue to be as useful as it has been up to now.

7.2 Empirical Parameters of Solvent Polarity from Equilibrium Measurements

The “rst attempt to introduce an empirical relationship between an equilibrium constant and solvent polarity was made in 1914 by K. H. Meyer [24]. Studying the solvent-dependent keto-enol tautomerism of 1,3-dicarbonyl compounds, he found a proportionality between the equilibrium constants of various tautomeric compounds in the same set of di¤erent solvents (cf. Table 4-2 in Section 4.3.1). He therefore split the tautomeric equilibrium constant K_T into two independent factors according to Eq. (7-9).

$$K_T = \frac{[\text{enol}]}{[\text{diketo}]} \equiv L \cdot E \qquad (7\text{-}9)$$

E is the so-called enol constant and measures the enolization capability of the diketo form ($E = 1$ for ethyl acetoacetate by de“nition). Thus, the so-called desmotropic constant L is a measure of the enolization power of the solvent. By de“nition, the values of L are equal to the equilibrium constants of ethyl acetoacetate ($E = 1$), determined in di¤erent solvents [24]. This desmotropic constant seems to be the “rst empirical solvent parameter. It describes the relative solvation power of a solvent for diketo and enol forms of 1,3-dicarbonyl compounds. It was measured only for a few solvents and was soon forgotten.

In contrast to this early empirical solvent scale, one of the more recent, introduced by Eliel and Hofer in 1973 [25] and based on the solvent-dependent conformational equilibrium of 2-i-propyl-5-methoxy-1,3-dioxane, should be mentioned (cf. Table 4-9 in Section 4.4.3). In general, polar solvents shift this conformational equilibrium towards the more dipolar axial cis isomer. The authors proposed calling the standard molar Gibbs energy changes associated with this equilibrium, $\Delta G^\circ_{OCH_3}$, the D_1 scale (D for dioxane, 1 for the number of carbons in the alkoxy group). They also recommended the use of this solvent scale, known for seventeen solvents, in the estimation of other solvent-dependent equilibria and reaction rates [25].

Another approach to a new solvent scale was introduced by Gutmann in 1966 [26, 27]. Based on the fact that many chemical reactions are in”uenced primarily by coordi-

nation interactions between an electron-pair acceptor (EPA) substrate and electron-pair donor (EPD) solvents (cf. Sections 2.2.6 and 2.6), he provided an empirical measure of the Lewis basicity of solvents using the so-called donor number DN (or donicity). Antimony pentachloride was chosen as a reference compound in order to obtain a measure of the nucleophilic properties of a solvent since the antimony is readily coordinated on the acceptance of an electron-pair from an EPD solvent. According to Eq. (7-10), the donor number is defined as the negative value of the molar enthalpy for adduct

$$D: + SbCl_5 \underset{\text{in Cl-CH}_2\text{CH}_2\text{-Cl}}{\overset{\text{room temp.}}{\rightleftharpoons}} \overset{\oplus}{D}—\overset{\ominus}{Sb}Cl_5$$

$$\text{Solvent Donor Number DN} \equiv -\Delta H_{D-SbCl_5}/(\text{kcal} \cdot \text{mol}^{-1}) \qquad (7\text{-}10)$$

formation between antimony pentachloride and EPD solvents, D, measured calorimetrically in highly diluted solutions of 1,2-dichloroethane as an inert solvent at room temperature [26]. Corresponding enthalpy measurements in the reference solvent tetrachloromethane have led to analogous results [28].

Donor numbers vary from 2.7 for nitromethane, which is a weak EPD solvent, to 38.8 for hexamethylphosphoric triamide, a strong EPD solvent (cf. Table 2-3 in Section 2.2.6. The donor number was measured directly (i.e. calorimetrically, for ca. fifty solvents [26…28, 128]. It has also been estimated by other means such as ^{23}Na [29, 129], ^{27}Al [130], and ^{1}H NMR [131] spectroscopy. Visual estimates of the varying donor numbers of EPD solvents can easily be made using the colour reactions with copper(II), nickel(II), or vanadyl(IV) complexes as acceptor solutes instead of $SbCl_5$ [132]. A selection of donor numbers has already been given in Table 2-3 in Section 2.2.6. A discussion of methods for the determination of donor numbers and critical compilations of DN values for 170 and 134 solvents can be found in references [133] and [294], respectively.

Since donor numbers are defined in the non-SI unit kcal mol^{-1}, Marcus has recommended the use of dimensionless normalized donor numbers DN^N, which are defined as $DN^N = DN/(38.8\ \text{kcal} \cdot \text{mol}^{-1})$ [133]. The non-donor solvent 1,2-dichloroethane ($DN = DN^N = 0.0$) and the strong donor solvent hexamethylphosphoric triamide (HMPT: $DN = 38.8\ \text{kcal} \cdot \text{mol}^{-1}$; $DN^N = 1.0$) have been used to fix the DN^N scale. Although solvents with higher donicity than HMPT are known, it is expedient to choose this EPD solvent with the highest directly (i.e. calorimetrically, determined DN value as the second reference solvent [133]. The normalized DN^N values are also included in Table 2-3 in Section 2.2.6.

Since the donor numbers were measured in dilute solution in inert 1,2-dichloroethane*), they reflect the donicity of single EPD solvent molecules. However, in neat, associated EPD solvents an increase in the donicity occurs [134]. For such highly

* 1,2-Dichloroethane is obviously not a chemically inert solvent under all circumstances. For example, the EPD solvent triethylamine is rapidly quaternized by 1,2-dichloroethane under the catalytic action of $SbCl_5$, leading to an overestimated donor number for triethylamine [128]. Even solutions of HMPT and $SbCl_5$ in 1,2-dichloroethane contain non-negligible amounts of charged species, the formation of which contributes to the measured enthalpy [138].

structured neat EPD solvents (e.g. water, alcohols, amines) the term bulk donicity has been introduced [135] in order to rationalize the deviations of these solvents in plots of $^{23}Na^{\oplus}$ NMR chemical shifts [136] and ESR parameters [137] vs. the donor number. Unfortunately, great discrepancies exist between the DN_{bulk} values given in the literature when estimated by different methods. For this reason, they are not included in Table 2-3 in Section 2.2.6; see reference [133] for a collection and discussion of bulk donicities, DN_{bulk}.

Donor numbers are considered as semiquantitative measures of solute/EPD solvent interactions, and are particularly useful in the prediction of other EPD/EPA interactions in coordination chemistry. Numerous examples of the application of donor numbers have been given by Gutmann [26, 27, 30]; cf. also [113, 133]. It has been shown that donor number correlations are parallel with correlations based on the highest occupied molecular orbital (HOMO) eigenvalues of EPD solvent molecules [139]. For non-HBD solvents, a fair correlation has been obtained between their donor numbers and their gas-phase proton affinities PA, indicating that the DN values do indeed reflect the intrinsic molecular properties of EPD solvents [140].

The donor number approach has been criticized for both conceptual [141] and experimental reasons [28, 133, 138, 265]. Therefore, the search for other empirical Lewis-basicity parameters has continued.

Another remarkable Lewis basicity scale for 75 non-HBD solvents has been established by Gal and Maria [138, 142]. This involved very precise calorimetric measurements of the standard molar enthalpies of 1:1 adduct formation of EPD solvents with gaseous boron trifluoride, $\Delta H^{\circ}_{D-BF_3}$, in dilute dichloromethane solutions at 25°C, according to Eq. (7-11).

$$D: + BF_3 \underset{\text{in } CH_2Cl_2}{\overset{25\,^{\circ}C}{\rightleftharpoons}} \overset{\oplus}{D}-\overset{\ominus}{B}F_3 \qquad (7\text{-}11)$$

A selection of $\Delta H^{\circ}_{D-BF_3}$ values has already been given in Table 2-4 in Section 2.2.6. This new Lewis basicity scale is more comprehensive and seems to be more reliable than the donor number scale. Analogously, a Lewis basicity scale for 88 carbonyl compounds (esters, carbonates, aldehydes, ketones, amides, ureas, carbamates) has been derived from their standard molar enthalpies of complexation with gaseous boron trifluoride in dichloromethane solution [143]. The corresponding $\Delta H^{\circ}_{CO-BF_3}$ values range from 33 kJ · mol^{-1} for di-t-butyl ketone to 135 kJ · mol^{-1} for 3-diethylamino-5,5-dimethylcyclohexen-2-one.

Solution calorimetry of the molecular probes pyrrole, N-methylpyrrole, benzene, and toluene in 35 solvents has been used by Cáceres et al. to determine a solvent HBA basicity scale, ranging from the gas phase to HMPT [31a]. Analogously, a solvent HBD acidity scale was derived calorimetrically using N-methylimidazole and N-methylpyrrole as probe molecules in 36 solvents, ranging from the gas phase to 2,2,2-trifluoroethanol [31b].

Another, more general approach for the estimation of EPD/EPA interactions between Lewis acids and bases, not restricted to the solvent as a reaction partner, was given by Drago [32]; cf. Eq. (2-12) and Table 2-6 in Section 2.2.6.

A compilation and critical comparison of various Lewis basicity scales for EPD solvents has been given by Persson et al. [144, 292].

The solvent-dependent tautomerization of a pyridoxal 5′-phosphate Schiff base has been shown to be another appropriate model process for the measurement of solvent polarities [32a]. This model process seems to be particularly useful for the determination of the polarities of sites of proteins at which pyridoxal 5′-phosphate is bound [32a].

Furthermore, an empirical hydrophobicity parameter derived from measurements of the distribution of a solute between two immiscible liquids should be mentioned (cf. Section 2.2.7 dealing with hydrophobic interactions). The hydrophobic or lipophilic character of organic compounds plays an important role in their ability to interfere with biochemical systems. Therefore, systematic efforts have been made to obtain numerically defined constants to assess the hydrophobic character of organic compounds. A hydrophobicity parameter which has proven quite valuable in the fields of toxicology, pharmacology, and environmental science is the Hansch–Leo 1-octanol/water partition coefficient $K_{o/w}$ or $P_{o/w}$ as defined in Eq. (7-12),

$$K_{o/w} = \frac{c_i(\text{1-octanol})}{c_i(\text{water})} \equiv P_{o/w} \qquad (7\text{-}12)$$

where c_i(1-octanol) and c_i(water) are the molar equilibrium concentrations of the solute i in the two immiscible phases 1-octanol and water, respectively [145–147]. The 1-octanol/water partition coefficients are often used as hydrophobicity parameters in the Gibbs energy-based form of $\lg P_{o/w}$. They are known for a vast number of organic compounds, particularly for those compounds normally used as organic solvents. A $\lg P_{o/w}$ value of 3.90 for n-hexane means that this hydrophobic solvent is preferably found in the 1-octanol phase, whereas a $\lg P_{o/w}$ value of -1.35 for dimethyl sulfoxide reveals the hydrophilicity of this particular solvent. A compilation of $\lg P_{o/w}$ values for 102 organic solvents can be found in reference [149], which also deals with correlations of $\lg P_{o/w}$ with some solvatochromic solvent parameters. 1-Octanol/water partition coefficients have served as the basis for many papers in a field of biochemistry known as quantitative structure/activity relationship (QSAR), inspired by Hansch's original observation that distribution of a solute between these two solvents provides an accurate measure of lipophilic/hydrophilic interactions. For example, $\lg P_{o/w}$ is a good measure of the ease with which drugs penetrate membranes and bind to hydrophobic surfaces. The interrelation between the narcotic potencies of various substances and the partition coefficient has long been known. Many applications of this hydrophobicity parameter in the framework of linear Gibbs energy relationships have been reported [145–147].

It should be mentioned that water and 1-octanol are not completely immiscible. The solubility of water in 1-octanol is $s = 2.46$ mol/L at 25 °C [= mole fraction $x(H_2O) = 0.289$], and that of 1-octanol in water is $s = 3.29 \cdot 10^{-3}$ mol/L [= x(1-octanol) $= 9.32 \cdot 10^{-4}$] [148]. In the application of $K_{o/w}$ values, it is tacitly assumed that the solvent properties of 1-octanol saturated with water are not different from those of neat 1-octanol. In practice, the presence of water in 1-octanol should increase the concentration of polar and H-bonding solutes and decrease the concentration of nonpolar solutes. However, solvatochromic studies by Carr et al. have shown that the water saturation of 1-octanol has only a very small effect on the properties of bulk 1-octanol. The water is almost completely associated with the hydroxy group of 1-octanol and scarcely affects the properties of this medium [148].

Using 1-octanol/water partition coefficients for aromatic solutes, Hansch also defined a hydrophobicity parameter p_X for organic substituents X according to Eq. (7-12a) [145…147].

$$p_X = \lg P_{o/w}^{C_6H_5X} - \lg P_{o/w}^{C_6H_6} \qquad (7\text{-}12a)$$

$P_{o/w}^{C_6H_5X}$ and $P_{o/w}^{C_6H_6}$ are the partition coefficients between 1-octanol and water for C_6H_5X and C_6H_6, respectively. A positive p_X, such as $+0.56$ for $X = CH_3$, means that these substituents favour the organic phase relative to the hydrogen atom ($X = H$). If substituents X have negative values of p_X, such as -0.67 for OH, then they cause partitioning into water. The p_X substituent parameters have also been widely used in quantitative structure/activity relationships, particularly in pharmacology.

Another microscopic hydrophobicity substituent parameter MH was evaluated by Menger et al. [150]. Addition of ammonium salts $R\text{-}NMe_3^{\oplus}\ X^{\ominus}$ (R = alkyl, aryl; $X^{\ominus}$ = halide ions) to 10^{-5} M 4-nitrophenyl laurate in water destroys or disrupts the ester aggregates formed in aqueous solution. The corresponding shift of the aggregation equilibrium deshields the laurate ester group, resulting in enhanced rate constants for the basic hydrolysis of 4-nitrophenyl laurate. The more hydrophobic the R group of the ammonium salt, the greater its disaggregation power, and the greater the rate enhancement [150].

A further quantitative measure of the solvent solvophobic effect* has been introduced by Abraham et al. [282]. It has been shown that the standard molar Gibbs energies of transfer of nonpolar, hydrophobic solutes X (X = argon, alkanes, and alkane-like compounds) from water (W) to other solvents (S) can be linearly correlated through a set of equations such as Eq. (7-12b),

$$\Delta G_t^{\circ}(X, W \rightarrow S) = M \cdot R_T + D \qquad (7\text{-}12b)$$

where R_T is a solute parameter, and M and D characterize the solvent. M values are referred to water, i.e. $M = 0$ for water by definition. With another fixed point, i.e. the M value of the most hydrophobic solvent n-hexadecane ($M = -4.2024$), a scale of solvophobic power Sp has been defined according to Eq. (7-12c),

$$Sp = 1 - \frac{M\,(\text{solvent})}{M\,(\text{n-hexadecane})} = 1 + \frac{M\,(\text{solvent})}{4.2024} \qquad (7\text{-}12c)$$

where the Sp values of water and n-hexadecane are arbitrarily taken as unity and zero. The Sp values provide a simple quantitative measure of the solvophobic power of solvents, relative to the two reference solvents water ($Sp = 1$) and n-hexadecane ($Sp = 0$).

---

* The solvophobic or hydrophobic effect is simply regarded as the experimental observation of the relative insolubility in water (or other highly structured liquids) of certain organic solutes, as compared to their solubility in nonaqueous solvents. The so-defined solvophobic (hydrophobic) effect should be distinguished from the solvophobic (hydrophobic) interaction between two or more solute molecules in solution; cf. Section 2.2.7. The solvophobic (hydrophobic) effect is only related to solute/solvent interactions [282].

at 298 K. For pure solvents, the following order of decreasing solvophobic power has been obtained [282]: water $\gg$ formamide $>$ 1,2-ethanediol $>$ methanol $>$ ethanol $>$ 1-propanol $>$ 2-propanol $>$ 1-butanol $\gg$ n-hexadecane. Good correlations between S_p values and the rate constants as well as stereoselectivities of several Diels…Alder cycloaddition reactions, carried out in hydrophobic media, have been reported [283, 297]; also Section 5.5.8.

In this context, another empirical solvent parameter called SI should be mentioned. SI stands for Solvent Influence (in Russian, BP for Влияние Растворителя). This parameter was introduced by Shmidt et al. in 1967 and was derived from the study of many different extraction equilibria, i.e. of the distribution of organic and inorganic compounds between two immiscible liquid phases [298…301]. It was found that in the extraction of metal salts using various extraction reagents, the distribution coefficients of the extractable compound depend on the specific electrophilic and/or nucleophilic properties of the solvents used as diluent. From a large number of well-studied extraction systems, Eq. (7-12d) has been derived,

$$\lg(K/K_0) = a \cdot SI + b \qquad (7\text{-}12d)$$

where K is the extraction constant for the extraction under consideration with a given solvent as diluent, K_0 is the extraction constant for the same system with a standard diluent (e.g. benzene), and a is a constant coefficient for a series of extraction systems differing only in the nature of the solvent used as diluent. For numerous extraction systems, linear correlations have been found between $\lg K$ and the diluent SI parameter [298…301]. This and other linear Gibbs energy relationships applicable in the correlation analysis of distribution and extraction constants have been reviewed by Makitra [302].

Finally, a parameter describing the softness of solvents should be mentioned. In terms of Pearson's principle of hard and soft acids and bases, the hardness of an ion or molecule is understood as resistance to a change or deformation in the electronic cloud (cf. Section 3.3.2). This property, or its complement, the softness of solvents, is expected to play a role in the solvation of hard and soft solutes. Marcus defined a scale of solvent softness (from *malakos* = soft in Greek) according to Eq. (7-12e) as the difference between the mean of the standard molar Gibbs energies of transfer of sodium and potassium ions from water (W) to a given solvent (S) and the corresponding transfer energy for silver ions, divided by 100 [285].

$$m = \left\{ \frac{1}{2} [\Delta G_t^\circ(Na^\oplus, W \rightarrow S) + \Delta G_t^\circ(K^\oplus, W \rightarrow S)] - \Delta G_t^\circ(Ag^\oplus, W \rightarrow S) \right\} \cdot \frac{1}{100} \qquad (7\text{-}12e)$$

Since water is a hard solvent, the Gibbs energy of transfer of ions from water as a reference solvent to other solvents should depend on the softness of these solvents in a different manner for hard and soft ions. For ions of equal charge and size, hard ions should prefer water and soft ions the softer solvents. The definition of m by Eq. (7-12e) has been given because the size of the soft $Ag^\oplus$ ion is intermediate between those of hard $Na^\oplus$ and $K^\oplus$. The m values for 34 organic solvents have been determined, e.g. $m = -0.12$ for 2,2,2-trifluoroethanol, 0.00 for water (by definition), 0.64 for pyridine,

and 1.35 for N,N-dimethylformamide. The degree of softness among solvents with oxygen, nitrogen, and sulfur donor atoms increases in the series O-donor (alcohols, ketones, amides) < N-donor (nitriles, pyridines, amines) < S-donor solvents (thioethers, thioamides). Applications of this softness scale of solvents have been reported [285].

The donor properties of soft EPD solvents have also been described by the softness parameter σ_P of Gritzner [290, 303]. This parameter is based on the standard molar Gibbs energies of transfer of soft Ag^+ ions from benzonitrile as a reference solvent to other soft solvents and should only be used for soft solute/soft solvent interactions. Further solvent softness parameters based on the Raman IR absorption of the symmetrical stretching vibration of the Hg–Br bond in $HgBr_2$ have been developed by Persson et al. [287, 292]; cf. also Section 3.3.2. The relationships between these solvent softness scales have recently been reviewed [304].

7.3 Empirical Parameters of Solvent Polarity from Kinetic Measurements

Since reaction rates can be strongly affected by solvent polarity (Chapter 5), the introduction of solvent scales using suitable solvent-sensitive chemical reactions was obvious [33, 34]. One of the most ambitious attempts to correlate reaction rates with empirical parameters of solvent polarity has been that of Winstein and his co-workers [35…37]. They found that the S_N1 solvolysis of 2-chloro-2-methylpropane (t-butyl chloride, t-BuCl) is strongly accelerated by polar, especially protic solvents (cf. Eq. (5-13) in Section 5.3.1. Grunwald and Winstein [35] defined a solvent "ionizing power" parameter Y using Eq. (7-13),

$$Y = \lg k^{t\text{-BuCl}} - \lg k_0^{t\text{-BuCl}} \qquad (7\text{-}13)$$

where $k_0^{t\text{-BuCl}}$ is the first-order rate constant for the solvolysis of t-butyl chloride at 25 °C in aqueous ethanol (80 cL/L ethanol and 20 cL/L water; $Y = 0$) as reference solvent, and $k^{t\text{-BuCl}}$ is the corresponding rate constant in another solvent. This reaction was selected as the model process because it was believed to occur by an essentially pure S_N1 mechanism, with ionization of the C–Cl bond as the rate-determining step. Choosing a standard reaction and a reference solvent, a linear Gibbs energy relationship is then written in the familiar form of Eq. (7-14):

$$\lg(k/k_0)_{RX} = m \cdot Y + c \qquad (7\text{-}14)$$

In Eq. (7-14), k and k_0 are the specific rate constants for the S_N1 solvolysis of RX (in this case t-BuCl) in a given solvent and in the standard solvent, respectively, m is the sensitivity of the specific rate of solvolysis of RX to changes in the solvent "ionizing power" (Y), Y is a parameter characteristic of the given solvent, and c is the intercept (zero for an ideally behaved solvolysis). Eq. (7-14) is expected to be applicable to reactions very similar to the standard reaction, that is, S_N1 substitutions. The similarity between Y and m of Eq. (7-14), and σ and ϱ of the Hammett equation (7-6) is obvious. Y values are known for some pure, mainly protic solvents and for various binary mixtures of organic solvents with water or a second organic solvent [35, 36]. Typical Y values are

indicated in Table 7-1. It should be noted that the Y value of the standard solvent lies about midway between the extremes. Y values of binary mixtures are not related to solvent composition in a simple manner.

The solvolysis of t-butyl chloride at 120 °C was also chosen as a solvent-dependent model reaction by Koppel and Pal•m [38] in order to calculate a solvent polarity scale which is restricted to nonspeci“c interactions only. In order to con“rm the limiting S_N1 mechanism for the solvolysis of t-butyl chloride, Schleyer and his co-workers [39] compared the solvolysis rates of t-butyl chloride and 1-bromoadamantane in a large series of solvents. They reasoned that any rate-determining elimination or nucleophilic solvent assistance in t-butyl chloride solvolysis will result in a failure to correlate with 1-bromoadamantane. In 1-bromoadamantane, backside nucleophilic solvent attack and elimination are impossible. An excellent correlation between solvolysis data for t-butyl chloride and 1-bromoadamantane has been found, indicating that t-butyl chloride seems to solvolyze by a limiting S_N1 mechanism, free from nucleophilic solvent participation and from rate-determining elimination. Similarly good correlations have been observed between the solvolysis rates of t-butyl chloride and 1-adamantyl tosylate [40] as well as 2-adamantyl tosylate [41]. The absence of any S_N2 character in the solvolysis of t-butyl hepta“uorobutyrates in aqueous ethanol and similar solvents has been con“rmed [151]. In most cases, the solvolysis rate data of t-butyl substrates are best explained by an S_N1 process with electrophilic solvent assistance of the leaving group in protic solvents through the formation of strong hydrogen bonds. That is, the ionization rates of t-butyl halides should be mainly dependent on solvent polarity and solvent electrophilicity, but not on solvent nucleophilicity [152]; cf. Eqs. (5-135) and (5-136) in Section 5.5.7. However, according to Bentley et al. [153], Kevill et al. [155], and Bunton et al. [156] it is now thought that t-butyl chloride reacts solvolytically with weak but signi“cant nucleophilic solvent assistance. Only 1- and 2-adamantyl substrates seem to react by an S_N1 (limiting) mechanism due to their cage structure, which absolutely precludes nucleophilic attack by solvents from the rear. This has been concluded from the observation that the relative reactivities of t-butyl chloride and 1-chloroadamantane are di¤erent in nucleophilic solvents and in weakly nucleophilic solvents such as tri“uoroacetic acid, hexa“uoro-2-propanol, or tri“uoroethanol, in which t-butyl chloride is abnormally unreactive [153]. See also reference [157] for a nice picture of the solvolysis of 1-bromoadamantane as compared to that of t-butyl bromide with and without backside solvent assistance.

The Grunwald…Winstein equation (7-14) is fairly successful in a large number of cases. Good linear relationships between $\lg k$ and Y are found for the solvolysis of various tertiary haloalkanes and secondary alkyl sulfonates, i.e. reactions which proceed by an S_N1 mechanism, as found for the standard reaction. For reactions involving borderline mechanisms (e.g. solvolysis of secondary haloalkanes) and for S_N2 reactions (e.g. solvolysis of primary haloalkanes), the application of Eq. (7-14) is less satisfactory. When solvolysis rates for di¤erent binary mixtures of solvents are correlated using Eq. (7-14), the well-known phenomenon of dispersion is often observed, that is, lines or curves of slightly di¤erent slopes are observed for each solvent system [35, 36]. In other words, the substrate parameter m is solvent dependent in these cases. If Eq. (7-14) were strictly obeyed, all the points should lie on a single straight line. These observations indicate that the reaction rate depends not only on the ionizing power of the solvent (for

Table 7-1. Empirical parameters based on kinetic measurements for solvents of decreasing polarity.

Solvents	Y [35, 36]	lgk_1[a)]	X [51]	$\mathscr{S}$[b)]	Ω [52]
Water	3.493	−1.180	…	…	0.869[k)]
Formic Acid	2.054	−0.929	…	…	…
2,2,2-Tri"uoroethanol	1.045[c,d)]	…	…	…	…
Formamide	0.604	…	…	…	0.825[k)]
Methanol/Water (80 cL/L + 20 cL/L)	0.381	…	…	…	…
Ethanol/Water (80 cL/L + 20 cL/L)	0.000	−2.505	…	…	…
Tetra-n-hexylammonium benzoate	−0.39[e)]	…	…	…	…
Acetone/Water (80 cL/L + 20 cL/L)	−0.673	…	…	…	…
1,4-Dioxane/Water (80 cL/L + 20 cL/L)	−0.833	…	…	…	…
Methanol	−1.090	−2.796	0.91	−1.89	0.845
Acetic acid	−1.675	−2.772	0.00	…	0.823
Ethanol	−2.033	−3.204	…	−2.02	0.718
2-Propanol	−2.73	…	…	…	…
t-Butanol	−3.26	…	…	…	…
Dimethyl sulfoxide	…	−3.738	1.6	…	…
Nitromethane	…	−3.921	…	0.041	0.680
Acetonitrile	…	−4.221	0.04	−0.328	0.692
N,N-Dimethylformamide	−3.5[f)]	−4.298	0.8	−0.222	0.620
Acetic anhydride	…	−4.467	…	…	…
1,2-Dichloroethane	…	…	…	−0.420	0.600
Dichloromethane	…	…	…	−0.553	…
Pyridine	…	−4.670	…	…	0.595
Acetone	…	−5.067	…	−0.824	0.619
Trichloromethane	…	…	…	−0.886	…
Chlorobenzene	…	…	−1.9	−1.15	…
1,2-Dimethoxyethane	…	…	…	…	0.543
1,4-Dioxane	…	…	…	−1.43	…
Ethyl acetate	…	−5.947	…	−1.66	…
Tetrahydrofuran	…	−6.073	…	−1.54	…
Benzene	…	…	…	−1.74	0.497[g)]
Tetrachloromethane	…	…	−4.8	−2.85	…
Diethyl ether	…	−7.3	…	−2.92	0.466[h)]
Cyclohexane	…	…	…	−4.15	0.595[i)]
Decalin	…	…	…	…	0.537[i)]
Triethylamine	…	…	…	…	0.445
n-Hexane	…	…	…	ca. −5	…

[a)] Ionization of 4-methoxyneophyl tosylate at 75°C [37].
[b)] S_N2 reaction of tris(n-propyl)amine with iodomethane at 20°C [49, 50].
[c)] V. J. Shiner, W. Dowd, R. D. Fisher, S. R. Hartshorn, M. A. Kessick, L. Milakofsky, and M. W. Rapp, J. Am. Chem. Soc. 91, 4838 (1969).
[d)] D. E. Sunko, I. Szele, and M. Tomić, Tetrahedron Lett. 1972, 1827, 3617.
[e)] C. G. Swain, A. Ohno, D. K. Roe, R. Brown, and T. Maugh, J. Am. Chem. Soc. 89, 2648 (1967).
[f)] S. D. Ross and M. M. Labes, J. Am. Chem. Soc. 79, 4155 (1957).
[g)] W. M. Jones and J. M. Walbrick, J. Org. Chem. 34, 2217 (1969).
[h)] J. F. King and R. G. Pews, Can. J. Chem. 43, 847 (1965).
[i)] R. Braun and J. Sauer, Chem. Ber. 119, 1269 (1986); value at 20°C.
[k)] A. Lattes, I. Rico, A. de Savignac, and A. A. Samii, Tetrahedron 43, 1725 (1987).

which Y is a measure), but also on the nucleophilicity of the solvent. Obviously, nucleophilic solvent assistance may be relevant in reactions with increasing S_N2 characteristics. Therefore, the trends in m values suggest their usefulness in determining the extent of solvent nucleophilic participation. Those reactions which are S_N1 exhibit m values near 1.00, while values for S_N2 substrates range from 0.25 to 0.35. Values of m between these extremes are typical for secondary substrates lying in the borderline area between limiting S_N1 and pure S_N2 mechanisms. The term "S_N2(intermediate) mechanism" has been suggested for reactions where there is evidence of nucleophilic solvent assistance and evidence for a nucleophilically solvated ion pair reaction intermediate (cf. Eq. (5-136) in Section 5.5.7. Thus, a gradual change in mechanism going from S_N2 (one-stage) through S_N2(intermediate) to an S_N1(limiting) mechanism has been postulated [41, 153]; cf. Section 5.5.7 for a discussion of this solvent-influenced S_N2/S_N1 mechanistic spectrum.

To account for nucleophilically solvent-assisted processes, Grunwald, Winstein et al. [42] later provided a four-parameter equation of the type shown in Eq. (7-15)*

$$\lg(k/k_0)_{RX} = m \cdot Y + l \cdot N + c \qquad (7\text{-}15)$$

In Eq. (7-15), the additional term $l \cdot N$ involves the sensitivity l, this indicating the substrates response to changes in solvent nucleophilicity (N). As in Eq. (7-14), Y denotes the solvent ionizing power, and ethanol/water (80:20 cL/L) is again chosen as the standard solvent with $N = 0$ (and $Y = 0$) for this binary solvent mixture. The intercept c should be zero for well-behaved solvolyses. It is somewhat unfortunate that m has been retained as the sensitivity to solvent ionizing power, since, in reactions where $l \neq 0$, different values form will be obtained depending on whether the correlation analysis is in terms of Eq. (7-14) or (7-15). Obviously, the simple Grunwald…Winstein equation (7-14) can only be used when the $l \cdot N$ term of the extended equation (7-15) makes either a negligible $(l = 0)$ or constant contribution, which is then included in c. The extended Grunwald…Winstein equation (7-15) has been explicitly evaluated by both Peterson et al. [43] and Bentley, Schleyer et al. [44], using two reference substrates, one with a high sensitivity to solvent nucleophilicity (e.g. bromomethane [43] or methyl tosylate [44]; $l = 1.00$), and the other with a low sensitivity to solvent nucleophilicity (e.g. t-butyl chloride [43] or 2-adamantyl tosylate [44]; $l = 0.00$ and $m = 1.00$). Eq. (7-15) correlates solvolysis data for such widely varying substrates as methyl and 2-adamantyl tosylates [44].

The evaluation of the bulk nucleophilicity of solvents (N) in terms of Eq. (7-15) is more difficult than the establishment of the Y scale of solvent ionizing power according to Eq. (7-14) [305]. As already mentioned, for substrates RX which ionize without

---

* The original equation was expressed as the partial differential equation [42]:

$$d \lg k = \left(\frac{\partial \lg k}{\partial Y}\right)_N dY + \left(\frac{\partial \lg k}{\partial N}\right)_Y dN$$

If the partial derivatives are constant and equal to m and l, respectively, this expression leads to Eq. (7-15).

nucleophilic solvent assistance, one can assume that $m = 0$ for solvolysis of this substrate and Eq. (7-15) simplifies to Eq. (7-14). This requirement is closely satisfied by substrates RX with R = 1- or 2-adamantyl. However, for the solvolysis of a neutral substrate RX with an anionic leaving group X^-, there is no corresponding system where $m \to 0$. In order to establish an N scale for the solvolysis of a standard substrate RX, Eq. (7-15) can be rearranged to give Eq. (7-16a), if one takes $l = 1$ by definition for this substrate:

$$N_{RX} = \lg(k/k_0)_{RX} - m_{RX} \cdot Y_x \qquad (7\text{-}16a)$$

Y_x values are available for the common leaving groups X [306] and the value of $\lg(k/k_0)_{RX}$ can be determined experimentally. The difficulty is to obtain the appropriate sensitivity value m_{RX} for use in Eq. (7-16a).

Based on the reasonable assumption that primary methyl substrates should be subject to the most pronounced nucleophilic solvent assistance towards their solvolyses, methyl tosylate (MeOTs) was chosen by Bentley, Schleyer et al. as a standard substrate ($l = 1$) for the evaluation of N_{RX} [44, 154]. Methyl tosylate solvolyzes cleanly in a wide variety of solvents without steric hindrance. From a plot of $\lg(k/k_0)_{MeOTs}$ against Y_{OTs}, a value of $m_{RX} = 0.3$ was chosen to give Eq. (7-16b) [44, 154, 305, 307]:

$$N_{OTs} = \lg(k/k_0)_{MeOTs} - 0.3 \cdot Y_{OTs} \qquad (7\text{-}16b)$$

Values of N derived in this way are designated as N_{OTs} [158] or N_{BS} [305] values; they range from -5.56 (CF_3CO_2H) to $+0.2$ (2-propanol). The values of Y in this equation are based on 2-adamantyl tosylate (2-AdOTs) instead of t-butyl chloride solvolysis in order to retain the same leaving group (TsO). Although it is a secondary substrate, 2-adamantyl tosylate solvolyzes according to an S_N1 (limiting) or k_c mechanism* through the rate-determining formation of contact ion pairs without detectable nucleophilic solvent assistance [154]. The Y_{OTs} scale of solvent ionizing power spans a rate range of over eight orders of magnitude, from $Y_{OTs} = +4.57$ (CF_3CO_2H) to -3.74 (t-butanol) [158], and can even be extended to aqueous sulfuric acid as the ionizing medium [159]. The Y_{OTs} scale correlates nonlinearly with the original $Y_{t\text{-}BuCl}$ values [158]. Using the extended Eq. (7-16c) N_{OTs} values in conjunction with Y_{OTs} values gave good to excellent correlations for the solvolyses of a great variety of sulfonates [44, 154].

$$\lg(k/k_0)_{ROTs} = m \cdot Y_{OTs} + l \cdot N_{OTs} + c \qquad (7\text{-}16c)$$

In spite of these successful correlations, the choice of $m_{RX} = 0.3$ for use in Eq. (7-16b) was criticized for some reasons; see references [160, 305] for details. Therefore, Kevill et al. have introduced a considerable improvement by making a fundamental

---

* According to Winstein, real solvolyses are either nucleophilically solvent assisted (designated k_s and including both substitution and elimination processes) or anchimerically assisted (designated k_Δ), with k_c representing the hypothetical limit which is reached when nucleophilic solvent assistance and anchimeric assistance approach zero. Anchimeric assistance (from the Greek anchi + meros, neighbouring parts) means neighbouring group participation during the ionization step (cf. Eq. (7-20) for an example.

change in the type of substrate undergoing S_N1 solvolysis; see [160, 305] and references cited therein. Using substrates of the type $R{-}X^+$ (i.e. sulfonium and oxonium salts), the anionic leaving group X^- of RX is replaced by a neutral molecule leaving group according to $R{-}X^+ \rightarrow R^+ + X$. With a neutral leaving group X, the mY term in Eq. (7-16b) can be ignored. According to the Hughes…Ingold rules, the positive charge is dispersed during the activation process and a small but signi"cant rate decrease with increasing solvent polarity is to be expected; see reaction type (b) in Table 5-4 in Section 5.3.1. Among the •onium salts studied (e.g. triethyloxonium hexa"uorophosphate [160], etc.), the S-methyl-dibenzothiophenium ion (MeDBTh$^+$; as its tri"uoromethanesulfonate) has been found to be the most useful because of its structural analogy to methyl tosylate (MeX compared to MeX$^+$); cf. Eq. (7-16b). With this new standard substrate, a new solvent nucleophilicity scale (N_T) was de"ned by Kevill et al. [305, 308], as expressed in Eq. (7-17):

$$N_T = \lg(k/k_0)_{\mathrm{MeDBTh}^+} \qquad (7\text{-}17)$$

In Eq. (7-17), k and k_0 are the speci"c rate constants of S_N1 solvolysis of the S-methyl-dibenzothiophenium ion in a given solvent and in a standard solvent (EtOH/H_2O, 80:20 cL/L; $N_T = 0$), respectively. N_T values are known for 37 pure and binary solvents and span a range from $N_T = -5.26$ [$(CF_3)_2CHOH/H_2O$, 97:3 cg/g] to +0.37 (EtOH) [305, 308]. The N_T solvent nucleophilicity scale can be used directly to correlate the solvolytic behaviour of other RX^+ substrates with a neutral nucleofuge, and also in applications of the extended Grunwald…Winstein equation (7-15). Many examples of the use of N_T values in the correlation of speci"c rates of solvolysis of a variety of RX and $R{-}X^+$ substrates are given in reference [305].

Interestingly, an analysis of the speci"c rate constants for t-butyl chloride solvolysis in 46 solvents based on the extended Grunwald…Winstein equation (7-15), using Y_{Cl} and N_T, gives a very good correlation (correlation coe‹cient $r = 0.988$) with values of $m = 0.86$, $l = 0.38$, and $c = 0$ [309]. This result con"rms the involvement of nucleophilic solvation of the developing carbocation in the S_N1 solvolysis of t-butyl chloride, in addition to the strong electrophilic solvation of the anionic leaving group [305].

Very precise experimental studies clearly show that Winstein•s original Y values, as de"ned by Eq. (7-14), do not apply to leaving groups other than the chloride ion [153, 158…166], most probably due to the variable amount of speci"c solvation of di¤erent leaving groups by solvent molecules. If allowance is made for di¤erent leaving groups X, the solvolytic rate data of substrates RX can be correlated with data for a reference compound according to the general Eq. (7-18).

$$\lg(k/k_0)_{RX} = m \cdot Y_X \qquad (7\text{-}18)$$

Y_X values for di¤erent leaving groups X have been de"ned using $m = 1.00$ for the solvolyses of 1-adamantyl substrates (X = Cl [153], Br [153], I [161], picrate [162], and $^{\oplus}SMe_2$ [163]) and 2-adamantyl substrates (X = OTs [44, 158], $OClO_3$ [162, 166], OSO_2CF_3 = OTf [164, 166], and picrate [166]). Additional Y_{OTs} values have been obtained from solvolyses of 1-adamantyl tosylate (X = OTs [158, 162]). Further Y_{OTf} values for tri"uoromethanesulfonates (tri"ates) have been determined from the sol-

volysis of 7-norbornyl tri”ate [165]. This range of Y_X values ensures that speci“c leaving group solvation e¤ects do not interfere with mechanistic deductions from the solvolysis rate data. Su‹cient data are now available to evaluate trends in the various Y_X scales. A comparison of the Y_X scales and a discussion of the correlations between them can be found in references [166, 306]. For example, the relative rates for the S_N1 solvolyses of di¤erent bridgehead substrates having the same leaving group are found to be almost solvent-independent [167].

An analysis of Winstein•s Y and N values and their modi“cations in terms of Kamlet and Taft•s linear solvation energy relationships (cf. Section 7.7) has been given by Abraham et al. [288].

The medium in”uence on the solvolyses of a variety of further substrates, i.e. benzylic [310…313], allylic and propargylic compounds [312], as well as very crowded tertiary haloalkanes [314], has been studied and analyzed in the framework of the extended Grunwald…Winstein equation (7-15).

A four-parameter relationship (7-19) similar to Eq. (7-15) was proposed by Swain et al. in 1955 [45].

$$\lg(k/k_0) = c_1 \cdot d_1 + c_2 \cdot d_2 \qquad (7\text{-}19)$$

d_1 and d_2 are measures of solvent nucleophilicity and electrophilicity, respectively, and c_1 and c_2 measure the substrate•s sensitivity to these solvent properties. The rate constant k_0 refers to the reaction in the standard solvent, 80:20 cL/L ethanol/water. This approach is essentially statistical since, in contrast to Eq. (7-15), four rather than two parameters (m and l) are varied. Numerous values of d_1, d_2, c_1, and c_2 were calculated, using scales based on certain standard systems, in such a way as to “t a large number of experimental rate constants [45]. Although a satisfactory correlation was achieved for a wide range of solvents and substrates, this approach has been criticized for a lack of connection with chemical reality [42, 46]. Thus, the substrate factors c_1 and c_2 are of little mechanistic signi“cance, e.g. t-butyl chloride has a higher c_1 value than bromomethane, suggesting that bromomethane (S_N2 substrate) should be less sensitive to solvent nucleophilicity than t-butyl chloride (S_N1 substrate). Overall, this treatment was not very successful and Swain et al. later conceded that the quantitative justi“cation for the critical subsidiary conditions chosen for the iterative calculation technique used was questionable [265].

An alternative model system, potentially superior to t-butyl chloride in the evaluation of solvent ionizing power, is based on the speci“c rate of anchimeric (p-aryl assisted solvolysis or decomposition of 4-methoxyneophyl tosylate at 75 °C according to Eq. (7-20) [37]. Use of this compound avoids some practical disadvantages of t-butyl



(7-20)

chloride: t-butyl chloride, being rather volatile, is troublesome to weigh accurately; it reacts extremely rapidly in highly polar solvents, and its solvolysis rate constant cannot usually be determined in nonhydroxylic solvents. Therefore, Winstein et al. [37] put forward the values of $\lg k_1$ for the ionization of 4-methoxyneophyl tosylate as a reference reaction suitable for measuring solvent polarity even in fairly nonpolar and non-HBD solvents (cf. Table 7-1)*. The $\lg k_1$ (or $\lg k_{ion}$) scale is equivalent to the Y scale for protic solvents and has been extended to non-HBD solvents, for which Y values are not available. Although a primary substrate, 4-methoxyneophyl tosylate is believed to solvolyze through concerted anchimeric b-aryl participation without nucleophilic solvent interference and internal ion-pair return [37]. A more sophisticated picture of solvent effects on b-arylalkyl tosylate solvolyses has been given by Schleyer et al. He demonstrated that primary and secondary b-arylalkyl substrates solvolyze through competition between discrete b-aryl assisted (k_Δ) and b-aryl non-assisted reaction pathways (k_s), each of which leads to distinct sets of products [168]. This is due to the fact that the neighbouring aryl group and the solvent molecules are in competition to displace the leaving group. By decreasing the solvent nucleophilicity as well as introducing electron-releasing substituents in the aryl group, the kinetic influence of the b-aryl group becomes more noticeable. Therefore, the $\lg k_1$ (or $\lg k_{ion}$) values, as measures of the solvent ionizing power, are subject to the same limitations as found for the Y values; cf. the preceding discussion of Eq. (7-14).

In addition to the application of S_N1 reactions as model reactions for the evaluation of solvent polarity, Drougard and Decroocq [48] suggested that the value of $\lg k_2$ for the S_N2 Menschutkin reaction of tri-n-propylamine and iodomethane at 20°C … termed "$\mathscr{S}$" according to Eq. (7-21) … should also be used as a general measure of solvent polarity.

$$\mathscr{S} = \lg k_2[(CH_3CH_2CH_2)_3N + CH_3I] \qquad (7\text{-}21)$$

The second-order rate constants of this quaternization reaction have been determined for seventy-eight solvents by Lassau and Jungers [49, 50]. A selection from this relatively extensive solvent scale is given in Table 7-1.

Analogous to the Y scale and based on a nucleophilic substitution reaction, Gielen and Nasielski [51] suggested that a solvent polarity scale could be based on electrophilic aliphatic substitution reactions such as the reaction of bromine with tetramethyltin shown in Eq. (7-22).



(7-22)

This model reaction probably passes through a non-cyclic dipolar activated complex (S_E2 mechanism) in polar solvents, while in less polar solvents a cyclic activated complex gains importance (S_i2 mechanism) [51]. The linear Gibbs energy relationship

* In his appreciation of Winstein's scientific work, Bartlett [47] proposed the term W for "Winstein parameter" for these values commonly referred to as "k_1" or "$\lg k_{ion}$".

given by these authors is as follows:

$$\lg(k/k_0) = p \cdot X \tag{7-23}$$

k and k_0 are the rate constants of reaction (7-22) in a given solvent and in glacial acetic acid, respectively. The latter is used as the reference solvent ($X = 0$ by definition). X is a parameter characteristic of the given solvent, and p is a parameter characteristic of the given reaction; $p = 1$ for Eq. (7-22) by definition. X values have only been determined for seven solvents to date (see Table 7-1). The two solvent scales X and Y, are not quite in harmony, since the reaction rate is influenced by the electrophilic and nucleophilic character of the solvent to differing degrees. The electrophilic substitution reaction (7-22) is probably supported by a nucleophilic pull of the solvent on the leaving group ($-SnMe_3$). The coupling constants $^2J_{^{117}Sn-^1H}$ obtained from ^1H NMR spectra of $(CH_3)_3SnBr$ in various solvents, which can be taken as a measure of the nucleophilic character of the solvent with respect to tin, show a correlation with the X values, in agreement with this assumption [51].

The fact that the rate of some Diels…Alder [4+2] cycloaddition reactions is affected, albeit only slightly, by the solvent was used by Berson et al. [52] in establishing an empirical polarity parameter called W. These authors found that, in the Diels…Alder addition of cyclopentadiene to methyl acrylate, the ratio of the endo product to the exo product depends on the reaction solvent. The endo addition is favoured with increasing solvent polarity; cf. Eq. (5-43) in Section 5.3.3. Later on, Pritzkow et al. [53] found that not only the endo/exo product ratio but also the absolute rate of the Diels…Alder addition of cyclopentadiene to acrylic acid derivatives increases slightly with increasing solvent polarity. The reasons for this behaviour have already been discussed in Section 5.3.3. Since reaction (5-43) is kinetically controlled, the product ratio [endo]/[exo] equals the ratio of the specific rate constants, and Berson et al. [52] define

$$\lg(k_{endo}/k_{exo}) = \lg[endo]/[exo] \equiv W \tag{7-24}$$

The values of W lie between 0.445 (triethylamine) and 0.869 (water), and are known for fourteen solvents (cf. Table 7-1). Owing to the low solubility of the reactants in polar media, an extension of this scale is limited.

Related to Diels…Alder [2+2] cycloadditions are 1,3-dipolar cycloadditions, which are known to be far less solvent-dependent (cf. Eq. (5-44) in Section 5.3.3. Nagai et al. [169] found that the 1,3-dipolar cycloaddition reaction of diazo-diphenylmethane to tetracyanoethene (TCNE) is an exception; it is 180 times faster in nonbasic trichloromethane than in the EPD solvent 1,2-dimethoxyethane (cf. Eq. (7-25). The second-order



(7-25)

rate constant decreases with increasing solvent basicity. This solvent effect can be interpreted in terms of the solute/solvent interaction between the soft Lewis acidic p-acceptor TCNE and basic EPD solvents. Thus, one of the educts is stabilized through specific solvation resulting in a corresponding rate deceleration. Using reaction (7-25) as the reference process, an empirical parameter D_p of solvent Lewis basicity for interactions with soft p-acceptors has been proposed according to Eq. (7-26):

$$\lg(k_0/k) = D_p \qquad (7\text{-}26)$$

k_0 and k are the rate constants of reaction (7-25) at 30°C in benzene (reference solvent with $D_p = 0$) and other solvents, respectively [169]. D_p values are known for 34 solvents [169]. They are relatively large for aromatic solvents (i.e. soft EPD solvents) compared to other empirical Lewis basicity parameters that have been determined by employing rather hard Lewis acid probes, e.g. $SbCl_5$-derived DN values [cf. Eq. (7-10)], CH_3OD-derived B values [cf. Eq. (7-34)], and 4-nitrophenol-derived b values [cf. Eq. (7-51)]. The D_p values have been applied successfully to reactions of diazo-diphenylmethane with various soft p-acceptors such as other cyano-substituted alkenes as well as quinones [170].

Finally, it should be mentioned that the solvent-dependent intramolecular Beckman…Chapman rearrangement of anti-methylketoxime O-picrates, $H_3C(R)C{=}N{-}OX$ (e.g. with R = 1-adamantyl and X = 2,4,6-trinitrophenyl), has been proposed as a standard reaction for measuring solvent polarities [315]. This anionotropic sextet rearrangement proceeds 62 times faster in dimethyl sulfoxide than in tetrahydrofuran. The solvent-dependent rate-determining step is the migration of R to nitrogen with simultaneous loss of the phenolate XO^- as a leaving group, to yield a nitrilium ion as an intermediate. The advantages of this probe reaction have been discussed [315].

7.4 Empirical Parameters of Solvent Polarity from Spectroscopic Measurements

Spectroscopic parameters of solvent polarity have been derived from solvent-sensitive standard compounds absorbing radiation in spectral ranges corresponding to UV/Vis, IR, ESR, and NMR spectra (cf. Chapter 6) [1…9]. The first suggestion that solvatochromic dyes should be used as indicators of solvent polarity was made by Brooker et al. [54] in 1951, but Kosower [5, 55] in 1958 was the first to set up a comprehensive solvent scale.

Since then, solvent-dependent absorptions of a great variety of compounds have been studied as potential reference processes for establishing empirical scales of solvent polarity. Most approaches include negatively or positively solvatochromic dyes because they are experimentally easy to handle. The use of solvatochromic dyes as solvent polarity indicators has been reviewed [1, 2, 293…296, 316], see, in particular, reference [293]; only a few of these dyes can be mentioned in this Section.

Kosower [5, 55] has taken the longest-wavelength intermolecular charge-transfer (CT) transition of 1-ethyl-4-(methoxycarbonyl)pyridinium iodide as a model process. It exhibits a marked negative solvatochromism (cf. the formula of this dye and its UV/Vis

absorption spectra in Fig. 6-3 in Section 6.2.1. A solvent change from pyridine to methanol causes a hypsochromic shift of the longest-wavelength CT band of $\Delta\tilde{\nu} = -88$ nm ($\Delta\tilde{\nu} = +5960\ cm^{-1}$). This is due to stabilization of the electronic ground state, which is an ion pair, relative to the "rst excited state, which is a radical pair, with increasing solvent polarity. The general conditions for the appearance of solvatochromism have already been discussed in Section 6.2.2. Kosower de"ned his polarity parameter Z, as the molar transition energy E_T, expressed in kcal/mol, for the CT absorption band of 1-ethyl-4-(methoxycarbonyl)pyridinium iodide in the appropriate solvent according to Eq. (7-27):

$$E_T/(kcal \cdot mol^{-1}) = h \cdot c \cdot \tilde{\nu} \cdot N_A = 2.859 \cdot 10^{-3} \cdot \tilde{\nu}/cm^{-1} \equiv Z \qquad (7\text{-}27)$$

h is Planck•s constant, c is the velocity of light, $\tilde{\nu}$ is the wavenumber of the photon which produces the electronic excitation, and N_A is Avogadro•s number. A Z value of 83.6 for methanol means that a transition energy of 83.6 kcal is necessary to promote one mole of the standard dye, dissolved in methanol, from its electronic ground state to its "rst excited state*. The stronger the stabilizing e¤ect of the solvent on the ground-state ion-pair as compared with the less dipolar radical pair in the excited state, the higher this transition energy and, thus, the Z value. A high Z value corresponds to high solvent polarity. The basis for the use of Z as a measure of solvent polarity has already been given in Section 6.2.1; cf. also [5, 55, 171, 172]. An alternative explanation for the negative solvatochromism of substituted pyridinium iodides has been given [171].

Z values cover a range from 94.6 (water) to about 60 kcal/mol (1-octane) and were originally measured for 21 pure solvents and 35 binary solvent mixtures [5, 56], as well as some electrolytes [57] and surfactant solutions [58]. Various authors have since gradually extended this to include 45 pure solvents. Z values for a further 41 pure solvents have been determined by Gri<ths and Pugh [172], who also compiled all available Z values and their relationships with other solvent polarity scales. A selection of Z values together with some other spectroscopic solvent polarity parameters is given in Table 7-2.

There are some serious limitations in the determination of Z values. First, Z values can only be obtained by direct measurement over the solvent range trichloromethane ($Z = 63.2$ kcal/mol) to 70:30 cL/L ethanol/water ($Z = 86.4$ kcal/mol). In highly polar solvents, the long-wavelength charge-transfer band moves to such short wavelengths that it cannot be observed underneath the much stronger $\pi \rightarrow \pi^*$ absorption band of the pyridinium ion. Therefore, the Z value for water was obtained by extra-

---

* For conversion according to the International System of Units, the following equations can be used:

$$1\ kcal \cdot mol^{-1} = 4.184\ kJ \cdot mol^{-1};$$

$$E_T/(kJ \cdot mol^{-1}) \quad \text{or} \quad Z/(kJ \cdot mol^{-1}) = 1.196 \cdot 10^{-2} \cdot \tilde{\nu}/cm^{-1}.$$

The choice of $kcal \cdot mol^{-1}$ as unit for the Z scale is related to the usefulness of this quantity for comparison to chemical reactions and their activation energies. Since this and other parameters have been widely used in the literature, conversion of the units into $kJ \cdot mol^{-1}$ has been avoided in order to avert confusion.

polating the Z values measured for acetone/water, ethanol/water, and methanol/water mixtures to zero organic component in a plot against Winstein-Y values. Because the lines were extrapolated a considerable distance, the original Z value for water (94.6 kcal/mol) has been reexamined and a lower Z value (91.8 kcal/mol) is advocated by Griffiths and Pugh [172]. Secondly, the standard pyridinium iodide is not soluble in many nonpolar solvents. By using the more soluble 4-(butoxycarbonyl)-1-ethylpyridinium iodide and pyridine-1-oxide as secondary standards, it was possible to calculate Z values of nonpolar solvents [5, 55].

Z values are both temperature- and pressure-dependent. The CT absorption band of substituted pyridinium halides is shifted hypsochromically as the temperature of the solution is decreased [59]. Thus Z values decrease with increasing temperature due to a lowering of the solute/solvent interactions at the higher temperature. Furthermore, it has been shown that the CT absorption band of 1-ethyl-4-(methoxycarbonyl)-pyridinium iodide is shifted bathochromically for solutions in methanol and ethanol with increasing pressure (up to 1920 bar), while for other solvents such as acetone and N,N-dimethylformamide it is shifted hypsochromically [60]. Except for the lower alcohols, the bulk solvent polarity generally increases with pressure [60].

Z values have been widely used to correlate other solvent-sensitive processes with solvent polarity, e.g. the $n \rightarrow \sigma^*$ absorption of haloalkanes [61], the $n \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$ absorption of 4-methyl-3-penten-2-one [62], the $\pi \rightarrow \pi^*$ absorption of phenol blue [62], the CT absorption of tropylium iodide [63], as well as many kinetic data (Menschutkin reactions, Finkelstein reactions, etc. [62]). Copolymerized pyridinium iodides, embedded in the polymer chain, have also been used as solvatochromic reporter molecules for the determination of microenvironment polarities in synthetic polymers [173]. No correlation was observed between Z values and the relative permittivity ε_r or functions thereof [317]. Measurement of solvent polarities using empirical parameters such as Z values has already found favour in textbooks for practical courses in physical organic chemistry [64].

Brownstein suggested Eq. (7-28) analogous to the Hammett equation (7-6), for the general description of solvent effects [65]:

$$\lg k - \lg k_0 = \lg(k/k_0) = S \cdot R \qquad (7\text{-}28)$$

k is the rate constant, equilibrium constant, or the function of a spectral shift for a reaction or absorption in the relevant solvent, and k_0 is the corresponding quantity for dry ethanol as reference solvent. S ($= 0.00$ for dry ethanol by definition). S is characteristic of the solvent and R gives the susceptibility of the given property towards a change of solvent. As a standard process, Brownstein [65] chose the CT absorption of Kosower's dye [55], and assigned to it an R value of 1.00. Having chosen a standard solvent and a standard reaction, it was then possible to calculate R and S values for other reactions and solvents, respectively. From Kosower's work, 58 S values were used to determine R values for 9 reactions. In a continuation of this process, 155 S values and 78 R values were deduced, including R values for solvent-dependent UV/Vis, IR, and NMR absorptions, rates of reactions, and positions of equilibria [65]. The S values represent statistical averages of a variety of different solvent polarity parameters, including Z and Y, and therefore, cannot be related to a specific model process. In principle, this is

Table 7-2. Empirical parameters based on spectroscopic measurements for solvents of decreasing polarity (25°C, 1013 hPa).

Solvents	Z/(kcal · mol^{-1}) [5, 55, 172]	S [65]	w_R/(kcal · mol^{-1}) [77]	w_B/(kcal · mol^{-1}) [77]	F [79]	G [85]	a(^{14}N)/mT [89]
Water	94.6	0.1540	…	68.9	0.545	…	1.7175
2,2,2-Tri"uoroethanol	…	…	…	…	0.475	…	…
2,2,3,3-Tetra"uoro-1-propanol	86.3	…	…	…	…	…	…
1,2-Ethanediol	85.1	0.0679	40.4	…	0.295	…	1.6364
Ethanol/Water 80:20 cL/L	84.8	0.0650	…	…	…	…	…
2-Aminoethanol	84.4[a]	…	…	…	…	…	…
Methanol	83.6	0.0499	43.1	63.0	0.285	…	1.6210
Formamide	83.3	0.0463	…	…	0.245	…	…
Ethanol	79.6	0.0000	43.9	60.4	0.255	…	1.6030
Acetic acid	79.2	0.0050	…	…	0.37	…	1.6420
2-Octanol	78.6[b]	…	…	…	…	…	…
1-Propanol	78.3	−0.0158	44.1	…	…	…	…
Diethyleneglycol monomethyl ether	78.1[c]	…	…	…	…	…	…
N-Methylacetamide	77.9[b]	…	…	…	…	…	…
1-Butanol	77.7	−0.0240	44.5	56.8	…	…	1.6018
Tetrahydrothiophene-1,1-dioxide	77.3[d]	…	…	…	…	…	…
2-Propanol	76.3	−0.0413	44.5	56.1	0.19	…	1.5973
Propylene carbonate	72.4[e]	…	…	…	…	…	…
2-Methyl-2-propanol, t-Butanol	71.3	−0.1047	…	…	0.130	…	1.5860
Acetonitrile	71.3	−0.1039	45.7	53.7	0.135	93	1.5666
Nitromethane	71.2[e]	−0.134	44.0	…	…	99	1.5759
Dimethyl sulfoxide	70.2	…	42.0	…	0.115	…	1.5692
N,N-Dimethylformamide	68.4	−0.1416	43.7	51.5	0.135	…	1.5635
N,N-Dimethylacetamide	66.9[b]	…	43.0	…	…	…	…
Tri-n-hexyl-n-heptylammonium iodide (mp. 95.7°C; in molten form)	66.4[f]	…	…	…	…	…	…
Acetone	65.5	−0.1748	45.7	50.1	…	…	1.5527
Nitrobenzene	…	−0.218	42.6	…	…	100[g]	…
Diethyleneglycol diethyl ether	65.2[c]	…	…	…	…	…	…
Benzonitrile	65.0[b]	…	43.3	…	…	…	…
Dichloromethane	64.7	−0.1890	44.9	47.5	…	100	1.5752

1,1,2,2-Tetrachloroethane	64.2[f]	−0.083	...	...	...	...	...
Pyridine	64.0	−0.1970	43.9	50.0	...	94	1.5608
cis-1,2-Dichloroethene	63.9[g]	...	...	...	...	...	...
1,2-Dichloroethane	63.8[d]	−0.151	...	...	...	95	1.5655
Trichloromethane	63.2	−0.2000	44.2	...	0.095	106	1.5863
Hexamethylphosphoric triamide	62.8[a]	...	...	...	0.115	...	...
1,1-Dichloroethane	62.1[d]	...	...	...	...	...	...
Triethyleneglycol dimethyl ether	61.3[a]	...	...	...	...	[c] 90	...
Fluorobenzene	60.2[a]	...	...	...	...	...	...
Di-n-butyl ether	60.1[b]	−0.286	48.6	...	...	61	...
1,2-Dichlorobenzene	60.0[a]	...	...	...	...	...	...
Ethyl acetate	59.4[b]	−0.210	47.2	...	...	...	...
Bromobenzene	59.2[a]	−0.164	44.6	...	...	...	1.5479
1,2-Dimethoxyethane	59.1	...	...	...	...	...	1.5424
Ethoxybenzene, Phenetole	58.9[a]	...	...	...	...	[c] 89	...
Tetrahydrofuran	58.8[a]	...	46.6	...	0.10	...	1.5373
Chlorobenzene	58.0[a]	−0.182	45.2	...	...	...	1.5472
2-Methyltetrahydrofuran	55.3	...	...	...	...	...	...
1,4-Dioxane	...	−0.179	48.4	...	0.115	86	1.5452
Diethyl ether	...	−0.277	48.3	...	0.08	64	1.5334
Benzene	54	−0.215	46.9	...	0.02	80	1.5404
Toluene	...	−0.237	47.2	41.7	...	74	1.5347
Carbon disul“de	...	−0.240	...	...	...	74	1.5289
Tetrachloromethane	...	−0.245	48.7	...	−0.01	69	1.5331
Triethylamine	...	−0.285	49.3	...	...	62	...
n-Hexane	...	−0.337	50.9	...	0.07	44	1.5134
Cyclohexane	...	−0.324	50.0	...	0.035	49	...
Gas phase	...	−0.556	...	...	...	0	...

[a] W. N. White and E. F. Wolfarth, J. Org. Chem. 35, 2196 (1970).
[b] R. S. Drago, D. M. Hart, and R. L. Carlson, J. Am. Chem. Soc. 87, 1900 (1965); R. S. Drago and K. F. Purcell, in T. C. Waddington (ed.): Non-Aqueous Solvent Systems, Academic Press, London, New York, 1965, p. 211 ¤.
[c] J. F. King and R. G. Pews, Can. J. Chem. 43, 847 (1965).
[d] J. E. Gordon, J. Am. Chem. Soc. 87, 4347 (1965).
[e] C. Walling and P. J. Wagner, J. Am. Chem. Soc. 86, 3368 (1964).
[f] P. H. Emslie and R. Foster, Rec. Trav. Chim. Pays-Bas 84, 255 (1965).
[g] J. E. Dubois and A. Bienvenüe, Tetrahedron Lett. 1966, 1809.

an interesting attempt at generalization but many of the correlations used to calculate R and S values are rather poor. It would seem that too many different solvent-dependent processes are being mixed-up and treated in an oversimplified way.

The practical limitations in the Z-value approach can be overcome by using pyridinium N-phenolate betaine dyes such as (44) as the standard probe molecule. They exhibit a negatively solvatochromic $\pi \rightarrow \pi^*$ absorption band with intramolecular charge-transfer character (cf. discussion of this dye in Section 6.2.1, its UV/Vis spectrum in Fig. 6-2, and its dipole moment in the electronic ground and excited states mentioned in Table 6-1, dye no. 12.



(44)



(45): R= -C(CH$_3$)$_3$
(46): R= -CF$_3$



(47)

Dimroth and Reichardt [66] have proposed a solvent polarity parameter, $E_T(30)$, based on the transition energy for the longest-wavelength solvatochromic absorption band of the pyridinium N-phenolate betaine dye (44) (dye no. 30 in reference [66]). According to Eq. (7-27), the $E_T(30)$ value for a solvent is simply defined as the transition energy of the dissolved betaine dye (44) measured in kcal/mol [2, 66…68, 293] (for conversion into SI units, see the footnote on page 412). The major advantage of this approach is that the solvatochromic absorption band is at longer wavelengths for (44) than for Kosower's dye, generating an extraordinarily large range for the solvatochromic behaviour: from $\lambda = 810$ nm, $E_T(30) = 35.3$ kcal/mol, for diphenyl ether, to $\lambda = 453$ nm, $E_T(30) = 63.1$ kcal/mol, for water. Since the greater part of this solvatochromic range lies within the visible region of the spectrum, it is even possible to make a visual estimation of solvent polarity. For example, the solution colour of (44) is red in methanol, violet in ethanol, green in acetone, blue in isoamyl alcohol, and greenish-yellow in anisole [66]. A remarkable feature of these solution colour changes is that nearly every colour of the visible spectrum can be obtained by employing suitable binary mixtures of solvents of different polarity. A description of nice illustrative test-tube experiments and visual classroom demonstrations of solvent polarities by means of betaine dye (44) (which is commercially available) is given in a series of papers published in the Journal of Chemical Education [318…322]. To date, the betaine dye (44) holds the world record in solvatochromism with a direct experimentally observed hypsochromic shift of $\Delta\lambda = -357$ nm (ΔE_T ca. 28 kcal/mol = 117 kJ/mol) for a solvent change from diphenyl ether to water. Owing to this exceptionally large displacement of

the solvatochromic absorption band, the $E_T(30)$ values provide an excellent and very sensitive characterization of the polarity of solvents, high $E_T(30)$ values corresponding to high solvent polarity.

$E_T(30)$ values have been determined for more than 360 pure organic solvents [66, 174…176, 293] and for a great number of binary solvent mixtures [68…72, 124, 177…192, 293]. A collection of $E_T(30)$ values for pure organic solvents, representing the most comprehensive empirical solvent polarity scale so far known, is given in Table 7-3.* Some discrepancies between these $E_T(30)$ values given in Table 7-3 and those published in

concentrations of the more polar solvent ($c^* \gg c_p$), a good approximation of Eq. (7-30a) is $E_T(30) = E_D \cdot (c_p/c^*) + E_T^\circ(30)$, and the $E_T(30)$ values increase linearly with the molar concentration of the more polar solvent c_p. For high concentrations of the more polar solvent ($c^* \ll c_p$), Eq. (7-30a) can be written as $E_T(30) = E_D \cdot \ln(c_p/c^*) + E_T^\circ(30)$, and values of $E_T(30)$ correlate linearly with the logarithm of the molar concentration of the more polar solvent $\ln c_p$. Eq. (7-30a) is valid with high precision for about 80 investigated solvent mixtures and can even be applied to more complicated solvent systems than those mentioned above [191, 192].

In contrast to Eq. (7-30a), which describes the $E_T(30)$ behaviour in binary solvent mixtures in a rather empirical way, a more rational preferential solvation model has been developed by Connors et al. [327] as well as by Bosch and Rósset al. [328], based on the following simple two-step solvent exchange model:

$$I(S1)_2 + 2\ S2 \rightleftharpoons I(S2)_2 + 2\ S1$$

$$I(S1)_2 + S2 \rightleftharpoons I(S12)_2 + S1$$

where S1 and S2 indicate the two pure solvents being mixed, and S12 represents a solvent formed by interaction of solvents 1 and 2. $I(S1)_2$, $I(S2)_2$, and $I(S12)_2$ represent the solvatochromic indicator betaine dye (44) solvated by solvents S1, S2, and S12. The two solvent-exchange processes can be characterized by two preferential solvation parameters, $f_{2/1}$ and $f_{12/1}$. They measure the tendency of the indicator dye to be solvated by solvents S2 and S12 with reference to solvation by solvent 1. The preferential solvation parameters are defined as $f_{2/1} = (x_2^S/x_1^S)/(x_2^0/x_1^0)^2$ and $f_{12/1} = (x_{12}^S/x_1^S)/(x_2^0/x_1^0)$, where x_1^S, x_2^S, and x_{12}^S are the mole fractions of solvents S1, S2, and S12, respectively, in the solvation shell of the indicator dye, and x_1^0 and x_2^0 are the mole fractions of the two solvents S1 and S2 in the bulk binary solvent mixture. The normalized E_T^N polarity of the binary solvent mixture is then calculated as an average of the E_T^N values of solvents S1, S2, and S12 as follows:

$$E_T^N = x_1^S \cdot E_{T1}^N + x_2^S \cdot E_{T2}^N + x_{12}^S \cdot E_{T12}^N$$

From these equations, a general equation that relates the E_T^N value of a binary solvent mixture to the E_T^N values of the two pure solvents, the preferential solvation parameters, and the solvent composition can be derived, viz.:

$$E_T^N = \frac{E_{T1}^N(1 - x_2^0)^2 + E_{T2}^N f_{2/1}(x_2^0)^2 + E_{T12}^N f_{12/1}(1 - x_2^0)x_2^0}{(1 - x_2^0)^2 + f_{2/1}(x_2^0)^2 + f_{12/1}(1 - x_2^0)x_2^0} \qquad (7\text{-}30b)$$

The general equation (7-30b) has been successfully applied by Bosch, Rósset al. to a great variety of binary solvent mixtures (> 70), even to solvent mixtures showing synergistic behaviour due to HBD solvent/HBA solvent interactions [328].

It should be mentioned that an equation analogous to Eq. (7-30a) has been successfully applied to salt effects on reaction rates arising from variations in solvent polarity on the addition of electrolytes (ionophores) [197] (cf. also Eq. (5-99) in Section 5.4.5. For electrolyte solutions, the added salt can be treated as a more polar ••co-solvent•• [197].

The strong dependence of $E_T(30)$ values on the composition of binary mixtures of

solvents with different polarity can be used for the quantitative UV/Vis spectroscopic determination of water in organic solvents [68, 192, 198, 199, 321, 322].

In general, the $E_T(30)$ values exhibit a good, mostly linear correlation with a large number of other solvent-sensitive processes such as light absorption, reaction rates, and chemical equilibria [2]. Applications of $E_T(30)$ values to chemical reactivity [124, 200…202] and analytical chemistry [203] have been reviewed. Their application to photochemical processes has been discussed [204, 205]. Its extreme sensitivity to small changes in the surrounding medium has made the betaine dye (44) a useful molecular probe in the study of micellar systems, microemulsions, phospholipid bilayers, model liquid membranes, polymers, organic…inorganic polymer hybrids, sol…gel matrices, surface polarities, and the retention behaviour in reversed-phase liquid chromatography; see page 333 and references [298…302, 443…451] of Chapter 6. Even standard molar Gibbs energies of transfer of chloride ions, ΔG_t° ($Cl^\ominus$, $H_2O \rightarrow S$), could be linearly correlated to the $E_T(30)$ values [206]. $E_T(30)$ values have also been used to calculate acceptor numbers, AN_E, which are not otherwise available by direct measurements [207]; Table 2-5 in Section 2.2.6.

It should be mentioned that the pyridinium *N*-phenolate betaine dye (44) is not only very sensitive to changes in solvent polarity, but in addition its longest-wavelength solvatochromic absorption band also depends on changes in temperature [73, 175, 180, 208] and pressure [74, 182, 208], on the addition of electrolytes (ionophores) [209…213], as well as on the introduction of substituents in the peripheral phenyl groups; Fig. 7-2 in Section 7.1 and reference [332] for a review.

The thermo-solvatochromism of (44) can be easily observed by means of a betaine solution in ethanol: at $-75\,^\circ C$ the solution is red-colored, while at $+75\,^\circ C$ it is blue-violet, corresponding to absorption maxima of $\lambda_{max} = 513$ and 568 nm, respectively [73]. The reason for this thermo-solvatochromism is the increased stabilization of the dipolar electronic ground state of (44) relative to the less dipolar excited state with decreasing temperature, due to better solute/solvent interactions at low temperature. It can be stated that, the lower the temperature, the higher the $E_T(30)$ value.

Tamura and Imoto [74] and Hendrickson, Drickamer et al. [182] observed pressure effects on the solution spectra of (44). In all solvents used they found a hypsochromic shift of the longest-wavelength absorption band with increasing pressure. The observed hypsochromic shift of a betaine solution in methanol amounts to $\Delta\lambda = -15$ nm ($\Delta\tilde{\nu} = +600\ cm^{-1}$; $\Delta E_T = +1.7$ kcal/mol) on raising the pressure from 1 to 10 kbar [182]. On the supposition that this piezo-solvatochromism results from better solute/solvent interactions with increasing pressure, it can be stated that, the higher the pressure, the more polar the solvent, and the higher the $E_T(30)$ value.

The addition of electrolytes (ionophores) to solutions of (44) causes hypsochromic shifts of its solvatochromic absorption band [197, 209…213]. This phenomenon can be designated as halo-solvatochromism*. For example, the addition of KI, NaI,

---

* The halo-solvatochromism of (44) can be considered as the only genuine halochromism, i.e. a colour change with increasing ionic strength of the medium without chemically changing the chromophore. The term halochromism, as introduced by Baeyer et al. [214], denotes the trivial colour change of a dye on addition of acids or bases. This is simply caused by the creation of a new chromophore in an acid/base reaction whereby a colourless compound is rendered coloured on salt formation, e.g.

$$(C_6H_5)_3C{-}Cl\ (\text{colourless}) + AlCl_3 \rightleftharpoons (C_6H_5)_3C^\oplus\ (\text{yellow}) + AlCl_4^\ominus.$$

LiI, BaI_2, $Ca(SCN)_2$, and $Mg(ClO_4)_2$ to solutions of (44) in acetonitrile leads to a differential hypsochromic band shift which increases with this ionophore order, i.e. with increasing charge density of the cation [211]. Obviously, salts act similarly to other polar compounds (solvents) when added to solutions of (44). The polarity of binary ionophore/solvent mixtures as a function of composition can be quantitatively described in a manner similar to other binary solvent/solvent mixtures [197, 213].

It should be noted that the polarity of the medium also influences the ^1H and ^{13}C NMR chemical shifts of dye (44) [215, 216]; cf. Section 6.5.1. The sites in the betaine molecule most influenced by the solvent are those nearest to the positive and negative charges within the dye molecule, and this is reflected in the NMR chemical shifts.

The solvent, temperature, pressure, ionophore, and substituent effects on the UV/Vis spectra of the pyridinium-N-phenolate betaine dyes indicate the extreme sensitivity of this class of compounds to small changes in the environment. Their behaviour may be compared to that of the Princess and the Pea in one of H. C. Andersen's fairy-tales [76]*). Their utility for setting up linear Gibbs energy relationships is demonstrated by the fact that the same betaine dye can be used for establishing kinetic and spectroscopic scales of substituent constants (cf. Figs. 7-1 and 7-2) as well as a spectroscopic scale of solvent polarity (cf. Table 7-3).

Further solvent polarity scales based on UV/Vis absorption as well as fluorescence spectra have been proposed by Brooker et al. [77], Dähne et al. [78], de Mayo et al. [217], Dubois et al. [79], Mukerjee et al. [218] and Wrona et al. [219], Walter et al. [220], Walther [81] and Lees et al. [82], Zelinskii et al. [80], Winnik et al. [222], Kamlet and Taft [84, 84a, 224, 226]. Buncel et al. [333], and Catalán et al. [296, 334…337]. In addition to these scales, a great variety of further positively and negatively solvatochromic dyes have been recommended as solvent polarity indicators. A review describes about 60 organic and inorganic compounds, the solvatochromism of which is sufficiently large for their potential application as empirical solvent polarity probes [293].

The solvent dependence of the $\pi \rightarrow \pi^*$ transition energies of two merocyanine dyes was used by Brooker et al. [77] to establish the solvent polarity parameters w_R and w_B (cf. Table 7-2). w_R is based on the positively solvatochromic merocyanine dye no. 1 in Table 6-1 of Section 6.2.1 (red shift with increasing solvent polarity), while w_B represents the transition energies of the negatively solvatochromic merocyanine dye no. 13 in Table 6-1.

Dähne et al. [78] proposed the positively solvatochromic 5-(dimethylamino)pentadien-2,4-al (dye no. 3 in Table 6-1) as a solvent polarity indicator and recommended a relative polarity function RPM (from the German "Relatives Polaritätsmaß").

The $\pi \rightarrow \pi^*$ transition energy E_{sp} of a spiropyran zwitterion of the type described in Section 4.4.2 [(27a) $\rightleftharpoons$ (27b)] has been used by de Mayo et al. to characterize the polarity of solid oxide surfaces such as that of silica gel [217].

Dubois et al. [79] formulated F values (formerly F values) as solvent polarity parameters based on the position of the solvent-sensitive $\pi \rightarrow \pi^*$ transition of certain saturated aliphatic ketones, as shown in Eq. (7-31)

---

* The princess was so sensitive to her surroundings that she was able to feel a pea through 20 mattresses and 20 eider-down quilts in her bed. This extreme sensitivity corresponds in some way to the sensitivity of the pyridinium-N-phenolate betaine dye (44).

$$\Delta\tilde{\nu}_H^S = \tilde{\nu}^S - \tilde{\nu}^H = F(\tilde{\nu}^H - 32637) - 174 \qquad (7\text{-}31)$$

Solvents are characterized by deviations from unity of the slope of plots of $\tilde{\nu}^S$ values (wavenumber of absorption for various ketones in solvent S) against $\tilde{\nu}^H$ (for various ketones in n-hexane as reference solvent); cf. Table 7-2.

Mukerjee et al. [218] and Wrona et al. [219] have used the highly solvatochromic $n \rightarrow p^*$ transition energy of the stable 2,2,6,6-tetramethylpiperidine-1-oxide radical (TMPNO) for the development of a solvent polarity scale. So-called E_B^N values, as empirical measures of solvent Lewis acidity, have been determined for 53 pure organic solvents and some binary solvent/water mixtures [219].

Using the negative solvatochromism of the $n \rightarrow p^*$ absorption of N,N-(dimethyl)-thiobenzamide S-oxide, an E_T^{SO} solvent scale has been proposed by Walter et al. [220]. This scale comprises of 36 solvents and three binary solvent/water mixtures and is thought to be particularly useful for characterizing protic solvents.

Furthermore, the E_K scale of Walther [81] and the E^*_{MLCT} scale of Lees et al. [82] should be mentioned. Both rather comprehensive solvent scales are based on the negatively solvatochromic metal-to-ligand charge-transfer absorption (MLCT; $d \rightarrow p^*$) of the two zero-valent group VI metal complexes (46) and (47) of the common formula $M(CO)_4$(diimine). A consistent explanation for the solvatochromic behaviour of such coordination compounds with MLCT absorption has been given by Kaim et al. [83]; cf. also Section 6.2.1.



(46)



(47)

For a review of the use of solvatochromic metal complexes as visual indicators of solvent polarity, see reference [221].

Solvatochromic fluorescent probe molecules have also been used to establish solvent polarity scales. The solvent-dependent fluorescence maximum of 4-amino-N-methylphthalimide was used by Zelinskii et al. to establish a "universal scale for the effect of solvents on the electronic spectra of organic compounds" [80, 213]. More recently, a comprehensive Py scale of solvent polarity including 95 solvents has been proposed by Winnik et al. [222]. This is based on the relative band intensities of the vibronic bands I and III of the $p^* \rightarrow p$ emission spectrum of monomeric pyrene; cf. Section 6.2.4. A significant enhancement is observed in the 0–0 vibronic band intensity I_I relative to the $0 \rightarrow 2$ vibronic band intensity I_{III} with increasing solvent polarity. The ratio of emission intensities for bands I and III serves as an empirical measure of solvent polarity: $Py = I_I/I_{III}$ [222]. However, there seems to be some difficulty in determining precise Py values, as shown by the varying Py values from different laboratories; the reasons for these deviations have been investigated [223].

An interesting approach, called the *solvatochromic comparison method*, was used to evaluate a b scale of solvent hydrogen-bond acceptor (HBA) basicities, an a scale of solvent hydrogen-bond donor (HBD) acidities, and a p^* scale of solvent dipolarity/polarizability using UV/Vis spectral data of solvatochromic compounds, was employed and further developed by Kamlet, Taft et al. [84, 84a, 224, 226]. Magnitudes of enhanced solvatochromic shifts, $\Delta\Delta\tilde{\nu}$, in HBA solvents*) have been determined for 4-nitroaniline relative to homomorphic**) N,N-diethyl-4-nitroaniline (compound no. 2, Table 6-1, Section 6.2.1). Both standard compounds are capable of acting as HBA substrates (at the nitro oxygens) in HBD solvents, but only 4-nitroaniline can act as a HBD substrate in HBA solvents. Taking the $\Delta\Delta\tilde{\nu}$ value of 2800 cm^{-1} for hexamethylphosphoric triamide (a strong HBA solvent) as a single "xed reference point ($b_i = 1.00$), a b scale of solvent HBA basicities for HBA solvents was developed [84]. Using the same solvatochromic comparison method, i.e. the enhanced solvatochromic shifts, $\Delta\Delta\tilde{\nu}$, in HBD solvents for 4-nitroanisole and the pyridinium-*N*-phenolate betaine dye (44), an a scale of HBD acidities was evaluated. Taking the $\Delta\Delta\tilde{\nu}$ value of 6240 cm^{-1} for methanol (a strong HBD solvent) as a "xed reference point ($a_i = 1.00$), an a scale of solvent HBD acidities for HBD solvents was established [84].

The same authors also introduced a p^* scale of solvent dipolarity/polarizability [84a]. This p^* scale is so named because it is derived from solvent e¤ects on the p^* electronic transitions of a selection of seven positively solvatochromic nitroaromatics of the type D–C_6H_4–A, where D and A stand for electron-donor (e.g. NMe_2) and electron-acceptor (e.g. NO_2) groups, respectively: 4-nitroanisole, *N*,*N*-diethyl-3-nitroaniline, 4-methoxy-β-nitrostyrene, 1-ethyl-4-nitrobenzene, *N*-methyl-2-nitro-*p*-toluidine, *N*,*N*-diethyl-4-nitroaniline, and 4-(dimethylamino)benzophenone. Given a solvatochromic indicator compound, the p^* value for a solvent S was de"ned according to Eq. (7-32):

$$p^*(S) = [\tilde{\nu}(S) - \tilde{\nu}(c\text{-}C_6H_{12})]/[\tilde{\nu}(DMSO) - \tilde{\nu}(c\text{-}C_6H_{12})] \qquad (7\text{-}32)$$

where $\tilde{\nu}(S)$ corresponds to the wavenumber of the maximum of the long-wavelength solvatochromic absorption band of the indicator measured in the various solvents. Cyclohexane and dimethyl sulfoxide were used as reference solvents by taking $p^*(c\text{-}C_6H_{12}) = 0.00$ and $p^*(DMSO) = 1.00$ by de"nition. Solvent e¤ects on the $\tilde{\nu}_{max}$ values of these seven primary solvatochromic indicators were employed in the initial construction of the p^* scale, which was then expanded and re"ned by multiple least-squares correlations with additional solvatochromic indicators. In this way, an averaged p^* scale of solvent dipolarity/polarizability was established, which measures the ability of the solvent to stabilize a charge or a dipole by virtue of its dielectric e¤ect [84a]. A normalized range of 0.00 (for cyclohexane) to 1.00 (for dimethyl sulfoxide) for the p^* values of common solvents has been chosen so that, taken together with the a scale of

---

* The term hydrogen-bond acceptor (HBA) refers to the acceptance of the proton of a hydrogen-bond. Therefore, HBA solvents are also electron-pair donor (EPD) solvents. Hydrogen-bond donor (HBD) refers to the donation of the proton. Therefore, HBD solvents behave as protic solvents.

** The term *homomorphic* molecules was introduced by Brown et al. [225]. Molecules having the same or closely similar molecular geometry are *homomorphs*, e.g. ethane is a homomorph of methanol, toluene of phenol, 4-aminobenzenesulfonamide of 4-aminobenzoic acid, and *N*,*N*-diethyl-4-nitroaniline of 4-nitroaniline.

solvent HBD acidities and the b scale of solvent HBA basicities (which have also been scaled to range from 0.00 to 1.00), these parameters can be used in a multiparameter equation (cf. Section 7.7).

In 1994, a review on the further development and improvement of the π^* scale was given by Laurence, Abboud et al. [227]. They redetermined π^* values for a total of 229 solvents, this time using only two (instead of seven) solvatochromic nitroaromatics as indicator compounds, i.e. 4-nitroanisole and N,N-dimethylamino-4-nitroaniline, for good reasons; see later and reference [227] for a more detailed discussion. A thermodynamic analysis of the π^* scale [and the $E_T(30)$ scale] has been reported by Matyushov et al. [228]. Using six novel diaza merocyanine dyes of the type R–N=N–R′ (R = N-methylpyridinium-4-yl or N-methylbenzothiazolium-2-yl, and R′ = 2,6-disubstituted 4-phenolates or 2-naphtholate) instead of nitroaromatics as positively solvatochromic probe compounds, an analogous π^*_{azo} scale was developed by Buncel et al., which correlates reasonable well with the π^* scale, but has some advantages; for a detailed discussion, see references [333]. Another π^* scale, based solely on naphthalene, anthracene, and b-carotene, was constructed by Abe [338]. π^* values are mixed solvent parameters, measuring the solvent dipolarity and polarizability. The differences in the various π^* scales are caused by the different mixture of dipolarity and polarizability measured by the respective indicator. The π^* scale of Abe is practically independent of the solvent dipolarity, whereas Kamlet…Taft's π^* and Buncel's π^*_{azo} reflect different contributions of both solvent dipolarity and polarizability.

A selection of Kamlet and Taft's solvatochromic parameters a, b, and π^* for 40 organic solvents, taken from a more recent comprehensive and improved collection [227, 294], is given in Table 7-4.

The parameters in Table 7-4 were arrived at by averaging the multiple normalized solvent effects on a variety of solvent-dependent properties involving various types of solvatochromic indicator dyes. Therefore, the solvatochromic parameters of Table 7-4 are no longer directly based on the solvent effects indicated by a distinct single solvatochromic indicator dye. Rather, they are statistically averaged values resulting from a series of successive approximations [226]. Kamlet and Taft's solvatochromic parameters have been used in different combinations in one-, two-, and three-parameter correlations known as linear solvation energy relationships (LSERs). An impressive series of more than 40 articles entitled "Linear solvation energy relationships" has been published: Part 1 [229]…Part 41 [230]; see Section 7.7 for a further discussion of such multiparameter correlations.

---

* There is a discussion in the literature concerning the use of averaged and statistically optimized "constant" solvent polarity parameters (e.g. a, b, π^*) instead of experimentally derived parameters which are based on a distinct, single, and well-understood solvent-dependent reference process [e.g. Y_{OTs}, Z, $E_T(30)$]. There are some practical reasons in favour of the experimentally derived solvent parameters. They are related to a distinct chemical or physical reference process and the corresponding solvent scale can easily be enlarged by new precise measurements. Averaged and statistically optimized solvent parameters are no longer directly related to a distinct reference process and are thus ill-defined. New data may be difficult to incorporate into the existing framework, and even if they are, then this can lead to a modification of the already calculated "constant" solvent parameters. It has been pointed out "that it is better to study one good model with precision than to take the average of results obtained from many poor models" [237]; also [153].

Table 7-4. A selection of Kamlet and Taft•s solvatochromic parameters a, b, and p^* for 40 organic solvents, taken mainly from references [227, 294] as well as from references [226, 295].

Solvents	a	b	p^*
Gas phase	0.00	0.00	−1.23
n-Hexane	0.00	0.00	−0.11
Cyclohexane	0.00[a)]	0.00[a)]	0.00[a)]
Carbon disul“de	0.00	0.07	0.51
Dichloromethane	0.13	0.10	0.73
Trichloromethane, Chloroform	0.20	0.10	0.69
Tetrachloromethane	0.00	0.10	0.21
Benzene	0.00	0.10	0.55
Methylbenzene, Toluene	0.00	0.11	0.49
Chlorobenzene	0.00	0.07	0.68
Nitrobenzene	0.00	0.30	0.86
Pyridine, Azine	0.00	0.64	0.87
Diethyl ether	0.00	0.47	0.24
Tetrahydrofuran	0.00	0.55	0.55
1,4-Dioxane	0.00	0.37	0.49
Triethylamine	0.00	0.71	0.09
Propanone, Acetone	0.08	0.48	0.62
Cyclohexanone	0.00	0.53	0.68
Ethyl acetate	0.00	0.45	0.45
Ethyl benzoate	0.00	0.41	0.68
(±)-Propylene carbonate	0.00	0.40	0.83
Dimethyl sulfoxide	0.00	0.76	1.00[a)]
Tetrahydrothiophene-1,1-dioxide, Sulfolane	0.00	0.39	0.90
Acetonitrile	0.19	0.40	0.66
Nitromethane	0.22	0.06	0.75
Hexamethylphosphoric acid triamide, HMPT	0.00	1.00[a)]	0.87
N,N-Dimethylformamide, DMF	0.00	0.69	0.88
N-Methylformamide	0.62	0.80	0.90
Formamide	0.71	0.48	0.97
t-Butanol	0.42	0.93	0.41
Cyclohexanol	0.66	0.84	0.45
1-Butanol	0.84	0.84	0.47
2-Propanol	0.76	0.84	0.48
1-Propanol	0.84	0.90	0.52
Ethanol	0.86	0.75	0.54
Ethane-1,2-diol, Glycol	0.90	0.52	0.92
Methanol	0.98	0.66	0.60
Acetic acid	1.12	0.45	0.64
2,2,2-Tri”uoroethanol	1.51	0.00	0.73
1,1,1,3,3,3-Hexa”uoro-2-propanol	1.96	0.00	0.65
Water	1.17	0.47	1.09

[a)] By de“nition.

An extension of the b scale of solvent HBA basicity, using only 4-nitroaniline and N,N-diethyl-4-nitroaniline as solvatochromic indicators (β_{KT} scale), has been described by Krygowski et al. [231, 232]. Some further improvements of Kamlet and Taft•s solvatochromic parameters have been proposed by Kolling [233] and Bekárek [234]. According to Bekárek, better correlations are obtained using modi“ed b_n, a_n, and p_n^*

values, which are derived from the original b, a, and p^* values by dividing them by the refractive index function $(n^2 - 1)/(2n^2 + 1)$. This is in order to eliminate the polarization contribution of the solvent molecules in the cybotactic solvation sphere during the electronic excitation of the solvatochromic indicators. According to Bekárek, the original b, a, and p^* values are adequate for correlating certain types of spectral properties (e.g., HFS constants in ESR spectra), but their applicability to other types of solvent effects could be improved by using the modified b_n, a_n, and p_n^* values [234]. However, this modification has been strongly criticized by Kamlet, Taft, and Abboud for conceptual and computational reasons. This leads to the conclusion that in all correlations the original solvatochromic parameters are in fact best [235].

Interestingly, a statistical principal component analysis of the solvatochromic shift data sets previously used by Kamlet and Taft in defining the p^* scale has shown that, rather than one (p^*), two solvent parameters (y_{1k} and y_{2k}) are necessary to describe the solvent-induced band shifts of the studied solvatochromic indicators [236]. This is not unexpected since the p^* parameters are assumed to consist of a blend of dipolarity and polarizability contributions to the solute/solvent interactions.

Laurence et al. [237…239] have tried to improve Kamlet and Taft's solvatochromic comparison method [224] by introducing a new thermosolvatochromic comparison method. In doing this, they tried to eliminate various shortcomings including sometimes insufficiently precisely determined solvatochromic parameters. According to the solvatochromic comparison method, the p^* value of a solvent S is measured by the bathochromic shift relative to cyclohexane, $-\Delta\tilde{\nu}_S$, of the $p \rightarrow p^*$ transition of a non-HBD indicator dye (e.g. 4-nitroanisole); cf. Eq. (7-32). The basicity parameter b of the same solvent is measured by the supplementary bathochromic shift $\Delta\Delta\tilde{\nu}_S$ obtained using a second HBD indicator dye (e.g. 4-nitrophenol), which is a homomorph of the first dye. Plotting the absorption wavenumbers of the HBD indicator dye i against the wavenumbers of its non-HBD homomorph j, for non-HBD and non-HBA solvents, leads to the linear reference line given in Fig. 7-3.

Because of the specific HBD solute/HBA solvent interaction, all HBA solvents fall below this line, the higher the basicity the lower their position. The Kamlet…Taft values are the means of the normalized values $\Delta\Delta\tilde{\nu}_S(i-j)$ for several indicator dyes j



Fig. 7-3. Illustration of the solvatochromic comparison method according to [237]. Reference line equation: $\tilde{\nu}(i) = a_{ij} \cdot \tilde{\nu}(j) + b_{ij}$. Reference solvent: cyclohexane. ○ non-HBA and non-HBD solvents; ● an HBA solvent.

and various homomorph pairs i/j, according to Eq. (7-33) [237].

$$-\Delta\Delta\tilde{\nu}_S(i-j) = [a_{ij} \cdot \tilde{\nu}_S(j) + b_{ij}] - \tilde{\nu}_S(i) \tag{7-33}$$

It is obvious that the accuracy of the determination of $-\Delta\Delta\tilde{\nu}_S(i-j)$ depends strongly on the precision with which the reference line of Fig. 7-3 is established.

Laurence et al. [237, 238] have shown that some of the absorption bands of the solvatochromic indicator dyes previously used by Kamlet and Taft to "x the reference lines exhibit a solvent- and temperature-dependent vibrational "ne structure, which makes the determination of the band maxima $\tilde{\nu}_{max}$, di‹cult. Furthermore, the indicator dyes used are thermo-solvatochromic, i.e. $\tilde{\nu}_{max}$ is temperature-dependent. No allowance for this temperature-dependence of the a, b, and p^* values has been made previously.

Finally, the right choice of solvent on going from the gas phase to the most polar non-HBA and non-HBD solvents has to be made in order to "x the reference line as given in Fig. 7-3. Taking all these di‹culties into account, and using 4-nitrophenol/4-nitro-anisole and 4-nitroaniline/N,N-dimethyl-4-nitroaniline as distinct solvatochromic indicator couples i and j, as well as a carefully selected set of non-HBA and non-HBD solvents, Laurence et al. [237, 238] have determined more precise reference lines. From these, it was possible to derive new solvent dipolarity/polarizability and solvent HBA basicity parameters, which they believe to be more correct than the original a and b parameters; see also reference [227]. In particular, the b scale seems to be mainly a scale of solvent HBA basicity against NH donor solutes and does not have the degree of general validity as originally claimed [84].

Other spectroscopic measurements of solvent polarity use as standard processes solvent-sensitive infrared stretching absorptions of groups such as OH and X—H · · · B, where X may be C, S, N, O, or P, and B is a hydrogen-bond acceptor (HBA) [cf. Eqs. (6-6) and (6-7) in Section 6.3]. Schleyer et al. [85] proposed the relationship (7-34) for the correlation of solvent-sensitive IR vibrations.

$$\frac{\tilde{\nu}^0 - \tilde{\nu}^s}{\tilde{\nu}^0} = a \cdot G \tag{7-34}$$

If X—H · · · B vibrations are being examined, $\tilde{\nu}^0$ refers to its gas-phase value. The value for a is a measure of solvent susceptibility of a particular IR vibration, and G* is a function of the solvent only. Since solvent shifts of $\tilde{\nu}_{C=O}$ and $\tilde{\nu}_{S=O}$ are proportional to solvent shifts of $\tilde{\nu}_{X-H\cdots B}$, G values are calculated from the solvent shifts of the carbonyl bands of N,N-dimethylformamide and benzophenone and the sulfonyl band of dimethyl sulfoxide. An arbitrary value of 100 was assigned to dichloromethane to "x the scale ($G = 0$ for the gas phase) [85]. Values of G are given in Table 7-2. Further G values have been determined by Somolinos et al. [240] and their relationships to other solvent polarity parameters have been investigated [241].

Kagiya et al. [86] have measured the IR wavenumbers of the O—D and C=O stretching vibrations of monodeuteriomethanol (CH_3OD) and acetophenone, respec-

* The abbreviation G was chosen from the name of a close aquaintance of one of the authors [85]; private communication to C. R.

tively, in various solvents. Taking benzene as the reference solvent, they used the wavenumber shift relative to the maximum found in benzene as a measure of the electron-pair donating power and the electron-pair accepting power of a given solvent. Measurements of the O–D stretching band of CH_3OD, $\tilde{\nu}_{MeOD}$, have been greatly extended (up to 55 aprotic solvents) and improved by Shorter *et al.* [87]. Based on the measurements of Kagiya *et al.* [86], Koppel and Pal•m [6] have de“ned a Lewis basicity parameter B_{MeOD} using Eq. (7-35),

$$B_{MeOD}/cm^{-1} = \Delta\tilde{\nu}_{OD} = \tilde{\nu}^{0}_{MeOD} - \tilde{\nu}_{MeOD\cdots B} \tag{7-35}$$

where $\tilde{\nu}^{0}_{MeOD}$ and $\tilde{\nu}_{MeOD\cdots B}$ refer to the O—D stretching vibration of CH_3OD measured in the gas phase ($\tilde{\nu}^{0}_{MeOD} = 2720\ cm^{-1}$ [87]) and in a given HBA solvent B. A comprehensive collection of B_{MeOD} values can be found in reference [87].

In a similar way, another set of solvent Lewis-basicity parameters, B_{PhOH}, based on band shifts of the O—H stretching vibration of phenol in tetrachloromethane induced by hydrogen-bond formation with added HBA solvents B, was used by Koppel and Paju [88] to classify 198 solvents, according to Eq. (7-36) ($B_{PhOH} = 0$ for CCl_4):

$$B_{PhOH}/cm^{-1} = \Delta\tilde{\nu}^{CCl_4}_{OH} = \tilde{\nu}^{CCl_4}_{PhOH} - \tilde{\nu}^{CCl_4}_{PhOH\cdots B} \tag{7-36}$$

This B_{PhOH} scale was further extended by Makitra *et al.* [242]. It has been shown that the spectroscopically determined B_{PhOH} values correlate well with Gutmann•s calorimetrically measured donor numbers, DN [243]; cf. Eq. (7-10).

Another remarkable IR spectroscopic parameter of solvent Lewis basicity has been introduced by Laurence *et al.* [239], using the so-called *infrared comparison method* analogous to Kamlet and Taft•s solvatochromic comparison method [224]. The band maxima wavenumbers of the C=O stretching vibration of the two homomorphs CCl_3CO_2H and $CCl_3CO_2CH_3$ have been measured in the gas phase as well as in non-HBD and non-HBA solvents in order to establish a reference line that conforms to Eq. (7-37); cf. Fig. 7-3.

$$\tilde{\nu}^{CCl_3CO_2H}_{C=O} = 1.171 \cdot \tilde{\nu}^{CCl_3CO_2CH_3}_{C=O} - 293 \tag{7-37}$$

This linear relationship demonstrates the similarity of the polarity e¤ects on both homomorphic vibrators, trichloroacetic acid and its methyl ester. For HBA solvents B, however, the $\tilde{\nu}_{C=O}$ data points are displaced below the reference line of Eq. (7-37). These deviations are caused by the formation of solute/solvent hydrogen bonds $CCl_3CO_2H\cdots B$, resulting in a decrease in the C=O vibration wavenumber. The hydrogen-bond induced wavenumber shift $\Delta\tilde{\nu}_{C=O}$ for a HBA solvent is then calculated by Eq. (7-38); cf. Eq. (7-33).

$$\begin{aligned}\Delta\tilde{\nu}_{C=O} &= \tilde{\nu}^{CCl_3CO_2H}_{C=O}\ [\text{calc. from Eq. (7-37)}] - \tilde{\nu}^{CCl_3CO_2H}_{C=O}\ [\text{observed}] \\ &= [1.171 \cdot \tilde{\nu}^{CCl_3CO_2CH_3}_{C=O} - 293] - \tilde{\nu}^{CCl_3CO_2H}_{C=O}\end{aligned} \tag{7-38}$$

Due to this subtraction, the nonspecific polarity effect of the solvent is disentangled from its basicity effect on the C=O vibrator of CCl_3CO_2H in HBA solvents. $\Delta\tilde{\nu}_{C=O}$ values range from 1.4 cm^{-1} for benzene to 29.3 cm^{-1} for tri-n-butyl phosphate [239]. The $\Delta\tilde{\nu}_{C=O}$ displacements correlate well with the $\Delta\tilde{\nu}_{OD}$ values of Eq. (7-35). However, this $\Delta\tilde{\nu}_{C=O}/\Delta\tilde{\nu}_{OD}$ relationship is family-dependent, i.e. different linear correlations are obtained for different classes (families) of organic HBA solvents such as S, O, and N bases [239]. Unfortunately, $\Delta\tilde{\nu}_{C=O}$ values are not available for carbonyl-containing solvents because of insufficient solvent transparency, and for more basic solvents such as water, DMSO, and HMPT because of decarboxylation and proton-transfer reactions. An excellent review on solvatochromic scales derived from various similarity models employed in IR and UV/Vis spectroscopy has been given by Laurence [339]. The similarity principle simply states that similar changes in molecular structure or medium should produce similar changes in the solvatochromic properties of the solutes under study.

Electron spin resonance (ESR) and nuclear magnetic resonance (NMR) measurements have also been used to establish solvent polarity scales. Knauer et al. [89] determined the nitrogen hyperfine splitting constants, $a(^{14}N)$, of several aminyloxide radicals (nitroxides), such as di-t-butyl aminyloxide, in 31 solvents. It has been known for some time that the ^{14}N isotropic HFS constant in the ESR spectrum of an aminyloxide free radicals is sensitive to the polarity of the solvent in which it is dissolved (cf. Table 6-5 and the discussion in Section 6.4). Since $a(^{14}N)$ is easily measured in almost all solvents, it may prove to be useful as an empirical solvent polarity parameter, especially in cases where the other parameters are difficult to determine owing to limited solubility. Only very low radical concentrations are usually needed to obtain an ESR spectrum. The $a(^{14}N)$ values of di-t-butyl aminyloxide are included in Table 7-2. Further $a(^{14}N)$ values of di-t-butyl aminyloxide have been determined by Kolling [244], Reddoch et al. [245], and Symonset al. [246], who also discussed their solvent-dependence, their applicability as solvent parameters, and their correlations with other solvent parameters in more detail. The $a(^{14}N)$ values of 2,2,6,6-tetramethylpiperidine-1-oxide (TMPNO) [247, 248] and 4-acetyl-1-methylpyridinyl free radicals [244] have also been discussed as potential solvent polarity probes. A critical comparison of the TMPNO $a(^{14}N)$ values with other physical and empirical solvent parameters has been given by Kecki et al. [248]. For example, the $a(^{14}N)$ values of aminyl oxides correlate well with the $E_T(30)$ and AN values, but not with the solvent relative permittivities or dipole moments or functions thereof [245, 248].

Interesting solvent scales based on NMR measurements have been proposed by Taft et al. [90] and by Gutmann, Mayer et al. [91]. A solvent polarity parameter, designated as P, has been defined by Taft et al. [90] as the ^{19}F chemical shift (in ppm) of 4-fluoro-nitrosobenzene in a given solvent, relative to the same quantity in the reference solvent cyclohexane (cf. Table 6-6 and the discussion in Section 6.5.1). These parameters define a scale ranging from $P = 0.0$ in cyclohexane to $P = 2.7$ in sulfolane, and can easily be measured in a wide variety of solvents. The P values appear to be related to the ability of the solvents to form specific 1:1 complexes with the nitroso group of the standard compound. A compilation of P values can be found in reference [92]. In addition, ^{13}C chemical shifts of (trifluoromethyl)benzene and phenylsulfur pentafluoride have been used by Taft et al. to study nonspecific dipolar interactions with HBD solvents and utilized to define p^* values of solvent dipolarity/polarizability for protic solvents [249].

Complementary to the donor numbers DN [cf. Eq. (7-10)], Gutmann, Mayer et al. introduced so-called acceptor numbers AN as measures of the Lewis acidity or EPA property of organic solvents [91, 134, 207, 251]. Acceptor numbers are derived from the relative ^{31}P NMR chemical shift values, δ_{corr}, of triethylphosphane oxide, related to those of the 1:1 adduct $Et_3PO—SbCl_5$. δ_{corr} is the observed ^{31}P chemical shift corrected for concentration effects and for differences in magnetic volume susceptibility.

$$(Et_3P{=}O \longleftrightarrow Et_3\overset{\oplus}{P}{-}\overset{\ominus}{O}) + A \rightleftharpoons Et_3\overset{\delta\oplus}{P}{=}\overset{\delta\ominus}{O}{-}A$$

With n-hexane as the reference solvent, values of AN for a given solvent A are calculated according to Eq. (7-39).

$$AN = \frac{\delta_{corr}(A) - \delta_{corr}(n\text{-}C_6H_{14})}{\delta_{corr}(Et_3PO—SbCl_5) - \delta_{corr}(n\text{-}C_6H_{14})} \cdot 100 \qquad (7\text{-}39)$$

The AN scale was set up by defining $AN = 0$ for n-hexane, and $AN = 100$ for $Et_3PO—SbCl_5$ dissolved in 1,2-dichloroethane. The term $\delta_{corr}(Et_3PO—SbCl_5) - \delta_{corr}(n\text{-}C_6H_{14})$ was experimentally found to be 42.58 ppm; Eq. (7-39) reduces therefore to Eq. (7-40).

$$AN = \frac{\delta_{corr}(A) - \delta_{corr}(n\text{-}C_6H_{14})}{42.58\ ppm} \cdot 100 = \Delta\delta_{corr} \cdot 2.348/ppm \qquad (7\text{-}40)$$

A selection of AN values has already been given in Table 2-5 of Section 2.2.6; also Table 6-6 in Section 6.5.1. The observed solvent-dependent ^{31}P chemical shifts result mainly from the polarization of the dipolar P—O group, induced by the interaction with electrophilic solvents A, particularly HBD solvents. The decrease in electron density at the phosphorus atom results in a deshielding proportional to the strength of the probe/solvent interaction. In solutions of protic acids, the ^{31}P chemical shift of the O-protonated triethyl hydroxyphosphonium salt is observed. Since Et_3PO is very hygroscopic and therefore not very suitable from an experimental point of view, the use of $(n\text{-}Bu)_3PO$ instead of Et_3PO as probe molecule has been recommended [250].

The AN values represent dimensionless numbers expressing the Lewis acidity properties of a given solvent A relative to those of $SbCl_5$, which is also the standard compound for setting up the donor number scale. Owing to the good solubility properties of triethylphosphane oxide, acceptor numbers are available for many types of coordinating and non-coordinating solvents. They are particularly useful in characterizing the Lewis acidities of protic solvents and protonic acids. Relationships have been found between the acceptor number and other empirical parameters such as Z, $E_T(30)$, and Y values, as well as many thermodynamic solvation quantities and other solvent-dependent processes [91, 139, 140, 207, 250…253]. The fairly good linear correlations of AN with Z and $E_T(30)$ reveal that the latter represent, to a large extent, measures of the electrophilic properties of the solvents, and therefore of necessity fail when applied to reactions which are mainly influenced by the nucleophilic properties of the solvents. The acceptor numbers are linearly correlated with the $E_T(30)$ parameter according to

$AN = 1.850 \cdot E_T(30) - 59.5$, as established for 51 solvents ($r = 0.944$) by Marcus [294]. This satisfactory linear correlation has been used to calculate AN values from $E_T(30)$ values for such solvents, for which the former are difficult to measure [207].

Finally, the ambitious approach of Catalán et al. to introduce complete new comprehensive scales of solvent dipolarity/polarizability (PP scale), solvent basicity (SB scale), and solvent acidity (SA scale) must be mentioned [296, 335…337]. These three UV/Vis spectroscopic scales are based on carefully selected positively solvatochromic and homomorphic pairs of probe dyes and include values for about 200 organic solvents; for a recent review, see reference [296]. The molecular structures of the three pairs of homomorphic indicator dyes proposed are as follows:



and



(48) (DMANF) *(49)* (FNF)



and



(50) (NI) *(51)* (MNI)



and



(52) (TBSB) *(53)* (DTBSB)

Based on the well-known strong solvent-dependent UV/Vis absorption and fluorescence of 2-amino-7-nitrofluorene (see Section 6.2.4), the donor/acceptor-substituted fluorenes 2-dimethylamino-7-nitrofluorene (48) (DMANF) and its homomorph 2-fluoro-7-nitrofluorene (49) (FNF) were selected as positively solvatochromic indicator dyes for measuring the solvent dipolarity and polarizability. The long-wavelength intramolecular charge-transfer absorption of (48) is connected with an increase in dipole moment of $\Delta m \approx 60 \cdot 10^{-30}$ Cm (18 D; $m_g < m_e$), and is thus sensitive to solute/solvent dipole/dipole and dipole/induced dipole interactions. The rigid and highly conjugated 16 p electron system of (48), a phenylogous nitramide, is readily polarizable, and is thus capable of registering solute/solvent polarizability interactions. It is further assumed that the basicity of (48) does not change very much upon electronic excitation. In order to remove specific solute/solvent interactions, the homomorph (49) was used as a second-

ary probe dye and it is believed that the use of differences of the absorptions of (48) and (49) will cancel many spurious effects. Both compounds exhibit a significant positive solvatochromism, that of (49) being smaller than that of (48). A solvent change from perfluoro-n-hexane to dimethyl sulfoxide results in bathochromic band shifts of $\Delta\lambda = +67$ nm ($\Delta\tilde{\nu} = -4038$ cm^{-1}) and $\Delta\lambda = +26$ nm ($\Delta\tilde{\nu} = -2380$ cm^{-1}), for (48) and (49) respectively [335]. The difference $\Delta\tilde{\nu} = \tilde{\nu}(49) - \tilde{\nu}(48)$, taken as a measure of solvent dipolarity/polarizability (SPP scale), varies from $\Delta\tilde{\nu} = 4692$ cm^{-1} for the gas phase to 6811 cm^{-1} for dimethyl sulfoxide. Using these two media as references to fix the scale from 0.000 (gas phase) to 1.000 (dimethyl sulfoxide, DMSO), the SPP scale is then defined as [296, 335]:

$$\mathrm{SPP(solvent)} = \frac{\Delta\tilde{\nu}(\mathrm{solvent}) - \Delta\tilde{\nu}(\mathrm{gas})}{\Delta\tilde{\nu}(\mathrm{DMSO}) - \Delta\tilde{\nu}(\mathrm{gas})} = \frac{\Delta\tilde{\nu}(\mathrm{solvent}) - 4692}{2119} \qquad (7\text{-}40a)$$

A selection of SPP values is collected in Table 7-5 for nearly the same set of solvents as given in Table 7-4. Whereas cyclohexane is often used as a nonpolar reference solvent in other solvent scales, the SPP scale shows a considerable gap between cyclohexane and the gas phase (0.557 units), which is nearly as wide as that between cyclohexane and polar hexafluoro-2-propanol (0.457 units). Therefore, the choice of cyclohexane as a reference solvent to define a solvent scale has been called into question [335]. For a comparison of the SPP scale with Kamlet and Taft's π^* scale, see reference [338].

For the construction of a solvent hydrogen-bond acceptor (HBA) basicity scale (SB scale), the positive solvatochromism of 5-nitroindoline (50) (NI) and its homomorph 1-methyl-5-nitroindoline (51) (MNI) has been used by Catalán et al. [296, 336]. 5-Nitroindoline (50) can act as a hydrogen-bond donor (HBD), forming H-bonds to HBA solvents, whereas its N-methyl derivative (51) no longer has this property. The two homomorphs have the same HBA properties as well as similar dipole moments and polarizabilities. Thus, they exhibit the same sensitivity to changes in solvent dipolarity/polarizability and should only register a change in the solvent's HBA basicity. A solvent change from tetramethylguanidine (TMG; a strong HBA solvent) to hexafluoro-2-propanol (a non-HBA solvent) leads to bathochromic band shifts of $\Delta\lambda = +22$ nm ($\Delta\tilde{\nu} = -1253$ cm^{-1}) and $\Delta\lambda = +56$ nm ($\Delta\tilde{\nu} = -2963$ cm^{-1}) for (50) and (51), respectively [336]. In order to cancel spurious effects accompanying solute-specific HBD solute/HBA solvent interactions, the difference $\Delta\tilde{\nu} = \tilde{\nu}(50) - \tilde{\nu}(51)$ is taken as a measure of the solvent HBA basicity. This difference varies from $\Delta\tilde{\nu} = 1570$ cm^{-1} for the gas phase to -165 cm^{-1} for TMG. Using these two media as references to normalize the scale between 0.000 (gas phase) and 1.000 (TMG), the SB scale is then defined as [296, 336]:

$$\mathrm{SB(solvent)} = \frac{\Delta\tilde{\nu}(\mathrm{solvent}) - \Delta\tilde{\nu}(\mathrm{gas})}{\Delta\tilde{\nu}(\mathrm{TMG}) - \Delta\tilde{\nu}(\mathrm{gas})} = \frac{\Delta\tilde{\nu}(\mathrm{solvent}) - 1570}{1735} \qquad (7\text{-}40b)$$

A selection of SB values is collected in Table 7-5. Not unexpectedly, a satisfactory linear correlation exists between Catalán's SB values and Kamlet and Taft's β values (see Table 7-4) for 98 solvents ($r = 0.928$), with deviations for some aliphatic amines and ethers with long alkyl chains. For comparisons of the SB scale with further solvent basicity scales, see reference [336].

Table 7-5. A selection of Catalán's solvatochromic parameters SA, SB, and SPP for 40 solvents, taken from reference [296].

Solvents	SA	SB	SPP
Gas phase	0[a)]	0.000[b)]	0.000[b)]
Perfluoro-n-hexane	0[a)]	0.057	0.214
n-Hexane	0[a)]	0.056	0.519
Cyclohexane	0[a)]	0.073	0.557
Dichloromethane	0.040	0.178	0.876
Trichloromethane, Chloroform	0.047	0.071	0.786
Tetrachloromethane	0[a)]	0.044	0.632
Benzene	0[a)]	0.124	0.667
Methylbenzene, Toluene	0[a)]	0.128	0.655
Chlorobenzene	0[a)]	0.182	0.824
Nitrobenzene	0.056	0.240	0.986
Pyridine	0.033	0.581	0.922
Diethyl ether	0[a)]	0.562	0.694
Tetrahydrofuran	0.0	0.591	0.838
1,4-Dioxane	0.0	0.444	0.701
Triethylamine	0[a)]	0.885	0.617
Propanone, Acetone	0.0	0.475	0.881
Cyclohexanone	0.0	0.482	0.874
Ethyl acetate	0.0	0.542	0.795
Ethyl benzoate	0.0	0.417	0.835
(±)-Propylene carbonate	0.106	0.341	0.930
Dimethyl sulfoxide	0.072	0.647	1.000[b)]
Tetrahydrothiophene-1,1-dioxide, Sulfolane	0.052	0.365	1.003
Acetonitrile	0.044	0.286	0.895
Nitromethane	0.078	0.236	0.907
Hexamethylphosphoric acid triamide, HMPT	0[a)]	0.813	0.932
N,N-Dimethylformamide, DMF	0.031	0.613	0.954
N-Methylformamide	0.444	0.590	0.920
Formamide	0.674	0.414	0.833
2-Methyl-2-propanol, t-Butanol	0.145	0.928	0.829
Cyclohexanol	0.258	0.854	0.847
1-Butanol	0.341	0.809	0.837
2-Propanol	0.283	0.762	0.848
1-Propanol	0.367	0.727	0.847
Ethanol	0.400	0.658	0.853
Ethane-1,2-diol, Glycol	0.565	0.534	0.932
Methanol	0.605	0.545	0.857
Acetic acid	0.689	0.390	0.781
2,2,2-Trifluoroethanol	0.893	0.107	0.912
Hexafluoro-2-propanol	1.00	0.014	1.014
Water	1.062	0.025	0.962

[a)] Assumed value.
[b)] By definition.

Finally, a solvent hydrogen-bond donor (HBD) acidity scale (SA scale) was developed by Catalán et al., using the homomorphic merocyanines (52) and (53) as positively solvatochromic indicator dyes, i.e. 1-methyl-4-[(4-oxocyclohexadienylidene)-ethylidene]-1,4-dihydropyridine with one (TBSB) and two t-butyl groups (DTBSB) in the positions a- and a,a′- to the carbonyl group, respectively [296, 337]. The acronyms TBSB and DTBSB stand for o-t-butyl- and o,o′-di-t-butyl-stilbazolium betaine, respectively, i.e. names valid for the zwitterionic mesomeric ground-state structures of these dyes. The positive solvatochromism observed is more in agreement with the less dipolar 1,4-dihydropyridine structure ($m_g < m_e$). At least one of the two lone-pairs of the carbonyl group in dye (52) can interact with HBD solvents by hydrogen bonding. In dye (53), the carbonyl group is shielded on both sides by bulky t-butyl groups and thus specific solute/solvent interactions are hindered. A solvent change from tetrahydrofuran (a non-HBD solvent) to methanol (a strong HBD solvent) results in bathochromic band shifts of $\Delta\lambda = +84$ nm ($\Delta\tilde{\nu} = -2537$ cm^{-1}) and $\Delta\lambda = +15$ nm ($\Delta\tilde{\nu} = -379$ cm^{-1}) for dyes (52) and (53), respectively. As expected, the solvatochromic shift observed for the sterically hindered dye (53) is much less than that for (52), the latter resulting from non-specific solute/solvent interactions only.

The SA scale was established using the solvatochromic comparison method of Kamlet and Taft [224, 226]. A good linear correlation ($r = 0.961$) has been found between the wavenumbers of the absorption maxima of dyes (52) and (53), measured in fifty non-acidic or non-HBD solvents, according to Eq. (7-40c):

$$\tilde{\nu}(52) = 1.405 \cdot \tilde{\nu}(53) - 6289 \qquad (7\text{-}40c)$$

It describes a reference line that can be used to quantify the deviations from this line found for acidic or HBD solvents. These deviations, $\Delta\tilde{\nu}$ (solvent), are then given by Eq. (7-40d) [cf. Fig. 7-3 and Eq. (7-33)]:

$$\Delta\tilde{\nu}(\text{solvent}) = \tilde{\nu}(52) - [1.405 \cdot \tilde{\nu}(53) - 6289] \qquad (7\text{-}40d)$$

In going from tetrahydrofuran to ethanol, these $\Delta\tilde{\nu}$ values increase from –39 to 1300 cm^{-1}. In order to obtain a normalized scale of near-zero HBD acidity for non-HBD solvents and a value of roughly unity for HBD solvents, the SA scale was defined as follows [296, 337a]:

$$SA(\text{solvent}) = \frac{\Delta\tilde{\nu}(\text{solvent})}{\Delta\tilde{\nu}(\text{ethanol})} \cdot 0.4 = \frac{\Delta\tilde{\nu}(\text{solvent})}{1300} \cdot 0.4 \qquad (7\text{-}40e)$$

The direct determination of SA values for more acidic solvents was not possible because in such solvents the indicator dyes (52) and (53) are protonated at the carbonyl oxygen atom. For such acidic solvents, 3,6-diethyl-1,2,4,5-tetrazine (DETZ) was introduced as a third probe dye [337b]. This aromatic tetrazine exhibits a solvent-dependent $n \rightarrow p^*$ absorption in the visible region: a change from methylcyclohexane ($\lambda_{max} = 550$ nm) to hexafluoro-2-propanol ($\lambda_{max} = 517$ nm) leads to a hypsochromic band shift of $\Delta\lambda = -33$ nm ($\Delta\tilde{\nu} = +1160$ cm^{-1}), mainly due to the HBD acidity of the

solvent (see Section 6.2.3). Analogous application of Kamlet and Taft's solvatochromic comparison method made it possible to extend the SA scale to more acidic solvents; see reference [337b] for details. A selection of SA values is included in Table 7-5. For HBD solvents with known α and SA values, a rather good linear correlation between Kamlet and Taft's α values and Catalán's SA values has been found ($n = 7$, $r = 0.951$) [337a].

The general SPP scale of solvent dipolarity/polarizability and the specific SB and SA scales of solvent HBA basicity and HBD acidity, respectively, are orthogonal to one another and they can be used in the correlation analysis of solvent effects in single- or, in combination with the others, in two- or three-parameter correlation equations, depending on the solvent-influenced process under consideration; see also Section 7.7. Examples of the correlation analysis of a variety of other solvent-dependent processes by means of SPP, SB, and SA values, including those used for the introduction of other solvent polarity parameters, can be found in references [335…337, 340…342]. In particular, comparisons with Kamlet and Taft's π^* scale [340] and Winstein and Grunwald's Y scale [341] have been made.

7.5 Empirical Parameters of Solvent Polarity from Other Measurements

An important measure of the total molecular cohesion per unit volume of liquid is the cohesive pressure (also called cohesive energy density), which characterizes the energy associated with all the intermolecular solvent/solvent interactions in a mole of the solvent. The cohesive pressure is defined as the molar energy of vapourization to a gas at zero pressure, ΔU_v, per molar volume of the solvent, V_m, according to Eqs. (3-5) and (5-76) in Sections 3.2 and 5.4.2, respectively [93, 94]. The cohesive pressure is related to the internal pressure p; cf. Eq. (3-6) and Table 3-2 in Section 3.2.

The square root of the cohesive pressure has been termed by Hildebrand and Scott the solubility parameter d because of its value in correlating and predicting solubility behaviour of non-electrolytes (ionogens) [93, 94]; cf. Eqs. (2-1) and (5-77) in Sections 2.1 and 5.4.2, respectively. Comprehensive lists of d values are given in references [94, 95, 343]; cf. Table 3-3 in Section 3.2 for a selection. Correlations of d values with other solvent polarity parameters have been attempted, but in most cases only relatively poor mutual relationships were found [96, 97, 255, 256]. Since d values, which are characterized by the energy needed to separate molecules of the liquid, give only a measure of the attractive forces between the solvent molecules, they need not necessarily be a measure of the solute/solvent interaction forces. It could be that some of the solvent/solvent interaction forces are also of relevance in particular solute/solvent interactions. Rather, the d values are related to the energy necessary to form a cavity in the solvent, which can then accommodate the solute molecule. In accordance with this is the experimental observation that a good solvent for a certain non-electrolyte solute should have a d value close to that of the solute [93, 94]. Depending on the solvent, creation of a cavity in an organic solvent requires about 20 . . . 40 kJ/mol (5 . . . 10 kcal/mol).

Sometimes, Hildebrand's solubility parameter d has been incorrectly used in linear Gibbs energy relationships; cf. for example [96, 97, 226, 255]. Since in linear Gibbs energy relationships the correlated solvent-dependent solute properties (e.g., lg K, lg k, $h \cdot n$) are proportional to Gibbs energy changes of reaction or activation (ΔG, $\Delta G^{\neq}$) and

excitation energies (E_T), all the terms of a regression equation should include an energy dimension; cf. Eqs. (7-2) and (7-3) in Section 7.1. Therefore, it is unjustified to use δ values in such regression equations because they have the dimension of the square root of the energy ($J^{1/2} \cdot cm^{-3/2}$). Instead of δ, the cohesive pressure c, which is equal to δ^2, should be used, as has been demonstrated in work dealing with the inclusion of cavity terms in multiparameter regression equations [256…258].

It has been shown that gas-liquid chromatographic methods are particularly suitable for a quantitative characterization of the polarity of solvents. In gas-liquid chromatography it is possible to determine the solvent power of the stationary liquid phase very accurately for a large number of substances [98, 99, 259, 260]. Many groups of substances exhibit a certain dependence of their relative retention parameters on the solvation characteristics of the stationary phase or of the separable components. In determining universal gas-chromatographic characteristics, the so-called retention index I, introduced by Kováts [100], is frequently used. The elution maxima of individual members of the homologous series of n-alkanes (C_nH_{2n+2}) form the fixed points of the system of retention indices. The retention index is defined by means of Eq. (7-41),

$$I = 100 \cdot \frac{\lg V_x - \lg V_n}{\lg V_{n+1} - \lg V_n} + 100 \cdot n \qquad (7\text{-}41)$$

where V_x is the specific retention volume of solute x and V_n and V_{n+1} are the retention volumes of two n-alkanes with n and $n+1$ carbon atoms, respectively; hence, $V_n < V_x < V_{n+1}$. The retention index is independent of the gas-chromatographic equipment used, and depends only on the solute x, the temperature, and the stationary phase. The retention index specifies with which n-alkane a solute leaves the separation column, whereby the number of carbon atoms of the respective n-alkane is multiplied by 100. In other words, a retention index of 800 or 1100 means that the solute leaves the column simultaneously with either n-octane or n-undecane, respectively. A retention index of 732 implies that the solute exhibits the same retention time as a hypothetical n-alkane with 7.32 carbon atoms.

Kováts and Weiß [101] used these retention indices to examine the polarity of stationary liquid phases. If the total Gibbs energy of dissolution is determined by the work of separating the solvent molecules to form a hole, by a dispersion term, a polar term, and the energy of the donor/acceptor interaction, the sum of the last two terms is proportional to the difference between the retention indices of the solute on the column with a given liquid phase X (I_T^X) and with a standard nonpolar stationary phase (i.e. a hydrocarbon; $I_T^{St\,Ap}$) at temperature T. This difference, $\Delta I_T^X = I_T^X - I_T^{St\,Ap}$, is then proposed as the polarity parameter. Using 1-chloro- and 1-bromo-n-hexadecane as the standard dipolar stationary phase, and n-hexadecane as the standard nonpolar stationary phase, $\Delta I_{50}^{Cl/Br}$ was defined as the new polarity parameter according to Eq. (7-42) [101].

$$\Delta I_{50}^{Cl/Br} = I_{50}^{\text{1-chloro or 1-bromo-n-hexadecane}} - I_{50}^{\text{n-hexadecane}} \qquad (7\text{-}42)$$

The $\Delta I_{50}^{Cl/Br}$ values show correlations with other empirical solvent polarity parameters, such as Z, $E_T(30)$, and $\lg k_1$ of 4-methoxyneophyl tosylate solvolysis [101].

Another more rigorous approach based on retention indices, taking into account the fact that the polarity of the column depends not only on the nature of the stationary phase but also on the type of substance being analyzed, has been suggested by Rohrschneider [98, 102]. The polarity of a stationary phase must therefore be assessed simultaneously with respect to a whole group of compounds possessing varying donor/acceptor properties. Rohrschneider proposed the "ve-term equation (7-43) for the solute retention, which has found widespread use in gas-chromatography [98, 99, 259, 260].

$$\Delta I = a \cdot x + b \cdot y + c \cdot z + d \cdot u + e \cdot s \qquad (7\text{-}43)$$

a, b, c, d, and e are polarity factors of the stationary liquid phase, and x, y, z, u, and s are polarity factors of the substances undergoing analysis. Having determined the polarity factors of 22 di¤erent stationary phases with respect to chosen standard substances (benzene, ethanol, butan-2-one, nitromethane, and pyridine), and those of 30 other solutes, Rohrschneider calculated 660 retention indices with only small limits of error.

Using a di¤erent set of standard substances, i.e. substituting 1-butanol, pentan-2-one, and 1-nitropropane for the rather volatile ethanol, butan-2-one, and nitromethane, McReynolds developed an analogous approach [103]. Altogether, he characterized over 200 liquid stationary phases using a total of 10 probes. A statistical analysis of the McReynolds retention index matrix using the principal component analysis method has shown that only three components are necessary to reproduce the experimental data matrix [262]. The "rst component is related to the polarity of the liquid phase, the second depends almost solely on the solute, and the third is related to speci"c interactions with solute hydroxy groups [262].

Both of these approaches used in the characterization of stationary liquid-phase polarities by means of retention indices have been further explored and expanded [104, 259…261]. For a review on the characterization of solvent properties of phases used in gas-liquid chromatography by means of the retention index system, see reference [344]. Similar methods for the characterization of solvent polarity in liquid-liquid and liquid-solid chromatography can be found in references [105…107]; also Section A-7 and Tables A-10 and A-11 in the Appendix.

7.6 Interrelation and Application of Solvent Polarity Parameters

From the previous Sections, we can conclude that there are many empirical solvent scales, the most comprehensive of which are the solvatochromic ones, see for example Table 7-3. Unfortunately, too many solvent scales have been proposed during the last decades. Around 35 di¤erent solvent scales are known.* Only about ten of them have found wider application in the correlation analysis of solvent e¤ects, i.e. Y, Z, $E_T(30)$, a,

* At present, the situation is not quite as bad as in the correlation analysis of substituent e¤ects, where even more substituent parameters than common substituents seem to be known. It has been suggested that new solvent polarity scales should only be introduced into the literature if they exhibit signi"cant advantages over existing solvent scales [235].

b, p^*, DN and AN, as well as SPP, SA, and SB. The application of most solvent scales is restricted by the fact that they are known only for an insufficient number of solvents. The catalogue of common organic solvents available to the chemist numbers about 300, not to speak of the infinite number of solvent mixtures! The extension of most solvent scales is restricted by the inherent properties of the selected reference process, which exclude the determination of solvent parameters for certain, often important solvents (e.g. chemical reactions between solute and solvent, solubility problems, etc.). For this reason, the most comprehensive solvent polarity scales are those derived from spectroscopic reference processes, which are the most easily measured for a large set of organic solvents.

In general, all these parameters constitute more comprehensive measures of solvent polarity than the relative permittivity or any other single physical characteristic, since they reflect more reliably the complete picture of all intermolecular forces acting between solute and solvent molecules. The solvent-dependent processes used to define solvent polarity parameters may be regarded as probes that permit a purely empirical investigation of solvent effects. In applying these parameters, however, it is tacitly assumed that the intermolecular interactions in the reference system used to develop a particular solvent scale are similar to those in the system the prediction of whose solvent effects is being raised. This is obviously true only for closely related solvent-dependent processes. Therefore, the use of single solvent parameters to predict solvent effects on equilibria, reaction rates, and spectral absorptions should be very restricted. One cannot expect a parameter to be universally useful for all kinds of solvent-sensitive processes, since any correlation of solvent effects of a particular process with a solvent polarity parameter is, in fact, a comparison with the effect of solvent on a reference process. However, if one compares the various empirical solvent scales (cf. Tables 7-1 to 7-5), based on very different solvent-sensitive reference processes, varying strongly in the energies involved, one finds, surprisingly, that most of the existing empirical solvent scales agree with each other very well qualitatively and even sometimes quantitatively. In spite of the large energy changes connected with a solvent change from methanol to hydrocarbon solvents for Z (ca. 100 kJ/mol), $E_T(30)$ (ca. 105 kJ/mol), $\lg k_1$ of 4-methoxyneophyl tosylate solvolysis ($-\ln 10 \cdot RT \cdot \lg k_1$ ca. 29 kJ/mol), and W values ($-\ln 10 \cdot RT \cdot W$ ca. 2 kJ/mol), in all four cases one obtains similar polarity orders for the solvents. This led Berson to observe that "... in this respect a set of solvents behaves like an elephant, which can lift a log or a peanut with equal dexterity" [52].

In particular, there are good linear correlations between the $E_T(30)$ values and some other empirical solvent polarity parameters according to Eq. (7-44),

$$y = a \cdot E_T(30) + b \quad \text{or} \quad y = a \cdot E_T^N + b \tag{7-44}$$

where a and b were determined by the method of least squares. Compilations of such linear correlations between $E_T(30)$ values and other solvent polarity parameters can be found in references [124, 294b]. For example, there is a satisfactory linear correlation between the values of $E_T(30)$ and Z according to Eq. (7-45) ($n = 60$ solvents; correlation coefficient $r = 0.972$) [294b].

$$Z/(\text{kcal} \cdot \text{mol}^{-1}) = 1.27 \cdot E_T(30)/(\text{kcal} \cdot \text{mol}^{-1}) + 13.0 \tag{7-45}$$

The excellent $Z/E_T(30)$ correlation for a selected set of 15 solvents common to both scales has been used to calculate $E_T(30)$ values from Z values for acidic solvents for which $E_T(30)$ values are not available [172]; cf. footnote [f] of Table 7-3. A similar satisfactory linear correlation between $E_T(30)$ values and acceptor numbers allows the calculation of AN values that are not directly available [207, 294b].

Of particular interest are the correlations between $E_T(30)$ or E_T^N values and the solvatochromic parameters p^*, a, and b of Kamlet and Taft [84a, 226, 235] as well as the SPP, SA, and SB parameters of Catalán et al. [335…337].

For 166 solvents, a satisfactory correlation between $E_T(30)$ or E_T^N values and Kamlet and Taft's solvatochromic parameters p^* and a has been obtained by Marcus [294b], according to Eqs. (7-46a) and (7-46b) ($n = 166, r = 0.979$):

$$E_T(30)/(\text{kcal}\cdot\text{mol}^{-1}) = 11.5\cdot p^* + 15.2\cdot a + 31.2 \qquad (7\text{-}46a)$$

$$E_T^N = 0.36\cdot p^* + 0.47\cdot a + 0.01 \qquad (7\text{-}46b)$$

An analogous satisfactory correlation has been found by Catalán [296] between $E_T(30)$ or E_T^N values and the solvatochromic parameters SPP, SA, and SB for 138 solvents, according to Eqs. (7-47a) and (7-47b) ($n = 138, r = 0.965$):

$$E_T(30)/(\text{kcal}\cdot\text{mol}^{-1}) = 20.1\cdot \text{SPP} + 24.9\cdot \text{SA} + 3.9\cdot \text{SB} + 20.7 \qquad (7\text{-}47a)$$

$$E_T^N = 0.62\cdot \text{SPP} + 0.77\cdot \text{SA} + 0.12\cdot \text{SB} - 0.31 \qquad (7\text{-}47b)$$

Eqs. (7-46) and (7-47) are in agreement with the observation that $E_T(30)$ and E_T^N values measure not only a blend of solvent dipolarity and solvent polarizability (as described by p^* and SPP, respectively), but for HBD solvents the solvent HBD acidity as well. In HBD solvents, the HBD acidity can even be the dominating solvent property: the quotients of the corresponding regression coefficients in Eqs. (7-46b) and (7-47b) are $0.47/0.36 = 1.31$ and $0.77/0.62 = 1.24$, respectively, i.e. greater than unity, which would be expected for equal contributions of both types of solute/solvent interactions. For non-HBD solvents, for which a and SA values are equal to zero, the $E_T(30)$ and E_T^N values essentially reflect only a blend of solvent dipolarity and polarizability. Not unexpectedly, the solvent HBA (or EPD) basicity (as described by b and SB, respectively) is not measured by the $E_T(30)$ or E_T^N values: the b and SB parameters make little or no contribution in the correlation equations (7-46) and (7-47). This is in agreement with the molecular structure of the pyridinium N-phenolate indicator dye (44), which has no functional groups capable of interaction with HBA solvents. In the correlation analysis of solvent effects with significant contributions of HBD solute/HBA solvent interactions, a combination of $E_T(30)$ values and solvent donor numbers DN can considerably improve the correlations; see Eq. (7-60) in Section 7.7.

For attempts to improve the $E_T(30)/p^*$ correlations using modified Kamlet…Taft parameters (p_n^* and a_n), see references [234, 235].

The solvatochromic polarity parameter $E_T(30)$ or E_T^N is also related to some physical solvent properties beyond the solvent dipole moment, provided that solvents capable of specific solute/solvent interactions are excluded. In particular, Brédas et al.

[194] and Makitra et al. [345] have tried to correlate $E_T(30)$ values with various functions of the solvent relative permittivities ε_r and solvent refractive indices n. For a selection of 50 aliphatic non-HBD solvents, the correlation equations (7-48a,b) have been established by Makitra et al. [345] ($n = 50, r = 0.952$):

$$E_T(30)/(\text{kcal} \cdot \text{mol}^{-1}) = 43.70 \cdot f(\varepsilon_r) + 4.12 \cdot f(n^2) + 20.63 \qquad (7\text{-}48a)$$

$$E_T^N = 1.349 \cdot f(\varepsilon_r) + 0.127 \cdot f(n^2) - 0.311 \qquad (7\text{-}48b)$$

where $f(\varepsilon_r)$ is equal to the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$ and $f(n^2)$ represents the electronic polarizability term $(n^2 - 1)/(n^2 + 2)$, taken from the Lorenz…Lorentz equation. From the magnitudes of the respective regression coefficients, it can easily be seen that the $f(\varepsilon_r)$ term describes the predominant influence on the $E_T(30)$ or E_T^N parameter, provided that only non-specific solute/solvent interactions are considered [194, 345]. Somewhat different correlation equations have been found for aromatic solvents [194, 345]. For further correlation equations, including cross-terms such as $f(\varepsilon_r, n^2) = (\varepsilon_r - 1)(n^2 - 1)/(2\varepsilon_r + 1)(2n^2 + 1)$, see references [194, 345]. However, it was not possible to find relationships between $E_T(30)$ and various functions of permittivity and polarizability that are statistically valid for all groups of solvents.

Further single- and multi-parameter equations for correlations between $E_T(30)$ values and other empirical solvent polarity parameters, between various other empirical solvent parameters (e.g. a/Z, a/AN, b/DN, etc.), and between empirical solvent parameters and macroscopic physical solvent properties have been collected in the reviews of Marcus [294b], Abboud and Notario [295], and Catalán [296].

The application of single solvent parameters such as the $E_T(30)$ values in correlating other solvent-dependent processes has proved to be surprisingly successful. This means that the blend of different solute/solvent interaction forces, as measured by the solvatochromic pyridinium N-phenolate betaine dye (44), is nearly the same as in numerous other solvent-dependent chemical reactions and spectral absorptions. It seems, therefore, that the blend of solute/solvent interactions as measured by the $E_T(30)$ values represents a kind of "mean solvent polarity", representative of many solvent-sensitive processes. As indicated by Eqs. (7-46) and (7-47), $E_T(30)$ values represent a combined measure of solvent dipolarity/polarizability and solvent HBD acidity (Lewis acidity), the latter property only being significant for protic (HBD) solvents. $E_T(30)$ correlations for more than 100 solvent-sensitive processes have been collected in references [2, 124, 200…203, 294b]. Three typical examples are given in Figs. 7-4 to 7-6.

Fig. 7-4 shows an excellent correlation between $E_T(30)$ values and the $\pi \rightarrow \pi^*$ transition energies of the open-chain form of a merocyanine dye obtained by irradiation of the corresponding photochromic benzthiazolospiropyran [108] (cf. the related spiropyran (27a)/merocyanine (27b) equilibrium in Section 4.4.2).

The structure of the open-chain form was assigned on the basis of its negative solvatochromic behaviour, which is similar to that of other meropolymethines such as the pyridinium N-phenolate betaines [108]. The correlation shown in Fig. 7-4 allows one to calculate absorption maxima of the merocyanine dye in other solvents for which $E_T(30)$ values are known.



Fig. 7-4. Correlation between $E_T(30)$ [66] and the $p.p^*$ transition energy, E_T', of a merocyanine dye [108]. $E_T' = 0.42 \cdot E_T(30) + 33.5$ ($n = 14$; $r = 0.986$; $s = 0.482$).

A correlation between $E_T(30)$ values and the rate of thermal racemization of chiral allyl p-tolyl sulfoxide according to Eq. (5-38) in Section 5.3.2 is shown in Fig. 7-5 [109].

The linear decrease in reaction rate with increasing solvent polarity has been considered as evidence in support of the proposed reaction mechanism, involving a less dipolar cyclic activated complex (cf. discussion of this reaction in Section 5.3.2).

Fig. 7-6 demonstrates the correlation between $E_T(30)$ and the relative rates for the S_N2 Menschutkin reaction between a tertiary amine and a haloalkane in non-HBD solvents. The values of the second-order rate constants are taken from the compilation made by Abraham and Grellier [110].

A comparison of Fig. 7-6 with Fig. 5-11 in Section 5.4.3 reveals some improvement of the poor correlation between $\lg(k/k_0)$ and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$. If HBD solvents are included in the $E_T(30)/\lg(k/k_0)$ correlation, two lines are obtained as in Fig. 5-12, one for non-HBD and one for HBD solvents. This demonstrates the utility of such correlations in discovering speci“c solute/solvent interactions such as hydrogen-bonding.



Fig. 7-5. Correlation between $E_T(30)$ [66] and $\lg k_1$ for the thermal racemization of chiral allyl p-tolyl sulfoxide at 60.7°C [109]; cf. Eq. (5-38). $\lg k_1 = -0.078 \cdot E_T(30) - 0.55$ ($n = 7$; $r = -0.976$; $s = 0.222$).

It has been stated that, when speci“c hydrogen-bonding e¤ects are excluded, and di¤erential polarizability e¤ects are similar or minimized, the solvent polarity scales derived from UV/Vis absorption spectra ($Z, S, E_T(30), p^*, w_R, E_K$), ”uorescence spectra (Py), infrared spectra (G), ESR spectra ($a(^{14}N)$), ^{19}F NMR spectra (P), and ^{31}P NMR spectra (AN) are linear with each other for a set of select solvents, i.e. non-HBD aliphatic solvents with a single dominant group dipole [263]. This result can be taken as con“rmation that all these solvent scales do in fact describe intrinsic solvent properties and that they are to a great extent independent of the experimental methods and indicators used in their measurement [263]. That these empirical solvent parameters correlate linearly with solvent dipole moments and functions of the relative permittivities (either alone or in combination with refractive index functions) indicates that they are a measure of the solvent dipolarity and polarizability, provided that speci“c solute/solvent interactions are excluded.

Applying a quantitative structure…property relationship (QSPR) analysis of 45 di¤erent empirical solvent scales and 350 solvents, a direct calculation of predicted



Fig. 7-6. Correlation between $E_T(30)$ [66] and $\lg(k/k_0)$ for the reaction between triethylamine and iodoethane in apolar and dipolar non-HBD solvents at 25°C (rate constants relative to *n*-hexane as "slowest" solvent) [110].
$\lg(k/k_0) = 0.248 \cdot E_T(30) - 6.54$ ($n = 28$; $r = 0.920$). (1) *n*-hexane, (2) cyclohexane, (3) tetrachloromethane, (4) diethyl ether, (5) toluene, (6) benzene, (7) 1,4-dioxane, (8) 1,1,1-trichloroethane, (9) iodobenzene, (10) bromobenzene, (11) chlorobenzene, (12) tetrahydrofuran, (13) ethyl acetate, (14) trichloromethane, (15) 1,1-dichloroethane, (16) 2-butanone, (17) dichloromethane, (18) acetophenone, (19) nitrobenzene, (20) benzonitrile, (21) 1,2-dichloroethane, (22) acetone, (23) propionitrile, (24) *N*,*N*-dimethylformamide, (25) dimethyl sulfoxide, (26) acetonitrile, (27) nitromethane, and (28) propylene carbonate.

values of empirical solvent parameters for any scale and for any previously unmeasured solvent is possible by means of the CODESSA program (i.e. comprehensive descriptors for structural and statistical analysis), developed by Katritzky et al. [346]. The QSPR models for each of the solvent scales are constructed using only theoretical descriptors, derived solely from the molecular solvent structure. This QSPR study has enabled a classification of the various empirical solvent scales, which provides considerable insight into the physicochemical background of these solvent scales [346]. Furthermore, a principal component analysis (PCA) has been carried out with 40 solvent scales as variables, each of them having 40 data points for 40 solvents as objects [347]. This PCA treatment allows a rational classification and grouping of both solvents and solvent scales. For 36

of the solvent polarity scales, an average of 88% of the variance is already described by the “rst three principal components, which can be roughly correlated to the solvent relative permittivity and dipolarity, as well as to their electrophilicity (HBD acidity) and nucleophilicity (HBA basicity) [347]; see also the end of Section 3.5. Another statistical treatment of a set of 32 solvent polarity scales for 45 solvents, using the program SMIRC (i.e. selection of a set of minimally interrelated columns), has been carried out by Palm et al. [348], who introduced the “rst multi(four)-parameter equation for the correlation analysis of solvent e¤ects; see Section 7.7.

There has been some criticism of the method of linear solvation Gibbs energy relationships. Using simulated solvatochromic correlation analyses by modeling dipole/dipole and dipole/induced-dipole interactions with a single combined parameter, Carr et al. have shown that, although good correlations can be obtained, the regression coe‹cients can be incorrect and not representative of the system under study [264]. Therefore, caution against overinterpretation of solvatochromic regression equations has been strongly recommended [264].

In conclusion, it is fair to say that the method of linear Gibbs energy correlations is still the most practical and simple method for predicting solvent e¤ects on reaction rates, equilibria, and spectral absorptions, as well as for predicting substituent e¤ects for reactions in solution. In so far as one understands why the model process responds to a solvent or substituent change, something can be learnt about the particular process under study. This kind of procedure has been criticized as being too empirical. However, one should take into account that not only the basic postulates of linear Gibbs energy relationships (i.e. additivity and separability) are of theoretically acceptable form, but the choice of a suitable reference process also requires intensive application of theory since the solute/solvent interaction of the model process and the process under investigation must be closely related.

7.7 Multiparameter Approaches

In spite of the observation that single empirical parameters, such as those mentioned in the foregoing Sections, may serve as good approximations of solvent polarity in the sense de“ned in Section 7.1 [1, 3], there are many examples of solvent-sensitive processes known which cannot be correlated to only one empirical solvent parameter. It has been repeatedly found that the simple concept of ••polarity•• as a universally determinable and applicable solvent characteristic is a gross oversimpli“cation. The solvation capability or solvation power of a solvent, which has been roughly divided into non-speci“c and speci“c solute/solvent interactions, is the result of many di¤erent kinds of interaction mechanisms between the molecules of the solute and the solvent (cf. Section 2.2). Solvent e¤ects are basically more complicated and often more speci“c than substituent e¤ects. In the latter case, linear Gibbs energy relationships such as the Hammett equation are well-established and are known to work very well. In order to take into account two or more aspects of solvation, a multiparameter approach of the general form

$$A = A_0 + b \cdot B + c \cdot C + d \cdot D + \cdots \qquad (7\text{-}49)$$

has been tried, where A is the value of a solvent-dependent physicochemical property ($\lg K$, $\lg k$, $h \cdot n$, etc.) in a given solvent, and A_0 is the statistical quantity corresponding to the value of this property in the gas phase or in an inert solvent. $B, C, D, \ldots$ represent independent but complementary solvent parameters, which account for the different solute/solvent interaction mechanisms. $b, c, d, \ldots$ are the regression coefficients describing the sensitivity of property A to the different solute/solvent interaction mechanisms. Such an equation can be applied only to data for a large number of well-chosen solvents, and its success must be examined by proper statistical methods [14, 15]. The separation of solvent polarity into various solute/solvent interaction mechanisms is purely formal and may not even be theoretically valid as the interactions could be coupled, i.e. not operate independently of each other. Nevertheless, if this separation can be reasonably accomplished, the resultant parameters may be used to interpret solvent effects through such multiple correlations, thus providing information about the type and magnitude of interactions with the solvent. On the basis of Eq. (7-49), the often observed failure of single solvent parameters can readily be understood. Any single empirical solvent parameter must have a fixed relative sensitivity to each of the various interaction mechanisms implied in Eq. (7-49). Thus, only application to processes that have the same relative sensitivity to various interaction mechanisms as the single solvent parameter will give a good correlation. Studies of multiparameter approaches to solvent effects on physical and chemical properties based on the general Eq. (7-49) have been made by Katritzky et al. [111], Koppel and Palm [6, 112], Kamlet and Taft [84a, 224, 226], Krygowski and Fawcett [113], Swain et al. [265], Mayer [266, 251], Dougherty [114], Pytela [349], Famini and Wilson [350, 364…369], Drago [351], and Catalán [296, 335…337]. Multiparameter treatments of solvent effects have been reviewed [293, 294, 296, 316, 339, 352].

Katritzky et al. [111] tested various multiparameter equations using linear combinations of existing empirical solvent parameters. The most successful treatment combines the $E_T(30)$ values (cf. Table 7-3) with functions of relative permittivity and index of refraction. Using $E_T(30)$ and the Kirkwood function $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, a two-parameter equation was constructed which allows independent variation of dipole/dipole and hydrogen-bonding forces. This equation is based on the assumptions that the Kirkwood function adequately represents dipole/dipole interactions and that $E_T(30)$ values are sensitive to both dipolar interactions and the interaction between solute and hydrogenbond donor (HBD) solvents. It could be shown that correlations of rates, equilibria, and spectral properties are indeed significantly improved by such multiparameter treatment, which implicitly allows for various independent interaction mechanisms between solvent and solute ground, transition, and excited states [111]. For a recent quantitative structure…property relationship (QSPR) analysis and a principal component analysis (PCA) of a large variety of different solvent scales by Katritzky et al., see references [346, 347] and the end of Section 3.5.

A more rigorous approach has been suggested by Koppel and Palm [6, 112], who argue that a complete description of all solute/solvent interactions must include both non-specific and specific effects. They proposed the general four-parameter equation (7-50), which relates the variation of a given property A to two non-specific (Y and P) and two specific characteristics of the solvent (E and B).

$$A = A_0 + y \cdot Y + p \cdot P + e \cdot E + b \cdot B \qquad (7\text{-}50)$$

A and A_0 are defined as in Eq. (7-49); by definition, A is equal to A_0 for the gas phase. The non-specific parameters Y (*) and P measure solvent polarization and polarizability, respectively, according to classical dielectric theory. E and B are specific parameters measuring the Lewis acidity (electrophilic solvating power) and Lewis basicity (nucleophilic solvating power) of the solvent, and y, p, e, and b are the corresponding regression coefficients indicating the sensitivity of A to the four different solvent parameters. Relative permittivities, ε_r, are the basis of Y, and were used in the form of the Kirkwood function, $(\varepsilon_r - 1)/(2\varepsilon_r + 1)$, or as $(\varepsilon_r - 1)/(\varepsilon_r + 2)$, a function based on the expression for molar polarization. The functions $(n^2 - 1)/(2n^2 + 1)$ or $(n^2 - 1)/(n^2 + 2)$ of the refractive index for sodium light were used for the polarizability parameter, P. In the effective numerical range, these pairs of functions are approximately co-linear, and the choice between the two functions $f(\varepsilon_r)$ and the two functions $f(n^2)$, respectively, is largely arbitrary.

A scale of Lewis acidity, E, was based on the $E_T(30)$ values discussed in Section 7.4 (cf. Table 7-3), but these were corrected for the influence of non-specific effects, and adjusted to an origin $E = 0$ for the gas phase by means of Eq. (7-51) [6, 112, 115].

$$E = E_T(30) - E_T^0(30) - y \cdot Y - p \cdot P$$

$$= E_T(30) - 25.10 - 14.84 \cdot Y - 9.59 \cdot P \qquad (7\text{-}51)$$

The subtraction of the polarization $(y \cdot Y)$ and polarizability $(p \cdot P)$ contributions from the total solvent effect allows an estimation of the contribution from specific solute…solvent interactions. This correction of $E_T(30)$ values was made using least-squares regression analysis by correlating the data for suitably selected non-specifically and specifically interacting solvents. E values derived in this way from $E_T(30)$ values are presented in references [6, 115]; they range from zero (gas phase, saturated hydrocarbons) to about 22 kcal/mol for water. By definition, $e = 1$ in Eq. (7-50) for the reference process, i.e. the $p \rightarrow p^*$ transition of the pyridinium N-phenolate betaine dye (44). The reason for assuming that $E_T(30)$, and thus E, largely relates to Lewis acidity in protic solvents has already been mentioned.**

A scale of Lewis basicity, B, was based on the O–D infrared stretching band of CH_3OD, according to Eq. (7-35) in Section 7.4 [6, 87, 112]. EPD solvents reduce $\tilde{\nu}_{O-D}$ through hydrogen-bonding and the wavenumber shift measures the strength of the HBD solute/EPD solvent interaction. For use in Eq. (7-50) the shifts were adjusted to an origin $B = 0$ for the gas phase. By definition, $b = 1$ in Eq. (7-50) for the reference process, the IR wavenumber shifts of CH_3OD.

A detailed analysis of solvent effects on various solvent-sensitive processes by means of Eq. (7-50) has been presented by Koppel and Palm [6, 112]. If Eq. (7-50)

---

* Not to be confused with the Y values of Grunwald and Winstein in Eq. (7-14).

** The procedure for parameterization of solvent electrophilicity has been criticized, mainly because it was found that the use of $E_T(30)$ instead of E in the multiple regression treatment of solvent effects is often quite successful; see reference [15, 116] for examples. It has been shown that values of $E_T(30)$ and E are linearly correlated, at least for solvents with an $E_T(30)$ value of greater than ca. 40 kcal/mol [178]. This calls into question the value of Koppel and Palm•s division of $E_T(30)$ into pure electrophilicity effects and non-specific effects by means of Eq. (7-51).

quantitatively re”ects the in”uence of all basic types of solute/solvent interactions, it should be possible to correlate any solvent-dependent kinetic or spectral data in terms of Y, P, E, and B solvent parameters. This has been done for more than sixty solvent-sensitive processes [6]. Surprisingly, the majority of the processes investigated by Koppel and Palm [6] depend only on a single speci“c solvent parameter: in “fty cases, only electrophilic solvation is important (thus con“rming the wide applicability of single $E_T(30)$ values), while in seven cases, nucleophilic solute/solvent interaction is the predominant solvation mechanism. Only one representative example will be mentioned in more detail in order to demonstrate what conclusions for any particular case can be drawn from Eq. (7-50). For a critical discussion of the Koppel…Palm treatment of solvent e¤ects and valuable comments on its applicability, see reference [15].

The rate of reaction between benzoic acid and diazo-diphenylmethane in 44 non-HBD solvents correlates signi“cantly with all four Koppel…Palm parameters of Eq. (7-50), according to Eq. (7-52) with $n = 44$, $r = 0.976$, and standard deviation $s = 0.188$ [15, 116];cf. also [117].

$$\lg k_2 = -3.13 + 4.58f(\varepsilon_r) + 11.96f(n^2) + 0.195E - 0.018B \qquad (7\text{-}52)$$

The regression coe‹cients a and b are of opposite sign, and the order of decreasing signi“cance of the four terms is $B > f(\varepsilon_r) > E > f(n^2)$, as shown by the stepwise regression. This result supports the mechanism given in the following reaction scheme:



The rate-determining step involves a proton-transfer from the carboxylic acid to form a diphenylmethyldiazonium/carboxylate ion-pair, which rapidly reacts in subsequent product-determining steps to give esters (or ethers in the case of alcoholic solvents). The negative sign of the basicity term of Eq. (7-52) indicates nucleophilic stabilization of the initial carboxylic acid, i.e. rate deceleration with increasing solvent basicity. The positive sign of the acidity term, however, indicates electrophilic stabilization of the activated complex, resulting in rate acceleration with increasing solvent acidity. Thus, the two counteracting solvent e¤ects are nicely unravelled by this kind of treatment.

Recently, a statistical treatment of 31 di¤erent solvent parameter scales was carried out by Palm et al. [348], in order to “nd the minimum number of solvent parameters necessary for an adequate multiparameter description of solvent-dependent processes. Numerous applications of the Koppel…Palm Eq. (7-50) and modi“cations thereof to the correlation analysis of many solvent-dependent processes, taken from di¤erent areas of

chemistry (e.g. kinetics [353, 354], extraction [302], partitioning [355], and solubility [356]), have been described by Makitra et al.

Another important treatment of multiple interacting solvent effects, in principle analogous to Eq. (7-50) but more precisely elaborated and more generally applicable, has been proposed by Kamlet, Abboud, and Taft (KAT) [84a, 224, 226]. Theirs and Koppel and Palm's approaches have much in common, i.e. that it is necessary to consider non-specific and specific solute/solvent interactions separately, and that the latter should be subdivided into solvent Lewis-acidity interactions (HBA solute/HBD solvent) and solvent Lewis-basicity interactions (HBD solute/HBA solvent). Using the solvatochromic solvent parameters s, b, and p^*, which have already been introduced in Section 7.4 (cf. Table 7-4), the multiparameter equation (7-53) has been proposed for use in so-called linear solvation energy relationships (LSER).

$$A = A_0 + s(p^* + dd) + aa + bb \qquad (7\text{-}53)$$

The solute property A can represent, for example, the logarithm of a rate or equilibrium constant, as well as a position of maximal absorption in a UV/Vis, IR, NMR, or ESR spectrum*; A_0 is the regression value of this solute property in cyclohexane as reference solvent.

p^* is an index of solvent dipolarity/polarizability, which measures the ability of the solvent to stabilize a charge or a dipole by virtue of its dielectric effect. For a set of select solvents, i.e. non-HBD aliphatic solvents with a single dominant group dipole, the p^* values are proportional to the dipole moment of the solvent molecule. The p^* scale was selected to run from 0.00 for cyclohexane to 1.00 for dimethyl sulfoxide. The p^* values correspond to the use of relative permittivity and refractive index in the Koppel–Palm equation (7-50) as measures of polarization and polarizability interactions. Therefore, functions of ε_r and n^2 are not included in the KAT equation (7-53). The advantage of using p^* values instead of the functions $f(\varepsilon_r)$ and $f(n^2)$ is that the latter terms are properties of the bulk solvent, whereas the p^* values are derived from electronic transitions occuring on a molecular-microscopic level in solute-organized cybotactic regions, i.e. within the solvation shell of the solute [84a]. For later improvements of the p^* scale, see reference [227].

d is a discontinuous polarizability correction term equal to 0.0 for non-chloro-substituted aliphatic solvents, 0.5 for poly-chloro-substituted aliphatics, and 1.0 for aromatic solvents. The d values reflect the observation that differences in solvent polarizability are significantly greater between the three solvent classes than within the individual classes. Thus, the sign and magnitude of the d term is related to the variable dipolarity/polarizability blend observed in the solvent influence on the solute property A.

a is a measure of the solvent hydrogen-bond donor (HBD) acidity and corresponds to Koppel and Palm's Lewis-acidity parameter E. It describes the ability of a solvent to donate a proton in a solvent-to-solute hydrogen bond. The a-scale was

* In the original papers, the symbols XYZ and $(XYZ)_0$ were used instead of A and A_0 [84a]. For the sake of consistency with the general Eq. (7-49), A and A_0 are used in Eq. (7-53).

selected to extend from zero for non-HBD solvents (e.g. cyclohexane) to about 1.0 for methanol.

b is a measure of the solvent hydrogen-bond acceptor (HBA) basicity and corresponds to Koppel and Palm's Lewis-basicity parameter B. It describes the solvent's ability to accept a proton (or, vice versa, to donate an electron pair) in a solute-to-solvent hydrogen bond. The b-scale was selected to extend from zero for non-HBD solvents (e.g. cyclohexane) to about 1.0 for hexamethylphosphoric acid triamide (HMPT).

The regression coefficients s, d, a, and b in Eq. (7-53) measure the relative susceptibilities of the solvent-dependent solute property A to the indicated solvent parameters. Due to the normalization of the a, b, and p^* scale (from ca. 0.0 to ca. 1.0), the a/s, b/s, and a/b ratios are assumed to provide quantitative measures of the relative contribution of the indicated solvent parameters.

Eq. (7-53) has been used in the correlation analysis by multiple regression of numerous reaction rates and equilibria, spectroscopic data, and various other solvent-dependent processes. An impressive series of 46 articles entitled "Linear Solvation Energy Relationships" (LSER) has been published: Part 1 [229] ... Part 46 [230]; see also the summarizing articles [127, 224, 227] and the critical compilation of solvent parameters (e.g. p^* and b) by Abboud and Notario [295].

Eq. (7-53) was subsequently extended to Eq. (7-54) by the introduction of two further solvent parameters, the d_H^2 term*) and the x term [226]

$$A = A_0 + s(p^* + dd) + aa + bb + hd_H^2 + ex \qquad (7\text{-}54)$$

d_H^2 represents Hildebrand's solubility parameter squared and corresponds to the cohesive pressure c, which characterizes the energy associated with the intermolecular solvent/solvent interactions; cf. Eqs. (3-5) and (5-77) in Sections 3.2 and 5.4.2, respectively, for the definition of c and d. Thus, d_H^2 is considered as a measure of the enthalpy or Gibbs energy input required to separate solvent molecules to provide a suitably sized cavity for the solute. This solvent cavity term is important for multiple correlations dealing with enthalpies or Gibbs energies of solution, Gibbs energies of transfer between two solvents, or gas-liquid chromatographic partition coefficients. In most cases, the hd_H^2 term is only significant for highly structured solvents such as water, formamide, and 1,2-ethanediol.

In addition to the b parameter of solvent Lewis basicity, the coordinate covalency parameter x has been found to be useful in correlating certain types of so-called family-dependent solute basicity properties [226, 267]. Family-independent (FI) basicity properties are defined as those which have a linear relationship with b when all solute bases are considered together. Family-dependent (FD) basicity properties are those which exhibit a linear relationship with b only when different families of solutes having similar HBA sites are considered separately. Thus, FD properties can be correlated to FI properties if an empirical coordinate covalency parameter x is used in correlation equa-

---

* The subscript H is used to distinguish the Hildebrand solubility parameter d from the polarizability correction term d in Eq. (7-54). In earlier formulations of Eq. (7-54), d_H instead of d_H^2 was wrongly used [96, 97, 226, 255]. In linear Gibbs energy relationships, all the terms of the regression equation should have the dimension of an energy, but d_H has the dimension of the square root of the energy ($J^{1/2} \cdot cm^{-3/2}$); cf. Section 7.5 and references [256...258].

tions such as $A = A_0 + bb + ex$. Values of x are equal to -0.2 for P=O bases (e.g. HMPT), 0.0 for C=O and S=O bases (arbitrary reference value), 0.1 for triple-bonded nitrogen bases (e.g. nitriles), 0.2 for single-bonded oxygen bases (e.g. ethers), 0.6 for sp^2-hybridized nitrogen bases (e.g. pyridines), and 1.0 for sp^3-hybridized nitrogen bases (e.g. amines). The x values are interpreted as being approximate measures of the relative coordinate covalencies of the bonds which are formed between solute and solvent at the base centres of the solvent. Coordinate covalencies of solute/solvent adducts of a given solute acid decrease in strength (x decreases) as the electronegativity of the solvent base centre increases, because the positive charge created by coordinate covalency on an increasingly electronegative atom is unfavourable. The x values have been found to be useful in correlating the basicity behaviour of neutral oxygen and nitrogen bases of widely di¤ering properties [267]. It has been stressed that the correlation equation $A = A_0 + bb + ex$ formally and conceptually resembles the Drago E/C treatment, with b and b corresponding to E_A and E_B, and e and x corresponding to C_A and C_B; cf. Eq. (2-12) in Section 2.2.6.

The multiparameter equation (7-54) seems to be rather di‹cult to apply. However, in practice, most of the linear solvation energy relationships that have been reported are simpler than indicated by Eq. (7-54) since one or more terms are inappropriate. For example, if the solute property A does not involve the creation of a cavity or a change in cavity volume between initial and activated or excited states (as is the case for solvent e¤ects on spectral properties), the d_H^2 term is dropped from Eq. (7-54). If the solvent-dependent process under study has been carried out in non-HBD solvents only, the a term drops out. On the other hand, if the solutes are not hydrogen-bond donors or Lewis acids, the b term drops out of Eq. (7-54). Thus, for many solvent-dependent processes, Eq. (7-54) can be reduced to a more manageable one-, two- or three-parameter correlation equation by a judicious choice of solutes and solvents [226].

The KAT equations (7-53) and (7-54) have been widely and successfully applied in the correlation analysis of all kinds of solvent-dependent processes; a review on their use in chromatography has been given by Carr [357]. Only two representative examples will be mentioned. Multiple regression of $\lg k_2$ for the reaction between benzoic acid and diazo-diphenylmethane using the KAT equation (7-53), for the same set of 44 non-HBD solvents as used in the Koppel…Palm treatment of this reaction given in Eq. (7-52), leads to Eq. (7-55a), with $n = 44$, $r = 0.980$, and $s = 0.171$ [15, 268].

$$\lg k_2 = 0.20 + 1.21p^* + 2.71a - 3.70b \qquad (7\text{-}55a)$$

A comparison of Eqs. (7-52) and (7-55a) shows that both treatments give similar results, with opposite signs of the solvent Lewis-acidity and basicity parameters. This is in agreement with the given reaction mechanism, involving simultaneous nucleophilic and electrophilic solvent assistance.

Multiple regression analysis of $\lg k_1$ of the strongly solvent-dependent solvolysis/dehydrohalogenation of 2-chloro-2-methylpropane (t-BuCl) for $n = 21$ solvents using the KAT equation (7-54) leads to Eq. (7-55b), with $r = 0.997$, and $s = 0.242$ [288]:

$$\lg k_1(\text{t-BuCl}) = -14.60 + 5.10 \cdot p^* + 4.17 \cdot a + 0.73 \cdot b + 0.0048 \cdot d_H^2 \qquad (7\text{-}55b)$$

From this correlation equation, it is immediately obvious that the "rst-order rate constant k_1 is increased by dipolar solvents (with large p^* values) and HBD solvents (with large a values). The small but statistically signi"cant $b \cdot b$ term in Eq. (7-55b) is consistent with a weak nucleophilic solvent assistance in the solvolysis of 2-chloro-2-methylpropane. Increasing cohesive pressure (as given by δ_H^2) increases k_1 to a very small extent because of the electrostriction of the substrate solvation shell in going from the educt to the more dipolar activated complex, cf. Eq. (5-13) in Section 5.3.1. The complex solvent in"uence on 2-chloro-2-methylpropane solvolysis as described by Eq. (7-55b) is of relevance for its use as the standard reaction for the determination of Grunwald…Winstein•s Y values as measures of solvent ionizing power, cf. the discussion of this topic and the question of nucleophilic solvent assistance in S_N1 reactions in the "rst part of Section 7.3.

A few comments of the KAT treatment of solvent e¤ects should be made. Some shortcomings and the sometimes insu‹cient precision in the determination of the solvatochromic parameters have been discussed by Laurence et al. [167, 237, 238], who also recommended improvements by virtue of a new thermosolvatochromic comparison method; cf. Section 7.4 for a further discussion of this method. A study of linear Gibbs energy relationships (LGER) in a homologous series of n-alkane and n-alkylnitrile solvents, using Kamlet and Taft•s solvatochromic indicator solutes, has shown that a single-lumped parameter of solvent dipolarity/polarizability such as the p^* values cannot be applied simultaneously to n-alkanes and n-alkylnitriles [269]. Therefore, it has been concluded by Carr et al. [269] that the hypothesis that solvent dipolarity and polarizability can be represented by a single parameter is certainly not generally valid. Kamlet, Abboud, and Taft have tried to consider this observation by incorporating the solvent polarizability correction term $d\delta$ in Eq. (7-53), which has di¤erent values for groups of aliphatic, chlorine-substituted aliphatic, and aromatic solvents,* but the important distinction to be made is that the solute sensitivity to the two types of polarization can be quite di¤erent [269]. In this context, reference has to be made to Sjöström and Wold•s results [236], who recommended, on statistical grounds, the use of a two-parameter equation of the form $A = A_0 + s_0 p_0^* + s_d p_d^*$, where p_0^* and p_d^* represent solvent orientational and distortional polarizability scales, respectively.

Furthermore, there has been considerable discussion on the question as to whether or not linear Gibbs energy relationships (LGER), such as linear solvation energy relationships (LSER), are really fundamental laws of chemistry, re"ecting simple physicochemical relationships, or rather local empirical rules of similarity, i.e. only locally valid linearizations of more complicated relationships [126, 127]; cf. Section 7.1 for a further discussion of this dialogue. Both parties, the chemometricians and the physical organic chemists, have tried to exemplify the merits of the two types of treatment of solvent e¤ects, i.e. principal component analysis (PCA) and linear solvation energy relationships (LSER), by means of the correlation and rationalization of solvent e¤ects on the ^{13}C NMR chemical shifts of lithium indenide, a planar delocalized carbanion,

* The description of di¤erent solvent polarizabilities by means of three discontinuous polarizability correction terms $d\delta$ ($d = 0.0$, 0.5, or 1.0 for three solvent groups) as in Eqs. (7-53) and (7-54) is surely an oversimpli"cation [167].

measured in a set of 13 solvents; see references [270, 271] as well as [126, 127] for this lengthy discussion.

Kamlet, Taft, Abraham et al. have further modified Eq. (7-54) in order to correlate the solubility and distribution behaviour of nonelectrolyte solutes with solvent properties, according to Eq. (7-56) [271, 272],

$$A = A_0 + A'(d_H^2)_1(V_2/100) + Bp_1^*p_2^* + Ca_1b_2 + Db_1a_2 \qquad (7\text{-}56)$$

where subscript 1 refers to the solvent and subscript 2 to the solute, and A′, B, C, and D are the regression coefficients for the endoergic cavity term, the exoergic dipolarity/polarizability term, and the exoergic hydrogen-bonding terms of adduct formation between HBD solvents and HBA solutes (measured by a_1 and b_2) as well as between HBA solvents and HBD solutes (measured by b_1 and a_2), respectively. V_2 is the molar volume of the solute, taken as its molecular mass divided by its liquid density at 20°C. Whereas $(d_H^2)_1$ represents the solvent's contribution to the cavity term, $V_2/100$ represents the solute's contribution to the cavity term. Instead of V_2, $V_2/100$ is used so that the parameter describing the cavity term will roughly cover the same numerical range as the other independent variables a, b, and p^* (ca. 0.0 . . . 1.0). This simplifies the evaluation of the relative contributions of the various terms of Eq. (7-56) to the property A.

When dealing with the effects of different solvents on the properties of a single solute (e.g. solvent effects on reaction rates), the factors relating to the solute can be subsumed into the regression coefficients of Eq. (7-56), and the following equation results:

$$A = A_0 + h(d_H^2)_1 + sp_1^* + aa_1 + bb_1 \qquad (7\text{-}57)$$

In Eq. (7-57), the dependence of the solute property A on each term is now given solely by solvent parameters. Furthermore, when the solute property A does not involve cavity formation or a cavity change, the cavity term drops out, and Eq. (7-57) takes the form of Eq. (7-53) (with $d = 0$).

Conversely, when dealing with solubilities or other properties of a set of different solutes in a single solvent, or with distributions of different solutes between a certain pair of solvents, the resulting Eq. (7-58) relates property A only to the solute parameters V_2, p_2^*, a_2, and b_2, and the solvent parameters are now subsumed into the regression coefficients.

$$A = A_0 + m(V_2/100) + sp_2^* + aa_2 + bb_2 \qquad (7\text{-}58)$$

For example, this equation has been successfully used to correlate the Hansch-Leo 1-octanol/water partition coefficient, $K_{o/w}$, of 102 aliphatic and aromatic solutes according to Eq. (7-59) with the indicated solute properties ($n = 102$, $r = 0.989$, $s = 0.175$) [149]; cf. Eq. (7-12) in Section 7.2 for the definition of $K_{o/w}$ [145…148].

$$\lg K_{o/w} = 2.74(V_2/100) - 0.92p_2^* - 3.49b_2 + 0.20 \qquad (7\text{-}59)$$

Dealing with this type of multiparameter correlation analysis, a series of twelve articles entitled ••Solubility Properties in Polymers and Biological Media•• was published by Kamlet, Taft, Abraham et al. (Part 1 [273]. . . Part 12 [274]), as well as another series entitled ••Solute…Solvent Interactions in Chemical and Biological Systems•• (Part 4 [358]. . . Part 7 [359]). The application of the LSER equation (7-58) to the prediction of solubilities of organic nonelectrolytes in water, blood, and other body tissues has been reviewed [286].

The KAT equation (7-58) for the correlation analysis of solute properties (e.g. solubility, partitioning, gas-chromatographic behaviour, etc.) has been further developed and re“ned by Abraham [360]. Abraham•s multiparameter LSER equation consists of “ve solute descriptors, speci“ed as follows: a combined dipolarity/polarizability descriptor (p_2^H), two descriptors for the e¤ective HBD acidity (a_2^H) and HBA basicity (b_2^H), the excess molar refraction (R_2), and the solute gas/n-hexadecane partition coef“cient $(\lg L)$ or McGowan•s characteristic volume (V_x). This multiparameter equation describes the properties of a series of solutes in a given solvent. That is, the solvent properties remain constant and only the solute is varied. Because solvent parameters are not needed, a more detailed consideration of this widely applied and highly successful approach is beyond the scope of this book. For some leading references, see [361] (correlation of gas-liquid chromatographic data), [362] (water/1-octanol partitioning, brain penetration, skin permeability), and [363] (solubility of C_{60} fullerene).

In formal analogy to the KAT equation (7-53), Catalán has combined his solvatochromically derived solvent parameters SPP, SA, and SB in the three-parameter equation (7-60) [296, 335…337]:

$$A = A_0 + a \cdot SPP + b \cdot SA + c \cdot SB \qquad (7\text{-}60)$$

where SPP corresponds to the solvent dipolarity/polarizability, SA to the solvent HBD acidity, and SB to the solvent HBA basicity; for their precise de“nition, see Eqs. (7-40a), (7-40b), and (7-40e) as well as Table 7-5 in Section 7.4. This equation has been used to investigate some other reported solvent polarity scales [296]. For example, it has been shown that the $E_T(30)$ scale correlates well with Eq. (7-60), which gives some insight into the various solute/solvent interactions contributing to this empirical solvent polarity scale; see Eqs. (7-47a,b) in Section 7.6.

Multiple regression analysis of $\lg k_1$ for the solvent-dependent solvolysis/dehydrohalogenation of 2-chloro-2-methylpropane (t-BuCl) by means of Eq. (7-60) leads, for $n = 148$ solvents and binary solvent mixtures to Eq. (7-61), with $r = 0.990$ and $s = 0.40$ [341]:

$$\lg k_1(\text{t-BuCl}) = -20.07 + 10.62 \cdot SPP + 7.89 \cdot SA + 1.71 \cdot SB \qquad (7\text{-}61)$$

In analogy to Eq. (7-55b), all three individual solvent parameters contribute to the overall solvent e¤ect, increasing the “rst-order rate constant k_1 with increasing SPP, SA, and SB values. Again, a small but signi“cant accelerating e¤ect results from the solvent HBA basicity term $c \cdot SB$, in agreement with a weak nucleophilically assisted S_N1 reaction; cf. also Eq. (7-55b) and the discussion of the mechanism of this S_N1 reaction in connection with the Y values of Grunwald and Winstein.

A more simpli“ed but likewise successful empirical two-parameter approach for the description of solvent e¤ects has been proposed by Krygowski and Fawcett [113]. They assume that only speci“c solute/solvent interactions need to be considered. These authors postulated that the solvent e¤ect on a solute property A can be represented as a linear function of only two independent but complementary parameters describing the Lewis acidity and Lewis basicity of a given solvent. Again, for reasons already mentioned, the $E_T(30)$ values were chosen as a measure of Lewis acidity. In addition, Gutmann•s donor numbers DN [26, 27] were chosen as a measure of solvent basicity (cf. Table 2-3 and Eq. (7-10) in Sections 2.2.6 and 7.2, respectively). Thus, it is assumed that the solvent e¤ect on A can be described in terms of Eq. (7-62)*

$$A = A_0 + a \cdot E_T(30) + b \cdot DN \qquad (7\text{-}62)$$

a and b are regression coe‹cients describing the sensitivity of the solute property A to electrophilic and nucleophilic solvent properties, respectively. Since a and b are not necessarily on the same scale, due to the fact that $E_T(30)$ and DN do not vary over the same range for a given set of data, the regression coe‹cients were normalized to give $\bar{a}$ and $\bar{b}$. Application of Eq. (7-62) involves the supposition that non-speci“c solute/solvent interactions are negligible or nearly constant and can be included in the solvent Lewis acidity and basicity terms. Obviously, this is a serious simpli“cation. However, in spite of this simpli“cation, the Krygowski…Fawcett treatment of solvent e¤ects has been successfully applied in many cases [113]. Satisfactory correlations were obtained in 90% of the cases involving ion/solvent and ion/ion interactions, and in 75% of those involving dipole/dipole interactions [113]; see references [118…120, 232] for applications of Eq. (7-62) and its modi“cations.

A further interesting two-parameter treatment of solvent e¤ects has been given by Swain et al. [265]. It is based on a computer calculation involving 1080 data sets for 61 solvents and 77 solvent-sensitive reactions and physicochemical properties, taken from the literature (e.g. rate constants, product ratios, equilibrium constants, UV/Vis, IR, ESR, and NMR spectra). According to these calculations, all solvent e¤ects can be rationalized in terms of two complementary solvent property scales, i.e., A_j, measuring the solvent•s anion-solvating tendency or acity**), and B_j, measuring the solvent•s cation-solvating tendency or basity**), which are combined in Eq. (7-63)*** [265].

---

* In the original paper, Q and Q_0 were used instead of A and A_0 [113]. For the sake of consistency with the general Eq. (7-49), A and A_0 are used in Eq. (7-62).

** The new names acity and basity were chosen because, although they are obviously kinds of acidity and basicity, they are neat (bulk) solvent properties involved in solute solvations. Such solvent properties cause speci“c local electrostatic solute/solvent interactions without major covalency changes and, therefore, are usually omitted from equations describing chemical reactions [265].

*** In the original publication, Eq. (7-63) takes the form

$$p_{ij} = c_i + a_i \cdot A_j + b_i \cdot B_j$$

with p_{ij} representing the solvent-dependent solute property i in solvent j, and c_i representing the predicted value for a reference solvent for which $A_j = B_j = 0$ [265]. For the sake of consistency with the general Eq. (7-49), A and A_0 instead of p_{ij} and c_i are used in Eq. (7-63), in spite of the unfortunate cumulation of the letter A.

$$A = A_0 + a_i \cdot A_j + b_i \cdot B_j \qquad (7\text{-}63)$$

A_j and B_j characterize the solvent j. A and A_0 as well as the multiple regression coefficients a_i and b_i depend only on the solvent-sensitive solute property i under study. Constants a_i and b_i represent the sensitivity of solute property i to a solvent change. A nonlinear least-squares procedure, using equal statistical weighting of the 1080 data, has been used to evaluate and to optimize all 353 $[= (2 \cdot 61) + (3 \cdot 77)]$ constants A_j, B_j, a_i, b_i, and A_0 in order to get the best possible "fit consistent with Eq. (7-63). In order to obtain values of A_j and B_j which represent physically significant solvent influences that are cleanly separated, some scale-setting subsidiary conditions have to be fixed. As trivial, arbitrary conditions that set zeros and scale factors but do not affect rank orders, $A_j = B_j = 0.00$ for n-heptane and $A_j = B_j = 1.00$ for water have been chosen. As two non-trivial, critical conditions, $A_j = 0.00$ for hexamethylphosphoric acid triamide (HMPT) and $B_j = 0.00$ for trifluoroacetic acid (TFA), have been chosen. This choice is equivalent to the assumption that HMPT is almost as poor an anion solvator, and TFA is almost as poor a cation solvator, as n-heptane.

Table 7-6 lists a selection of A_j and B_j values in order of their sum $(A_j + B_j)$, which may be considered as a measure of "solvent polarity" in terms of the overall solvation capability of a solvent; cf. Section 7.1 and references [1, 3] for this definition of solvent polarity. A plot of A_j against B_j shows that both parameters are highly variable, but there is no correlation between A_j and B_j, an essential condition for the application of Eq. (7-63). Interestingly, the overall correlation coefficient between the 1080 input data and the predictions made by Eq. (7-63) is excellent, at $r = 0.991$. That means that the two unrelated solvent parameters A_j and B_j alone account for over 98% of the solvent effects in the set of 77 solvent-sensitive processes that have been examined by Swain et al. [265]. Since, for the 1080 diverse input data, the solvent properties can be adequately represented by only two parameters A_j and B_j, there must be a correlation among the three neat solvent properties, i.e. anion-solvating tendency, hydrogen-bonding acidity (HBD acidity), and electrophilicity, and likewise cation-solvating tendency, hydrogen-bonding basicity (HBA basicity), and nucleophilicity, respectively.

The coefficients a_i and b_i, calculated for the 77 solvent-sensitive processes used to establish Eq. (7-63) as well as for 11 further solvent-dependent reactions (including another 75 data), have been discussed in detail in reference [265]. For example, $a_i = 1.87$ and $b_i = -0.05$ for the $n \rightarrow p^*$ absorption of benzophenone. With the high value of a_i and the negligible value of b_i, this solvent-dependent UV/Vis absorption comes close to measuring A_j in pure form. The $E_T(30)$ values (cf. Table 7-3) are characterized by $a_i = 30.36$ and $b_i = 4.45$ $(a_i/b_i = 6.8)$, in agreement with other observations showing that the $E_T(30)$ values are mainly related to the solvent Lewis acidity, and not to the solvent Lewis basicity. Application of Eq. (7-63) to several hundred increasingly diverse additional solvent-dependent reactions has led to satisfactory correlations (with $r > 0.965$); four explanations for non-agreement have been found and discussed by Swain et al. [265].

In view of the success in correlating so many solvent effects by only two solvent parameters according to Eq. (7-63), Swain et al. concluded that not all four solvent parameters of the KAT Eq. (7-53) are necessary; additional parameters do not improve the "ts already observed with Eq. (7-63), and, for example, the t parameter of Eq. (7-53)

Table 7-6. Selection of values of solvent acity A_j, and solvent basity, B_j, for 34 solvents, calculated according to Eq. (7-63) [265].

Solvents j	A_j	B_j	$(A_j + B_j)$[a]
n-Heptane	0.00	0.00	0.00
Cyclohexane	0.02	0.06	0.09
Triethylamine	0.08	0.19	0.27
Tetrachloromethane	0.09	0.34	0.43
Diethyl ether	0.12	0.34	0.46
Carbon disul“de	0.10	0.38	0.48
Toluene	0.13	0.54	0.67
Benzene	0.15	0.59	0.73
Ethyl acetate	0.21	0.59	0.79
Tetrahydrofuran	0.17	0.67	0.84
Chlorobenzene	0.20	0.65	0.85
1,4-Dioxane	0.19	0.67	0.86
t-Butanol	0.45	0.50	0.95
Methoxybenzene	0.21	0.74	0.96
2-Butanone	0.23	0.74	0.97
2-Propanol	0.59	0.44	1.03
Acetone	0.25	0.81	1.06
Acetic acid	0.93	0.13	1.06
Hexamethylphosphoric acid triamide	0.00	1.07	1.07
1-Propanol	0.63	0.44	1.08
Ethanol	0.66	0.45	1.11
1,2-Dichloroethane	0.30	0.82	1.12
Dichloromethane	0.33	0.80	1.13
Trichloromethane	0.42	0.73	1.15
Pyridine	0.24	0.96	1.20
Acetonitrile	0.37	0.86	1.22
N,N-Dimethylformamide	0.30	0.93	1.23
Methanol	0.75	0.50	1.25
Nitromethane	0.39	0.92	1.31
Dimethyl sulfoxide	0.34	1.08	1.41
Ethane-1,2-diol	0.78	0.84	1.62
Formamide	0.66	0.99	1.65
Tri”uoroacetic acid	1.72	0.00	1.72
Water	1.00	1.00	2.00

[a] The solvents are listed in order of their sum $(A_j + B_j)$, which is considered as reasonable measure of ••solvent polarity•• in terms of the overall solvation capability of a solvent [265].

is rendered super”uous [265]. These conclusions, as well as Swain et al.•s general approach, have been criticized by Taft, Abboud, and Kamlet [275], leading to a reply by Swain [276] dismissing this criticism. The reader is referred to references [275, 276] and to the June issue of the Journal of Organic Chemistry in 1984 for this interesting discussion (issue no. 11, p. 1989…2010).

Both Kamlet, Abboud, and Taft et al.•s [224, 226] and Swain et al.•s [265] multiparameter solvent e¤ect treatments have an inherent weakness in so far as the solvent parameters a, b, and p^* as well as A_j and B_j are averaged and statistically optimized parameters: the former are derived from various types of solvatochromic indicator dyes,

while the latter are calculated from a selection of 77 solvent-sensitive processes. Thus, they are no longer directly related to a distinct, carefully selected, well-understood single reference process [as, for example, Y_{OTs}, Z, and $E_T(30)$]; cf. footnote on page 379. The signi“cance of such averaged and statistically optimized solvent parameters depends above all on (i) the right choice of the various solvent-dependent processes used in the averaging procedure, which has to be done with a skillful hand, and (ii) the right choice of the critical subsidiary conditions, which have to be de“ned and justi“ed in order to get solvent parameters with a clear physical meaning at the molecular-microscopic level*. For example, concerning point (i), according to Taft, Abboud, and Kamlet [275], most of the 77 solvent-dependent processes selected by Swain et al. [265] involve only non-HBD solutes (reactants or indicator dyes). Therefore, Swain et al.•s “nding that Kamlet, Abboud, and Taft•s b parameter of solvent HBA basicity is super-”uous may have resulted from a somewhat unfortunate data selection. As far as point (ii) is concerned, Swain criticizes that in three-parameter equations such as (7-53) the six critical subsidiary conditions needed to assure that the derived solvent parameters are physically signi“cant are often not properly de“ned. It has been stressed that optimization of the correlations through adjustment of the solvent parameters is not enough and has nothing to do with the physical signi“cance of these calculated solvent parameters [276].

Another semiempirical multiparameter relationship for the description of solvent e¤ects on the statics and kinetics of chemical reactions, according to Eqs. (7-64) and (7-65), has been introduced by Mayer [266].

$$\Delta G^S - \Delta G^R = a \cdot (DN^S - DN^R) + b \cdot (AN^S - AN^R) + c \cdot (\Delta G_{vp}^{0S} - \Delta G_{vp}^{0R}) \quad (7\text{-}64)$$

$$\Delta\Delta G = a \cdot \Delta DN + b \cdot \Delta AN + c \cdot \Delta\Delta G_{vp}^0 \quad (7\text{-}65)$$

ΔG represents the Gibbs energy of reaction or activation ($\Delta G^{\neq}$), DN the donor number [26, 27], AN the acceptor number [91], and ΔG_{vp}^0 the standard molar Gibbs energy of vapourization of a solvent S and a reference solvent R, respectively. Acetonitrile ($\Delta G_{vp}^0 = 5.31$ kJ/mol) has been used as a reference solvent [266]. The coe‹cients a and b are correlated to the donor and acceptor strengths of the reaction partners relative to those of the reference compounds $SbCl_5$ and $(C_2H_5)_3PO$, respectively. This approach is based on a model developed for calculating the Gibbs energy change associated with the creation of cavities in the solvent, the size of which corresponds to the volume occupied by the solute molecules. Experimental equilibrium and rate data, including solubility measurements, complex-formation equilibria, ion-pair equilibria, and an S_NAr reaction,

* For a two-parameter treatment of solvent e¤ects (with two independent solvent vectors), only two critical subsidiary conditions must be de“ned in order to force the two solvent parameters to represent physically signi“cant solvent properties. Four other trivial arbitrary conditions have to be de“ned in order to “x zero reference points and scale-unit sizes. However, for a three-parameter treatment (with three independent solvent vectors), already six critical subsidiary conditions must be de“ned, in addition to the six trivial reference or scale-factor conditions. On the contrary, single-parameter treatments require no de“nition of critical subsidiary conditions, but only one reference (zero) condition and one standard (unit) condition, whose arbitrary assignment changes only the reference solvent and the scale-unit size [265, 276].

have been successfully used to test Eqs. (7-64) and (7-65) by the method of multiple linear regression analysis [266]. In a remarkable application of Eq. (7-65) (with $d = 0$), it has been shown that for non-HBD solvents the logarithm of their relative permittivities can be represented by a linear combination of the acceptor number and the donor number [134, 207]; cf. also [139].

Finally, a multiparameter correlation equation based solely on theoretically determined solvent descriptors, introduced by Famini and Wilson, deserves attention [350]. Linear solvation energy relationships (LSERs), such as the KAT equation (7-54) and its successors, can be summarized by the general form shown in Eq. (7-66):

$$A = A_0 + \text{dipolarity/polarizability term(s)} + \text{hydrogen-bonding term(s)} + \text{bulk/cavity term} \qquad (7\text{-}66)$$

where A and A_0 are the values of the solvent-dependent physicochemical property in a given solvent and in the gas phase or in an inert solvent, respectively; cf. also Eq. (7-49). As mentioned before, the dipolarity/polarizability terms are measures of solute/solvent dipole and induced dipole interactions (exoergic). The hydrogen-bonding terms give a measure of the energy of interaction (exoergic) when hydrogen-bonded HBA solute/HBD solvent and/or HBD solute/HBA solvent complexes are formed. The bulk or cavity term is a measure of the energy needed to overcome the cohesive solvent/solvent interactions (endoergic) to form cavities for the solute molecules. The regression coefficients of the LSER equation provide chemical insight into the specific molecular properties or solute/solvent interactions that impact most strongly on the solvent-dependent process under study. Usually, not all terms of an LSER equation are statistically significant.

The solvent descriptors in LSER equations mentioned so far have been experimentally obtained using carefully selected solvent-dependent reference processes, in most cases the UV/Vis absorption of solvatochromic indicator dyes. In addition to these experimentally based LSER equations, a new theoretical set of solvent parameters has been developed by Famini and Wilson [350]. Termed the theoretical linear solvation energy (TLSER) descriptors, these parameters are determined solely from computational methods permitting an a priori prediction of solvent properties. The TLSER descriptors were developed so as to give TLSER equations with correlation coefficients and standard deviations close in value to those for normal LSERs, and to be as widely applicable to solvent-dependent processes as the normal LSER set. A set of six TLSER descriptors has been developed by Famini and Wilson and the resulting generalized TLSER equation for solutes in a given solvent is shown in Eq. (7-67) [350]:

$$A = A_0 + a \cdot V_{mc} + b \cdot p_I + c \cdot e_a + d \cdot e_b + e \cdot q_- + f \cdot q_+ \qquad (7\text{-}67)$$

In this equation, V_{mc} represents the molecular van der Waals volume (in $Å^3$) and indicates the cavity size needed in the solvent matrix to accommodate the solute. The polarizability index, p_I, is obtained by dividing the polarizability volume by the molecular volume, to give a alternatively size-independent quantity that indicates the ease with which electrons may move throughout the solvent molecule. The solvent HBD

acidity and HBA basicity are modeled by covalent terms e_a and e_b, and electrostatic terms q_- and q_+. The e_a and e_b terms are derived from linear functions of the highest occupied molecular orbital (HOMO) energy and the lowest unoccupied molecular orbital (LUMO) energy, respectively. The covalent HBD acidity term, e_a, is de“ned as a linear transformation of the energy di¤erence between E_{HOMO} of water and E_{LUMO} of the solute, according to $0.30 - [0.01|(E_{LUMO\ (solute)} - E_{HOMO\ (water)})|]$. This transformation provides a •zero-point• reference for the scale and corrects the scale to be positive with increasing acidity. The covalent HBA basicity term, e_b, is likewise de“ned, but uses the LUMO energy of water and the HOMO energy of the solute, according to $0.30 - [0.01|(E_{LUMO\ (water)} - E_{HOMO\ (solute)})|]$. The water energies have been included for •aesthetic reasons• [350c]: the smaller these di¤erences are, the greater is the ability to form a hydrogen bond with water. The electrostatic contribution to the HBA basicity, q_-, is the magnitude of the largest negative formal charge on a nonhydrogen atom of the solvent molecule. The electrostatic contribution to the HBD acidity q_+, is de“ned as the magnitude of the charge on the most positive hydrogen atom in the solvent molecule. A table of TLSER descriptors for a selection of 30 solvents can be found in reference [351].

The TLSER methodology has been successfully applied to develop correlation equations for a wide variety of solvent-dependent properties and processes [350, 364…369]. Some examples are the characterization of other solvent polarity, acidity, and basicity scales [364], the acidities of substituted acetic acids in various solvents [365], the basicities of substituted dimethylamines in various solvents [366], the decarboxylation kinetics of 3-carboxybenzisoxazole [367], the C=O stretching frequencies of substituted pyrrolidin-2-ones [368], and gas…water distribution coe‹cients [369].

Of particular interest is the analysis of other, experimentally determined empirical solvent polarity scales by means of the theoretically derived TLSER equation (7-67). For example, correlation of the solvatochromic E_T^N scale [see Eq. (7-29) in Section 7.4] with the TLSER descriptors of Eq. (7-67) leads to Eq. (7-68) for $n = 29$ solvents, with $r = 0.937$ and $s = 0.075$ [351]:

$$E_T^N = 0.8585 - 0.3662 \cdot V_{mc} - 4.0 \cdot p_I + 0.3202 \cdot q_- + 1.5164 \cdot q_+ \qquad (7\text{-}68)$$

According to this equation, the E_T^N values include both the molecular volume and the polarizability index, which is indicative of solute/solvent dispersion interactions. The negative sign of a and b means that increasing dispersion interactions decrease the E_T^N values, in agreement with the red shift usually observed in UV/Vis spectra with increasing dispersion interactions. The covalent acidity and basicity terms drop out ($c = d = 0$). However, E_T^N has signi“cant contributions from both the electrostatic acidity and basicity terms q_+ and q_-, which is indicative of dipolar interactions and a permanent dipolarity contribution. That the electrostatic acidity term q_+ is clearly more signi“cant than q_- further indicates that there are additional acidity factors being included in this empirical solvent polarity scale. This is in agreement with the molecular structure of the solvatochromic indicator dye (see betaine dye (44) in Section 7.4) and other observations that have been already mentioned.

A more detailed discussion of this E_T^N correlation and the TLSER correlations of other experimentally derived solvent scales can be found in reference [351]. Obviously,

the computationally based TLSER parameters can be used in the same way as the experimentally based LSER parameters. The advantage of the TLSER descriptors is that they are obtained from computation and not from experiment. This permits the prediction of properties and correlations for compounds for which the solvatochromic or other solvent parameters are not readily available.

Finally, some special multiparameter correlations of solvent e¤ects will be mentioned. A common multiple regression equation with seven “tting constants has been derived for the relationship between the Gibbs energy of transfer of ions from water to polar solvents, $\Delta G_t^\circ(X, W \rightarrow S)$, and the properties of solvents and ions. Almost 200 data points can be described in terms of four solvent properties (SN, $E_T(30)$, α_1, and δ_H^2) and three ionic solute properties (charge, size, and softness) [277]. The substituent and solvent in”uence on the cationic polymerization of 4-substituted styrenes can be described by four-parameter correlation equations involving terms for the substituent and solvent in”uence as well as the initiator activity [278, 279]. Such correlations are especially useful for controlling industrial polymerization processes as well as speci“c product formation [278, 279]. Modi“ed multiparameter equations involving solvent viscosity parameters have been used to correlate the solvent in”uence on quantum yields of radiative electronic deactivation processes [280]. The dependence of the optical resolution of phenylglycine derivatives with L-(+)-tartaric acid on racemate structure and solvent polarity has been successfully described with multiparameter correlation equations [281]. The solubility of fullerene C_{60} in 113 organic solvents has been successfully correlated with a four-parameter equation using the LSER approach [370]. Increasing molar volume and solvent polarity [as measured by $E_T(30)$] diminished the solubility of C_{60}, whereas solvent polarizability and EPD ability enhanced its solubility. Using an analogous LSER approach, the rate constants for a Menschutkin reaction ($Et_3N + EtI$) and solvolyses of several tertiary haloalkanes have been successfully correlated by multiparameter equations; see reference [371] for details.

Many di¤erent solvent parameters and multiparameter equations have been introduced in this Chapter 7. Certainly, only a few of them will survive the test of applicability and acceptance by organic chemists. Indeed, the preference for certain time-tested solvent scales and multiparameter treatments is already clearly discernible. Amongst the one-parameter scales, the $E_T(30)$ or E_T^N scale and the DN scale have frequently been used, while the Kamlet…Abboud…Taft (KAT) LSER approach seems to be the most widely applied multiparameter approach.

The multiparameter treatment of solvent e¤ects can be criticized from at least three complementary points of view. First, the separation of solvent e¤ects into various additive contributions is somewhat arbitrary, since di¤erent solute/solvent interaction mechanisms can cooperate in a non-independent way. Second, the choice of the best parameter for every type of solute/solvent interaction is critical because of the complexity of the corresponding empirical solvent parameters, and because of their susceptibility to more than one of the multiple facets of solvent polarity. Third, in order to establish a multiparameter regression equation in a statistically perfect way, so many experimental data points are usually necessary that there is often no room left for the prediction of solvent e¤ects by extrapolation or interpolation. This helps to get a sound interpretation of the observed solvent e¤ect for the process under study, but simultaneously it limits the value of such multiparameter equations for the chemist in its daily laboratory work.

In this context, one should be aware of the important remark made by the Austrian philosopher Karl Popper in his autobiography [284]: ••It is always undesirable to make an e¤ort to increase precision for its own sake . . . since this usually leads to loss of clarity. . . . One should never try to be more precise than the situation demands. . . . Every increase in clarity is of intellectual value in itself; an increase in precision or exactness has only a pragmatic value as a means to some de“nite end.••

Appendix

A. Properties, Purification, and Use of Organic Solvents

A.1 Physical Properties

A selection of one-hundred useful organic solvents are listed according to decreasing polarity in Table A-1. Also given are their physical constants, *viz.* melting point, boiling point, relative permittivity, dipole moment, and index of refraction. The measure of polarity used is the empirical solvent parameter E_T^N, derived from the solvatochromism of a pyridinium *N*-phenolate betaine (*cf.* Section 7.4). Further physical data, including those of technically useful solvents, are found in references [1–8, 97–100]. Particularly useful collections of all kinds of solvent properties, relevant for a great variety of solvent applications, can be found in the handbooks of Marcus [8] and Wypych [99].

In Table A-2, a selection of chiral organic solvents is given, together with some physical data. Such chiral solvents have received much attention because of their use in determining optical purities [9, 134], as media for stereoselective syntheses [10–12, 135, 136], and as NMR shift reagents [13].

Another important property of organic solvents is their miscibility with other organic liquids (*cf.* Fig. 2-2 in Chapter 2). The farther two solvents are located from each other in Hecker's "mixotropic" series of solvents [51], given in Table A-10, the less miscible they are.

According to Hildebrand's solubility parameter approach [101], two liquids are miscible if their solubility parameters δ differ by no more than 3.4 units [101, 102]; *cf.* Eqs. (2-1) and (5-77) for the definition of δ. That is, mutual miscibility decreases as the δ values of two solvents become farther apart. Higher mutual solubility will follow if the δ values of the solvents are closer. A comprehensive collection of δ values has been given by Barton [100].

An alternative, more empirical but more accurate method for predicting miscibility has been given by Godfrey [103], using so-called *miscibility numbers* (*M*-numbers). These are serial numbers of 31 classes of organic solvents, ordered empirically by means of their lipophilicity (*i.e.* their affinity for oil-like substances), using simple test-tube miscibility experiments. All pairs of solvents whose *M*-numbers differ by 15 units or less are miscible in all proportions at 25 °C; a difference of $\geq$17 corresponds to immiscibility, and an *M*-number difference of 16 units indicates borderline behaviour (limited mutual miscibility) [103]. The central class of solvents with *M*-number equal to 16 (*e.g.* 2-*n*-butoxyethanol) comprises "universal" solvents, which are miscible with less lipophilic as well as with more lipophilic solvents.

A.2 Purification of Organic Solvents

Normally it is necessary to purify a solvent before use. Naturally, the purity that can be achieved depends on the nature of the impurities [14, 15] and the desired purity is

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

Table A-1. Compilation of one-hundred important organic solvents together with their physical constants[a], arranged in order of decreasing E_T^N value, as empirical parameter of solvent polarity[b].

Solvents	t_{mp}/°C[c]	t_{bp}/°C[d]	ε_r[e]	$\mu \cdot 10^{30}$/Cm[f]	n_D^{20}[g]	E_T^N[h]
(1) Water	0.0	100.0	78.36	6.2	1.3330	1.000
(2) 1,2-Ethanediol[i]	−12.6	197.5	37.70	7.7	1.4318	0.790
(3) Formamide	2.55	210.5	109.50	11.2	1.4475	0.775
(4) Methanol[k]	−97.7	64.5	32.66	9.6	1.3284	0.762
(5) *N*-Methylformamide	−3.8	200	182.40	12.9	1.4319	0.722
(6) Diethylene glycol	−7.8	245.7	31.69 (20 °C)	7.7	1.4475	0.713
(7) Triethylene glycol	−4.3	288.0	23.69 (20 °C)	10.0	1.4558	0.682
(8) Tetraethylene glycol	−6.2	327.3	19.7	10.8	1.4577	0.664
(9) 2-Methoxyethanol	−85.1	124.6	16.93	6.8	1.4021	0.657
(10) *N*-Methylacetamide[l]	30.6	206.7	191.3 (32 °C)	12.8	1.4253 (35 °C)	0.657
(11) Ethanol[k]	−114.5	78.3	24.55	5.5	1.3614	0.654
(12) 2-Aminoethanol	10.5	170.9	37.72	7.6	1.4545	0.651
(13) Acetic acid	16.7	117.9	6.17 (20 °C)	5.6	1.3719	0.648
(14) 1-Propanol[m]	−126.2	97.2	20.45	5.5	1.3856	0.617
(15) Benzyl alcohol	−15.3	205.4	12.7	5.5	1.5404	0.608
(16) 1-Butanol	−88.6	117.7	17.51	5.8	1.3993	0.586
(17) 1-Pentanol	−78.2	138.0	13.9	5.7	1.4100	0.568
(18) 3-Methyl-1-butanol	−117.2	130.5	15.19	6.1	1.4072	0.565
(19) 2-Methyl-1-propanol	−108.2	107.9	17.93	6.0	1.3959	0.552
(20) 2-Propanol[m]	−88.0	82.2	19.92	5.5	1.3772	0.546
(21) Cyclohexanol	25.15	161.1	15.0	6.2	1.4648 (25 °C)	0.509
(22) 2-Butanol	−114.7	99.5	16.56	5.5	1.3971	0.506
(23) 2-Pentanol	−50	119.0	13.71	5.5	1.4064	0.488
(24) Nitromethane[n]	−28.55	101.2	35.87	12.0	1.3819	0.481
(25) Propylene carbonate[l]	−54.5	241.7	64.92	16.5	1.4215	0.472
(26) 3-Pentanol	−8	115.3	13.35	5.5	1.4104	0.463
(27) Acetonitrile[l]	−43.8	81.6	35.94	13.0	1.3441	0.460
(28) Dimethyl sulfoxide[l]	18.5	189.0	46.45	13.5	1.4793	0.444
(29) Aniline	−6.0	184.4	6.98	5.0	1.5863	0.420
(30) Sulfolane[l]	28.45	287.3	43.3 (30 °C)	16.0	1.4816 (30 °C)	0.410
(31) Acetic anhydride	−73.1	140.0	20.63	9.4	1.3904	0.407
(32) Propanenitrile	−92.8	97.3	28.26	13.4	1.3658	0.398
(33) 2-Methyl-2-propanol[m]	25.6	82.3	12.47	5.5	1.3877	0.389
(34) *N*,*N*-Dimethylformamide (DMF)[l]	−60.4	153.1	36.71	12.7	1.4305	0.386

Table A-1. (Continued)

Solvents	t_{mp}/°C[c]	t_{bp}/°C[d]	ε_r[e]	$\mu \cdot 10^{30}$/Cm[f]	$n_D{}^{20}$[g]	E_T^N[h]
(35) *N*,*N*-Dimethylacetamide (DMAC)	−20.1	166.1	37.78	12.4	1.4384	0.377
(36) 1,3-Dimethylimidazolidin-2-one, (DMEU)[o]	8.2	225.5	37.60	13.6	1.4707 (25 °C)	0.364
(37) 1-Methylpyrrolidin-2-one[p]	−24.4	202	32.2	13.6	1.4700	0.355
(38) Acetone[n]	−94.7	56.1	20.56	9.0	1.3587	0.355
(39) 1,3-Dimethyl-2-oxohexahydropyrimidine (DMPU)[o]	−23.8	246.5	36.12	14.1	1.4881 (25 °C)	0.352
(40) 1,2-Diaminoethane[l]	11.3	116.9	12.9	6.3	1.4568	0.349
(41) Cyanobenzene[q]	−12.7	191.1	25.20	13.9	1.5282	0.333
(42) 1,2-Dichloroethane[q]	−35.7	83.5	10.36	6.1	1.4448	0.327
(43) 2-Butanone	−86.7	79.6	18.11	9.2	1.3788	0.327
(44) Nitrobenzene	5.8	210.9	34.78	14.0	1.5562	0.324
(45) 2-Pentanone	−76.9	102.3	15.38 (20 °C)	9.0	1.3908	0.321
(46) 2-Methyl-2-butanol	−8.8	102.0	5.78	5.7	1.4050	0.318
(47) Morpholine	−4.8	129.0	7.42	5.2	1.4542	0.318
(48) Tetramethylurea[o]	−1.2	175.3	23.60	11.6	1.4493 (25 °C)	0.315
(49) Hexamethylphosphoric triamide (HMPT)[l]	7.3	233	29.30	18.5	1.4588	0.315
(50) 3-Methyl-2-butanone	−92.2	94.9	15.87 (30 °C)	9.2	1.3880	0.315
(51) Dichloromethane[q]	−94.9	39.6	8.93	3.8	1.4242	0.309
(52) Acetophenone	19.7	202.0	17.39	9.8	1.5342	0.306
(53) Pyridine[l]	−41.5	115.3	12.91	7.9	1.5102	0.302
(54) Cyclohexanone	−32.1	155.7	15.50	10.3	1.4510	0.281
(55) 4-Methyl-2-pentanone	−84.2	117.5	13.11 (20 °C)	9.0	1.3958	0.269
(56) 1,1-Dichloroethane[q]	−97.0	57.4	10.0 (18 °C)	6.1	1.4164	0.269
(57) Quinoline	−14.9	237.2	8.95	7.3	1.6273	0.269
(58) 3-Pentanone	−39.0	102.0	17.00 (20 °C)	9.4	1.3923	0.265
(59) Trichloromethane	−63.6	61.2	4.89	3.8	1.4459	0.259
(60) 3,3-Dimethyl-2-butanone	−52.5	105.9	12.60	9.2	1.3952	0.256
(61) Methyl acetate	−98.1	56.9	6.68	5.6	1.3614	0.253
(62) Triethylene glycol dimethyl ether	−45	216	7.6	7.4	1.4224	0.253
(63) 2,4-Dimethyl-3-pentanone	−69.0	125.3	17.2 (20 °C)	9.1	1.3999	0.247
(64) Diethylene glycol dimethyl ether	−64.0	162.0	5.8	6.6	1.4078	0.244
(65) 1,2-Dimethoxyethane[r]	−69.2	84.6	7.20	5.7	1.3796	0.231
(66) Ethyl acetate	−83.6	77.2	6.02	5.9	1.3724	0.228
(67) 1,2-Dichlorobenzene	−17.1	180.5	9.93	8.3	1.5515	0.225
(68) 2,6-Dimethyl-4-heptanone	−46.0	168.2	9.91 (20 °C)	8.9	1.4122	0.225
(69) Diethylene glycol diethyl ether	−44.3	188.9	5.70	6.6	1.4115	0.210

Table A-1. (Continued)

Solvents	t_{mp}/°C[c]	t_{bp}/°C[d]	ε_r[e]	$\mu \cdot 10^{30}$/Cm[f]	n_D^{20}[g]	E_T^N[h]
(70) Tetrahydrofuran[s]	−108.4	66.0	7.58	5.8	1.4072	0.207
(71) Methoxybenzene	−37.5	153.7	4.33	4.2	1.5170	0.198
(72) Fluorobenzene	−42.2	84.8	5.42	4.9	1.4684 (15 °C)	0.194
(73) 1,1-Dichloroethene	−122.6	31.6	4.82 (20 °C)	4.3	1.4247	0.194
(74) Chlorobenzene	−45.6	131.7	5.62	5.6	1.5248	0.188
(75) Diethyl carbonate	−43.0	126.9	2.82 (20 °C)	3.0	1.3837	0.185
(76) Bromobenzene	−30.9	156.0	5.40	5.2	1.5568	0.182
(77) Ethoxybenzene	−29.6	169.9	4.22 (20 °C)	4.5	1.5074	0.182
(78) Iodobenzene	−31.4	188.4	4.49 (20 °C)	4.7	1.6200	0.170
(79) 1,1,1-Trichloroethane	−30.4	74.1	7.25 (20 °C)	5.7	1.4380	0.170
(80) 1,4-Dioxane[s]	11.8	101.3	2.21	1.5	1.4224	0.164
(81) Trichloroethene	−86.4	87.2	3.42 (16 °C)	2.7	1.4773	0.160
(82) Piperidine	−10.5	106.3	5.8 (20 °C)	4.0	1.4525	0.148
(83) Diethylamine	−49.8	55.5	3.78	4.0	1.3846	0.145
(84) Diphenyl ether	26.9	258.1	3.60	3.9	1.5763 (30 °C)	0.142
(85) *t*-Butyl methyl ether	−108.6	55.2	4.5 (20 °C)	4.1	1.3690	0.124
(86) Diethyl ether	−116.3	34.5	4.20	3.8	1.3524	0.117
(87) Benzene[t]	5.6	80.1	2.27	0.0	1.5011	0.111
(88) Di-*n*-propyl ether	−123.2	90.1	3.39 (26 °C)	4.4	1.3805	0.102
(89) Toluene[t]	−95.0	110.6	2.38	1.0	1.4969	0.099
(90) 1,4-Dimethylbenzene	13.3	138.4	2.27 (20 °C)	0.0	1.4958	0.074
(91) Di-*n*-butyl ether	−95.2	140.3	3.08 (20 °C)	3.9	1.3992	0.071
(92) Carbon disulfide[u]	−111.6	46.3	2.64 (20 °C)	0.0	1.6275	0.065
(93) Tetrachloromethane	−22.9	76.7	2.24	0.0	1.4602	0.052
(94) Triethylamine	−114.7	88.9	2.42 (20 °C)	2.2	1.4010	0.043
(95) Tri-*n*-butylamine	−70.0	214.1	2.29	2.6	1.4291	0.043
(96) *cis*-Decahydronaphthalene	−43.1	195.8	2.20 (20 °C)	0.0	1.4810	0.015
(97) *n*-Heptane	−90.6	98.5	1.92 (20 °C)	0.0	1.3876	0.012
(98) *n*-Hexane	−95.4	68.7	1.88	0.0	1.3749	0.009
(99) *n*-Pentane	−129.8	36.1	1.84 (20 °C)	0.0	1.3575	0.009
(100) Cyclohexane	6.8	80.8	2.02 (20 °C)	0.0	1.4262	0.006

[a] The physical constants were taken from the following references: (1) R. C. Weast, M. J. Astle: *CRC Handbook of Data on Organic Compounds*, Vols. I and II, CRC Press, Boca Raton/FL, USA, 1985; (2) D. R. Lide (ed.): *Handbook of Chemistry and Physics*, 82nd edition, CRC Press, Boca Raton/FL, USA, 2001; (3) J. A. Riddick, W. B. Bunger, T. K. Sakano: *Organic Solvents, Physical Properties and Methods of Purification*. 4th edition, in: A. Weissberger (ed.): *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986; (4) A. L. McClellan: *Tables of Experimental Dipole Moments*, Vols. 1–3, Freeman, San Francisco, 1963; Rahara Enterprises, El Cerrito/CA, USA, 1974 and 1989; (5) A. A. Maryott, E. R. Smith: *Table of Dielectric Constants of Pure Liquids*, NBS Circular 514, Washington DC, 1951; (6) Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998; (7) G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), Chem Tec Publishing, Toronto, and William Andrew Publishing, New York, 2001; (8) M. J. O'Neil et al. (eds.): *The Merck Index*, 13th ed., Merck and Co., Whitehouse Station/NJ, USA, 2001.

[b] C. Reichardt, Chem. Rev. *94*, 2319 (1994); *cf.* also Table 7-3 in Chapter 7.

[c] Melting point.

[d] Boiling point at 1013 hPa.

[e] Relative permittivity ("dielectric constant") of the pure liquid at 25 °C, unless followed by another temperature in parentheses.

[f] Dipole moment in Coulombmeter (Cm), measured in benzene, tetrachloromethane, 1,4-dioxane, or *n*-hexane at 20...30 °C. 1 Debye = $3.336 \cdot 10^{-30}$ Cm.

[g] Refractive index at the average *D*-line of sodium (16969 cm^{-1}) at 20 °C, unless followed by another temperature in parentheses.

[h] Normalised E_T^N values, derived from the transition energy at 25 °C of the long-wavelength visible absorption of a standard pyridinium *N*-phenolate betaine dye, $E_T(30)$[b]; *cf.* Eqs. (7-27) and (7-29) in Section 7.4.

[i] Y. Marcus, Pure Appl. Chem. *62*, 139 (1990) (1,2-Ethanediol and 2,2,2-Trifluoroethanol).

[k] Y. Marcus, S. Glikberg, Pure Appl. Chem. *57*, 855, 860 (1985) (Methanol and Ethanol).

[l] J. F. Coetzee (ed.): *Recommended Methods for the Purification of Solvents and Tests for Impurities*, Pergamon Press, Oxford, 1982. (Acetonitrile, Sulfolane, Propylene carbonate, Dimethyl sulfoxide, *N*,*N*-Dimethylformamide, Hexamethylphosphoric triamide, Pyridine, 1,2-Diaminoethane, *N*-Methylacetamide, and *N*-Methylpropionamide).

[m] Y. Marcus, Pure Appl. Chem. *58*, 1411 (1986) (1-Propanol, 2-Propanol, and 2-Methyl-2-propanol).

[n] J. F. Coetzee, T.-H. Chang, Pure Appl. Chem. *58*, 1535, 1541 (1986) (Acetone and Nitromethane).

[o] B. J. Barker, J. Rosenfarb, J. A. Caruso, Angew. Chem. *91*, 560 (1979); Angew. Chem. Int. Ed. Engl. *18*, 503 (1979) (DMEU, DMPU, and Tetramethylurea).

[p] M. Bréant, Bull. Soc. Chim. Fr. 725 (1971) (1-Methylpyrrolidin-2-one).

[q] K. M. Kadish, J. E. Anderson, Pure Appl. Chem. *59*, 703 (1987) (Benzonitrile, Dichloromethane, 1,1-Dichloroethane, and 1,2-Dichloroethane).

[r] C. Agami, Bull. Soc. Chim. Fr. 1205 (1968) (1,2-Dimethoxyethane).

[s] J. F. Coetzee, T.-H. Chang, Pure Appl. Chem. *57*, 633 (1985) (THF, 1,4-Dioxane).

[t] K. M. Kadish, X. Mu, J. E. Anderson, Pure Appl. Chem. *61*, 1823 (1989) (Benzene and Toluene).

[u] A. D. Dunn, W.-D. Rudorf: *Carbon Disulfide in Organic Chemistry*, Ellis Horwood, Chichester, and Wiley, New York, 1989.

Table A-2. Selection of thirty-four chiral solvents and cosolvents (in alphabetical order)[a].

Solvents	t_{fp}/°C	t_{bp}/°C (hPa)	$[\alpha]_D^{20}$ [b]	Configuration
(1) 2-Amino-1-butanol	−2	178	−10 (neat)	(*R*)
		(1013)	+10 (neat)	(*S*)
(2) 2-Amino-1-propanol (Alaninol)	–	174.5	−18 (neat)	(*R*)
		(1013)	+18 (neat)	(*S*)
(3) 1-Amino-2-propanol	24 . . . 26	160	−18 (water)	(*R*)
		(1013)	+18 (water)	(*S*)
(4) 1,4-Bis(dimethylamino)-butane-2,3-diol (DBD)[c]	43	68 . . . 70	+34 (benzene)	(2*R*, 3*R*)
		(0.7)	−35 (benzene)	(2*S*, 3*S*)
(5) 1,4-Bis(dimethylamino)-2,3-dimethoxybutane (DDB)[d]	–	62 . . . 64	−15 (neat)	(2*R*, 3*R*)
		(4)	+15 (neat)	(2*S*, 3*S*)
(6) 1,3-Butanediol	< −50	207.5	−31 (ethanol)	(*R*)
		(1013)	+30 (ethanol)	(*S*)
(7) 2,3-Butanediol	16	183	−13 (neat; 23 °C)	(2*R*, 3*R*)
		(1013)	+13 (neat; 23 °C)	(2*S*, 3*S*)
(8) 2-Butanol	−114	99.5	−13 (neat)	(*R*)
		(1013)	+13 (neat)	(*S*)
(9) 2-Chlorobutane	−131	68	−31 (neat)	(*R*)
		(1013)	+31 (neat)	(*S*)
(10) Diethyl tartrate[e]	18	280	+8.5 (neat)	(2*R*, 3*R*)
		(1013)	−8.5 (neat)	(2*S*, 3*S*)
(11) 2,3-Dimethoxybutane[f,g]	−84	109 . . . 110	+3.7 (neat)[f]	(2*R*, 3*R*)
		(1000)		
(12) 1-Dimethylamino-1-phenylethane[h]	–	81	+62 (neat; 26 °C)	(*R*)
		(16)	−64 (neat)	(*S*)
(13) 2,3-Dimethylpentane	−135	90	−11 (neat)	(*S*)
		(1013)		
(14) Ethyl lactate	−25	154	+11 (neat)	(*R*)
		(1013)	−11 (neat)	(*S*)
(15) 2-Heptanol	–	160	−9.5 (neat)	(*R*)
		(1013)	+10 (neat; 24 °C)	(*S*)
(16) 3-Heptanol	−70	157	+5 (neat; 25 °C)	(*S*)
		(1013)		
(17) 2-(Hydroxymethyl)oxirane (Glycidol)	–	167 (dec.)	+15 (neat; 23 °C)	(*R*)
		(1013)	−15 (neat)	(*S*)
(18) 2-Methyl-1-butanol	−70	129	−6 (neat)	(*S*)
		(1013)		
(19) 3-Methyl-2-butanol	–	113	+5 (neat)	(*S*)
		(1013)		
(20) 3-Methylhexane	−119	92	+9 (neat)	(*S*)
		(1013)		
(21) 4-Methyl-2-pentanol	−90	132	−21 (neat)	(*R*)
		(1013)	+21 (neat)	(*S*)
(22) *N*-Methyl-(1-phenylethyl)amine	–	184	+70 ($CHCl_3$)	(*R*)
		(973)	−75 ($CHCl_3$)	(*S*)
(23) 2-Methyltetrahydrofuran[i,j]	−137	80	−27 (neat)	(*R*)
		(1013)	+27 (neat)	(*S*)
(24) 1-(1-Naphthyl)ethylamine[k]	–	153	+83 (neat; 17 °C)	(*R*)
		(11)	−81 (neat; 25 °C)	(*S*)
(25) 2-Octanol[l]	−32	180	−9.5 (neat; 17 °C)	(*R*)
		(1013)	+9.5 (neat)	(*S*)
(26) 2-Pentanol	−73	119	−13 (neat; 25 °C)	(*R*)
		(1013)	+13 (neat; 25 °C)	(*S*)

Table A-2. (Continued)

Solvents	t_{fp}/°C	t_{bp}/°C (hPa)	$[\alpha]_D^{20}$ [b]	Configuration
(27) 1-Phenylethanol[m,n]	20	204	+44 (neat)	(*R*)
		(1013)	−44 (neat)	(*S*)
(28) (1-Phenylethyl)amine[o,p]	−10	187	+39 (neat)	(*R*)
		(1013)	−40 (neat)	(*S*)
(29) *N*-(1-Phenylethyl)formamide[q,r]	46 . . . 47	175 . . . 178	+180 (neat)	(*R*)
		(20)	−172 (neat)	(*S*)
(30) 1-Phenyl-1-propanol	–	218 . . . 220	+48 (hexane)	(*R*)
		(1013)	−47 (hexane)	(*S*)
(31) 1,2-Propanediol	−60	188	−15 (neat; 24 °C)	(*R*)
		(1013)	+17 (neat)	(*S*)
(32) 1,2,3,4-Tetramethoxybutane[s]	–	70	−6 (neat)	(2*S*, 3*S*)
		(19)		
(33) 2,2,2-Trifluoro-1-(1-naphthyl)-ethanol[t]	52 . . . 53	83 . . . 85	−26 (ethanol)	(*R*)
		(0.03)		
(34) 2,2,2-Trifluoro-1-phenylethanol[u]	20	73 . . . 76	−41 (neat; 25 °C)	(*R*)
		(9)	+31 (neat)	(*S*)

[a] The physical constants are taken from the following references: (1) *Beilstein's Handbuch der organischen Chemie*, 4th ed., Springer, Berlin; (2) J. A. Riddick, W. B. Bunger, T. Sakano: *Organic Solvents, Physical Properties and Methods of Purification*, in: A. Weissberger (ed.): *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986; (3) Fluka Chemie AG: *Chiral Compounds Chemistry*, Buchs/Switzerland, 1994; (4) Sigma-Aldrich Co.: *Chiral Nonracemic Compounds*, Milwaukee/WI, USA, 1998; (5) Merck-Eurolab GmbH: *Der Laborkatalog Merck®*. Darmstadt, 2001.

[b] Specific rotation (dimension $10^{-1} \cdot \text{deg} \cdot \text{cm}^2 \cdot \text{g}^{-1}$), measured at the average *D*-line of sodium (16969 cm^{-1}) at 20 °C, unless followed by another temperature in parentheses. Because of different enantiomeric purities of the solvents studied, the literature values of $[\alpha]_D$ often vary; therefore, only rounded values are given.

[c] D. Seebach, H. Daum, Chem. Ber. *107*, 1748 (1974).

[d] D. Seebach, H.-O. Kalinowski, W. Langer, G. Crass, E.-M. Wilka, Org. Synth. *61*, 24, 42 (1983).

[e] H. Plieninger, H. P. Kraemer, Angew. Chem. *88*, 230 (1976); Angew. Chem. Int. Ed. Engl. *15*, 243 (1976).

[f] H. L. Cohen, G. F. Wright, J. Org. Chem. *18*, 432 (1953); N. Allentoff, G. F. Wright, ibid. *22*, 1 (1957).

[g] J. D. Morrison, R. W. Ridgeway, Tetrahedron Lett. 569 (1969).

[h] W. H. Pirkle, M. S. Hoekstra, J. Magn. Reson. *18*, 396 (1975).

[i] D. C. Iffland, J. E. Davis, J. Org. Chem. *42*, 4150 (1977).

[j] D. Gagnaire, A. Butt, Bull. Soc. Chim. Fr. 312 (1961); E. R. Novak, T. S. Tarbell, J. Am. Chem. Soc. *89*, 73 (1967).

[k] T. G. Burlingame, W. H. Pirkle, J. Am. Chem. Soc. *88*, 4294 (1966).

[l] E. Axelrod, G. Barth, E. Bunnenberg, Tetrahedron Lett. 5031 (1969).

[m] J. C. Jochims, G. Taigel, A. Seeliger, Tetrahedron Lett. 1901 (1967).

[n] A. J. H. Houssa, J. Kenyon, J. Chem. Soc. 2260 (1930); E. Downer, J. Kenyon, ibid. 1156 (1939).

[o] W. H. Pirkle, J. Am. Chem. Soc. *88*, 1837 (1966).

[p] W. Theilacker, H. G. Winkler, Chem. Ber. *87*, 690 (1954).

[q] P. Abley, F. J. McQuillin, J. Chem. Soc., Chem. Commun. 477 (1969); J. Chem. Soc., Part C 844 (1971).

[r] R. Huisgen, C. Rüchardt, Liebigs Ann. Chem. *601*, 21 (1956).

[s] D. Seebach et al., Helv. Chim. Acta *60*, 301 (1977).

[t] W. H. Pirkle, M. S. Hoekstra, J. Org. Chem. *39*, 3904 (1974); J. Am. Chem. Soc. *98*, 1832 (1976).

[u] W. H. Pirkle, S. D. Beare, T. G. Burlingame, J. Org. Chem. *34*, 470 (1969), W. H. Pirkle, P. L. Rinaldi, J. Org. Chem. *42*, 3217 (1977).

determined by the intended use [16]. The following is a practical definition of the purity of a solvent: "A material is sufficiently pure if it does not contain impurities of such nature and in such quantity as to interfere with the use for which it is intended" [1]. Detailed prescriptions for purification are available in standard texts [1, 17, 104, 105]. The most frequently found impurity in organic solvents is water. A water content of only 20 μg/g (20 ppm) is equivalent to the total amount of solute in a 10^{-3} molar solution! Since water interferes undesirably with many reactions, its removal is one of the basic laboratory operations. Drying agents may bind water either physically or chemically [18, 19]. The best method depends in each case on the chemical nature of the solvent and the desired degree of dryness [20]. All organic solvents possessing a relative permittivity of less than 15 can be freed almost completely from water, alcohols, peroxides, and traces of acid, by simple adsorptive filtration through aluminium oxide (activity I) or silica gel (activity I), *e.g.* using a chromatography column 2 . . . 5 cm in diameter and 40 . . . 150 cm long [19, 21–24].

Another safe and nonhazardous, general, large-scale procedure for the purification of solvents without distillation for their use in air- and moisture-sensitive reactions has been developed by Grubbs *et al.* [137]. The solvent, slightly pressurized with nitrogen, is passed through two sequential purification columns. The first contains activated alumina (removing water, peroxides, and inhibitors), and the second, optional, column contains a supported copper redox catalyst as an oxygen scavenger. A simplified modification of this distillation-free solvent purification system has also been described [138].

Comparative studies of the drying efficiencies of a number of common desiccants for different types of organic solvents have been carried out by Burfield and coworkers [106–117]. Using a new and very sensitive tritiated water tracer method for determining water content, they obtained rather unexpected results concerning the drying efficiencies of commonly used desiccants. The results of this study, together with other recommended physical and chemical drying methods [1, 17, 18, 104, 105], are compiled in Table A-3. In particular, Burfield's results draw attention to the almost universal applicability of zeolite molecular sieves as desiccants, which are capable of drying even the most difficult organic solvents [107].

The classical method for the determination of low water contents in organic solvents is the nonaqueous iodometric titration introduced by Karl Fischer in 1935, using a solution of sulfur dioxide, iodine, and pyridine in a benzene/methanol mixture [139, 140]. By replacing some of the toxic ingredients (pyridine, benzene, methanol), this titration method has more recently been developed into a simple and environmentally safe standard procedure [141].

Another UV/Vis spectroscopic method for the determination of water in organic solvents involves the use of solvatochromic dyes (such as the pyridinium *N*-phenolate betaine dye *(44)* in Chapter 7.4), and is based on the observation that water has a very high polarity compared with most organic solvents [142–145]. Even small amounts of water cause a strong hypsochromic band shift of the dissolved solvatochromic dye, which can be related to the water content by a calibration curve. A typical detection limit of this method is of the order of 1 mg water in 1 mL solvent for routine spectrophotometers [142]. An analogous solvatochromic method has been developed for the determination of aqueous ethanol mixtures [146].

A.3 Spectroscopic Solvents

Solvents used in ultraviolet, visible, infrared, microwave, and radiowave spectroscopy must meet the following requirements: transparency and stability toward the radiation used, solubility and chemical stability of the substance to be examined, and a high and reproducible purity (''optical constancy''). Normally, intermolecular interaction with the solute should be minimal. On the other hand, important information about the solute can be obtained from the changes in the absorption spectrum arising from such interactions.

A collection of UV/Vis, IR, as well as ^{1}H and ^{13}C NMR spectra of common organic solvents can be found in reference [147]. A comprehensive list with ''UV and IR spectroscopic windows'' of organic solvents is given in reference [8].

The number of solvents useful in UV/Vis spectroscopy decreases with decreasing wavelength (increasing wavenumber) since the absorption of all substances increases in this direction. The *cut-off point* depends on the chemical nature and to a large extent on the purity of the solvent. Hence, numerous procedures for the production of optically pure solvents have been developed [25–29]. Solvents for the measurement of fluorescence spectra must often be particularly pure [30]. The cut-off points of the solvents normally used in UV/Vis spectroscopy are collected in Table A-4. Saturated hydrocarbons are among the most useful because of the weak intermolecular interactions and the lack of excitable π-electrons. Perfluorinated hydrocarbons are recommended for the far-UV region (<200 nm) [31–33]. The UV spectra of the more important organic solvents are reproduced in the ''DMS UV Atlas of Organic Compounds'' [34].

Solvents for infrared spectroscopy must meet the additional requirement that they do not attack the absorption cells themselves (normally made from alkali metal halides such as NaCl, KBr, and CsBr) [35]. The transparency regions of the IR solvents within the mid-IR region ($2 \ldots 16$ μm; $5000 \ldots 625$ cm^{-1}) are given in Table A-5. Complete IR spectra of organic solvents can be found in the ''DMS Working Atlas of Infrared Spectroscopy'' [36], in the ''Sadtler IR Spectra Handbook of Common Organic Solvents'' [118], as well as in ''The Sprouse Collection of Infrared Spectra'' [148]. Transmission characteristics of organic solvents in the near-IR region ($1 \ldots 3$ μm; $1000 \ldots 3333$ cm^{-1}) are given in references [37, 38], and for the far-IR region ($15 \ldots 35$ μm; $667 \ldots 286$ cm^{-1}) in references [39, 40]. The IR spectra of deuterated organic solvents between 2.5 and 16.7 μm ($4000 \ldots 600$ cm^{-1}) have also been measured [41]. The number of IR absorption bands active in a covalent compound decreases with the number of atoms and with increasing symmetry of the molecule. Therefore, small molecules of high symmetry are particularly useful IR solvents, *e.g.* carbon disulfide (point group $D_{\infty h}$) and tetrachloromethane (point group T_d).

In ^{1}H NMR spectroscopy, one uses solvents which either contain no hydrogens (*e.g.* CS_2, CCl_4, $Cl_2C{=}CCl_2$, hexachlorobutadiene) or deuterated solvents (*e.g.* C_6D_6, $CDCl_3$, D_2O, CD_3SOCD_3). Table A-6 gives the characteristic ^{1}H NMR absorption bands of common organic solvents. More complete collections of ^{1}H NMR chemical shifts of organic solvents are available in the literature [42, 43]. Table A-7 contains the ^{13}C NMR absorption bands of compounds used as solvents as well as reference substances in ^{13}C NMR spectroscopy [44]. In order to use ^{1}H and ^{13}C NMR chemical shifts diagnostically for certain functional groups, all solvent effects should be eliminated

Table A-3. Some recommended simple physical and chemical drying methods for thirty-three common organic solvents: ++ method gives superdry solvents with less than 1 ppm water; + solvent sufficiently dry for most chemical applications; (+) often used but less efficient; − explosive hazard (!) or other chemical reaction; no entry means not recommended or no information in the literature. For extensive compilations of more sophisticated purification methods, see references[a–f].

Drying methods		Static drying[k] with															Dynamic[l] drying with					Fractional distillation[m] [Other methods]
Solvents[Ref.]	$S_{H_2O}^{g}$ 25 °C	M.S. 0.3 nm[n]	M.S. 0.4 nm[n]	M.S. 0.5 nm[n]	P_4O_{10}	Na	Al/Hg[a,c,d]	$LiAlH_4$[z)]	CaH_2	Al_2O_3 B–I	B_2O_3	KOH (powder)	$CaCl_2$	$CaSO_4 \cdot 0.5\ H_2O$[w]	Na_2SO_4	K_2CO_3	M.S. 0.3 nm[n]	M.S. 0.4 nm[n]	M.S. 0.5 nm[n]	Al_2O_3 B–I[l]	Al_2O_3 N–I[l]	
Acetic acid[a,b,d]	∞				+[d]	−	−	−	−	−		−	−			−				−	−	+(from Ac_2O or P_4O_{10})[bj] [fractional freezing]
Acetone[a–d,p,q,β)]	∞	(+)[q]	(+)[q]		−	−	−	−	−	−	+[q]	−	−	(+)	(+)	−				−	−	+(from $CaSO_4 \cdot 0.5\ H_2O$)
Acetonitrile[a–e,i,o,p,w]	∞	+[o,p]			+[o]	−		−	+[d]		+[o]	−	(+)[o]	(+)[w]	(+)[a]	(+)[o]	+[e,f]				+[e,i]	+(from P_4O_{10} and then from K_2CO_3)
Benzene[a–d,i,o,p]	(0.63)		++[o,p]	+[e]	+[a]	+[o]		++[o]	++[o]	++[o]			++[o]	+[d]	(+)			+[f]		+[i]		+(from Na)[aj] [fractional freezing]
tert-Butanol[a,b,d,γ)]	∞	+[u]*	(+)[u]		−	(+)[u]	+	−	(+)[u]				−	(+)[d]		(+)[d]						+(from Mg/I_2)[d)] [fractional freezing]
2-Butanone[a–d)]	120.0				−	−	−	−	−	−		−	−	+[e]	+[d]	+[e]				−	−	+
Chloroform[a–d,i,x,y)]	0.6		+[y]	+[x,y]	+[f]	−!	−			+[y]		−	+[y]	+[a]	+[d]	+[a]		+[f]		+[i]		+(from M.S. 0.4 nm[n])[d)]
Cyclohexane[a,b,d,i)]	0.1		+[d]		+[d]	+[d]				+			+[d]					+[f]		+[i]		+(from Na or $LiAlH_4$)[d,z)]
1,2-Diaminoethane[b,d,e)]	∞			+[e]	−	+[e]		−				+[e]	−									+(from Na or M.S. 0.5 nm under N_2)[d,e)]
1,2-Dichloroethane[b,d)]	1.5				+[d]	−!	−		+[d]	+[b]			+[d]	+[d]	+[b]	+[b]				+		+(from P_4O_{10})
Dichloromethane[a–d,i,p,r)]	2.0		++[p]		+[d]	−!	−		+[d]	+		−	+[c]	+[d]				+[f]		+[i]		+(from P_4O_{10})
Diethyl ether[a–d,i,p,t,w,y,z)]	13.0		+*[p,t,z]	+[y]	−	+[c]		+		+			(+)[c,t]	(+)[w]		(+)[t]		+[f]		+[i]		+(from Na or $LiAlH_4$ under N_2)[z)]
1,2-Dimethoxyethane[b–d)]	∞				−	+[c]		+[c]	+[c]	+			+							+		+(under N_2)[c)]
Di-*i*-propylamine[b,d,s)]	(400)	+[s]	+[s]		−	+[s]		−	+[s]			(+)[s]	−									+(from NaOH)[d)]
N,*N*-Dimethylformamide[a–e,p,q)]	∞	+[q]	+[p,q]		+[q]	−		−	+[q]			−	−					+[f]				+(i. vac. under N_2 or as benzene azeotr.)
Dimethyl sulfoxide[a–e,i,q)]	∞	+[q]	+[q]	+[e]		−	−	−	+[d,e]				−	+[c,d]				+[d]				+(i. vac. from CaH_2)[e,q)] [fractional freezing]
1,4-Dioxane[a–e,i,o,p,r,w,z)]	∞	+[w]	+*[p,z]	+[f]	−	+[o]			+[o]	+		+[o]	(+)[w]	(+)[w]			+[e]	+	+[f]	+[i]		+(from Na under N_2)[c,e)]
1,2-Ethanediol[a,b,d,u)]	∞	+[u]*			−	−		−					−	(+)[d]	(+)[f]			+[d]				+(from Mg/I_2 or as benzene azeotr.)[u)]
Ethanol[a–d,u,v)]	∞	+[u]*			−	−	+[a,d]	−	+[d,u]				−				+[f,v]					+(from Mg/I_2 or as benzene azeotr.)[u)]
Ethyl acetate[a–d,i,p)]	(29.4)		+[p]		+[d]	−	−	−	+[d]			−	−	+[d]	+[f]	+[e]		+[f]			+[i]	+(from P_4O_{10})[a)]
HMPT[b–e,q)]	∞	+[q]	+[q]						+[q]			+[q]					+[e]					+(i. vac. from CaH_2, under Ar)[e,q)]

Methanol[a–d,u)]	∞	+[u]	–	–	–	–	+[a]	–	+[u]				–	+[d]			+[f]					+(from Mg/I_2)[u)]
N-Methylacetamide[b,d,e)]	∞					+[e]	–		–				–	–								+(i. vac.)[e)] [fractional freezing][e)]
1-Methyl-2-pyrrolidinone[a–d)]	∞					–		–					–	–								+(i. vac. or as benzene azeotrope)[c)]
Nitromethane[a,b,d,i,δ)]	19.0			+[f]		–[d]	–	–	–				–	+[d]	+[d]	+[d]						+(from M.S. 0.4 nm)
Propylene carbonate[d,e)]	(83)			+[e]	+[e]		–		–				–							+[e,d]		+(i. vac.)[e,d)]
2-Propanol[a–d,γ)]	∞	+[f]		+[c]	–	–	+[a]	–	+[d]					–	+			+[f]				+(from Mg/I_2)[a)]
Pyridine[a–e,i,s)]	∞	+[s]	+[s]					+[d]	+[s]	+		+[e,s]	–						+[f]		+[i]	+(from M.S. 0.5 nm/KOH)[e)]
Sulfolane[a–e)]	∞		+[c]							+		+		+[e]							+[e]	+(i. vac. from CaH_2)[e)]
Tetrachloromethane[a–d,i)]	0.1			+[f]	+[c]	+[i]	–[!]	–			+			+[i]		+	+[a]		+[f]		+[i]	+(from P_4O_{10})[d)]
Tetrahydrofuran[a–e,i,p,z)]	∞			+[p,z]	+[f]	–	+[d]		+[e,d]	+[d]	+		–[!c]	+[c]	+[c]			+[e]	+	+[f]	+[l]	+(from $LiAlH_4$ or Na under N_2)[e,z)]
Toluene[a–d,p,r)]	0.5			++[p]	+[c]	+[d]	+[d]		+	+[d]	+			+[d]	+[d]					+[f]		+	+(from Na)[d)]
Trichloroethene[a,b,d,i)]	(0.25)						–[!]	–					–	+[d]		+[f]	+[d]					+[i]	+

[a)] Houben-Weyl-Müller: *Methoden der organischen Chemie*, 4th edition, Thieme, Stuttgart, 1959, Vol. I/2, p. 765–868 (W. Bunge: *Eigenschaften und Reinigung der wichtigsten organischen Lösungsmittel*) and p. 869–885 (H. Rickert and H. Schwarz: *Trockenmittel*).

[b)] J. A. Riddick, W. B. Bunger, T. K. Sakano: *Organic Solvents. Physical Properties and Methods of Purification*, 4th ed., in A. Weissberger (ed.): *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986.

[c)] B. S. Furniss, A. J. Hannaford, P. W. G. Smith, A. R. Tatchell: *Vogel's Textbook of Practical Organic Chemistry*, 5th ed., Longman, London, New York, 1989, p. 395–413.

[d)] W. L. F. Armarego, D. D. Perrin: *Purification of Laboratory Chemicals*, 4th ed., Butterworth-Heinemann, Oxford/UK, 1996.

[e)] J. F. Coetzee (ed.): *Recommended Methods for Purification of Solvents and Tests for Impurities*, Pergamon Press, Oxford 1982; Pure Appl. Chem. *57*, 634 (1985).

[f)] Merck–Eurolab GmbH: *Der Laborkatalog Merck® – Chemikalien und Reagenzien*. D-64293 Darmstadt/Germany, 2001, p. 1285 (Lösungsmittel – Eigenschaften und Trocknung).

[g)] Maximum solubility of water in solvent in g/L, except values in parentheses which are in g/kg; ∞ means completely miscible with water[b,h,i)].

[h)] K. Schwetlick et al.: *Organikum – Organisch-chemisches Grundpraktikum*, 21st ed., Wiley-VCH, Weinheim/Germany, 2000, p. 741–762.

[i)] B. P. Engelbrecht: *Adsorptives Reinigen von Lösungsmitteln für die Chromatographie und Spektroskopie*, GIT Fachz. Lab. *23*, 681 (1979); Chem. Abstr. *91*, 133523y (1979); see also ICN Biomedicals GmbH (formerly Woelm Pharma GmbH): *Lösungsmittel-Reinigung mit ICN Adsorbentien*, D-37269 Eschwege/Germany.

[k)] Treatment with *ca.* 50 . . . 100 g/L desiccant at ambient temperature for about 24 h. Gentle agitation or stirring has an accelerating effect on drying. Sequential drying, accomplished by decanting monosiccated solvent onto a fresh charge of *ca.* 50 g/L desiccant, is more effective.

[l)] Column drying by percolating the solvent through the desiccant contained in a glass column of 2 . . . 5 cm diameter and 40 . . . 150 cm long, and collecting the eluent in a storage container protected from atmospheric moisture by a drying tube filled with molecular sieve.

[m)] Fractional distillation is often combined with static drying before or after the distillation.

[n)] Zeolite molecular sieves (sodium and calcium aluminosilicates) of nominal pore size 0.3 . . . 0.5 nm, normally used as beads except in cases where the use of powdered molecular sieve is essential (marked with an asterisk).

[o)] D. R. Burfield, K.-H. Lee, R. H. Smithers, J. Org. Chem. *42*, 3060 (1977).

[p)] D. R. Burfield, G.-H. Gan, R. H. Smithers, J. Appl. Chem. Biotechnol. *28*, 23 (1978); Chem. Abstr. *89*, 12551f (1978).

[q)] D. R. Burfield, R. H. Smithers, J. Org. Chem. *43*, 3966 (1978).

[r)] D. R. Burfield, R. H. Smithers, J. Chem. Technol. Biotechnol. *30*, 491 (1980); Chem. Abstr. *94*, 66822s (1981).

[s)] D. R. Burfield, R. H. Smithers, A. S. C. Tan, J. Org. Chem. *46*, 629 (1981).

[t)] D. R. Burfield, R. H. Smithers, J. Chem. Educ. *59*, 703 (1982).

[u)] D. R. Burfield, R. H. Smithers, J. Org. Chem. *48*, 2420 (1983); Y. Marcus, S. Glikberg, Pure Appl. Chem. *57*, 855, 860 (1985).

[v)] D. R. Burfield, G. T. Hefter, D. S. P. Koh, J. Chem. Technol. Biotechnol., Chem. Technol. Part A *34*, 187 (1984); Chem. Abstr. *101*, 133117u (1984).

[w)] D. R. Burfield, J. Org. Chem. *49*, 3852 (1984).

[x)] D. R. Burfield, J. Chem. Educ. *56*, 486 (1979); D. R. Burfield, E. H. Goh, E. H. Ong, R. H. Smithers, Gazz. Chim. Ital. *113*, 841 (1983).

[y)] D. R. Burfield, R. H. Smithers, Chem. Ind. (London) 240 (1980).

[z)] D. R. Burfield, J. Org. Chem. *47*, 3821 (1982); Deperoxidation of ethers with self-indicating molecular sieves 0.4 nm.

[α)] *Caution*: solutions of $LiAlH_4$ in oxygen-containing solvents may decompose at elevated temperatures (≧160 °C). Therefore, distillation should never be carried out to dryness, and solvents boiling above 100 °C should be distilled under reduced pressure.

[β)] J. F. Coetzee, T.-H. Chang, Pure Appl. Chem. *58*, 1535 (1986).

[γ)] Y. Marcus, Pure Appl. Chem. *58*, 1411 (1986).

[δ)] J. F. Coetzee, T.-H. Chang, Pure Appl. Chem. *58*, 1541 (1986).



Table A-4. Ultraviolet "cut-off points"[a)] of spectroquality solvents commonly used in UV/Vis spectroscopy, for 1 cm pathlengths (accuracy *ca.* ±5 nm)[b)].

[a)] The "cut-off point" in the ultraviolet region is the wavelength at which the absorbance approaches 1.0 using a 1-cm cell path with water as the reference. Solvents should not be used for measurements below the cut-off point, even though a compensating reference cell is employed. The cut-off points are very dependent on the purity of the solvent used. Most of the solvents listed above are available in highly purified "spectrograde" quality.

[b)] Compiled from the following references:

(1) Eastman Kodak Company: *Spectrophotometric Solvents*, Dataservice Catalog JJ-282, Rochester, New York 14650, USA, 1977;

(2) E. Merck: *UVASOLE® – Lösungsmittel und Substanzen für die Spektroskopie*, D-64293 Darmstadt, Germany;

(3) and from the reviews of Gordon and Ford [4] (p. 167), Pestemer [25], and Hampel [34].

[c)] Values for solid, as used in a pellet for example.

Table A-5. Infrared transmission characteristics[a] of spectroquality solvents commonly used in infrared spectroscopy in the 2 . . . 16 μm region (5000 . . . 625 cm^{-1}) for 0.1 mm solvent thickness[b].



[a] ■ 80 . . . 100% Transmission; □ 60 . . . 80% Transmission. The black lines represent the useful regions. In the areas where the solvent has absorption bands that are totally absorbing, no information can be obtained about the sample, even though a compensating reference cell is employed. There is no solvent that is completely transparent over the entire wavenumber range. The most useful solvents are tetrachloromethane and carbon disulfide.

[b] N. L. Alpert, W. E. Keiser, and H. A. Szymanski: *IR – Theory and Practice of Infrared Spectroscopy*. 2nd edition, first paperback printing, Plenum Publishing Corporation, New York, 1973, p. 326.

Table A-6. Characteristic ^1H NMR absorption bands[a] of thirty spectroquality solvents, commonly used in ^1H nuclear magnetic resonance spectroscopy[b] (δ scale: Tetramethylsilane $\delta_H = 0$ ppm).

Solvents	δ/ppm (multiplicity)[a]
Acetic acid-d_4	11.53(1)[c], 2.03(5)
Acetone-d_6	2.04(5)
Acetonitrile-d_3	1.93(5)
Benzene-d_6	7.15(br)
Tribromomethane-d_1	6.85(1)
Trichloromethane-d_1	7.24(1)
Cyclohexane-d_{12}	1.38(br)
Deuterium oxide	*ca.* 4.8[c]
1,2-Dichloroethane-d_4	3.72(br)
Dichloromethane-d_2	5.32(3)
Diethyl ether-d_{10}	3.34(m), 1.07(m)
Diethylene glycol dimethyl ether-d_{14}	3.49(br), 3.40(br), 3.22(5)
1,2-Dimethoxyethane-d_{10}	3.40(m), 3.22(5)
N,*N*-Dimethylformamide-d_7	8.01(br), 2.91(5), 2.74(5)
Dimethyl sulfoxide-d_6	2.49(5)
1,4-Dioxane-d_8	3.53(m)
Ethanol-d_6 (anhydrous)	5.19(1)[c], 3.55(br), 1.11(m)
Hexafluoroacetone · 3/2 D_2O	*ca.* 9.0(br)[c], 5.26(1)
Hexamethylphosphoric triamide-d_{18}	2.53(2 × 5)
Methanol-d_4	4.78(1)[c], 3.30(5)
Nitrobenzene-d_5	8.11(br), 7.67(br), 7.50(br)
Nitromethane-d_3	4.33(5)
2-Propanol-d_8	5.12(1)[c], 3.89(br), 1.10(br)
Pyridine-d_5	8.71(br), 7.55(br), 7.19(br)
Sulfuric acid-d_2	*ca.* 11.0[c]
1,1,2,2-Tetrachloroethane-d_2	5.91(1)
Tetrahydrofuran-d_8	3.58(br), 1.73(br)
Toluene-d_8	7.09(m), 7.00(br), 6.98(m), 2.09(5)
Trifluoroacetic acid-d_1	11.50(1)[c]
2,2,2-Trifluoroethanol-d_3	5.02(1)[c], 3.88(4 × 3)



[a] The ^{1}H chemical shifts of the residual hydrogen atoms are in ppm relative to tetramethylsilane, determined with solutions containing 5 mL/100 mL tetramethylsilane as the internal standard. The multiplicity of the peaks is given in parentheses; br indicates a broad peak without resolvable fine structure, while m denotes one with fine structure. Partial deuterium labeling results in broadening or splitting of the residual peaks, due either to deuterium-hydrogen coupling or non-equivalence of the residual hydrogens. If undeuterated solvents are used, the region obscured might be as wide as 2 . . . 3 ppm.

[b] Compiled from the following references:

(1) Merck, Sharp & Dohme Canada Ltd., Isotope Division: *Deuterated Solvents – Handy Reference Data*, Quebec/Canada, 1990;

(2) H.-O. Kalinowski, S. Berger, S. Braun: *Carbon–13 NMR Spectroscopy*, Wiley, Chichester, 1988;

(3) E. Pretsch, P. Bühlmann, C. Affolter: *Structure Determination of Organic Compounds – Tables of Spectral Data*. 3rd ed., Springer, Berlin, 2000 (p. 240–243).

[c] These peak positions may vary considerably depending on solute, concentration, and temperature.

Table A-7. Characteristic ^{13}C NMR absorption bands[a] of thirty spectroquality solvents, commonly used in ^{13}C nuclear magnetic resonance spectroscopy[b] (δ_C scale: tetramethylsilane $\delta_C = 0$ ppm).

Solvents	δ/ppm (multiplicity)[a]
Acetic acid-d_4	178.4(br), 20.0(7)
Acetone-d_6	206.0(13), 29.8(7)
Acetonitrile-d_3	118.2(br), 1.3(7)
Benzene-d_6	128.0(3)
Tribromomethane-d_1	12.4(3)
Carbon disulfide	192.7(1)
Trichloromethane-d_1	77.0(3)
Cyclohexane-d_{12}	26.4(5)
Dichlorodifluoromethane	126.2(3)
1,2-Dichloroethane-d_4	43.6(5)
Dichloromethane-d_2	53.8(5)
Diethyl ether-d_{10}	65.3(5), 14.5(7)
Diethylene glycol dimethyl ether-d_{14}	70.7(5), 70.0(5), 57.7(7)
1,2-Dimethoxyethane-d_{10}	71.7(5), 57.8(7)
N,N-Dimethylformamide-d_7	162.7(3), 35.2(7), 30.1(7)
Dimethyl sulfoxide-d_6	39.5(7)
1,4-Dioxane-d_8	66.5(5)
Ethanol-d_6 (anhydrous)	56.8(5), 17.2(7)
Hexafluoroacetone · 3/2 D_2O	122.5(4), 92.9(7)
Hexamethylphosphoric triamide-d_{18}	35.8(7)
Methanol-d_4	49.0(7)
Nitrobenzene-d_5	148.6(1), 134.8(3), 129.5(3), 123.5(3)
Nitromethane-d_3	62.8(7)
2-Propanol-d_8	62.9(3), 24.2(7)
Pyridine-d_5	149.9(3), 135.5(3), 123.5(3)
Tetrachloroethene	121.4(1)
Tetrahydrofuran-d_8	67.4(5), 25.3(br)
Toluene-d_8	137.5(1), 128.9(3), 128.0(3), 125.2(3), 20.4(7)
Trifluoroacetic acid-d_1	164.2(4), 116.6(4)
2,2,2-Trifluoroethanol-d_3	126.3(4), 61.5(4 × 5)



[a] The ^{13}C chemical shifts are in ppm relative to tetramethylsilane, determined with solutions containing 5 mL/100 mL tetramethylsilane as the internal standard. The multiplicity of the peaks is given in parentheses; br indicates a broad peak without resolvable fine structure. The chemical shifts can be dependent on solute, concentration, and temperature.

[b] Compiled from the following references:

(1) Merck, Sharp & Dohme Canada Ltd., Isotope Division: *Deuterated Solvents – Handy Reference Data*, Quebec/Canada, 1990;
(2) H.-O. Kalinowski, S. Berger, S. Braun: *Carbon–13 NMR Spectroscopy.* Wiley, Chichester, 1988.
(3) E. Pretsch, P. Bühlmann, C. Affolter: *Structure Determination of Organic Compounds – Tables of Spectral Data*, 3rd ed., Springer, Berlin, 2000 (p. 157–160).

as far as possible. The ideal situation can be approached by extrapolation to infinite dilution from measurements at different concentrations. Solvent effects can generally be ignored in inert solvents such as CCl_4 or CS_2 at solute concentrations of $\leq$50 mg/g.

Concerning solvents for matrix isolation see Section A.4.

A.4 Solvents as Reaction Media

The following criteria can be used in the proper choice of solvents for chemical reactions and kinetic investigations: a maximum working range between melting and boiling point, chemical and thermal stability in this temperature range, good solubility of reactants and products (or sometimes insolubility of the products), compatibility with the analytical methods employed, and usually high degree of purity. The most useful solvents for some common reaction types are indicated in Table A-8. When a solvent is used for technical purposes, other factors may play an additional role [45]: price, flammability (ignition point, flash point), explosiveness (upper and lower explosion limit), viscosity, miscibility (*e.g.* blending with extenders or diluents), toxicity, corrosive action, and feasibility of recycling.

More recently, two-phase solvent systems, sometimes with temperature-dependent mutual miscibility of the two components, have gained interest as reaction media [149–156]. Having different solubilities for educts, products, reagents, and catalysts, biphasic solvent combinations can facilitate the separation of products from reaction mixtures. Since perfluorohydrocarbons [149–154] and room temperature ionic liquids [155, 156] are immiscible with many common organic solvents, they are particularly suitable for the formation of such biphasic solvent systems; see also Section 5.5.13.

Another interesting area concerning solvents as reaction media as well as for spectroscopy is the technique of *matrix isolation*, used when the molecule of interest is extremely unstable. Matrix isolation involves the trapping of the molecule in a rigid cage of a chemically inert solvent (the matrix) at low temperature. The main requirements are that the unstable molecule must have a stable precursor from which it can be liberated (usually by irradiation), and that the fragments so produced must not react with it or with the matrix. The reaction products can then be studied by IR or UV/Vis spectroscopy. Typical matrix materials are the noble gases (especially argon), the lighter hydrocarbons and halohydrocarbons, nitrogen, carbon dioxide, and other solvents and solvent mixtures that produce clear glasses when cooled to low temperatures. The properties of a number of matrix solvents can be found in references [93–96]. The temperature has to be sufficiently low to prevent diffusion of the solute into the matrix lattice. The useful temperature range is from 1.5 K to about 0.6 K of the matrix melting point. It should be remembered that the properties of a matrix-isolated molecule may be different from those in the commonly used solvents because of the low temperature and unusual environment.

A.5 Solvents for Recrystallization

The following requirements must be met by a solvent used for the recrystallization of an organic compound [46–48]: high solubility at high temperatures and low solubility at

Table A-8. Compilation of solvents commonly used for some important chemical reactions[a,b].

Solvents \ Reactions	S_N1 Reactions[c]	S_N2 Reactions[c]	Oxidation reactions[d]	Ozonization reactions	Epoxidation reactions	Catalytic Hydrogenation[d,e]	Metal Hydride Reductions[f]	Aldol Reactions[d,g]	Wittig Reactions[h]	Diels-Alder Cycloadditions	Grignard Reactions	Friedel-Crafts Acylation and Alkylation Reactions	Halogenation	Nitration	Sulfonation	Diazotization	Diazo Coupling
Acetic acid	+		+	+	+	+		+					+	+	+	+	+
Acetone		+			+					+							
Acetonitrile		+							+	+							
Benzene			+		+		+	+	+	+		(+)				+	
t-Butanol	+		+						+								
Carbon disulfide												+					
Trichloromethane			+	+	+					+					+		
Cyclohexane						+											
Dichloromethane			+	+					+	+		+		+			
Diethyl ether					+		+	+	+	+	+						
Di-*n*-butyl ether											+						
1,2-Dichloroethane				+						+		+					
1,2-Dichlorobenzene										+		+	+	+			
1,2-Dimethoxyethane							+	+	+								
N,*N*-Dimethylformamide		+				+			+							+	
Dimethyl sulfoxide		+						+	+								
1,4-Dioxane			+		+	+	+			+					+		
Ethanol	+					+		+	+	+						+	+
Ethyl acetate			+	+		+											
HMPT		+							+								
Methanol	+			+		+		+	+	+							+
Nitrobenzene			+									+	+	+	+		
Nitromethane		+										+					
Petroleum ether						+						+			+		
Pyridine			+					+									+
Sulfuric acid			+											+	+	+	
Tetrachloroethene										+							
Tetrachloromethane				+	+								+	+	+		
Tetrahydrofuran						+	+	+	+	+	+						
Tetramethylene sulfone		+	+									+					
Toluene							+	+	+	+							
Trichloroethene										+		+					
Water	+		+			+		+		+						+	+

[a] *Cf.* A. K. Doolittle: *The Technology of Solvents and Plasticizers*, Wiley, New York, 1954.
[b] Table A-8 contains a more arbitrary selection from a plurality of possibilities. If more than one solvent is given for a reaction, then binary mixtures of these solvents can also be used as reaction media.
[c] C. K. Ingold: *Structure and Mechanism in Organic Chemistry*, 2nd edition, Cornell University Press, Ithaca/N.Y., and London, 1969.
[d] H. O. House: *Modern Synthetic Reactions*, 2nd edition, Benjamin, Menlo Park/California, 1972.
[e] P. N. Rylander: *Solvents in Catalytic Hydrogenation*, in W. H. Jones (ed.): *Catalysis in Organic Synthesis*, Academic Press, New York, London, 1980.
[f] H. C. Brown: *Organic Syntheses via Boranes*, Wiley-Interscience, New York, 1985.
[g] A. T. Nielsen and W. J. Houlikan: *The Aldol Condensation*, Org. React *16*, 1 (1968); *cf.* p. 76–77. – C. H. Heathcock: *Stereoselective Aldol Condensations*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part B, Chapter 5, p. 177ff., Elsevier, Amsterdam, 1984; *cf.* p. 199.
[h] I. Gosney and A. G. Rowley in J. I. G. Cadogan (ed.): *Organophosphorus Reagents in Organic Synthesis*, Academic Press, London, 1979, p. 24–25 and 41.

low temperatures of the substance to be recrystallized; impurities should be either highly soluble or totally insoluble; the boiling point should be as high as possible; the solvent should be chemically inert; the solvent must favor crystal growth, and be easily separated from the pure crystals by washing or drying. Hence, it should be either volatile or very soluble in a more volatile solvent and not form clathrates or solvates. In the first instance, the choice of solvent can be made according to the old rule "like dissolves like". The following list gives some guidelines [49]; a more complete collection of organic solvents used for recrystallization is given in Table A-9. When a compound is too soluble in one, and not soluble enough in another solvent, a binary mixture of the two may be a useful medium for recrystallization (*e.g.* acetic acid/water, ethanol/water, ethanol/benzene, acetone/petroleum ether, or trichloromethane/petroleum ether). For solvent selection, see also references [119, 120, 157].

Crystallization of a solid from solution cannot only be effected by changing the temperature (decreasing solubility with decreasing temperature) but also by changing the solvent polarity (decreasing solubility with decreasing solvent polarity) at constant temperature. A "cold" crystallization technique, using pairs of solvents of different polarity (*e.g.* methanol/water, acetone/water) has been considered in reference [121].

Compound to be recrystallized		Well soluble in
Hydrocarbons	*hydrophobic*	Hydrocarbons, Ethers,
Halohydrocarbons		Halohydrocarbons
Ethers		
Amines		Carboxylic Esters
Esters		
Nitrohydrocarbons		
Nitriles		Alcohols, 1,4-Dioxane
Ketones		Acetic Acid
Aldehydes		
Phenols		Alcohols, Water
Amides		
Alcohols		
Carboxylic Acids		
Sulfonic Acids		
Salts	*hydrophilic*	Water

A.6 Solvents for Extraction and Partitioning (Distribution)

The partitioning of a substance between two liquid phases (multistage partitioning, partition chromatography) and the extraction of solids require similar properties of a solvent [50–55]. When a substance has to be partitioned, a solvent system with limited miscibility of the components is required in order that the substance dissolves to a different extent in the two phases. The greater the chemical differences between any two solvents, the more limited their miscibility. Other requirements that the solvent system must fulfil are, *inter alia*, a favorable partition coefficient (the average partition coefficient of the component mixture should be between *ca.* 0.2 and 5), as high a separation

Table A-9. Compilation of solvents commonly used for crystallization of organic compounds[a], listed in order of decreasing solvent polarity as measured by the E_T^N values[b].

Solvent	E_T^N	$t_{bp}/°C$	Suitable solvent for	Second solvent for mixture[c]
Water	1.000	100.0	Salts, amides, some carboxylic acids	Acetone, alcohols, 1,4-dioxane, acetonitrile
Methanol	0.762	64.5	General, esters, nitro and bromo compounds	Water, diethyl ether, benzene
2-Methoxyethanol	0.657	124.6	Sugars	Water, benzene, diethyl ether
Ethanol	0.654	78.3	General, esters, nitro and bromo compounds	Water, hydrocarbons, ethyl acetate
Acetic acid	0.648	117.9	Salts, amides, some carboxylic acids	Water
Acetonitrile	0.460	81.6	Dipolar compounds	Water, diethyl ether, benzene
Acetone	0.355	56.1	General, nitro and bromo compounds, osazones	Water, hydrocarbons, diethyl ether
Dichloromethane	0.309	39.6	General, low-melting compounds	Ethanol, hydrocarbons
Pyridine	0.302	115.3	High-melting, poorly soluble compounds	Water, methanol, hydrocarbons
Trichloromethane	0.259	61.2	General, acid chlorides	Ethanol, hydrocarbons
Methyl acetate	0.253	56.9	General, esters	Water, diethyl ether
Ethyl acetate	0.228	77.1	General, esters	Diethyl ether, benzene, hydrocarbons
1,4-Dioxane	0.164	101.3	Amides	Water, benzene, hydrocarbons
Diethyl ether	0.117	34.5	General, low-melting compounds	Acetone, hydrocarbons
Benzene	0.111	80.1	Aromatics, hydrocarbons, molecular complexes	Diethyl ether, ethyl acetate, hydrocarbons
Toluene	0.099	110.6	Aromatics, hydrocarbons	Diethyl ether, ethyl acetate, hydrocarbons
Tetrachloromethane	0.052	76.7	Apolar compounds, acid chlorides, anhydrides	Diethyl ether, benzene, hydrocarbons
Ligroin	–	90–110	Hydrocarbons	Ethyl acetate, benzene, dichloromethane
Petroleum ether	–	40–60	Hydrocarbons	Any solvent on this list from ethanol down
n-Hexane	0.009	68.7	Hydrocarbons	Any solvent on this list from ethanol down
Cyclohexane	0.006	80.8	Hydrocarbons	Any solvent on this list from ethanol down

[a] *Cf.* A. J. Gordon and R. A. Ford: *The Chemist's Companion – A Handbook of Practical Data, Techniques, and References*. Wiley, New York, London, Sydney, Toronto, 1972, p. 442.
[b] C. Reichardt, Chem. Rev. *94*, 2319 (1994).
[c] Trial and error is usually required in selecting a second solvent for a mixture. There are some generally successful mixtures, such as diethyl ether/methanol (or ethanol) for highly associated solids (especially amides and alcohols) and many natural products, and diethyl ether/petroleum ether (or benzene) for dipolar compounds (especially esters and alcohols) and hydrocarbons[a]. *Cf.* also J. B. Baumann: *Solvent Selection for Recrystallization*. J. Chem. Educ. *56*, 64 (1979); R. E. R. Craig: *Rapid, Efficient Determination of Recrystallization Solvents at the Microscale Level*, J. Chem. Educ. *66*, 88 (1989).

factor as possible (the ratio of the partition coefficients should not, in general, be smaller than 1.5), linearity of the partition isotherm (*i.e.* concentration independence of the partition isotherm), a sufficiently large capacity, high selectivity, no tendency for emulsion formation and rapid separation of the phases (this requires low viscosity, a large density difference, and a sufficient surface tension), absence of irreversible reactions between solvent and solute, and ease of recovery of the substance [51]. The optimization of these various requirements is difficult. Generally, a compromise between these, sometimes competing, factors must be made in the choice of phase partners. The solvents used most frequently in partitioning have been divided into five classes according to the intermolecular interactions between the solvent molecules. The difference lies predominantly in the number and strength of the intermolecular hydrogen bonds [51]. Typical representatives of these five classes of solvents are water, methanol, pyridine, trichloromethane, and *n*-heptane. Within each class the solvents can be ordered according to increasing solubility in water or decreasing solubility in *n*-heptane. Thus, a "mixotropic" series of solvents has been established [51], an extended version of which is presented in Table A-10. This series gives valuable information concerning both the miscibility of solvents and their use in partitioning methods (paper, column, and thin-layer chromatography). The farther that two solvents are from each other in Table A-10, the less miscible they are. More than 400 liquid stationary phases for gas-liquid chromatography are commercially available. The selection of the proper stationary phase for a separation problem is often done by "trial and error"; here again, the rule "like dissolves like" is often useful. Collections of established liquid stationary phases can be found in references [4, 56]. Methods for assessing the polarity and selectivity of stationary phases in gas-liquid chromatography have been reviewed [133].

For the optimization of solvent composition in high-performance liquid chromatography (HPLC), see references [122–125].

In addition to common organic solvents, supercritical fluids (scf's) can be used for a great variety of extraction processes [158–165]. Supercritical fluid extraction (SFE), mostly carried out with sc-CO_2 as eluant, has many advantages compared to extractions with conventional solvents. The solvent strength of a supercritical fluid can easily be controlled by the pressure and temperature used for the extraction: at a constant temperature, extraction at lower pressures will favour less polar analytes, while extraction at higher pressures will favour more polar and higher molar mass analytes. As supercritical fluids such as CO_2 and N_2O have low critical temperatures ($t_c = 31$ °C and 36 °C, respectively), SFE can be performed at moderate temperatures to extract thermolabile compounds. Typical industrial applications using sc-CO_2 include caffeine extraction from coffee beans [158] as well as fat and oil extraction from plant and animal tissues [165]. For some physical properties of supercritical solvents, see Section 3.2.

A.7 Solvents for Adsorption Chromatography

The accumulation of an organic substance on the surface of an adsorbent is determined by its dipolarity as well as its polarizability and molecular size. This is also true for solvents, which are adsorbed more strongly the more dipolar their molecules. The mole-

Table A-10. Mixotropic Solvent Series[a–e].

(1) Water
(2) Lactic acid
(3) Formamide
(4) Morpholine
(5) Formic acid
(6) Acetonitrile
(7) Methanol
(8) Acetic acid
(9) Ethanol
(10) 2-Propanol
(11) Acetone
(12) 1-Propanol
(13) 1,4-Dioxane
(14) Propanoic acid
(15) Tetrahydrofuran
(16) *t*-Butanol
(17) 2-Methylpropanoic acid
(18) 2-Butanol
(19) 2-Butanone
(20) Cyclohexanone
(21) Phenol
(22) *t*-Amyl alcohol
(23) 1-Butanol
(24) 3-Methylphenol
(25) Cyclohexanol
(26) *i*-Amyl alcohol
(27) 1-Pentanol
(28) Benzyl alcohol
(29) Ethyl acetate
(30) 1-Hexanol
(31) 2,4,6-Trimethylpyridine
(32) Pentanoic acid
(33) Ethyl formate
(34) 3-Methylbutanoic acid
(35) Furan
(36) Diethyl ether
(37) 1-Octanol
(38) Diethoxymethane
(39) Hexanoic acid
(40) Butyl acetate
(41) Di-*i*-propoxymethane
(42) Nitromethane
(43) 1-Bromobutane
(44) Di-*i*-propyl ether
(45) Butyl butyrate
(46) 1-Bromopropane
(47) Di-*n*-butyl ether
(48) Dichloromethane
(49) Trichloromethane
(50) Di-*i*-amyl ether
(51) 1,2-Dichloroethane
(52) Bromobenzene
(53) 1,1,2-Trichloroethane
(54) 1,2-Dibromethane
(55) Bromoethane
(56) Benzene
(57) 1-Chloropropane
(58) Trichloroethene
(59) Toluene
(60) Xylenes
(61) Tetrachloromethane
(62) Carbon disulfide
(63) Decalin
(64) Cyclopentane
(65) Cyclohexane
(66) *n*-Hexane
(67) *n*-Heptane
(68) Kerosene
(69) Petroleum ether
(70) Petroleum
(71) Paraffin oil

[a] This series applies generally to partition separations by paper-, column-, or thin-layer chromatography. The solvents listed go from most to least hydrophilic behaviour.
[b] E. Hecker, Chimia *8*, 229 (1954); E. Hecker: *Verteilungsverfahren in Laboratorien*, Verlag Chemie, Weinheim, 1955, p. 92 and 139.
[c] E. Heftmann (ed.): *Chromatography*, 2nd ed., Reinhold Publishing Company, New York, 1967.
[d] O. Mikeś: *Laboratory Handbook of Chromatographic Methods*, Van Nostrand, London, 1970.
[e] *Cf.* also A. J. Gordon and R. A. Ford: *The Chemist's Companion – A Handbook of Practical Data, Techniques, and References*, Wiley-Interscience, New York, 1972, p. 379.

cules of the adsorbed compound and the solvent compete for the active sites on the adsorbent surface. Thus, an organic compound will be adsorbed more strongly from an apolar than from a polar solvent. Conversely, a previously adsorbed material can only be replaced by a solvent when the latter has a higher affinity for the adsorbent [24, 52, 57–60, 126, 127].

The following requirements are desirable when choosing an eluant: high purity (in particular, the absence of water and other strongly polar compounds), solubility of the crude mixture, low viscosity, ease of regeneration, and suitability for analytical methods employed (*e.g.* when using UV/Vis detection during elution, the solvent itself must be transparent at the wavelength used). Furthermore, the eluant must be chemically inert to the adsorbate; *e.g.* acetone and ethyl acetate on an alkaline adsorbent such as aluminium oxide are readily transformed into diacetone alcohol and acetic acid, respectively. The success of a separation problem often depends more on the proper choice of solvent than on the choice of stationary phase. The solvents used as eluants can be arranged according to their increasing eluting power, the so-called "eluotropic" series of solvents, by an empirical determination of the retention times for a constant adsorbent and test mixture [24, 52, 57–61]. The shorter the retention time of the test sample, the higher the eluting power and hence the polarity of the solvent. Oxide adsorbents such as aluminium oxide and silica give almost identical eluotropic series. A generalised eluotropic series according to Pusey [61] is as follows: saturated hydrocarbons < aromatic hydrocarbons < halogenated hydrocarbons < ethers < esters < ketones < alcohols < carboxylic acids. Table A-11 gives the eluotropic series for standard solvents in conjugation with hydrophilic adsorbents according to Snyder [59, 60]. For hydrophobic adsorbents such as charcoal and polyamides, the eluotropic series is almost reversed. Often, mixtures of two or three solvents of different polarity lead to a better separation than a pure solvent. Again, the multicomponent eluants can be ordered in an eluotropic series [59, 60]. For example, the eluting power increases steadily in the series: petroleum ether, petroleum ether/benzene (100, 200, and 500 mL/L), benzene, benzene/ethanol (20, 50, 100, and 200 mL/L). Since the eluting power of a solvent is the result of interactions between adsorbent, solvent, and sample, eluotropic series are generally valid only for the substance class for which they have been established. The mixotropic (Table A-10) as well as the eluotropic series (Table A-11) reflect, approximately, the series of increasing polarity given by the empirical polarity parameter E_T^N (*cf.* Table A-1).

The most common approach to solvent characterization used by chromatographers is the 'solvent-triangle classification method' introduced by Snyder [166, 167]. Three solvent selectivity factors, x_i, representing the solvent's dipolarity, HBD acidity, and HBA basicity contributions to the total polarity, P', are plotted on a triangular diagram, allowing a classification of the many solvents used in liquid chromatography into eight classes with distinctly different selectivities.

In addition to conventional liquid chromatography, supercritical fluid chromatography (SFC), using a supercritical fluid as mobile phase (mostly scf-CO_2), has attracted attention in the last decades [58, 164, 168, 169]. Supercritical fluids provide a favourable medium for the transport of solutes through a chromatographic column because they resemble a gas in terms of viscosity, a liquid in terms of density, and are intermediate between these two phases in terms of diffusivity. For some physical properties of supercritical solvents, see Section 3.2.

Table A-11. Eluotropic solvent series for hydrophilic adsorbents such as alumina or silica, listed in order of increasing eluting power of the solvent[a–e], quantitatively measured by Snyder's empirical eluant strength parameter ε°[f].

Solvents	ε° (Al_2O_3)[f]	Solvents	ε° (Al_2O_3)[f]
(1) Fluoroalkanes (1,1,2-Trichloro-1,2,2-trifluoroethane)	−0.25	(20) 2-Butanone	0.51
		(21) Triethylamine	0.54
(2) *n*-Pentane	*0.00*	(22) Acetone	0.56
(3) *n*-Hexane	0.01	(23) 1,4-Dioxane	0.56
(4) 2,2,4-Trimethylpentane	0.01	(24) Tetrahydrofuran	0.57
(5) Cyclohexane	0.04	(25) Ethyl acetate	0.58
(6) Cyclopentane	0.05	(26) Diethylamine	0.63
(7) Carbon disulfide	0.15	(27) Nitromethane	0.64
(8) Tetrachloromethane	0.18	(28) Acetonitrile	0.65
(9) 1,1,1-Trichloroethane	0.19	(29) Pyridine	0.71
(10) *t*-Butyl methyl ether	0.20	(30) Dimethyl sulfoxide	0.75
(11) Xylene	0.26	(31) 1- and 2-Propanol	0.82
(12) Di-*i*-propyl ether	0.28	(32) Ethanol	0.88
(13) Toluene	0.29	(33) Methanol	0.95
(14) Chlorobenzene	0.30	(34) 1,2-Ethanediol	1.11
(15) Benzene	0.32	(35) Acetic acid	High ($\gg$1)
(16) Diethyl ether	0.38	(36) Water	Higher
(17) Trichloromethane	0.40	(37) Aqueous salt solutions, buffers	Highest
(18) Dichloromethane	0.42		
(19) 1,2-Dichloroethane	0.44		

[a] The term *eluotropic series* for solvents arranged in the order of increasing elution strength was introduced by W. Trappe, Biochem. Z. *305*, 150 (1940).

[b] As the relative elution power depends not only on the adsorbent, but in many cases on the compound types being separated, there exists no universal eluotropic series of solvent strengths. This series was given by L. R. Snyder[d]. For another eluotropic solvent series, see J. C. Touchstone: *Practice of Thin-Layer Chromatography*, 2nd ed., Wiley, Chichester, 1983.

[c] In binary solvent mixtures, addition of a small amount of one solvent to another less polar solvent rapidly increases the eluting power. The further away in the series the solvent pair is, the more drastic the change.

[d] For reversed-phase adsorbents such as charcoal or completely silanized silica, this eluotropic solvent series is valid in the reverse order. In this case, the eluting power increases in the following sequence: water < methanol < ethanol < acetone < 1-propanol < 1-butanol $\approx$ diethyl ether $\approx$ ethyl acetate < benzene < *n*-hexane[d].

[e] L. R. Snyder: *Principles of Adsorption Chromatography*, Dekker, New York, 1968; L. R. Snyder: *Solvent Selection for Separation Processes*, in E. S. Perry and A. Weissberger (eds.): *Techniques of Chemistry*, 3rd ed., Vol. XII, p. 25–75, Wiley-Interscience, New York, 1978; L. R. Snyder and J. J. Kirkland: *Introduction to Modern Liquid Chromatography*, 2nd ed., Wiley-Interscience, New York, 1979.

[f] The solvent strength ε° is a dimensionless number and indicates the free energy of adsorption of the solvent per unit area of adsorbent with unit activity, defined as $\varepsilon^{\circ} = 0$ for *n*-pentane on alumina[d]. For silica (SiO_2) as adsorbent, the ε° values are smaller by a factor of ca. 0.8 [ε° (SiO_2) $\approx 0.8 \cdot \varepsilon^{\circ}$ (Al_2O_3)], but the ranking of solvents remains the same.

The organization of solvents for analytical and preparative thin-layer chromatography also appears to be related to their viscosities [58, 126, 127].

A.8 Solvents for Acid/Base Titrations in Non-Aqueous Media

Many acids and bases are too weak to be titrated in aqueous solution (ionization constants $<10^{-8}$). The use of non-aqueous amphiprotic differentiating titration solvents (*cf.* Section 3.3.1), however, often permits one to reach a sharp titration end-point [62–70]. Non-aqueous solvents for the titration of weak acids and bases should be obtainable water-free and in a high state of purity, be chemically inert to the titrant and the substance under investigation, readily dissolve the substance and its titration product, or if the latter precipitates favour the formation of compact, crystalline, non-voluminous material. Weak bases are frequently titrated with perchloric acid in acetic acid, and weak acids with tetraalkylammonium hydroxide in 1,2-diaminoethane, alcohols, or pyridine [62–70]. The autoprotolysis constant is a particularly important criterion for solvent selection, since this constant determines the acidity or basicity region available in the solvent used. The smaller the autoprotolysis constant, the greater the range of acid or base strengths that can exist in a solvent and the greater the likelihood that it will be a differentiating solvent. Hence, acid-base titrations are best carried out in solvents with small K_{auto} values [64, 70]. The autoprotolysis constants, K_{auto}, of some amphiprotic solvents are given in Table A-12. These were often determined by conductivity measurements. It should be noted that the measured K_{auto} values often represent only a minimum value since, for extremely weakly basic or acidic solvents, it is difficult to distinguish between electrical conductivity caused by autoprotolysis and by impurities. A solvent is amphiprotic when it shows well-defined self-ionization, such that an autoprotolysis constant can be reproducibly measured. It is recommended that the term aprotic be used in place of amphiprotic for solvents with $K_{\text{auto}} < 10^{-20}$ ($\text{p}K_{\text{auto}} > 20$) [71]. In Table A-12 are also listed some solvents whose self-ionization cannot be due to protolysis, *e.g.* acetic anhydride. Just as it is possible to define pK areas, one can also evaluate intervals of electrochemical potential for various solvents useful for titrations with glass and calomel electrodes [72].

A.9 Solvents for Electrochemistry

Many electrochemical reactions, especially of organic compounds, are better carried out in non-aqueous solvents and may not even proceed in water. The following requirements should be met by these solvents [73–77]: sufficient solubility of the compounds to be examined and, of necessity, of the supporting electrolyte as well (usually tetraalkylammonium salts), chemical inertness towards the electrolyte and the reactive intermediates formed (*e.g.* the frequently formed radical anions would immediately be protonated by protic solvents), and as high a relative permittivity as possible (usually $\varepsilon_r > 10$). The latter will increase the electrical conductivity by favoring the dissociation of the electrolyte and hence decreasing the electrical resistance of the solution. Nevertheless, even solvents of low relative permittivity ($\varepsilon_r < 5$) can be used for electrochemical

Table A-12. Autoprotolysis constants (auto-ionization ionic products) of some amphiprotic solvents at 25 °C[a] according to 2 SH $\rightleftharpoons$ $SH_2^{\oplus} + S^{\ominus}$; $K_{auto} = [SH_2^{\oplus}] \cdot [S^{\ominus}]$[b].

Solvents	Lyonium Ion	Lyate Ion	$pK_{auto} = -\lg[K_{auto}/(mol^2 \cdot L^{-2})]$	References
Sulfuric acid	$H_3SO_4^{\oplus}$	$HSO_4^{\ominus}$	3.33	[c]
2-Aminoethanol	$HO—CH_2CH_2—NH_3^{\oplus}$	$H_2N—CH_2CH_2—O^{\ominus}$	5.7	[d]
Formic acid	$HC(OH)_2^{\oplus}$	$HCO_2^{\ominus}$	6.2	[e]
N-Methylformamide	$CH_3—\overset{\oplus}{N}H{=}CH—OH$	$CH_3—N{=}CH—O^{\ominus}$	10.74	[v]
Hydrogen fluoride	$H_2F^{\oplus}$	$F^{\ominus}$	12.5 (0 °C)	[w]
Water	$H_3O^{\oplus}$	$HO^{\ominus}$	14.00	[f]
Acetic acid	$CH_3C(OH)_2^{\oplus}$	$CH_3CO_2^{\ominus}$	14.45	[g]
Acetic anhydride	$CH_3CO^{\oplus}$	$CH_3CO_2^{\ominus}$	*ca.* 14.5 (20 °C)	[h]
Acetamide	$CH_3CO—NH_3^{\oplus}$	$CH_3CO^{\ominus}{=}NH$	14.6 (98 °C)	[i]
Deuterium oxide	$D_3O^{\oplus}$	$DO^{\ominus}$	14.96	[k]
1,2-Diaminoethane	$H_2N—CH_2CH_2—NH_3^{\oplus}$	$H_2N—CH_2CH_2—NH^{\ominus}$	15.2	[l]
1,2-Ethanediol	$HO—CH_2CH_2—OH_2^{\oplus}$	$HO—CH_2CH_2—O^{\ominus}$	15.84	[m]
Formamide	$HCO—NH_3^{\oplus}$	$HCO^{\ominus}{=}NH$	16.8 (20 °C)	[n]
Methanol	$H_3C—OH_2^{\oplus}$	$H_3C—O^{\ominus}$	17.20	[o]
Ethanol	$H_5C_2—OH_2^{\oplus}$	$H_5C_2—O^{\ominus}$	18.88	[p]
1-Propanol	$H_7C_3—OH_2^{\oplus}$	$H_7C_3—O^{\ominus}$	19.43	[p]
Hexamethylphosphoric triamide	$[(CH_3)_2N]_3\overset{\oplus}{P}—OH$		20.56	[o]
1-Pentanol	$H_{11}C_5—OH_2^{\oplus}$	$H_{11}C_5—O^{\ominus}$	20.65	[q]
2-Propanol	$(H_3C)_2C—OH_2^{\oplus}$	$(H_3C)_2C—O^{\ominus}$	20.80	[p]
1-Butanol	$H_9C_4—OH_2^{\oplus}$	$H_9C_4—O^{\ominus}$	21.56	[q]
Ethyl acetate	$H_3C—\overset{\overset{OH}{|}}{C}{=}\overset{\oplus}{O}C_2H_5$	$H_2C{=}\overset{\overset{O^{\ominus}}{|}}{C}—OC_2H_5$	22.83	[o]
N,*N*-Dimethylacetamide	$H_3C—\overset{\overset{OH}{|}}{C}{=}\overset{\oplus}{N}(CH_3)_2$	$H_2C{=}\overset{\overset{O^{\ominus}}{|}}{C}—N(CH_3)_2$	23.95	[o]
1-Methyl-2-pyrrolidinone			24.15	[o]
Tetrahydrothiophene-1,1-dioxide			25.45	[o]

Table A-12. (Continued)

Solvents	Lyonium Ion	Lyate Ion	$pK_{auto} = -\lg[K_{auto}/(mol^2 \cdot L^{-2})]$	References
2-Butanone			25.94	[o]
2-Methyl-2-propanol			26.8	[r]
N,*N*-Dimethylformamide			29.4	[r]
Ammonia			32.5 (−33 °C)	[s]
Acetone			32.5	[r]
Dimethyl sulfoxide			33.3	[t]
Acetonitrile			≥33.3	[u]

[a] Values at 25 °C unless stated otherwise.
[b] For more extensive compilations of autoprotolysis constants, see:
1. A. P. Kreshkov, Talanta *17*, 1029 (1970);
2. E. J. King: *Acid-Base Behaviour*, in A. K. Covington and T. Dickinson (eds.), *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, p. 342 and 391.
3. B. Trémillon: *Chemistry in Non-Aqueous Solvents*, Reidel Publishing Company, Dordrecht, Boston, 1974, p. 70ff.
4. S. Rondinini, P. Longhi, P. R. Mussini, and T. Mussini, Pure Appl. Chem. *59*, 1693 (1987);
5. Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998, Table 4.5 on p. 167–172.
[c] P. A. H. Wyatt, Trans. Faraday Soc. *65*, 585 (1969).
[d] P. W. Brewster, F. C. Schmidt, and W. B. Schaap, J. Phys. Chem. *65*, 990 (1961).
[e] M. Bréant, C. Beguin, and C. Coulombeau, Anal. Chim. Acta *87*, 201 (1976).
[f] H. S. Harned, J. Am. Chem. Soc. *47*, 930 (1925); H. S. Harned and B. B. Owen: *The Physical Chemistry of Electrolytic Solutions*, 3rd edition, Reinhold, New York, 1958; see also S. J. Hawkes, J. Chem. Educ. *72*, 799 (1995).

[g] S. Bruckenstein and I. M. Kolthoff, J. Am. Chem. Soc. *78*, 2974 (1956).
[h] G. Jander and H. Surawski, Z. Elektrochem., Ber. Bunsenges. Physik. Chem. *65*, 527 (1961).
[i] S. Guiot and B. Trémillon, J. Electroanal. Chem. *18*, 261 (1968).
[k] A. K. Covington, R. A. Robinson, and R. G. Bates, J. Phys. Chem. *70*, 3820 (1966).
[l] W. B. Schaap, R. E. Bayer, J. R. Siefker, J. Y. Kim, P. W. Brewster, and F. C. Schmidt, Record Chem. Progr. *22*, 197 (1961); Chem. Abstr. *56*, 12664 (1962).
[m] K. K. Kundu, P. K. Chattopadhyay, D. Jana, and M. N. Das, J. Phys. Chem. *74*, 2633 (1970).
[n] F. H. Verhoek, J. Am. Chem. Soc. *58*, 2577 (1936).
[o] A. P. Kreshkov, N. Sh. Aldarova, and B. B. Tanganov, Zh. Fiz. Khim. *44*, 2089 (1970); Russian J. Phys. Chem. *44*, 1186 (1970); Chem. Abstr. *73*, 124160 d (1970); *cf.* also A. P. Kreshkov, N. T. Smolova, N. Sh. Aldarova, and N. A. Gabidulina, Zh. Fiz. Khim. *46*, 667 (1972); Chem. Abstr. *77*, 33798 j (1972).
[p] R. Schaal and A. Tézé, Bull. Soc. Chim. Fr. 1783 (1961).
[q] A. P. Kreshkov, N. Sh. Aldarova, and N. T. Smolova, Zh. Fiz. Khim. *43*, 2846 (1969); Chem. Abstr. *72*, 54413 s (1970).
[r] L. N. Bykova and S. I. Petrov, Zh. Anal. Khim. *27*, 1076 (1972); Chem. Abstr. *77*, 93626 m (1972); *cf.* also [o].
[s] L. V. Coulter, J. R. Sinclair, A. G. Cole, and G. C. Roper, J. Am. Chem. Soc. *81*, 2986 (1959).
[t] J. Courtot-Coupez and M. Le Démézet, Bull. Soc. Chim. Fr. 1033 (1969); *cf.* also [r].
[u] I. M. Kolthoff and M. K. Chantooni, J. Phys. Chem. *72*, 2270 (1968); *cf.* also [r].
[v] T. Oncescu, A.-M. Oancea, and L. de Maeyer, J. Phys. Chem. *84*, 3090 (1980).
[w] R. Gut, J. Fluorine Chem. *15*, 163 (1980).

investigations [170]. Furthermore, the solvent should be of low viscosity in order to guarantee rapid mass transport of ions to the electrodes. An absence of conducting impurities, especially water, is particularly important when measuring electrical conductivities [78, 79]. In addition, the solvent should have a large anodic and cathodic voltage limit, which defines the "window" of accessible electronic levels available for electron-transfer processes. Although organic solvents have intrinsic limits based on their chemical oxidation-reduction properties, the practical working limits also depend on the nature of the working electrode material and the composition of the supporting electrolyte. Therefore, the voltage limits are a system property and not only a solvent property [77]. Practical working limits in organic solvents are given in the literature [8, 73, 74, 77, 80, 81]. The electrochemical windows of a range of ionic liquids, *i.e.* liquid at room temperature, can be found in reference [171].

The rather narrow electrochemical window of water, limited by the discharge of hydrogen and oxygen, has stimulated the use of nonaqueous solvents for electrochemical reactions. Procedures for measuring and reporting electrode potentials in nonaqueous solvents are presented in reference [128]. The solvent influence on the redox properties of cations and anions has been reviewed [172], as have applications of ion-selective electrodes in nonaqueous solvents [129] and the influence of nonaqueous solvents on the polarographic half-wave potentials of cations [173].

A.10 Toxicity of Organic Solvents

Organic solvents show varied physiological and toxicological properties, which all too often are neglected in the laboratory. All solvents influence the human organism to a greater or lesser extent [82–89, 130–132]. The extent of this influence depends on the time of exposure. High doses over short time intervals can lead to acute poisoning; small doses over prolonged periods can induce chronic damage.

Solvents can be ingested through the skin and the respiratory organs. Inhalation of solvent vapours affects not only the lungs but the whole circulatory system and hence the whole body. They accumulate principally in lipid and fat-rich cells in the nervous system, brain, bone marrow, liver, and body fat. Acute poisoning manifests itself by derangement of the central nervous system (euphoria, dizziness, unconsciousness). Chronic poisonings occur initially without any obvious symptoms and damage of the organs characteristic of the different solvents appears only later.

Many solvents are absorbed not only through the respiratory organs but also through the skin (*e.g.* tetrachloromethane, dimethyl sulfoxide, benzene). This leads to dehydration of the skin due to the removal of water and fat, thereby allowing the invasion of bacteria and dirt.

Hence, organic solvents should be handled with care. In the USA, the *threshold limit values* (*TL* values) are used as a measure of the inhalation toxicity for chronic interaction with solvent vapours [90]. In the Federal Republic of Germany, the *maximum concentration values at the workplace* (*MAK* values) are used [91, 92]. Threshold limit values refer to airborne concentrations of substances and represent conditions under which it is believed that workers may be repeatedly exposed daily without adverse effect. They refer to time-weighted average concentrations for a normal 8-hour workday

Table A-13. *TL* Values[a], *MAK* Values[b,c], and Vapour Pressures[b,c] for seventy-seven organic solvents.

Solvents[d]	*TL* Values (TWA)[e]		*MAK* Values[f]		Vapour pressure at 20 °C in hPa
	mL/m³	mg/m³	mL/m³	mg/m³	
Acetic acid	10	25	–	–	15.3
Acetic anhydride	5	21	5	21	4
Acetone	500	1200	500	1200	240
Acetonitrile (skin)	20	34	20	34	97
1-Aminobutane (skin)	–	–	5	15	122 (25 °C)
2-Aminoethanol	3	7.5	2	5.1	0.53
Ammonia	25	17	20	14	8570
Aniline (skin)[g]	2	7.6	2	7.7	0.4
Benzene (skin)[h]	0.5	1.6	–	–	101
n-Butane	800	1900	1000	2400	2137 (21 °C)
1-Butanol (skin)	20	61	100	310	6.3
2-Butanol	100	303	–	–	–
t-Butanol	100	303	20	62	40.8
2-Butanone (skin)	200	590	200	600	105
1-Butyl acetate	150	713	100	480	13.3
t-Butyl methyl ether[g]	50	180	50	180	270
Carbon disulfide (skin)	10	31	5	16	400
Chlorobenzene	10	46	10	47	12
Cyclohexane	100	344	200	700	104
Cyclohexanol (skin)	50	205	50	210	1.2
Cyclohexanone (skin)[g]	25	100	–	–	5
1,2-Diaminoethane (skin)	10	25	10	25	12
1,1-Dichloroethane	100	405	100	410	240
1,2-Dichloroethane[h]	10	40	–	–	87
1,1-Dichloroethene[g]	–	–	2	8	667
1,2-Dichloroethene (*cis* + *trans*)	200	793	200	800	220
Dichloromethane[g]	50	174	–	–	475
Diethylamine (skin)	5	15	5	15	260
Diethylene glycol	–	–	10	44	0.027
Diethylene glycol dimethyl ether (skin)	–	–	5	28	2
Diethyl ether	400	1213	400	1200	587
Dimethoxymethane	1000	3112	1000	3200	440
Di-*i*-propyl ether	250	1045	200	850	180
N,*N*-Dimethylacetamide (skin)	10	36	10	36	1.3
N,*N*-Dimethylformamide (skin)	10	30	10	30	3.5
2,6-Dimethyl-4-heptanone	25	145	–	–	2.2 (25 °C)
1,4-Dioxane (skin)	20	72	20	73	41
Diphenyl ether (vapour)	1	7	1	7.1	0.3 (25 °C)
1,2-Ethanediol (skin)	–	–	10	26	–
Ethanol	1000	1884	500	960	59
2-Ethoxyethanol (skin)	5	18	5	19	ca. 5
Ethyl acetate	400	1441	400	1500	97
Ethyl benzene (skin)[g]	100	434	–	–	9
Ethyl formate (skin)	100	303	100	310	256
Formic acid	5	9.4	5	9.5	42
n-Heptane	400	1640	500	2100	48

Table A-13. (Continued)

Solvents[d]	*TL* Values (TWA)[e]		*MAK* Values[f]		Vapour pressure at 20 °C in hPa
	mL/m³	mg/m³	mL/m³	mg/m³	
Hexamethylphosphoric triamide[h] (skin)	–	–	–	–	0.04
n-Hexane (skin)	50	176	50	180	160
Methanol (skin)	200	262	200	270	128
2-Methoxyethanol (skin)	5	16	5	16	ca. 11
Methyl acetate	200	638	200	610	220
3-Methyl-1-butanol	100	361	100	370	4
Methylcyclohexane	400	1606	200	810	48
Methyl formate (skin)	100	266	50	120	640
2-Methyl-1-propanol	50	152	100	310	11.7
1-Methylpyrrolidin-2-one (vapour) (skin)	–	–	19	80	0.3
Morpholine (skin)	20	72	10	36	10
Nitrobenzene (skin)[g]	1	5	–	–	0.2
Nitromethane (skin)[g]	20	50	–	–	37
n-Octane	300	1400	500	2400	15
n-Pentane	600	1770	1000	3000	573
3-Pentanone	200	705	–	–	–
Phenol (skin)[g]	5	19	–	–	0.47
2-Propanol	400	983	200	500	44
2-Propen-1-ol (skin)[g]	0.5	1.2	–	–	24
i-Propylbenzene, Cumene (skin)	50	246	50	250	4
Pyridine	5	14	5	16	20
1,1,2,2-Tetrachloroethane (skin)[g]	1	7	1	7	7
Tetrachloroethene (skin)[g]	25	170	–	–	19
Tetrachloromethane (skin)[g]	5	31	0.5	3.2	120
Tetrahydrofuran (skin)	200	590	50	150	200
Toluene (skin)	50	188	50	190	29
1,1,1-Trichloroethane (skin)	350	1910	200	1100	133
Trichloroethene[h]	50	269	–	–	77
Trichloromethane (skin)	10	49	0.5	2.5	210
Triethylamine (skin)	1	4.2	1	4.2	72
Xylene (all three isomers) (skin)	100	434	100	440	15

[a] *2002 Threshold Limit Values (TLVs®) for Chemical Substances and Physical Agents*, adopted by the American Conference of Governmental Industrial Hygienists (ACGIH®), available by ACGIH, 1330 Kemper Meadow Drive, Cincinnati/Ohio, 45240–1634, USA [90].

[b] *Maximale Arbeitsplatzkonzentration und Biologische Arbeitsstofftoleranzwerte 2002 (Maximum Concentrations and Biological Tolerance Values at the Workplace)*, Report No. 38 of the "Senatskommission zur Prüfung gesundheitsschädlicher Arbeitsstoffe" of the Deutsche Forschungsgemeinschaft (DFG), D-53170 Bonn/Germany; Wiley-VCH, Weinheim/Germany, 2002 [91].

[c] L. Roth, U. Weller: *Sicherheitsfibel Chemie*, 5th ed., Ecomed Verlagsgesellschaft, München, 2001.

[d] Where solvents are followed by the designation "skin", this refers to the potential significant contribution to the overall exposure by the cutaneous route, including mucous membranes and the eyes, either by contact with vapours or, of probable greater significance, by direct skin contact with

Table A-13. (Continued)

the solvent. This attention-calling designation is intended to suggest appropriate measures for the prevention of cutaneous absorption so that the threshold limit is not invalidated.
[e)] Adopted *T*hreshold *L*imit *V*alues – Time Weighted Average (TLV–TWA) – in parts of vapour or gas per million parts of contaminated air by volume (mL/m^3; ppm) at 25 °C and 1013 hPa, and in approximate milligrams of substance per cubic metre of air (mg/m^3).
[f)] *M*aximale *A*rbeitsplatz*k*onzentrationen (maximum concentrations at the workplace) in parts of vapour or gas per million of contaminated air by volume (mL/m^3; ppm) at 20 °C and 1013 hPa, and in approximate milligrams of substance per cubic metre of air (mg/m^3).
[g)] Solvent suspect of carcinogenic potential in humans.
[h)] Solvent recognized as having carcinogenic potential, *i.e.* capable of inducing malignant tumors as shown through experience with animals and/or humans. – See also L. Roth, G. Rupp: *Krebserzeugende Stoffe*, 3rd ed., Wissenschaftliche Verlagsgesellschaft, Stuttgart/Germany, 2000.

and a 40-hour working week. *MAK* values represent the highest allowed airborne concentrations of gaseous, vapourous, or dusty chemical substances within a work area, which will neither, as far as is known, adversely affect the health of the workers nor disturb them, even by repeated long-term exposure. They refer to time-weighted average concentrations for an 8-hour workday and a 40-hour working week [91, 92]. Table A-13 gives the *TL* and *MAK* values of frequently used organic solvents. These values are based on the best available information from industrial experience and from experimental human and animal studies. The basis on which the *MAK* values are established is given in reference [92]. Since the amount and nature of the information available for establishing *TL* and *MAK* values varies, the precision of the estimated *TL* and *MAK* values is also subject to variation and the latest information should be consulted. The *TL* and *MAK* values in Table A-13 refer to those issued in 2002.

The vapour pressures of organic solvents are also listed in Table A-13, since these give an additional indication of their potential dangers caused by their volatility.

The designation ''skin'' is used to indicate that there is also a danger through skin ingestion either by airborne or by direct contact with the chemical substance [90–92]. When handling such compounds, meticulous care in avoiding contact with the skin, hair, and clothing is imperative for health protection.

The *TL* and *MAK* values should be used as guides in the control of health hazards. They are not constants that can be used to draw fine lines between safe and dangerous concentrations. Nor is it possible to calculate the *TL* or *MAK* values of solvent mixtures from the data in Table A-13, because antagonistic action or potentiation may occur with some combinations. It should be noted that occupational exposure limits such as the *TL* and *MAK* values are not intended for use as a comparative measure of one solvent against another. The values set airborne concentration limits on chemical exposure, but do not describe the ease with which that airborne limit is achieved. In addition, the vapour pressure of the solvent must also be considered. The lower the vapour pressure, the lower the airborne concentration. In order to better compare the safety of volatile compounds such as organic solvents, the use of the *vapour hazard ratio (VHR)* has been recommended as a feasible measure [175]. The vapour hazard ratio is defined as the quotient of the saturation concentration of a solvent (in mg/m^3; at a given temperature and pressure) and its occupational exposure limit (in mg/m^3; *e.g. TL* or *MAK* values), according to:

Table A-14. Some recommendations for the substitution of hazardous solvents by safer ones.

New safer solvent	Substitute for	Chemical reactions[a]	References
Perfluoroalkanes [*e.g.* perfluorohexane (bp. 57 °C), perfluoro(methylcyclohexane) (bp. 76 °C), perfluorodecalin (bp. 142 °C)]	Tetrachloromethane	Alkene brominations, reactions in fluorous biphasic systems (FBS)	[b] ... [f]
(Trifluoromethyl)benzene (bp. 102 °C)	Benzene, dichloromethane	Friedel–Crafts reactions acylations, tosylations, silylations, Swern oxidations, radical reactions	[g]
Supercritical fluids (*e.g.* sc-CO_2: $t_c = 31$ °C, $p_c = 73$ bar)	Many other less polar solvents	Many types of organic reactions, *e.g.* isomerizations, hydroformylations, Diels–Alder reactions, oxidations	[h] ... [l]
t-Butyl methyl ether (bp. 55 °C)	Et_2O, 1,4-Dioxane, THF, other ethers, benzene	Grignard reactions; as an octane booster in gasoline	[m] ... [o]
Dimethoxymethane (bp. 42 °C), Diethoxymethane (bp. 88 °C)	Et_2O, 1,4-dioxane, THF, other ethers, benzene, $CHCl_3$	Organometallic reactions, preparation of Grignard reagents and organolithium derivatives, Schmidt reaction	[p] ... [r]
1,3-Dimethylimidazolidin-2-one (DMI, DMEU) (bp. 225 °C)	HMPT	Organometallic reactions, S_N reactions, Wittig-Horner reactions, Birch reductions	[s] ... [t]
1,3-Dimethylpropyleneurea (DMPU) (bp. 246 °C)	HMPT	Organometallic reactions, Enolate and Michael additions, S_N and Wittig-Horner reactions	[u]
N,*N*,*N*′,*N*′-Tetraethylsulfamide (TES) (bp. 85–88 °C/0.5 Torr)	HMPT, ethers	Many types of organometallic reactions, *e.g.* Grignard reactions	[v]
(−)-(*S*)-Ethyl lactate (bp. 154 °C)	Many solvents, *e.g.* xylene, glycol ethers, 1-methylpyrrolidone	Many types of reactions, compatible with the ester group	[w]
Water (bp. 100 °C)	Many other polar solvents	Many types of reactions, *e.g.* Diels–Alder reactions, Claisen rearrangements, carbonyl additions, transition metal-catalyzed reactions (Suzuki, Stille, and Heck coupling reactions), redox reactions	[x] ... [z] [α] ... [γ]
Ionic liquids [*e.g.* 1-ethyl-3-methylimidazolium triflate (mp. −9 °C); 1-*n*-butyl-3-methylpyridinium chloride (mp. 98 °C); tetra-*n*-butylammonium bromide (mp. 130 °C)]	Many other polar solvents	Many types of reactions, *e.g.* transition metal-catalyzed reactions, Heck, Diels–Alder, hydrogenation, hydroformylation and S_E reactions, alkene polymerizations	[δ] ... [θ]

Table A-14. (Continued)

New safer solvent	Substitute for	Chemical reactions[a]	References
No solvent at all (*i.e.* solvent-free reactions)	All solvents	Many types of reactions, using the following techniques: Reactions on solid mineral supports (alumina, silica, clays); without any solvent, support, or catalyst between neat reactants; solid-liquid phase-transfer-catalyzed and microwave-activated reactions	[*ι*] . . . [*ξ*]

[a] Only a few typical reactions can be mentioned; see references for the whole spectrum of possible reactions.
[b] S. M. Pereira, G. P. Savage, G. W. Simpson, Synth. Commun. *25*, 1023 (1995).
[c] B. Cornils, Angew. Chem. *109*, 2147 (1997); Angew. Chem. Int. Ed. Engl. *36*, 2057 (1997).
[d] U. Diederichsen, Nachr. Chem. Tech. Lab. *47*, 805 (1999).
[e] B. Betzemeier, P. Knochel, Top. Curr. Chem. 206, 61 (1999).
[f] A. Endres, G. Maas, Chem. Unserer Zeit *34*, 382 (2000).
[g] J. J. Maul, P. J. Ostrowski, G. A. Ublacker, B. Linclau, D. P. Curran, Top. Curr. Chem. *206*, 79 (1999).
[h] G. Kaupp, Angew. Chem. *106*, 1519 (1994); Angew. Chem. Int. Ed. Engl. *33*, 1452 (1994).
[i] W. Leitner, Top. Curr. Chem. *206*, 107 (1999); Acc. Chem. Res. *35*, 746 (2002).
[j] R. Noyori (ed.): *Supercritical Fluids*, Chem. Rev. *99*, 353–633 (1999).
[k] S. L. Wells, J. M. DeSimone, Angew. Chem. *113*, 534 (2001); Angew. Chem. Int. Ed. Engl. *40*, 518 (2001).
[l] A. K. Sunol, S. G. Sunol, *Supercritical Fluids*, in G. Wypych (ed.): *Handbook of Solvents*, ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001; Chapter 21.1, p. 1419ff.
[m] M. J. O'Neil (ed.): *The Merck Index*, 13th ed., Merck & Co., Whitehouse Station/N.J., 2001, p. 6056.
[n] D. Stoye: *Solvents*, in *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed., VCH, Weinheim, 1993, Vol. A24, p. 495.
[o] H. Quast, M. Witzel, Liebigs Ann. Chem. 699 (1993) (for azeotropic distillations).
[p] P. C. Kierkus, in L. A. Paquette (ed.): *Encyclopedia of Reagents for Organic Syntheses*, Wiley, Chichester, 1995, Vol. 3, p. 1977ff.
[q] G. Schilling, GIT Fachz. Lab. *36*, 1157 (1992).
[r] N. Gálvez, M. Moreno-Manas, R. M. Sebastián, A. Vallribera, Tetrahedron *52*, 1609 (1996).
[s] J. Malinowski, Specialty Chemicals *8*(3), 246 (1988); Chem. Abstr. *110*, 7144g (1989).
[t] E. M. Leahy, in L. A. Paquette (ed.): *Encyclopedia of Reagents for Organic Syntheses*, Wiley, Chichester, 1995, Vol. 3, p. 2086ff.
[u] A. Keck, D. Seebach, in L. A. Paquette (ed.): *Encyclopedia of Reagents for Organic Syntheses*, Wiley, Chichester, 1995, Vol. 3, p. 2123ff.
[v] H. G. Richey et al., J. Org. Chem. *46*, 2823 (1981); ibid. *52*, 479 (1987).
[w] E. J. J. Grabowski, in L. A. Paquette (ed.): *Encyclopedia of Reagents for Organic Syntheses*, Wiley, Chichester, 1995, Vol. 4, p. 2476ff.
[x] R. Breslow, Acc. Chem. Res. *24*, 159 (1991).
[y] P. A. Grieco, Aldrichimica Acta *24*(3), 59 (1991); P. A. Grieco: *Organic Syntheses in Water*, Blackie Academic and Professional, Hampshire/U.K., 1998.
[z] A. Lubineau, J. Augé, Y. Queneau, Synthesis 741 (1994); A. Lubineau, J. Augé, Top. Curr. Chem. *206*, 1 (1999).
[α] C.-J. Li, T.-H. Chan: *Organic Reactions in Aqueous Media*, Wiley, Chichester, 1997.

Table A-14. (Continued)

[β] B. Cornils, W. A. Herrmann: *Aqueous-Phase Organometallic Catalysis*, Wiley-VCH, Weinheim, 1998; B. Cornils, Top. Curr. Chem. *206*, 133 (1999).
[γ] D. Sinou, Top. Curr. Chem. *206*, 41 (1999). – N. Akiya, P. E. Savage, Chem. Rev. *102*, 2725 (2002).
[δ] Y. Chauvin, H. Olivier-Bourbigou, CHEMTECH *25*(9), 26 (1995). – M. Freemantle, Chem. Eng. News (May 15) 37–50 (2000).
[ε] T. Welton, Chem. Rev. *99*, 2071 (1999). – P. Wasserscheid, T. Welton, *Ionic Liquids in Synthesis*, Wiley-VCH, Weinheim, 2002.
[ζ] P. Wasserscheid, W. Keim, Angew. Chem. *112*, 3926 (2000); Angew. Chem. Int. Ed. Engl. *39*, 3772 (2000).
[η] D. W. Rooney, K. R. Seddon: *Ionic Liquids*, in G. Wypych (ed.): *Handbook of Solvents*. ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001, Chapter 21.2 on p. 1459ff.; M. J. Earle and K. R. Seddon, Pure Appl. Chem. *72*, 1391 (2000).
[θ] V. P. W. Böhm, W. A. Herrmann, Chem. Eur. J. *6*, 1017 (2000).
[ι] J. O. Metzger, Angew. Chem. *110*, 3145 (1998); Angew. Chem. Int. Ed. Engl. *37*, 2975 (1998).
[κ] M. Binnewies: *Chemische Transportreaktionen – Die Gasphase als Lösungsmittel*, Chem. Unserer Zeit *32*, 15 (1998).
[λ] A. Loupy, Top. Curr. Chem. *206*, 153 (1999). – A. Loupy (ed.): *Microwaves in Organic Syntheses*, Wiley-VCH, Weinheim, 2002.
[μ] R. S. Varma, Green Chem. *1*, 43 (1999); ACS Symp. Ser. *767*, 292 (2000).
[ν] K. Tanaka, F. Toda, Chem. Rev. *100*, 1025 (2000).
[ξ] J. M. Thomas, R. Raja, G. Sankar, B. F. G. Johnson, D. W. Lewis, Chem. Eur. J. *7*, 2973 (2001).

$$VHR = \frac{\text{Saturation concentration (mg/m}^3\text{)}}{\text{Exposure limit (mg/m}^3\text{)}}$$

$$= \frac{\text{Sättigungskonzentration (mg/m}^3\text{)}}{\text{Grenzwert (mg/m}^3\text{)}} = GZ$$

In Germany, the vapour hazard ratio is described and recommended as the *Gefährdungszahl (GZ)* in the Technical Guidelines of the German 'Gefahrstoffverordnung' (TRGS No. 420) [176]. According to this definition, a solvent with a lower *vapour hazard ratio* or *Gefährdungszahl* is the safer one. For example, the *GZ* values for methanol and toluene (with *MAK* values of 270 and 190 mg/m^3, respectively) are (168000 mg/m^3)/(270 mg/m^3) = *622* and (110000 mg/m^3)/(190 mg/m^3) = *579*, respectively. Thus, compared to methanol, toluene is the safer solvent, in spite of the fact that methanol has the higher exposure limit. This is because the *GZ* (and *VHR*) value reflects the potential health risk of a solvent not only in terms of the exposure limit, but also the potential ease with which that limit is achieved under a given set of conditions. *VHR* values can range from 50 (for 1-methylpyrrolidin-2-one; low exposure risk) to 17644 for dichloromethane (high exposure risk). Different *GZ* and *VHR* values can be calculated under different conditions since the vapour pressure changes with temperature and pressure. A collection of *VHR* values, calculated according to the national exposure limits of many countries, can be found at http://www.dbe.dupont.com.

So-called *odor threshold values* are not included in Table A-13 since these often show great discrepancy in the literature. They depend strongly on the experimental technique and individual sensitivity. Nevertheless, an unpleasant smell together with any irritation should be taken as a warning signal.

As organic solvents can be released into the environment by air emissions, industrial and waste-treatment effluents, accidental spillages, leaking tanks, and the land disposal of solvent-containing wastes, considerable emphasis has also been placed on the environmental impact of solvents. Investigations concerning the fate of organic solvents in water, soil, and air; the methods of solvent detection and solvent recycling; and the substitution of hazardous solvents by safer ones, as well as the national and international regulations concerning solvents as volatile organic compounds and hazardous air pollutants have recently been extensively reviewed [174].

The replacement of hazardous organic solvents by safer ones is of particular interest for chemists in their routine daily laboratory work. Table A-14 collects a selection of more recent recommendations for such solvent substitutions, together with relevant references.

Of special interest is the renaissance of water as a useful solvent for many organic reactions, as well as the increasing importance of supercritical fluids and ionic liquids. In the context of sustainable chemistry and clean technology, the best solvent is no solvent at all. In following this axiom, increasing efforts have recently been made to develop solvent-free synthetic reactions using various modern techniques. The results are interesting, but not so promising that this book on solvent effects will become superfluous in the foreseeable future.

References

Introduction

[1] Hermannus Boerhaave: *Elementa Chemiae*, Editio Altera, Leydensi multo correctior et accuratior, G. Cavelier, Parisii, 1733, Tomus Primus, p. 558. [2] G. A. Lindeboom: *Herman Boerhaave – The Man and his Work*, Methuen, London, 1968; *cf.* also Endeavour *28*, 2 (1969). [3] P. Walden: *Die Lösungstheorien in ihrer geschichtlichen Aufeinanderfolge*, in W. Herz (ed.): *Sammlung chemischer und chemisch-technischer Vorträge*, Enke, Stuttgart, 1910, Vol. XV, Heft 8–12; Chem. Zentralbl. *1910 II*, 1352; P. Walden: *Elektrochemie nichtwässriger Lösungen*, J. A. Barth, Leipzig, 1924. [4] F. Szabadváry: *Geschichte der Analytischen Chemie*, Vieweg, Braunschweig, and Akadémiai Kiadó, Budapest, 1966, p. 38. [5] J. H. van't Hoff: *Über die Theorie der Lösungen*, in F. W. Ahrens (ed.): *Sammlung chemischer und chemisch-technischer Vorträge*, Enke, Stuttgart, 1900, Vol. V, Heft 1; Chem. Zentralbl. *1900 I*, 696. [6] J. R. Partington: *A History of Chemistry*, MacMillan, London, New York, 1964, Vol. 4, Chapter XX, p. 637ff. [7] For a survey of older works on solvent effects on reaction rates, see M. Magat, Z. Phys. Chem. *A 162*, 432 (1932); Chem. Zentralbl. *1933 I*, 1566. [8] J. H. van't Hoff: *Die chemische Dynamik*, in *Vorlesungen über theoretische und physikalische Chemie*, Vieweg, Braunschweig, 1898, Heft 1; Chem. Zentralbl. *1899 II*, *278*. [9] M. Berthelot and L. Péan de Saint-Gilles, Ann. Chim. et Phys., 3. Ser., *65*, 385 (1862); *66*, 5 (1862); *68*, 255 (1863). Translated into German and edited by M. and A. Ladenburg in: *Ostwald's Klassiker der exakten Naturwissenschaften*, Nr. 173, Engelmann, Leipzig, 1910. [10] N. Menschutkin, Z. Phys. Chem. *1*, 611 (1887); ibid. *5*, 589 (1890); ibid. *6*, 41 (1890); ibid. *34*, 157 (1900).

[11] Th. Zincke and H. Bindewald, Ber. Dtsch. Chem. Ges. *17*, 3026 (1884); Th. Zincke and H. Thelen, ibid. *17*, 1809 (1884). [12] P. C. Laar, Ber. Dtsch. Chem. Ges. *18*, 648 (1885); ibid. *19*, 730 (1886). [13] C. K. Ingold: *Structure and Mechanism in Organic Chemistry*, 2nd edition, Cornell University Press, Ithaca, London, 1969, p. 794ff. [14] L. Claisen, Liebigs Ann. Chem. *291*, 25 (1896); especially pp. 30, 43, and 86. [15] W. Wislicenus, Liebigs Ann. Chem. *291*, 147 (1896); especially p. 176ff. [16] L. Knorr, Liebigs Ann. Chem. *293*, 70 (1896); especially p. 88. [17] A. Hantzsch and O. W. Schultze, Ber. Dtsch. Chem. Ges. *29*, 2251 (1896); especially p. 2256. [18] H. Stobbe, Liebigs Ann. Chem. *326*, 347 (1903); especially p. 357ff. [19] O. Dimroth, Liebigs Ann. Chem. *377*, 127 (1910); ibid. *399*, 91 (1913). [20] K. H. Meyer, Liebigs Ann. Chem. *380*, 212 (1911); Ber. Dtsch. Chem. Ges. *45*, 2843 (1912); ibid. *47*, 826 (1914).

[21] S. E. Sheppard, Rev. Modern Phys. *14*, 303 (1942); Chem. Abstr. *37*, 1654 (1943). [22] A. Hantzsch, Ber. Dtsch. Chem. Ges. *55*, 953 (1922). [23] A. Kundt: *Über den Einfluß des Lösungsmittels auf die Absorptionsspectra gelöster absorbierender Medien*, Poggendorfs Ann. Phys. Chem. N.F. *4*, 34 (1878); Chem. Zentralbl. *1878*, 498. [24] W. R. Brode, J. Phys. Chem. *30*, 56 (1926); Chem. Zentralbl. *1926 I*, 2775. [25] G. Scheibe, E. Felger, and G. Rößler, Ber. Dtsch. Chem. Ges. *60*, 1406 (1927). [26] N. Menschutkin: *Sur les conditions de l'acte de la combinaison chimique; modifications déterminées la présence des dissolvants, soi-disant indifférents* (Extrait d'une lettre de M. Menschutkin, professeur de chimie à Saint-Petersburg, à M. Louis Henry, Louvain), Bulletin de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique [3] *19*, 513–514 (1890); *Cf.* also A. Bruylants, Bulletin de l'Académie Royale de Belgique (Classe des Sciences) [5] *62*, 866–882 (esp. p. 877) (1976); Chem. Abstr. *87*, 166770x (1977). – I thank Prof. Bruylants, Louvain-La-Neuve, for drawing my attention to this important letter of Menschutkin. [27] M. V. Lomonosov: *O Deistvii Chimicheskikh Rastvoritelei Voobshche (On the Action of Chemical Solvents in General)*, in S. R. Mikulinskii (ed.): *M. V. Lomonosov – Izbrannye Proizvedeniya (Selected Works)*, Tom 1, p. 58–73; Estestvennye Nauki i Filosofiya, Nauka, Moskva, 1986. This paper was first published in 1747 in Latin and Lomonosov considered it as the first contribution to a pure physical chemistry. – I thank Prof. Möckel, Mühlhausen/Germany, for drawing my attention to this important publication of Lomonosov. [28] J. H. van't Hoff: *Raoult Memorial Lecture*, J. Chem. Soc. *81*, 969 (1902); see also M. Sutton, Chem. Br. *37*, 66 (2001) and J. van Houten, J. Chem. Educ. *78*, 1570 (2001). [29] K. J. Laidler: *A Century of Solution Chemistry*, Pure Appl. Chem. *62*, 2221 (1990); K. J. Laidler: *The World of Physical Chemistry*, Oxford University Press, Oxford/U.K., 1993. [30] H. Ohtaki: *Development of Modern Solution Chemistry: A Search of New Fields*, Coord. Chem. Rev. *185/186*, 735 (1999).

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

[31] O. Sterner: *Chemistry, Health and Environment*, Wiley-VCH, Weinheim/Germany, 1999. [32] C. Bolm, O. Beckmann and O. A. G. Dabard: *Die Suche nach neuen umweltfreundlichen Prozessen*, Angew. Chem. *111*, 957 (1999), Angew. Chem. Int. Ed. Engl. *38*, 907 (1999). [33] W. M. Nelson: *Choosing Solvents that Promote Green Chemistry*, ACS Symp. Ser. *767*, 313 (2000) (Green Chemical Syntheses and Processes); Chem. Abstr. *133*, 323232m (2000). [34] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001; Chapter 21 (Substitution of Solvents by Safer Products and Processes). [35] K. R. Seddon, Kinetika i Kataliz *37*, 743 (1996); Kinetics and Catalysis *37*, 693 (1996); Chem. Abstr. *125*, 285927s (1996). [36] K. R. Seddon, J. Chem. Technol. Biotechnol. *68*, 351 (1997); Chem. Abstr. *126*, 306898w (1997). [37] J. O. Metzger: *Lösungsmittelfreie organische Synthesen*, Angew. Chem. *110*, 3145 (1998); Angew. Chem. Int. Ed. Engl. *37*, 2975 (1998). [38] M. Binnewies: *Chemische Transportreaktionen – Die Gasphase als Lösungsmittel*, Chem. Unserer Zeit *32*, 15 (1998). [39] A. Loupy: *Solvent-free Reactions*, Top. Curr. Chem. *206*, 153 (1999). [40] R. S. Varma: *Solvent-free Organic Syntheses*, Green Chem. *1*, 43 (1999); ACS Symp. Ser. *767*, 292 (2000).

[41] K. Tanaka and F. Toda: *Solvent-free Organic Synthesis*, Chem. Rev. *100*, 1025 (2000). [42] J. M. Thomas, R. Raja, G. Sankar, B. F. G. Johnson, and D. W. Lewis: *Solvent-Free Routes to Clean Technology*, Chem. Eur. J. *7*, 2973 (2001).

Chapter 2

[1] D. H. Whiffen (ed.): *Manual of Symbols and Terminology for Physicochemical Quantities and Units, Appendix I: Solutions*, Pure Appl. Chem. *51*, 1 (1979). [2] T. S. Ree, T. Ree, and H. Eyring, Angew. Chem. *77*, 993 (1965); Angew. Chem. Int. Ed. Engl. *4*, 923 (1965); H. Eyring and M. S. Jhon: *Significant Liquid Structures*, Wiley, New York, 1969. [3] J. S. Rowlinson and F. L. Swinton: *Liquids and Liquid Mixtures*, 3rd ed., Butterworths, London 1982; Essays in Chemistry *1*, 1 (1970). [4] J. H. Hildebrand, J. M. Prausnitz, and R. L. Scott: *Regular and Related Solutions*, Van Nostrand-Reinhold, Princeton, 1970. [5] D. Henderson (ed.): *Liquid State*, Vol. 8 in "Physical Chemistry – An Advanced Treatise" (eds. H. Eyring, D. Henderson, and W. Jost), Academic Press, London, New York, 1971. [6] F. Kohler: *The Liquid State*, Vol. 1 in *Monographs in Modern Chemistry*, Verlag Chemie, Weinheim, 1972. [7] D. M. Heyes: *The Liquid State*, Wiley-VCH, Weinheim/Germany, 1997. [8] E. Wicke, Angew. Chem. *78*, 1 (1966); Angew. Chem. Int. Ed. Engl. *5*, 106 (1966). [9] R. A. Horne: *The Structure of Water and Aqueous Solutions*, in A. F. Scott (ed.): Survey of Progress in Chemistry *4*, 1 (1968). [10] D. Eisenberg and W. Kauzmann: *The Structure and Properties of Water*, Oxford University Press, New York, 1969.

[11] P. Krindel and I. Eliezer: *Water Structure Models*, in Coord. Chem. Rev. *6*, 217 (1971). [12] R. A. Horne (ed.): *Water and Aqueous Solutions – Structure, Thermodynamics, and Transport Processes*, Wiley, New York, 1972. [13] F. Franks (ed.): *Water – A Comprehensive Treatise*, Vols. 1–7, Plenum Press, London, 1972–1982. [14] W. A. P. Luck (ed.): *Structure of Water and Aqueous Solutions*, Verlag Chemie, Weinheim, 1974. [15] R. Ludwig: *Wasser – Von Clustern zur Flüssigkeit*, Angew. Chem. *113*, 1856 (2001); Angew. Chem. Int. Ed. Engl. *40*, 1808 (2001). [15a] M. Klose and J. I. Naberuchin: *Wasser – Struktur und Dynamik*, Akademie-Verlag, Berlin, 1986. [16] H. S. Frank and W.-Y. Wen, Discussions Faraday Soc. *24*, 133 (1957); H. S. Frank, Proc. Roy. Soc. London Ser. A *247*, 481 (1958). [17] D. T. Hawkins: *A Bibliography on the Physical and Chemical Properties of Water (1969–1974)*, in J. Sol. Chem. *4*, 625 (1975). [18] K. Ziegler, in Houben-Weyl-Müller, "Methoden der Organischen Chemie", 4th ed., Thieme, Stuttgart 1955, Vol. IV/2, p. 738. [19] F. Franks and D. J. G. Ives, Quart. Rev. Chem. Soc. *20*, 1 (1966). [20] O. Fuchs, Fortschr. Chem. Forsch. *11*, 74 (1968); Deutsche Farben-Zeitschrift *22*, 548 (1968); *23*, 17, 57, 111 (1969); Chem. Abstr. *71*, 70949f (1969).

[21] K. C. James, Education in Chemistry *9*, 220 (1972). [22] G. C. Pimentel (ed.): *Chemistry – An Experimental Science*, Freeman, San Francisco, 1963, p. 313 and 554. [23] G. Duve, O. Fuchs, and H. Overbeck: *Lösemittel Hoechst*, 6th ed., Hoechst AG, Frankfurt (Main), 1976. [24] A. F. M. Barton, *Handbook of Solubility Parameters and other Cohesion Parameters*, CRC Press, Boca Raton/Florida, 1983. [25] M. R. J. Dack: *The Importance of Solvent Internal Pressure and Cohesion*

to Solution Phenomena, Chem. Soc. Rev. (London) *4*, 211 (1975). [26] For recent reviews on intermolecular forces, see: (a) Reviews on van der Waals Molecules, Parts I–III, in: (I) Chem. Rev. *88*, 813–988 (1988); (II) Chem. Rev. *94*, 1721–2160 (1994); (III) Chem. Rev. *101*, 3861–4264 (2001); (b) J. Israelachvili: *Intermolecular and Surface Forces*, 2nd ed., Academic Press, London, 1991; (c) P. L. Huyskens, W. A. P. Luck, and T. Zeegers-Huyskens: *Intermolecular Forces*, Springer, Berlin, 1991; (d) P. W. Atkins: *Physical Chemistry*, 6th ed., Oxford University Press, Oxford/U.K., 1998, p. 659ff.; (e) K. Müller-Dethlefs and P. Hobza: *Noncovalent Interactions – A Challenge for Experiment and Theory*, Chem. Rev. *100*, 143 (2000); (f) R. Schmid: *Solvent Effects on Chemical Reactivity*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 13.1, p. 737–777. [27] For compilations of attempts to treat intermolecular forces by quantum mechanical methods, see: (a) A. D. Buckingham, P. W. Fowler, and J. M. Hutson: *Theoretical Studies of van der Waals Molecules and Intermolecular Forces*, Chem. Rev. *88*, 963 (1988); (b) A. J. Stone: *The Theory of Intermolecular Forces*, Oxford University Press, Oxford/U.K., 1996; (c) S. Scheiner (ed.): *Molecular Interactions – From van der Waals to Strongly Bound Complexes*, Wiley-VCH, Weinheim/Germany, 1997; (d) J. Tomasi, B. Mennucci, and C. Cappelli: *Solvents and Solutions as Assemblies of Interacting Molecules*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 8.1, p. 419–504. [28] H. H. Jaffé, J. Chem. Educ. *40*, 649 (1963). [29] W. H. Keesom, Physik. Z. *22*, 129, 643 (1921); *23*, 225 (1922). [30] R. L. Amey, J. Phys. Chem. *72*, 3358 (1968).

[31] M. Rabinowitz and A. Pines, J. Am. Chem. Soc. *91*, 1585 (1969). [32] P. Debye, Physik. Z. *21*, 178 (1920); *22*, 302 (1921). [33] F. London, Z. Physik. Chem. (B) *11*, 222 (1931); Z. Physik *63*, 245 (1930); Trans. Faraday Soc. *33*, 8 (1937). [34] R. Ulbrich, Chemiker-Ztg. *99*, 320 (1975). [35] E. F. Meyer and R. E. Wagner, J. Phys. Chem. *70*, 3162 (1966). [35a] C. H. Yoder, J. Chem. Educ. *54*, 402 (1977). [36] A. J. Parker, Quart. Rev. (London) *16*, 163 (1962); Usp. Khim. *32*, 1270 (1963). [37] M. L. Huggins, Angew. Chem. *83*, 163 (1971); Angew. Chem. Int. Ed. Engl. *10*, 147 (1971). [38] M. D. Joesten and L. J. Schaad: *Hydrogen Bonding*, Dekker, New York, 1974. [39] P. Schuster (ed.): *Hydrogen Bonds*, Top. Curr. Chem. *120*, 1–113 (1984) (particularly p. 35ff.). [40] F. Hibbert and J. Emsley: *Hydrogen Bonding and Chemical Reactivity*, Adv. Phys. Org. Chem. *26*, 255 (1990).

[41] G. A. Jeffrey and W. Saenger: *Hydrogen Bonding in Biological Structures*, Springer, Berlin, 1991. [42] C. B. Aakeröy and K. R. Seddon: *The Hydrogen Bond and Crystal Engineering*, Chem. Soc. Rev. *22*, 397 (1993). [43] C. L. Perrin and J. B. Nielson: *"Strong" Hydrogen Bonds in Chemistry and Biology*, Annu. Rev. Phys. Chem. *48*, 511 (1997). [44] M. S. Gordon and J. H. Jensen: *Understanding the Hydrogen Bond Using Quantum Chemistry*, Acc. Chem. Res. *29*, 536 (1996). [45] D. Hadzi (ed.): *Theoretical Treatments of Hydrogen Bonding*, Wiley-VCH, Weinheim/Germany, 1997. [46] G. A. Jeffrey: *An Introduction to Hydrogen Bonding*, Oxford University Press, Oxford/U.K., 1997. [47] E. M. Arnett, L. Jores, E. Mitchell, T. S. S. R. Murty, T. M. Gorrie, and P. v. R. Schleyer, J. Am. Chem. Soc. *92*, 2365 (1970). [48] D. Hadži and N. Kobilarov, J. Chem. Soc. *A 1966*, 439. [49] G. Kortüm: *Lehrbuch der Elektrochemie*, 5th ed., Verlag Chemie, Weinheim, 1972. [50] G. Briegleb, *Elektronen-Donator-Acceptor-Komplexe*, Springer, Berlin, 1961.

[51] R. S. Mulliken and W. B. Person: *Molecular Complexes – A Lecture and Reprint Volume*, Wiley-Interscience, New York, 1969. [52] O. Hassel, Angew. Chem. *82*, 821 (1970). [53] V. Gutmann: *Coordination Chemistry in Non-Aqueous Solvents*, Springer, Wien, New York, 1968; V. Gutmann: *Chemische Funktionslehre*, Springer, Wien, New York, 1971; V. Gutmann, *The Donor–Acceptor Approach to Molecular Interactions*, Plenum Publ. Corp., New York, 1978. [54] H. A. Bent, Chem. Rev. *68*, 587 (1968). [55] E. N. Gur'yanova, Usp. Khim. *37*, 1981 (1968); Russ. Chem. Rev. *37*, 863 (1968); E. N. Gur'yanova, I. P. Gol'dshtein, and I. P. Romm: *The Donor–Acceptor Bond*, Wiley, New York, 1975. [56] J. Yarwood (ed.): *Spectroscopy and Structure of Molecular Complexes*, Plenum Press, New York, 1973. [57] R. Foster: *Organic Charge-Transfer Complexes*, Academic Press, London, New York, 1969; R. Foster (ed.): *Molecular Complexes*, Vols. 1 and 2, Elek Science, London, 1973/74. [58] A. E. Reed, L. A. Curtiss, and F. Weinhold, Chem. Rev. *88*, 899 (1988). [59] P. Hobza and R. Zahradnik: *Intermolecular Complexes*, Elsevier, Amsterdam, 1988, and Mir, Moscow, 1989. [59a] W. B. Jensen: *The Lewis Acid-Base Definition – A Status Report*, Chem. Rev. *78*, 1 (1978). [59b] C. J. Bender: *Theoretical Models of Charge-Transfer*

Complexes, Chem. Soc. Rev. *15*, 475 (1986). [60] R. S. Drago, Structure and Bonding *15*, 73 (1973); J. Chem. Educ. *51*, 300 (1974).

[61] D. W. Turner: *Ionization Potentials*, Adv. Phys. Org. Chem. *4*, 31 (1966). [62] G. Briegleb: *Elektronenaffinitäten organischer Moleküle*, Angew. Chem. *76*, 326 (1964); Angew. Chem. Int. Ed. Engl. *3*, 617 (1964). [63] V. Kampars and O. Neilands: *The Electron Affinities of Organic Electron Acceptors*, Usp. Khim. *46*, 945 (1977); Russ. Chem. Rev. *46*, 503 (1977). [64] R. S. Mulliken and W. B. Person, J. Am. Chem. Soc. *91*, 3409 (1969). [65] D. W. Meek: *Lewis Acid-Base Interactions in Polar Non-Aqueous Solvents*, in J. J. Lagowski (ed.): *The Chemistry of Non-Aqueous Solvents*, Academic Press, London, New York, 1966, Vol. I, p. 1ff. [66] R. S. Drago and K. F. Purcell: *Coordinating Solvents*, in T. C. Waddington (ed.): *Non-Aqueous Solvent Systems*, Academic Press, London, New York, 1965, p. 211ff. [67] V. Gutmann, Coord. Chem. Rev. *2*, 239 (1967); ibid. *18*, 225 (1976); V. Gutmann and A. Scherhaufer, Monatsh. Chem. *99*, 335 (1968); V. Gutmann, Chimia *23*, 285 (1969); V. Gutmann, Electrochimica Acta *21*, 661 (1976). [67a] I. Lindqvist and M. Zackrisson, Acta Chem. Scand. *14*, 453 (1960). [68] U. Mayer, Pure Appl. Chem. *41*, 291 (1975). [69] V. Gutmann, Pure Appl. Chem. *27*, 73 (1971); Fortschr. Chem. Forsch. *27*, 59 (1972); Structure and Bonding *15*, 141 (1973). [70] U. Mayer, V. Gutmann, and W. Gerger, Monatsh. Chem. *106*, 1235 (1975); ibid. *108*, 489 (1977); U. Mayer, Pure Appl. Chem. *51*, 1697 (1979).

[71] R. S. Drago and B. B. Wayland, J. Am. Chem. Soc. *87*, 3571 (1965); R. S. Drago, Structure and Bonding *15*, 73 (1973); J. Chem. Educ. *51*, 300 (1974); Coord. Chem. Rev. *33*, 251 (1980); R. S. Drago: *Applications of Electrostatic-Covalent Models in Chemistry*, Surfside Publishers, Gainesville/Florida, 1994. [72] T. Kagiya, Y. Sumida, and T. Inoue, Bull. Chem. Soc. Jpn. *41*, 767 (1968). [73] For a review, see C. Agami, Bull. Soc. Chim. Fr. *1969*, 2183. [74] G. Nemethy and H. A. Scheraga, J. Chem. Phys. *36*, 3401 (1962). [75] G. Nemethy, Angew. Chem. *79*, 260 (1967); Angew. Chem. Int. Ed. Engl. *6*, 195 (1967). [76] E. M. Huque: *The Hydrophobic Effect*, J. Chem. Educ. *66*, 581 (1989). [77] T. P. Silverstein: *The Real Reason Why Oil and Water Don't Mix*. J. Chem. Educ. *75*, 116 (1998). [78] B. R. Baker, J. Chem. Educ. *44*, 610 (1967). [79] R. Cecil, Nature *214*, 369 (1967). [79a] R. D. Cramer, J. Am. Chem. Soc. *99*, 5408 (1977). [80] A. Ben-Naim, J. Chem. Phys. *54*, 1387 (1971); esp. footnote 5 on p. 1404.

[81] O. Wörz and G. Scheibe, Z. Naturforsch. *24*b, 381 (1969). [82] A. Ray, Nature *231*, 313 (1971). [83] S. N. Timasheff, Acc. Chem. Res. *3*, 62 (1970). [84] H. Zimmermann, Chemie in unserer Zeit *4*, 69 (1970). [85] O. Ya. Samoilov: *Structure of Aqueous Electrolyte Solutions and the Hydration of Ions*, Consultants Bureau, New York, Plenum Press, London 1965. [86] K. P. Mishchenko and G. M. Poltoratskii: *Problems of Thermodynamics and the Structure of Water and Non-Aqueous Electrolyte Solutions*, Khimiya, Leningrad, 1968; *Thermodynamics and Structure of Electrolyte Solutions*, Plenum Press, London, 1972. [87] G. Zundel: *Hydration and Intermolecular Interaction*, Academic Press, New York, London, 1969; Angew. Chem. *81*, 507 (1969); Angew. Chem. Int. Ed. Engl. *8*, 499 (1969). [88] J. F. Coetzee and C. D. Ritchie (eds.): *Solute–Solvent Interactions*, Vols. 1 and 2, Dekker, New York, London, 1969/1976. [89] E. S. Amis: *Solvent Composition and Chemical Phenomena*, Inorg. Chim. Acta Rev. *3*, 7 (1969). [90] L. P. Hammett: *Physical Organic Chemistry*, 2nd ed., McGraw-Hill, New York, 1970; *Physikalische Organische Chemie*, Verlag Chemie, Weinheim, 1973.

[91] H. G. Hertz: *Die Struktur der Solvathülle gelöster Teilchen*, Angew. Chem. *82*, 91 (1970); Angew. Chem. Int. Ed. Engl. *9*, 124 (1970). [92] M. J. Blandamer: *Structure and Properties of Aqueous Salt Solutions*, Quart. Rev. (London) *24*, 169 (1970). [93] A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973. [94] E. S. Amis and J. F. Hinton: *Solvent Effects on Chemical Phenomena*, Vol. 1, Academic Press, New York, London, 1973. [95] A. Ben-Naim: *Solvation Thermodynamics*, Plenum, New York, London, 1987. [96] J. E. Gordon: *The Organic Chemistry of Electrolyte Solutions*, Wiley, New York, 1975. [97] E. S. Amis: *Solvation of Ions*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. III, Part 1 of the series "Techniques of Chemistry", Wiley-Interscience, New York, 1975. [98] P. Schuster, W. Jakubetz, and W. Marius: *Molecular Models for the Solvation of Small Ions and Polar Molecules*, Top. Curr. Chem. *60*, 1 (1975). [98a] G. A. Krestov: *Thermodynamics of Solvation, Solution, and Dissolution; Ions and Solvents; Structure* and *Energetics*, Ellis Horwood,

New York, 1991; G. A. Krestov *et al.* (eds.): *Ionic Solvation*, Ellis Horwood, New York, 1994. [99] K. L. Wolf: *Theoretische Chemie*, 4th ed., J. A. Barth, Leipzig, 1959, p. 592, Fig. 283. [100] R. Daudel: *Quantum Theory of Chemical Reactivity*, Reidel, Dordrecht, Boston, 1973, p. 81ff.

[101] M. Davies, J. Chem. Educ. *48*, 591 (1971). [102] P. Sen and S. Basu, J. Chem. Phys. *48*, 4075 (1968). [103] S. F. Lincoln: *Solvent Coordination Numbers of Metal Ions in Solution*, Coord. Chem. Rev. *6*, 309 (1971). [104] J. F. Hinton and E. S. Amis: *Solvation Number of Ions*, Chem. Rev. *71*, 627 (1971). [105] J. A. Jackson, J. F. Lemons, and H. Taube, J. Chem. Phys. *32*, 553 (1960); M. Alei and J. A. Jackson, ibid. *41*, 3402 (1964). [106] J. F. Hinton and E. S. Amis: *NMR Studies of Ions in Pure and Mixed Solvents*, Chem. Rev. *67*, 367 (1967). [107] A. Fratiello: *Nuclear Magnetic Resonance Cation Solvation Studies*, Progr. Inorg. Chem. *17*, 57 (1972). [108] J. Burgess and M. C. R. Symons: *The Study of Ion–Solvent and Ion–Ion Interactions by Magnetic Resonance Techniques*, Quart. Rev. Chem. Soc. *22*, 276 (1968); M. C. R. Symons: *Spectroscopic Studies of Solvation*, Annu. Rep. Progr. Chem. Sect. A *73*, 91 (1976). [109] C. Deverell: *NMR Studies of Electrolyte Solutions*, Progr. Nucl. Magn. Reson. *4*, 235 (1969). [110] H. G. Hertz: *Magnetische Kernresonanzuntersuchungen zur Struktur von Elektrolytlösungen*, in H. Falkenhagen and W. Ebeling (eds.): *Theorie der Elektrolyte*, Hirzel, Stuttgart, 1971.

[111] A. I. Popov: *The Use of Alkali Metal Nuclear Magnetic Resonance in the Study of Solvation and Complexation of Alkali Metal Ions*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute–Solvent Interactions*, Dekker, New York, London, 1976, Vol. 2, p. 271ff.; A. I. Popov: *Alkali Metal, Magnesium-25, and Silver-109 NMR Studies of Complex Compounds in Nonaqueous Solvents*, in G. Mamantov (ed.): *Characterization of Solutes in Non-Aqueous Solvents*, Plenum Publ. Corp., New York, 1978. [111a] P. Laszlo: *Kernresonanzspektroskopie mit Natrium-23*, Angew. Chem. *90*, 271 (1978); Angew. Chem. Int. Ed. Engl. *17*, 254 (1978). [112] N. A. Matwiyoff, P. E. Darley, and W. G. Movius, Inorg. Chem. *7*, 2173 (1968). [113] J. C. Boubel, J. J. Delpuech, M. R. Khaddar, and A. Peguy, J. Chem. Soc. Chem. Commun. 1265 (1971). [114] J. F. Hinton and R. W. Briggs, J. Magn. Reson. *19*, 393 (1975). [115] Y. M. Cahen, P. R. Handy, E. T. Roach, and A. I. Popov, J. Phys. Chem. *79*, 80 (1975). [116] R. W. Gurney: *Ionic Processes in Solution*, Dover, New York, 1962. [116a] G. W. Stewart and R. M. Morrow, Proc. Nat. Acad. Sci. Washington *13*, 222 (1927); Chem. Zbl. *1927 II*, 371; G. W. Stewart, Phys. Rev. [2] *32*, 558 (1928); Chem. Zbl. *1929 I*, 17. [116b] J. R. Partington: *An Advanced Treatise on Physical Chemistry*, Longmans, Green, and Co., London, 1951, Vol. 2, p. 2. [117] G. Engel and H. G. Hertz, Ber. Bunsenges. Phys. Chem. *72*, 808 (1968). [118] H. Schneider: *The Selective Solvation of Ions in Mixed Solvents*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute–Solvent Interactions*, Dekker, New York, London, 1969/1976, Vol. 1, p. 301ff., Vol. 2, p. 155ff.; H. Schneider: *Ion Solvation in Mixed Solvents*, Top. Curr. Chem. *68*, 103 (1976). [119] H. Strehlow and H. Schneider: *Solvation of Ions in Pure and Mixed Solvents*, Pure Appl. Chem. *25*, 327 (1971); H. Strehlow, W. Knoche, and H. Schneider: *Ionensolvatation*, Ber. Bunsenges. Phys. Chem. *77*, 760 (1973). [120] J. I. Padova: *Solvation in Nonaqueous and Mixed Solvents*, Mod. Aspects Electrochem. *7*, 1 (1972); Chem. Abstr. *79*, 77566c (1973).

[121] H. Strehlow and H.-M. Koepp, Z. Elektrochem., Ber. Bunsenges. Phys. Chem. *62*, 373 (1958); H. Schneider and H. Strehlow, Z. Phys. Chem. (Frankfurt) *49*, 44 (1966). [122] H. Schneider and H. Strehlow, Z. Elektrochem., Ber. Bunsenges. Phys. Chem. *66*, 309 (1962). [123] H. Schneider and H. Strehlow, Ber. Bunsenges. Phys. Chem. *69*, 674 (1965). [123a] L. Rodehüser and H. Schneider, Z. Phys. Chem. (Frankfurt) *100*, 119 (1976). [124] K. Dimroth and C. Reichardt, Z. Anal. Chem. *215*, 344 (1966); Z. B. Maksimović, C. Reichardt, and A. Spirić, ibid. *270*, 100 (1974). [125] L. S. Frankel, T. R. Stengle, and C. H. Langford, Chem. Commun. 393 (1965); 393; Can. J. Chem. *46*, 3183 (1968); J. Phys. Chem. *74*, 1376 (1970). [126] R. H. Erlich, M. S. Greenberg, and A. I. Popov, Spectrochim. Acta, Part A *29*, 543 (1973). [127] R. D. Gillard, E. D. McKenzie, and M. D. Ross, J. Inorg. Nucl. Chem. *28*, 1429 (1966). [128] M. E. L. McBain and E. Hutchison: *Solubilization and Related Phenomena*, Academic Press, New York, 1955, Chapter 6. [129] E. M. Kosower: *An Introduction to Physical Organic Chemistry*, Wiley, New York, 1968. [130] F. M. Menger: *Aggregate organischer Moleküle mit Kollektiveigenschaften*, Angew. Chem. *103*, 1104 (1991); Angew. Chem. Int. Ed. Engl. *30*, 1086 (1991).

[131] Y. Moroi: *Micelles – Theoretical and Applied Aspects*, Plenum, New York, 1992. [132] J. H. Fendler: *Membrane Mimetic Chemistry*, Wiley-Interscience, New York, 1982. [133] S. H. Yalkow-

sky: *Solubility and Solubilization in Aqueous Media*, Oxford University Press, New York, 1999. [134] R. Fuoss, J. Chem. Educ. *32*, 527 (1955); R. M. Fuoss and F. Accascina: *Electrolytic Conductance*, Wiley-Interscience, New York, 1959. [135] E. Price: *Solvation of Electrolytes and Solution Equilibria*, in J. J. Lagowski (ed.): *The Chemistry of Non-Aqueous Solvents*, Academic Press, New York, London, 1966, Vol. I, p. 67ff. [136] J. Barthel: *Leitfähigkeit von Elektrolytlösungen*, Angew. Chem. *80*, 253 (1968); Angew. Chem. Int. Ed. Engl. *7*, 260 (1968); J. Barthel: *Ionen in nichtwäßrigen Lösungen*, Steinkopff, Darmstadt, 1976. [137] V. Gutmann, Chimia *23*, 285 (1969); Angew. Chem. *82*, 858 (1970); Angew. Chem. Int. Ed. Engl. *9*, 843 (1970); Chemie in unserer Zeit *4*, 90 (1970); Chemistry in Britain *7*, 102 (1971); Chimia *31*, 1 (1977). [138] M. Szwarc (ed.): *Ions and Ion Pairs in Organic Reactions*, Wiley-Interscience, New York, 1972/1974, Vols. 1 and 2; M. Szwarc, Acc. Chem. Res. *2*, 87 (1969). [139] U. Mayer: *Solvent Effects on Ion-Pair Equilibria*, Coord. Chem. Rev. *21*, 159 (1976). [140] D. N. Glew and D. A. Hames, Can. J. Chem. *49*, 3114 (1971).

[141] J. Smid: *Die Struktur solvatisierter Ionenpaare*, Angew. Chem. *84*, 127 (1972); Angew. Chem. Int. Ed. Engl. *11*, 112 (1972). [141a] W. Nernst, Nachrichten der Akademie der Wissenschaften zu Göttingen, Mathematisch-naturwissenschaftliche Klasse *1893*, 491; Z. Phys. Chem. *13*, 531 (1894). [141b] J. J. Thomson, Philosophical Magazine (London) *36*, 313 (1893). [141c] E. Beckmann, Z. Phys. Chem. *6*, 437 (1890); esp. p. 470. [142] M. B. Reynolds and C. A. Kraus, J. Am. Chem. Soc. *70*, 1709 (1948). [143] C. C. Evans and S. Sugden, J. Chem. Soc. *1949*, 270. [144] S. Winstein, L. G. Savedoff, S. Smith, I. D. R. Stevens, and J. S. Gall, Tetrahedron Lett. *1960*, No. 9, p. 24. [145] H. Normant, Angew. Chem. *79*, 1029 (1967); Angew. Chem. Int. Ed. Engl. *6*, 1046 (1967); Bull. Soc. Chim. Fr. *1968*, 791; Usp. Khim. *39*, 990 (1970). [146] F. Madaule-Aubry, Bull. Soc. Chim. Fr. *1966*, 1456. [147] P. Walden, Ber. Dtsch. Chem. Ges. *35*, 2018 (1902). [148] N. N. Lichtin: *Ionization and Dissociation Equilibria in Solution in Liquid Sulfur Dioxide*, Progr. Phys. Org. Chem. *1*, 75 (1963). [149] W. Karcher and H. Hecht: *Chemie in flüssigem Schwefeldioxid*, in G. Jander, H. Spandau, and C. C. Addison (eds.): *Chemie in nichtwäßrigen ionisierenden Lösungsmitteln*, Vieweg, Braunschweig, Pergamon Press, Oxford, 1967, Vol. III, Part 2, p. 79ff. [150] H. H. Freedman: *Arylcarbonium Ions*, in G. A. Olah and P. v. R. Schleyer (eds.): *Carbonium Ions*, Wiley-Interscience, New York, 1973, Vol. IV, p. 1501ff.

[151] E. Price and N. N. Lichtin, Tetrahedron Lett. *1960*, No. 18, p. 10. [152] M. Baaz, V. Gutmann, and O. Kunze, Monatsh. Chem. *93*, 1142 (1962). [153] Private communication from R. Waack and M. A. Doran to H. H. Freedman, cited in [150]. [154] A. G. Evans, A. Price, and J. H. Thomas, Trans. Faraday Soc. *52*, 332 (1956). [155] Y. Pocker, J. Chem. Soc. *1958*, 240. [156] A. G. Evans, A. Price, and J. H. Thomas, Trans. Faraday Soc. *51*, 481 (1955). [157] A. G. Evans, I. H. McEwan, A. Price, and J. H. Thomas, J. Chem. Soc. *1955*, 3098. [158] U. Mayer and V. Gutmann, Monatsh. Chem. *102*, 148 (1971). [159] B. Chevrier, J. M. Le Carpentier, and R. Weiss, J. Am. Chem. Soc. *94*, 5718 (1972); B. Chevrier and R. Weiss, Angew. Chem. *86*, 12 (1974); Angew. Chem. Int. Ed. Engl. *13*, 1 (1974). [160] D. Cook, Can. J. Chem. *37*, 48 (1959).

[161] For reviews, see: (a) I. P. Beletskaya: *Ions and Ion Pairs in Nucleophilic Aliphatic Substitution*, Usp. Khim. *44*, 2205 (1975); Russ. Chem. Rev. *44*, 1067 (1975); (b) K. Okamoto: *Generation and Ion-Pair Structures of Unstable Carbocation Intermediates in Solvolytic Reactions*, in X. Creary (ed.): *Advances in Carbocation Chemistry*, Vol. 1, p. 171ff., JAI Press, Greenwich/USA, and London/U.K., 1989; (c) M. B. Smith and J. March: *March's Advanced Organic Chemistry – Reactions, Mechanisms, and Structure*, 5th ed., Chapter 10, p. 397ff., Wiley-Interscience, New York, 2001. [162] S. Winstein, E. Clippinger, A. H. Fainberg, and G. C. Robinson, J. Am. Chem. Soc. *76*, 2597 (1954) ibid. *78*, 328 (1956). [163] H. Sadek and R. M. Fuoss, J. Am. Chem. Soc. *76*, 5897, 5905 (1954). [164] T. E. Hogen-Esch and J. Smid, J. Am. Chem. Soc. *88*, 307, 318 (1966). [165] G. A. Olah and P. v. R. Schleyer (eds.): *Carbonium Ions*, Vols. I–IV, Wiley-Interscience, New York, 1968–1973. [166] J. F. Garst: *Organoalkali Compounds in Ethers*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute–Solvent Interactions*, Dekker, New York, London, 1969, Vol. 1, p. 539ff. [167] H. F. Ebel: *Struktur und Reaktivität von Carbanionen und carbanionoiden Verbindungen*, Fortschr. Chem. Forsch. *12*, 387 (1969). [168] M. Schlosser: *Struktur und Reaktivität polarer Organometalle*, Springer, Berlin, 1973, p. 26ff. [168a] T. E. Hogen-Esch: *Ion-Pairing Effects in Carbanion Reactions*, Adv. Phys. Org. Chem. *15*, 153 (1978). [169] N. M. Atherton and S. I. Weissman, J. Am. Chem. Soc. *83*, 1330 (1961). [170] P. Chang, R. V. Slates, and M. Szwarc, J. Phys. Chem. *70*, 3180 (1966).

[171] G. Jancsó and D. V. Fenby: *Thermodynamics of Dilute Solutions*, J. Chem. Educ. *60*, 382 (1983). [172] M. D. Zeidler: *Struktur einfacher molekularer Flüssigkeiten*, Angew. Chem. *92*, 700 (1980); Angew. Chem. Int. Ed. Engl. *19*, 697 (1980); *cf.* also Z. Phys. Chem. N. F. *133*, 1 (1982). [173] J. A. Barker and D. Henderson: *The Fluid Phases of Matter*, Scientific American *254*, 94 (1981); *Einheitliche Theorie der Flüssigkeiten und Gase*, Spektrum der Wissenschaft, Januar 1982, p. 80ff. [174] J. N. Murrell and A. D. Jenkins: *Properties of Liquids and Solutions*, 2nd ed., Wiley, Chichester, 1994. [175] F. Franks: *Polywater*, MIT Press, Cambridge/Mass., London 1982; *Polywasser – Betrug oder Irrtum in der Wissenschaft*, Vieweg, Braunschweig, Wiesbaden, 1984. [176] F. Franks: *Water – A Matrix of Life*, 2nd ed., Royal Society of Chemistry, Cambridge/U.K., 2000. [177] W. A. P. Luck: *Modellbetrachtung von Flüssigkeiten mit Wasserstoffbrücken*, Angew. Chem. *92*, 29 (1980); Angew. Chem. Int. Ed. Engl. *19*, 28 (1980). [178] M. C. R. Symons: *Structure in Solvents and Solutions*, Chem. Soc. Rev. *12*, 1 (1983). [179] P. Huyskens: *Molecular Structure of Liquid Alcohols*, J. Mol. Struct. *100*, 403 (1983). [180] J. H. Hildebrand, J. Phys. Colloid Chem. *53*, 944 (1949); Chem. Abstr. *43*, 7285c (1949).

[181] S. L. Kittsley and H. A. Goeden, J. Am. Chem. Soc. *72*, 4841 (1950). [182] G. C. Maitland, M. Rigby, E. B. Smith, and W. A. Wakeham: *Intermolecular Forces – Their Origin and Determination*, Oxford University Press, Oxford, 1987. [183] H. Ratajczak and W. J. Orville-Thomas (eds.): *Molecular Interactions*, Vols. 1–3, Wiley, New York, 1980–1982. [184] H. Gnamm and O. Fuchs: *Lösungsmittel und Weichmachungsmittel*, 8th edition, Wissenschaftliche Verlagsgesellschaft, Stuttgart, 1980, Vol. I, Chapter 2, p. 9ff. [185] N. H. March and M. P. Tosi: *Coulomb Liquids*, Academic Press, New York, London, 1984. [186] J. H. Mahanty and B. W. Ninham: *Dispersion Forces*, Academic Press, New York, London, 1981. [187] S. Scheiner: *Hydrogen Bonding – A Theoretical Perspective*, Oxford University Press, Oxford/U.K., 1997. [188] I. Alkorta, I. Rozas, and J. Elguero: *Non-Conventional Hydrogen Bonds*, Chem. Soc. Rev. *27*, 163 (1998). [189] G. Desiraju and T. Steiner: *The Weak Hydrogen Bond in Structural Chemistry and Biology*, Oxford University Press, Oxford/U.K., 1999. [190] P. Hobza and Z. Havlas: *Blue-Shifting Hydrogen Bonds*, Chem. Rev. *100*, 4253 (2000); see also S. Scheiner and T. Kar, J. Phys. Chem. A *106*, 1784 (2002).

[191] W. M. Latimer and W. H. Rodebush, J. Am. Chem. Soc. *42*, 1419 (1920); *cf.* p. 1431. [192] N. Isenberg, J. Chem. Educ. *59*, 547 (1982). [193] W. Saenger, Nature (London) *279*, 343 (1979); W. Saenger and K. Lindner, Angew. Chem. *92*, 404 (1980); Angew. Chem. Int. Ed. Engl. *19*, 398 (1980); *cf.* also Nachr. Chem. Techn. Labor. *27*, 403 (1979). [194] R. Taylor and O. Kennard: *Hydrogen-Bond Geometry in Organic Crystals*, Acc. Chem. Res. *17*, 320 (1984). [195] M. Rospenk, J. Fritsch, and G. Zundel, J. Phys. Chem. *88*, 321 (1984). [196] M. J. Kamlet and R. W. Taft, J. Am. Chem. Soc. *98*, 377, 2886 (1976). [197] M. Mashima, R. T. McIver, R. W. Taft, F. G. Bordwell, and W. N. Olmstead, J. Am. Chem. Soc. *106*, 2717 (1984) (*cf.* footnote 12). [198] B. D. Gosh and R. Basu, J. Phys. Chem. *84*, 1887 (1980). [199] R. Schmid, J. Sol. Chem. *12*, 135 (1983). [200] Y. Marcus, J. Sol. Chem. *13*, 599 (1984).

[201] U. Mayer and V. Gutmann, Structure and Bonding *12*, 113 (1972); *cf.* p. 125. [202] A. I. Popov, Pure Appl. Chem. *41*, 275 (1975), and references cited therein. [203] T. Ogata, T. Fujisawa, N. Tanaka, and H. Yokoi, Bull. Chem. Soc. Jpn. *49*, 2759 (1976). [204] R. W. Soukop: *Farbreaktionen zur Klassifizierung von Lösungsmitteln*, Chemie in unserer Zeit *17*, 129, 163 (1983); R. W. Soukop and R. Schmid: *Metal Complexes as Color Indicators for Solvent Parameters*, J. Chem. Educ. *62*, 459 (1985). [205] R. Schmid, Rev. Inorg. Chem. *1*, 117 (1979); Chem. Abstr. *92*, 117098d (1980). [206] R. Schmid and V. N. Sapunov: *Non-Formal Kinetics in Search for Chemical Reaction Pathways*, Verlag Chemie, Weinheim, 1982. [207] V. Gutmann and G. Resch: *The Unifying Impact of the Donor–Acceptor Approach*, in N. Tanaka, H. Ohtaki, and R. Tamamushi (eds.): *Ions and Molecules in Solution*, Elsevier, Amsterdam, 1982, p. 203ff. [208] R. S. Drago, Coord. Chem. Rev. *33*, 251 (1980); Pure Appl. Chem. *52*, 2261 (1980). [209] R. W. Taft, N. J. Pienta, M. J. Kamlet, and E. M. Arnett, J. Org. Chem. *46*, 661 (1981). [210] G. Olofsson and I. Olofsson, J. Am. Chem. Soc. *95*, 7231 (1973).

[211] L. Elégant, G. Fratini, J.-F. Gal, and P.-C. Maria, J. Calorim. Anal. Therm. *11*, 3/21/1–3/21/9 (1980); Chem. Abstr. *95*, 121811k (1981). [212] P.-C. Maria and J.-F. Gal, J. Phys. Chem. *89*, 1296 (1985); P.-C. Maria, J.-F. Gal, J. de Franceschi, and E. Fargin, J. Am. Chem. Soc. *109*, 483

(1987). [213] U. Mayer: *NMR Spectroscopic Studies on Solute–Solvent and Solute–Solute Interactions*, in N. Tanaka, H. Ohtaki, and R. Tamamushi (eds.): *Ions and Molecules in Solution*, Elsevier, Amsterdam, 1983, p. 219ff. [214] J.-C. Bollinger, G. Yvernault, and T. Yvernault, Thermochim. Acta *60*, 137 (1983); Chem. Abstr. *98*, 118416s (1983). [215] R. S. Drago, G. C. Vogel, and T. E. Neddham, J. Am. Chem. Soc. *93*, 6014 (1971). [216] M. K. Kroeger and R. S. Drago, J. Am. Chem. Soc. *103*, 3250 (1981). [217] R. S. Drago: *The Coordination Model for Non-Aqueous Solvents*, Pure Appl. Chem. *52*, 2261 (1980). [218] R. S. Drago, M. K. Kroeger, and J. R. Stahlbush, Inorg. Chem. *20*, 306 (1981). [219] P. E. Doan and R. S. Drago, J. Am. Chem. Soc. *104*, 4524 (1982). [220] C. Tanford: *The Hydrophobic Effect*, 2nd ed., Wiley-Interscience, New York, 1980.

[221] A. Ben-Naim: *Hydrophobic Interactions*, Plenum, New York, 1980. [222] J.-Y. Huot and C. Jolicoeur: *Hydrophobic Effects in Ionic Hydration and Interactions*, in R. R. Dogonadze, E. Kálmán, A. A. Kornyshev, and J. Ulstrup (eds.): *The Chemical Physics of Solvation*, Part A, Elsevier, Amsterdam, 1985. [223] H. van de Waterbeemd: *Hydrophobicity of Organic Compounds*, Compudrug International, Wien, 1986. [224] L. R. Pratt: *Theory of Hydrophobic Effects*, Ann. Rev. Phys. Chem. *36*, 433 (1985). [225] W. Blokzijl and J. B. F. N. Engberts: *Hydrophobic Effects: Opinions and Facts*, Angew. Chem. *105*, 1610 (1993); Angew. Chem. Int. Ed. Engl. *32*, 1545 (1993). [226] H. S. Frank and M. W. Evans, J. Phys. Chem. *13*, 507 (1945). [227] D. F. Evans, S.-H. Chen, G. W. Schriver, and E. M. Arnett, J. Am. Chem. Soc. *103*, 481 (1981); D. Mirejovsky and E. M. Arnett, ibid. *105*, 1112 (1983). [228] F. A. Greco, J. Phys. Chem. *88*, 3132 (1984). [229] J. H. Hildebrand: *Is there a "hydrophobic" effect*? Proc. Natl. Acad. Sci. U.S.A. *76*, 194 (1979); Chem. Abstr. *90*, 134237z (1979). [230] M. H. Abraham, J. Am. Chem. Soc. *102*, 5910 (1980).

[231] A. Hvidt, Acta Chem. Scand., Part A *37*, 99 (1983); and references cited therein. [232] L. R. Pratt and A. Pohorille, Chem. Rev. *102*, 2671 (2002). [233] J. Burgess: *Ions in Solution – Basic Principles of Chemical Interaction*, 2nd ed., Horwood Publishing, Westergate/U.K., 1999. [234] B. G. Cox and W. E. Waghorne: *Thermodynamics of Ion-Solvent Interactions*, Chem. Soc. Rev. *9*, 381 (1980). [235] O. Popovych and R. P. T. Tomkins: *Nonaqueous Solution Chemistry*, Wiley-Interscience, New York, 1981. [236] B. E. Conway: *Ionic Hydration in Chemistry and Biophysics*, Elsevier, Amsterdam, 1981. [237] K. Burger: *Solvation, Ionic and Complex Formation Reactions in Non-Aqueous Solvents*, Elsevier, Amsterdam, 1983. [238] R. R. Dogonadze, E. Kálmán, A. A. Kornyshev, and J. Ulstrup (eds.): *The Chemical Physics of Solvation*, Part A: *Theory of Solvation*; Part B: *Spectroscopy of Solvation*; Part C: *Solvation Phenomena in Specific Physical, Chemical, and Biological Systems*. Elsevier, Amsterdam 1985–1988. [239] J. M. G. Barthel, H. Krienke, and W. Kunz: *Physical Chemistry of Electrolyte Solutions – Modern Aspects*, Top. Phys. Chem. *5*, 1 (1998). [240] N. Tanaka, H. Ohtaki, and R. Tamamushi (eds.): *Ions and Molecules in Solution*, Elsevier, Amsterdam, 1983.

[241] Y. Marcus: *Ion Solvation*, Wiley, Chichester, 1985. [242] Y. Marcus, Pure Appl. Chem. *54*, 2327 (1982). [243] R. Schurhammer and G. Wipff, J. Phys. Chem. A *104*, 11159 (2000). [244] Y. Marcus, Rev. Anal. Chem. *5*, 53 (1980); Pure Appl. Chem. *55*, 977 (1983); ibid. *57*, 1103 (1985); ibid. *58*, 1721 (1986); ibid. *62*, 899 (1990). [245] G. A. Vidulich and A. Fratiello, J. Chem. Educ. *55*, 672 (1978). [246] P. Laszlo and A. Stockis, J. Am. Chem. Soc. *102*, 7818 (1980) (^{59}Co NMR). [247] P. S. Pregosin: *Platinium-195 NMR*, Coord. Chem. Rev. *44*, 247 (1982). [248] B. E. Conway: *Local Changes of Solubility induced by Electrolytes: Salting-out and Ionic Hydration*, Pure Appl. Chem. *57*, 263 (1985). [249] H. Langhals: *Polarität binärer Flüssigkeitsgemische*, Angew. Chem. *94*, 739 (1982); Angew. Chem. Int. Ed. Engl. *21*, 724 (1982); H. Langhals: *Description of Properties of Binary Solvent Mixtures*, in R. I. Zalewski, T. M. Krygowski, and J. Shorter (eds.): *Similarity Models in Organic Chemistry, Biochemistry, and Related Fields*, Elsevier, Amsterdam, 1991, Chapter 6, p. 283ff. [250] S. Janardhanan and C. Kalidas: *Preferential Solvation of Ions in Binary Solvent Mixtures*, Rev. Inorg. Chem. *6*, 101 (1984).

[251] T. Ichikawa, H. Yoshida, A. S. W. Li, and L. Kevan, J. Am. Chem. Soc. *106*, 4324 (1984). [252] D. S. Gill, N. Kumari, and M. S. Chauhan, J. Chem. Soc., Faraday Trans. I *81*, 687 (1985). [253] A. J. Stace, J. Am. Chem. Soc. *106*, 2306 (1984). [254] H. Strehlow and G. Busse, Ber. Bunsenges. Phys. Chem. *88*, 467 (1984). [255] K. Remerle and J. B. F. N. Engberts, J. Phys. Chem. *87*, 5449 (1983). [256] G. González and N. Yutronic, Spectrochim. Acta Part A *39*, 269 (1983). [257] A.

Ben-Naim, J. Phys. Chem. *93*, 3809 (1989); Pure Appl. Chem. *62*, 25 (1990). [258] M. wa Muanda, J. B. Nagy, and O. B. Nagy, Tetrahedron Lett. *1974*, 3421; J. Chem. Soc., Faraday Trans. I *74*, 2210 (1978); J. Phys. Chem. *83*, 1961 (1979). [259] A. K. Covington and K. E. Newman, J. Chem. Soc., Faraday Trans. I *84*, 1393 (1988). [260] J. G. Dawber, S. Etemad, and M. A. Beckett, J. Chem. Soc., Faraday Trans. *86*, 3725 (1990).

[261] Y. Marcus, J. Chem. Soc., Faraday Trans. *87*, 1843 (1991), and previous papers. [262] K. L. Mittal *et al.* (eds.): *Surfactants in Solution*, Vols. 1–11, Plenum, New York, 1984–1992. [263] J. C. Brackmant and J. B. F. N. Engberts, Chem. Soc. Rev. *22*, 85 (1993). [264] J. N. Israelachvili: *Intermolecular and Surface Forces*, 2nd ed., Academic Press, London, New York, 1992; Chapters 16 and 17, p. 341ff. and p. 366ff. [265] K. Kosswig: *Surfactants*, in *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed., VCH, Weinheim, 1994, Vol. A 25, p. 747ff. [266] M. N. Jones and D. Chapman: *Micelles, Monolayers, and Biomembranes*, Wiley-Liss, New York, 1994. [267] D. O. Shah (ed.): *Micelles, Microemulsions and Monolayers – Science and Technology*, Dekker, New York, 1998. [268] L. J. Cline Love, J. G. Habarts, and J. G. Dorsey, Anal. Chem. *56*, 1132A (1984). [269] F. M. Menger: *On the Structure of Micelles*, Acc. Chem. Res. *12*, 111 (1979); F. M. Menger, R. Zana, and B. Lindman, J. Chem. Educ. *75*, 115 (1988). [270] C. W. von Nägeli and S. Schwendener: *Das Mikroskop. Theorie und Anwendung desselben*, 2nd edition, Verlag Engelmann, Leipzig, 1877; C. W. von Nägeli: *Theorie der Gärung*, Verlag Oldenbourg, München, 1879; C. W. von Nägeli: *Die Micellartheorie*, in A. Frey (ed.): *Ostwalds Klassiker der exakten Naturwissenschaften*, Vol. 227, Akademische Verlagsgesellschaft, Leipzig, 1928.

[271] F. Krafft, Ber. Dtsch. Chem. Ges. *29*, 1334 (1896). [272] A. Reychler, Zeitschrift für Chemie und Industrie der Kolloide *12*, 277 (1913); Chem. Zbl. *1913*, II 1, 491. [273] P. Fromherz, Ber. Bunsenges. Phys. Chem. *85*, 891 (1981). [274] J. K. Thomas, Chem. Rev. *80*, 283 (1980). [275] D. G. Whitten, J. C. Russell, and R. H. Schmehl, Tetrahedron *38*, 2455 (1982). [276] J. H. Fendler, J. Chem. Educ. *60*, 872 (1983). [277] C. A. Bunton: *Reactions in Micelles and Similar Self-Organized Aggregates*, in M. I. Page (ed.): *The Chemistry of Enzyme Action*, Chapter 13, p. 461ff., Elsevier, Amsterdam, 1984. [278] J. Smid (ed.): *Ions and Ion Pairs and Their Role in Chemical Reactions*, Pergamon Press, Oxford, 1979; *cf.* also Pure Appl. Chem. *51*, p. 49ff. (1979). [279] M. Simonetta: *Structure of Ion Pairs in Solution*. Int. Rev. Phys. Chem. *1*, 31 (1981); Chem. Abstr. *95*, 176553h (1981); *cf.* also Chem. Soc. Rev. *13*, 1 (1984). [280] N. Bjerrum: *Untersuchungen über Ionenassoziation*, Kong. Danske Vidensk. Meddelelser. Math.-phys. Kl. 7, Nr. 9, p. 3–48 (1926); Chem. Zentralbl. *1926* II, 1378; Svensk Kem. Tidskr. *38*, p. 2–18 (1926); Chem. Zentralbl. *1926* I, 2174.

[281] V. Gutmann: *The Donor–Acceptor Approach to Molecular Interactions*, Plenum Press, New York, 1978. [282] M. Feigel and H. Kessler, Chem. Ber. *112*, 3715 (1979); Acc. Chem. Res. *15*, 2 (1980). [283] M. Feigel, H. Kessler, D. Leibfritz, and A. Walter, J. Am. Chem. Soc. *101*, 1943 (1979). [284] E. B. Troughton, K. E. Molter, and E. M. Arnett, J. Am. Chem. Soc. *106*, 6726 (1984); E. M. Arnett and K. E. Molter, Acc. Chem. Res. *18*, 339 (1985); J. Phys. Chem. *90*, 383 (1986). [285] M. Born, Z. Phys. *1*, 45 (1920). [286] P. Müller (ed.): *Glossary of Terms Used in Physical Organic Chemistry – IUPAC Recommendations 1994*, Pure Appl. Chem. *66*, 1077 (1994). [287] E. Buncel and B. Menon, J. Org. Chem. *44*, 317 (1979). [288] G. Boche and F. Heidenhain, J. Am. Chem. Soc. *101*, 738 (1979); G. Boche, F. Heidenhain, W. Thiel, and R. Eiben, Chem. Ber. *115*, 3167 (1982). [289] W. Bauer and D. Seebach, Helv. Chim. Acta *67*, 1972 (1984). [290] H. J. Reich, J. P. Borst, R. R. Dykstra, and D. P. Green, J. Am. Chem. Soc. *115*, 8728 (1993).

[291] E. M. Arnett and R. A. Flowers, Chem. Soc. Rev. *22*, 9 (1993). [292] K. Okamoto, T. Kitagawa, K. Takeuchi, K. Komatsu, T. Kinoshita, S. Aonuma, M. Nagai, and A. Miyabo, J. Org. Chem. *55*, 996 (1990). [293] S. Hahn, W. M. Miller, R. N. Lichtenthaler, and J. M. Prausnitz, J. Sol. Chem. *14*, 129 (1985). [294] J. F. Hinton, K. R. Metz, and R. W. Briggs: *Thallium NMR Spectroscopy*, in G. A. Webb (ed.): Annual Reports on NMR Spectroscopy, Vol. 13, p. 211ff., Academic Press, London, New York, 1982. [295] J. J. Dechter: *NMR of Metal Nuclides*, Progr. Inorg. Chem. *29*, 285 (1982); Fig. 7 on p. 320. [296] C. Hansch, Acc. Chem. Res. *2*, 232 (1969); A. Leo, J. Chem. Soc., Perkin Trans. II *1983*, 825. [297] F. M. Menger and U. V. Venkataram, J. Am. Chem. Soc. *108*, 2980 (1986). [298] M. J. Kamlet and R. W. Taft, J. Org. Chem. *47*, 1734 (1982). [299] G. Heublein and D. Bauernfeind, J. Prakt. Chem. *326*, 81 (1984). [300] C. Sandorfy, R. Buchet, L. S. Lussier, P. Ménassa, and L. Wilson, Pure Appl. Chem. *58*, 1115 (1986).

[301] I. Persson, Pure Appl. Chem. *58*, 1153 (1986). [302] W. L. Jorgensen *et al.*, J. Am. Chem. Soc. *109*, 1891 (1987); ibid. *111*, 2507 (1989). [303] I. Persson, M. Sandström, and P. L. Goggin, Inorg. Chim. Acta *129*, 183 (1987); M. Sandström, I. Persson, and P. Persson, Acta Chem. Scand. *44*, 653 (1990). [304] M. Vedamuthu, S. Singh, and G. W. Robinson, J. Phys. Chem. *98*, 2222 (1994). [305] I. Ohmine and S. Saito: *Water Dynamics – Fluctuation, Relaxation, and Chemical Reactions in Hydrogen-Bonded Network Arrangements*, Acc. Chem. Res. *32*, 741 (1999). [306] J. C. Dore and J. Teixeira (eds.): *Hydrogen-Bonded Liquids*, Kluwer, Dordrecht/Netherlands, 1991. [307] P. Ball: *H_2O – A Biography of Water*, Weidenfeld and Nicolson, London, 1999. [308] H. Bertagnolli, M. Hoffmann, and M. Ostheimer, Z. Phys. Chem. N. F. *165*, 165 (1989). [309] B. G. Rao and U. C. Singh, J. Am. Chem. Soc. *112*, 3803 (1990). [310] P. Knops, N. Sendhoff, H.-B. Mekelburger, and F. Vögtle, Top. Curr. Chem. *161*, 1 (1991).

[311] R. S. Ruoff, D. S. Tse, R. Malhotra, and D. C. Lorents, J. Phys. Chem. *97*, 3379 (1993). [312] D. J. W. Grant and T. Higuchi: *Solubility Behaviour of Organic Compounds*, in W. H. Saunders and A. Weissberger (eds.): *Techniques of Chemistry*, Vol. 21, Wiley, New York, 1990. [313] S. H. Yalkowsky and S. Banerjee: *Aqueous Solubility – Methods of Estimation for Organic Compounds*, Dekker, New York, 1994. [314] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001; Chapter 5, p. 101ff.: *General Principles Governing Dissolution of Materials in Solvents*. [315] T. M. Letcher and R. Battino, J. Chem. Educ. *78*, 103 (2001). [316] W. G. Van der Sluys, J. Chem. Educ. *78*, 111 (2001). [317] J. Kochansky, J. Chem. Educ. *68*, 655 (1991). [318] M. Handa, M. Kataoka, M. Wakaumi, and Y. Sasaki, Bull. Chem. Soc. Jpn. *70*, 315 (1997). [319] J. H. Clark, C. J. Locke, and M. S. White, Spectrochim. Acta, Part A, *50A*, 891 (1994). [320] J. Emsley: *Very Strong Hydrogen Bonding*, Chem. Soc. Rev. *9*, 91 (1980).

[321] S. Scheiner and M. Cuma, J. Am. Chem. Soc. *118*, 1511 (1996). [322] N. S. Golubov, I. G. Shenderovich, S. N. Smirnov, G. S. Denisov, and H.-H. Limbach, Chem. Eur. J. *5*, 492 (1999). [323] H. Zimmermann, Angew. Chem. *76*, 1 (1964); Angew. Chem. Int. Ed. Engl. *3*, 157 (1964); Chem. Unserer Zeit *4*, 69 (1970). [324] P. P. Schmidt, J. Phys. Chem. *97*, 4249 (1993). [325] P. Jacques, J. Chem. Educ. *68*, 347 (1991). [326] V. A. Durov and I. Yu. Shilov, J. Chem. Soc., Faraday Trans. *92*, 3559 (1996). [327] D. C. Daniel and J. L. McHale, J. Phys. Chem. A *101*, 3070 (1997). [328] T. H. Grenby, Chemistry in Britain *27*, 342 (1991). [329] M. H. Abraham *et al.*, J. Chem. Soc., Perkin Trans. II, 699, 1355 (1989); ibid. 521 (1990); J. Phys. Org. Chem. *6*, 660 (1993). [330] M. H. Abraham: *Scales of Solute Hydrogen-Bonding*, Chem. Soc. Rev. *22*, 73 (1993).

[331] C. Laurence and M. Berthelot: *Observations on the Strength of Hydrogen Bonding*, Perspectives in Drug Discovery and Design *18*, 39 (2000); Chem. Abstr. *133*, 37623w (2000). [332] C. Ouvrard, M. Berthelot, and C. Laurence, J. Phys. Org. Chem. *14*, 804 (2001). [333] J.-L. M. Abboud, M. J. Kamlet, and R. W. Taft, Progr. Phys. Org. Chem. *13*, 485 (1981). [334] Y. Marcus: *The Effectiveness of Solvents as Hydrogen-Bond Donors*, J. Sol. Chem. *20*, 929 (1991). [335] C. A. Hunter, Angew. Chem. *105*, 1653 (1993); Angew. Chem. Int. Ed. Engl. *32*, 1584 (1993). [336] R. Rathore, S. V. Lindemann, and J. K. Kochi, J. Am. Chem. Soc. *119*, 9393 (1997). [337] J. C. Ma and D. A. Dougherty: *The Cation–π Interaction*, Chem. Rev. *97*, 1303 (1997). [338] J. B. Lambert, Y. Zhao, and S. M. Zhang, J. Phys. Org. Chem. *14*, 370 (2001). [339] Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998. [340] T. Chen, G. Hefter, and Y. Marcus, J. Sol. Chem. *29*, 201 (2000).

[341] W. R. Fawcett, J. Phys. Chem. *97*, 9540 (1993). [342] J.-L. M. Abboud and R. Notario, Pure Appl. Chem. *71*, 645 (1999). [343] R. Schmid: *Recent Advances in the Description of the Structure of Water, the Hydrophobic Effect, and the Like-Dissolves-Like Rule*, in W. Linert (ed.): *Highlights in Solute–Solvent Interactions*, Springer, Wien, New York, 2002, p. 59ff. [344] G. Hummer, S. Garde, A. E. Garcia, M. E. Paulaitis, and L. R. Platt: *Hydrophobic Effects on a Molecular Scale*, J. Phys. Chem. B *102*, 10469 (1998). [345] G. Graziano, Can. J. Chem. *76*, 437 (1998). [346] T. Lazaridis, Acc. Chem. Res. *34*, 931 (2001). [347] N. T. Southall, K. A. Dill, and A. D. J. Haymet, J. Phys. Chem. B *106*, 521 (2002). [348] A. M. Davis and S. J. Teague, Angew. Chem. *111*, 778 (1999), Angew. Chem. Int. Ed. Engl. *38*, 736 (1999). [349] C. Kalidas, G. Hefter, and Y. Marcus, Chem. Rev. *100*, 819 (2000). [350] G. Gritzner, Electrochim. Acta *44*, 73 (1998).

[351] H. Ohtaki: *Ionic Solvation in Aqueous and Nonaqueous Solutions*, in W. Linert (ed.): *Highlights in Solute–Solvent Interactions*, Springer, Wien, New York, 2002, p. 1ff. [352] K. Bhattacharyya and B. Bagchi, J. Phys. Chem. A *104*, 10603 (2000). [353] R. Argaman, T. Molotsky, and D. Huppert, J. Phys. Chem. A *104*, 7934 (2000). [354] A. Nitzan: *Polar Solvation Dynamics: Theory and Simulations*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001; Chapter 4.3, p. 132ff. [355] J. Tomasi and M. Persico: *Molecular Interactions in Solution*, Chem. Rev. *94*, 2027 (1994). [356] O. Tapia and J. Bertrán (eds.): *Solvent Effects and Chemical Reactivity*, Kluwer, Dordrecht/Netherlands, 1996. [357] C. J. Cramer and D. G. Truhlar: *Implicit Solvation Models: Equilibria, Structure, Spectra, and Dynamics*, Chem. Rev. *99*, 2161 (1999). [358] M. Orozco and F. J. Luque: *Theoretical Methods for the Description of the Solvent Effect on Biomolecular Systems*, Chem. Rev. *100*, 4187 (2000). [359] J. Li, T. Zhu, C. J. Cramer, and D. G. Truhlar, J. Phys. Chem. A *104*, 2178 (2000). [360] A. Papazyan and A. Warshel, J. Phys. Chem. B *102*, 5348 (1998).

[361] W. L. Jorgensen: *Computational Insights on Intermolecular Interactions and Binding in Solution*, CHEMTRACTS – Organic Chemistry *4*, 91 (1999). [362] J. M. Martinez, R. R. Pappalardo, and E. S. Marcos, J. Phys. Chem. A *101*, 4444 (1997). [363] F. J. Luque and M. Orozco, J. Phys. Chem. B *101*, 5573 (1997). [364] G. P. Das and D. S. Dudis, J. Phys. Chem. A *104*, 4767 (2000). [365] G. A. Krestov, V. N. Afanas'ev, and L. S. Efremova: *Physico-Chemical Properties of Binary Solvents*, Publisher Khimiya, Leningrad, 1988. [366] Y. Y. Fialkov and V. L. Chumak: *Mixed Solvents*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 9, p. 505ff. [367] S. J. Romanowski, C. M. Kinart, and W. J. Kinart, J. Chem. Soc., Faraday Trans. *91*, 65 (1995). [368] A. Wakisaka, H. Abdoul-Carime, Y. Yamamoto, and Y. Kiyozumi, J. Chem. Soc., Faraday Trans. *94*, 369 (1998). [369] Y. Marcus, J. Chem. Soc., Dalton Trans. 2265 (1991). [370] T. K. Varadarajan, T. V. Ramakrishna, and C. Kalidas, Ber. Bunsenges. Phys. Chem. *101*, 676 (1997).

[371] Y. Marcus, J. Chem. Soc., Perkin Trans. II, 1015, 1751 (1994). [372] A. Bagno, M. Campulla, M. Pirana, G. Scorrano, and S. Stiz, Chem. Eur. J. *5*, 1291 (1999). [373] C. Lerf and P. Suppan, J. Chem. Soc., Faraday Trans. *88*, 963 (1992); P. Suppan and N. Ghoneim: *Solvatochromism*, Royal Society of Chemistry, Cambridge/U.K., 1997, Chapter 6, p. 163ff. [374] R. D. Skwierczynski and K. A. Connors, J. Chem. Soc., Perkin Trans. II, 467 (1994). [375] W. A. Acree jr., J. R. Powell, and S. A. Tucker, J. Chem. Soc., Perkin Trans. II, 529 (1995). [376] A. K. Laha, P. K. Das, and S. Bagchi, J. Phys. Chem. A *106*, 3230 (2002). [377] J. Ortega, C. Ráfols, E. Bosch, and M. Rosés, J. Chem. Soc., Perkin Trans. II, 1607 (1995); ibid. 1497 (1996). [378] P. M. Mancini, C. Adam, A. del C. Pérez, and L. R. Vottero, J. Phys. Org. Chem. *13*, 221 (2000). [379] P. Jacques, J. Chem. Educ. *68*, 347 (1991). [380] I. A. Koppel and J. B. Koppel, Org. React. (Tartu) *20*, 523, 547 (1983); Chem. Abstr. *101*, 110180v, 110274d (1984).

[381] P. K. Kipkemboi and A. J. Easteal, Aust. J. Chem. *47*, 1771 (1994). [382] K. Herodes, I. Leito, I. Koppel, and M. Rosés, J. Phys. Org. Chem. *12*, 109 (1999). [383] E. B. Tada, L. P. Novaki, and O. A. El Seoud, J. Phys. Org. Chem. *13*, 679 (2000). [384] S. Niqam, A. de Juan, R. J. Stubbs, and S. C. Rutan, Anal. Chem. *72*, 1956 (2000). [385] M. Kemell and P. Pirilä, J. Solution Chem. *29*, 87 (2000). [386] H. Langhals, Anal. Lett. *23*, 2243 (1990); GIT Fachz. Lab. *35*, 766 (1991). [387] M. F. Vitha, J. Chem. Educ. *78*, 370 (2001). [388] C. Machado and V. G. Machado, J. Chem. Educ. *78*, 649 (2001). [389] J. Van Stam, S. Depaemelaere, and F. C. De Schryver, J. Chem. Educ. *75*, 94 (1998). [390] G. S. Hartley: *Aqueous Solutions of Paraffin-Chain Salts. A Study of Micelle Formation*, Herman, Paris, 1936; particularly Fig. 11A.

[391] Y. Moroi, K. Mitsonobu, T. Morisue, Y. Kadobayashi, and M. Sakai, J. Phys. Chem. *99*, 2372 (1995). [392] R. V. Bensasson, E. Bienvenue, M. Dellinger, S. Leach, and P. Seta, J. Phys. Chem. *98*, 3492 (1994). [393] D. T. Lai, M. A. Neumann, M. Matsumoto, and J. Sunamoto, Chem. Lett. (Tokyo) 64 (2000). [394] P. L. Luisi, J. Chem. Educ. *78*, 380 (2001). [395] M. P. Pileni: *Reversed Micelles as Microreactors*, J. Phys. Chem. 97, 6961 (1993). [396] F. M. Menger and H. Park: *Microemulsions as Reaction Media*, Recl. Trav. Chim. Pays-Bas *113*, 176 (1994). [397] E. Schultz: *Ionization or Dissociation?* J. Chem. Educ. *74*, 868 (1997). [398] S. J. Hawkes: *Salts are mostly NOT ionized*, J. Chem. Educ. *73*, 421 (1996). [399] A. Abbott: *Electrochemistry in Media of Low Dielec-*

tric Constant, Chem. Soc. Rev. *22*, 435 (1993). [400] J. Mattay and M. Vondenhof: *Contact and Solvent-Separated Radical Ion Pairs in Organic Photochemistry*, Top. Curr. Chem. *159*, 219 (1991).

[401] T. M. Bockman and J. K. Kochi: *From Solvolysis to Electron Transfer: Direct Observation of Ion-Pair Dynamics*, J. Chem. Soc., Perkin Trans. II, 1901 (1994). [402] G. Boche, Angew. Chem. *104*, 742 (1992); Angew. Chem. Int. Ed. Engl. *31*, 731 (1999). [403] A. P. Abbott and D. J. Schiffrin, J. Chem. Soc., Faraday Trans. *86*, 1453 (1990). [404] G. B. Schuster *et al.*, J. Am. Chem. Soc. *114*, 793 (1992); ibid. *115*, 11628 (1993). [405] G. H. Peslherbe, B. M. Ladanyi, and J. T. Hynes, J. Phys. Chem. A *104*, 4533 (2000). [406] L. M. Jackman and J. Bortiatynski: *Structure of Lithium Enolates and Phenolates in Solution*, in V. Snieckus (ed.): *Advances in Carbanion Chemistry*, Vol. 1, p. 45ff., JAI Press, Greenwich/USA, and London/U.K., 1992. [407] M. T. Reetz, S. Hütte, and R. Goddard, J. Am. Chem. Soc. *115*, 9339 (1993); J. Prakt. Chem. *341*, 297 (1999). [408] K. Seppelt: *'Non-coordinating' Anions*, Angew. Chem. *105*, 1074 (1993); Angew. Chem. Int. Ed. Engl. *32*, 1025 (1993). [409] R. Gschwind, P. R. Rajamohanan, M. John, and G. Boche, Organometallics *19*, 2868 (2000); G. Boche *et al.*, Chem. Eur. J. *6*, 3060 (2000). [410] C. D. Stevenson, R. M. Fico, and E. C. Brown, J. Phys. Chem. B *102*, 2841 (1998).

[411] M. Khajehpour and J. F. Kauffman, J. Phys. Chem. A *105*, 10316 (2001). [412] N. A. Izmailov, Zh. Fiz. Khim. *24*, 321 (1950); Chem. Abstr. *44*, 6706 (1950); ibid. *28*, 2047 (1954); Chem. Abstr. *50*, 7549 (1956); ibid. *30*, 2164 (1956); Chem. Abstr. *51*, 9266 (1957).

Chapter 3

[1] L. F. Audrieth and J. Kleinberg: *Non-Aqueous Solvents: Applications as Media for Chemical Reactions*, Wiley, New York, and Chapman and Hall, London, 1953. [2] H. H. Sisler: *Chemistry in Non-Aqueous Solvents*, Reinhold, New York, and Chapman and Hall, London, 1961. [3] G. Charlot and B. Trémillon: *Réactions Chimiques dans les Solvants et les Sels Fondus*, Gauthier-Villars, Paris, 1963; *Chemical Reactions in Solvents and Melts*, Pergamon Press, Oxford, 1969. [4] G. Jander, H. Spandau, and C. C. Addison (eds.): *Chemie in nichtwäßrigen Lösungsmitteln*, Vols. I–IV, Vieweg, Braunschweig, Wiley-Interscience, New York, and Pergamon Press, Oxford, 1963–1967. [5] T. C. Waddington (ed.): *Non-Aqueous Solvent Systems*, Academic Press, London, New York, 1965. [6] A. K. Holliday and A. G. Massey: *Non-Aqueous Solvents*, Pergamon Press, Oxford, 1965. [7] J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvents*, Vols. I–V, Academic Press, New York, London, 1966–1978. J. J. Lagowski: *Non-Aqueous Ionizing Liquids*, Rev. Chim. Miner. *15*, 1 (1978). [8] J. Jander and C. Lafrenz: *Wasserähnliche Lösungsmittel*, Verlag Chemie, Weinheim, 1968; *Ionizing Solvents*, Wiley, New York, London, 1968. [9] R. A. Zingaro: *Nonaqueous Solvents*, Raytheon Education Company, Lexington/Mass., 1968. [10] T. C. Waddington: *Non-Aqueous Solvents*, 2nd ed., Nelson, London, 1973; *Nichtwäßrige Lösungsmittel*, Hüthig, Heidelberg, 1972.

[11] B. P. Whim and B. P. Johnson: *Directory of Solvents*, Blackie Academic and Professional, London, 1995. [12] J. A. Riddick, W. B. Bunger, and T. Sakano: *Organic Solvents*, 4rd ed., in A. Weissberger (ed.): *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986. [13] D. R. Lide: *Handbook of Organic Solvents*, CRC Press, Boca Raton/FL, USA, 1995. [14] W. L. Archer: *Industrial Solvents Handbook*, Dekker, New York, 1996. [15] A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973. [16] V. Gutmann: *Coordination Chemistry in Non-Aqueous Solutions*, Springer, Wien, New York, 1968, p. 2 (and footnote 1). [17] A. Saupe: *Neuere Ergebnisse auf dem Gebiet der flüssigen Kristalle*, Angew. Chem. *80*, 99 (1968); Angew. Chem. Int. Ed. Engl. *7*, 97 (1968). [18] G. H. Brown: *Liquid Crystals and some of their Applications in Chemistry*, Anal. Chem. *41*, 26 A (1969); Chemie in unserer Zeit *2*, 42 (1968). [19] R. Steinsträßer and L. Pohl: *Chemie und Verwendung flüssiger Kristalle*, Angew. Chem. *85*, 706 (1973); Angew. Chem. Int. Ed. Engl. *12*, 617 (1973). [20] E. Sackmann: *Scientific Applications of Liquid Crystals*, in G. Meier, E. Sackmann, and J. G. Grabmaier (eds.): *Applications of Liquid Crystals*, Springer, Berlin, Heidelberg, New York, 1975, p. 21ff.

[21] D. Demus: *Eigenschaften, Theorie und Molekülbau flüssiger Kristalle*, Z. Chem. *15*, 1 (1975); D. Demus and H. Zaschke: *Flüssige Kristalle in Tabellen, I und II*, Deutscher Verlag für Grundstoffindustrie, Leipzig, 1976 and 1987. [22] E. I. Kovshev, L. M. Blinov, and V. V. Titov: *Thermotropic*

Liquid Crystals and Their Application, Usp. Khim. *46*, 753 (1977); Russ. Chem. Rev. *46*, 395 (1977). [22a] H. Kelker and R. Hatz, *Handbook of Liquid Crystals*, Verlag Chemie, Weinheim, 1980. [23] H. Kelker and B. Scheurle, Angew. Chem. *81*, 903 (1969); Angew. Chem. Int. Ed. Engl. *8*, 884 (1969). [24] B. R. Sundheim (ed.): *Fused Salts*, McGraw-Hill, New York, 1964. [25] W. Sundermeyer: *Salzschmelzen und ihre Verwendung als Reaktionsmedien*, Angew. Chem. *77*, 241 (1965); Angew. Chem. Int. Ed. Engl. *4*, 222 (1965); Chemie in unserer Zeit *1*, 150 (1967). [26] H. Bloom and J. W. Hastie: *Molten Salts as Solvents*, in T. C. Waddington (ed.): *Non-Aqueous Solvent Systems*, Academic Press, London, New York, 1965, p. 353ff. [27] G. J. Janz: *Molten Salts Handbook*, Academic Press, London, New York, 1967. [28] J. E. Gordon: *Applications of Fused Salts in Organic Chemistry*, in D. B. Denney (ed.): *Techniques and Methods of Organic and Organometallic Chemistry*, Dekker, New York, London, 1969, p. 51ff. [29] J. Braunstein, G. Mamantov, and G. P. Smith (eds.): *Advances in Molten Salt Chemistry*, Vols. 1–4, Plenum Publ. Corp., London, New York, 1971–1981. [30] D. H. Kerridge: *Molten Salts as Nonaqueous Solvents*, in J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvents*, Vol. 5B, Academic Press, London, New York, 1978.

[31] W. Sundermeyer, O. Glemser, and K. Kleine-Weischede, Chem. Ber. *95*, 1829 (1962). [31a] E. Channot, A. K. Sharma, and L. A. Paquette, Tetrahedron Lett. *1978*, 1963. [32] G. Duve, O. Fuchs, and H. Overbeck: *Lösemittel HOECHST*, 6th ed., Hoechst AG, Frankfurt (Main) 1976. [33] E. sz. Kováts: *Zu Fragen der Polarität – Die Methode der Linearkombination der Wechselwirkungskräfte*, Chimia *22*, 459 (1968). [34] C. Reichardt: *Optische Aktivität und Molekülsymmetrie*, Chemie in unserer Zeit *4*, 188 (1970). [35] D. Seebach: *Neue links- und rechtshändige Werkzeuge für den Chemiker*, in "25 Jahre Fonds der Chemischen Industrie 1950–1975", Frankfurt (Main) 1975, p. 13ff. [36] H. Pracejus: *Asymmetrische Synthesen*, Fortschr. Chem. Forsch. *8*, 493 (1967). [37] T. D. Inch: *Asymmetric Syntheses*, Synthesis *1970*, 466. [38] J. D. Morrison and H. S. Mosher: *Asymmetric Organic Reactions*, Prentice Hall, Englewood Cliffs/New Jersey, 1971, p. 411ff. [39] R. v. Ammon and R. D. Fischer, Angew. Chem. *84*, 737 (1972); Angew. Chem. Int. Ed. Engl. *11*, 675 (1972). [40] M. Raban and K. Mislow, Topics in Stereochemistry *2*, 199 (1967).

[41] R. Charles, U. Beither, B. Feibush, and E. Gil-Av, J. Chromatogr. *112*, 121 (1975). [42] H. Plieninger and H. P. Kraemer, Angew. Chem. *88*, 230 (1976); Angew. Chem. Int. Ed. Engl. *15*, 243 (1976); Tetrahedron *34*, 891 (1978). [43] G. B. Kauffman: *The Brönsted–Lowry Acid–Base Concept*, J. Chem. Educ. *65*, 28 (1988). [44] R. G. Bates: *Medium Effects and pH in Nonaqueous Solvents*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York, London, 1969, Vol. 1, p. 46ff. [45] R. P. Bell: *Acids and Bases – Their Quantitative Behaviour*, Methuen, London, 1969; *Säuren und Basen* (translated into German by F. Frickel). Verlag Chemie, Physik Verlag, Weinheim, 1974. [46] R. P. Bell: *The Proton in Chemistry*, 2nd ed., Chapman and Hall, London, 1973. [47] M. M. Davies: *Brønsted Acid-Base Behaviour in "Inert" Organic Solvents*, in J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvent Systems*, Academic Press, London, New York, 1970, Vol. III, p. 1ff. [48] L. N. Bykova and S. I. Petrov: *Acid-Base Equilibria in Amphiprotic Solvents and Potentiometric Titration*, Usp. Khim. *41*, 2065 (1972); Russ. Chem. Rev. *41*, 975 (1972). [49] E. J. King: *Acid-Base Behaviour*, in A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, p. 331ff. [50] R. G. Bates: *Acidity Functions in Aqueous and Nonaqueous Media*, Bull. Soc. Chim. Belg. *84*, 1139 (1975).

[51] J. N. Brønsted, Rec. Trav. Chim. Pays-Bas *42*, 718 (1923); Chem. Rev. *5*, 231 (1928); Angew. Chem. *43*, 229 (1930). [52] T. M. Lowry, Chem. Ind. (London) *42*, 43 (1923); Trans. Faraday Soc. *20*, 13 (1924). [53] For a compilation of important pK_a values of acids and bases, see A. J. Gordon and R. A. Ford: *The Chemist's Companion*, Wiley-Interscience, New York, 1972, p. 54ff. [54] J. N. Brønsted, Ber. Dtsch. Chem. Ges. *61*, 2049 (1928). [55] I. M. Kolthoff, E. B. Sandell, E. J. Meehan, and S. Bruckenstein: *Quantitative Chemical Analysis*, 4th ed., MacMillan Company, London, 1969 (especially Chap. 5 on Acid-Base Equilibria in Nonaqueous Solvents). [56] I. M. Kolthoff: *Acid-Base Equilibria in Dipolar Aprotic Solvents*, Anal. Chem. *46*, 1992 (1974). [57] V. I. Dulova, N. V. Lichkova, and L. P. Ivleva: *The Differentiating Action of Oxygen-containing Solvents on the Strengths of Acids*, Usp. Khim. *37*, 1893 (1968); Russ. Chem. Rev. *37*, 818 (1968). [58] L. N. Bykova and S. I. Petrov: *Differentiating Effect of Amphiprotic Solvents in Relation to Phenol and Benzoic Acid Derivatives*, Usp. Khim. *39*, 1631 (1970); Russ. Chem. Rev. *39*, 766 (1970). [59] I. M. Kolthoff, S. Bruckenstein, and M. K. Chantooni, J. Am. Chem. Soc. *83*, 3927 (1961). [60] R. H.

Boyd: *Acidity Functions*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Vol. 1, Dekker, New York, London, 1969, p. 98ff.

[61] C. Agami and M. Caillot, Bull. Soc. Chim. Fr. *1969*, 1990; C. Agami, ibid. *1969*, 2183. [62] M. L. Martin, Annales de Physique *7*, 35 (1962); G. J. Martin and M. L. Martin, J. Chim. Physique *61*, 1222 (1964). [63] G. N. Lewis: *Valence and the Structure of Atoms and Molecules*, Chemical Catalog Company, New York, 1923, p. 141ff.; J. Franklin Inst. *226*, 293 (1938). [64] M. R. Leach: *Lewis Acid/Base Reaction Chemistry*, Meta-Synthesis, Brighton/U.K., 1999; Chem. Abstr. *132*, 333992c (2000). [65] V. Gutmann: *Coordination Chemistry in Non-Aqueous Solvents*, Springer, Wien, New York, 1968; V. Gutmann: *Chemische Funktionslehre*, Springer, Wien, New York, 1971. [65a] W. B. Jensen: *The Lewis Acid-Base Concepts – An Overview*, Wiley, Chichester 1980. [66] R. G. Pearson, J. Am. Chem. Soc. *85*, 3533 (1963); Coord. Chem. Rev. *100*, 403 (1990); J. Chem. Educ. *64*, 561 (1987); ibid. *76*, 267 (1999). [67] R. G. Pearson: *Chemical Hardness – Applications from Molecules to Solids*, Wiley-VCH, Weinheim, 1997. [68] H. Werner: *Harte und weiche Säuren und Basen – ein neues Klassifizierungsprinzip*, Chemie in unserer Zeit *1*, 135 (1967). [69] R. D. Hancock and A. E. Martell, J. Chem. Educ. *73*, 654 (1996). [70] T.-L. Ho: *Hard and Soft Acids and Bases Principle in Organic Chemistry*, Academic Press, New York, 1977; T.-L. Ho, Chem. Rev. *75*, 1 (1975); T.-L. Ho, Tetrahedron *41*, 3 (1985).

[71] R. Th. Myers, Inorg. Chem. *13*, 2040 (1974). [72] R. S. Drago, J. Chem. Educ. *51*, 300 (1974). [73] A. J. Parker, Quart. Rev. (London) *16*, 163 (1962); Usp. Khim. *32*, 1270 (1963); Adv. Org. Chem. *5*, 1 (1965); Adv. Phys. Org. Chem. *5*, 173 (1967); Usp. Khim. *40*, 2203 (1971); Chem. Rev. *69*, 1 (1969); Pure Appl. Chem. *25*, 345 (1971). [74] J. Miller and A. J. Parker, J. Am. Chem. Soc. *83*, 117 (1961); A. J. Parker, J. Chem. Soc. *1961*, 1328. [75] J. F. Coetzee: *Ionic Reactions in Acetonitrile*, Progr. Phys. Org. Chem. *4*, 45 (1967). [76] R. S. Kittila: *A Review of Catalytic and Synthetic Applications for DMF and DMAC*, Bulletin DuPont de Nemours and Co., Wilmington/USA, 1960 (and Supplement 1962). [77] J. W. Vaughn: *Amides*, in J. J. Lagowski (ed.): *The Chemistry of Non-Aqueous Solvents*, Academic Press, New York, London, 1967, Vol. II, p. 192ff. [78] S. S. Pizey: *Dimethylformamide*, in: *Synthetic Reagents*, Wiley, New York, London, 1974, Vol. I, p. 1ff. [79] R. Sowada: *Darstellung, Eigenschaften und Verwendung von Dimethylsulfon*, Z. Chem. *8*, 361 (1968). [80] H. L. Schläfer and W. Schaffernicht: *Dimethylsulfoxid als Lösungsmittel für anorganische Verbindungen*, Angew. Chem. *72*, 618 (1960).

[81] C. Agami: *Le Diméthylsulfoxyde en Chimie Organique*, Bull. Soc. Chim. Fr. *1965*, 1021. [82] D. Martin, A. Weise, and H.-J. Niclas: *Das Lösungsmittel Dimethylsulfoxid*, Angew. Chem. *79*, 340 (1967); Angew. Chem. Int. Ed. Engl. *6*, 318 (1967). [83] D. Martin and H. G. Hauthal: *Dimethylsulfoxid*, Akademie-Verlag, Berlin, 1971; *Dimethyl Sulphoxide*, Van Nostrand Reinhold, London, 1975. [84] S. W. Jacob, E. E. Rosenbaum, and D. C. Wood (eds.): *Dimethyl Sulfoxide*, Dekker, New York, 1971, Vol. 1. [85] H. Normant: *Hexamethylphosphorsäuretriamid*, Angew. Chem. *79*, 1029 (1967); Angew. Chem. Int. Ed. Engl. *6*, 1046 (1967); Bull. Soc. Chim. Fr. *1968*, 791; Usp. Khim. *39*, 990 (1970); Russ. Chem. Rev. *39*, 457 (1970). [86] M. Bréant: *Propriétés chimiques et électrochimiques dans la N-méthylpyrrolidone*, Bull. Soc. Chim. Fr. *1971*, 725. [87] M. Bréant and G. Demange-Guerin: *Propriétés chimiques et électrochimiques dans la nitromethane*, Bull. Soc. Chim. Fr. *1975*, 163. [88] W. H. Lee: *Cyclic Carbonates*, in J. J. Lagowski (ed.): *The Chemistry of Non-aqueous Solvents*, Academic Press, New York, 1976, Vol. IV, p. 167ff. [89] E. M. Arnett and C. F. Douty: *Sulfolane – A Weakly Basic Aprotonic Solvent of High Dielectric Constant*, J. Am. Chem. Soc. *86*, 409 (1964). [90] T. Yamamoto: *Sulfolane*, Yuki Gosei Kagaku Kyokai Shi *28*, 853 (1970); Chem. Abstr. *73*, 109587c (1970). [90a] J. Martinmaa: *Sulfolane*, in J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvents*, Academic Press, New York, 1976, Vol. IV, p. 248ff.

[91] A. Lüttringhaus and H.-W. Dirksen: *Tetramethylharnstoff als Lösungsmittel und Reaktionspartner*, Angew. Chem. *75*, 1059 (1963). [91a] B. J. Barker, J. Rosenfarb, and J. A. Caruso: *Harnstoffe als Lösungsmittel in der chemischen Forschung*, Angew. Chem. *91*, 560 (1979); Angew. Chem. Int. Ed. Engl. *18*, 503 (1979). [91b] R. J. Lemire and P. G. Sears: *N-Methylacetamide as a Solvent*, Top. Curr. Chem. *74*, 45 (1978). [92] F. Madaule-Aubry: *Le rôle en chimie de certains solvants dipolaires aprotiques*, Bull. Soc. Chim. Fr. *1966*, 1456. [93] C. Agami: *Nouvelle classification des solvants aprotoniques*, Bull. Soc. Chim. Fr. *1967*, 4031. [94] C. D. Ritchie: *Interactions in Dipolar*

Aprotic Solvents, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York, London, 1969, Vol. 1, p. 219ff. [95] H. Liebig: *Präparative Chemie in aprotonischen Lösungsmitteln*, Chemiker-Ztg. *95*, 301 (1971). [96] E. S. Amis and J. F. Hinton: *Solvent Effects on Chemical Phenomena*, Academic Press, New York, London, 1973, Vol. 1, p. 271ff. [97] P. K. Kadaba: *Role of Protic and Dipolar Aprotic Solvents in Heterocyclic Syntheses via 1,3-Dipolar Cycloaddition Reactions*, Synthesis *1973*, 71. [98] J. H. Hildebrand and R. L. Scott: *The Solubility of Nonelectrolytes*, 3rd ed., Reinhold, New York, 1950; Dover, New York, 1964; J. H. Hildebrand and R. L. Scott: *Regular Solutions*, Prentice-Hall, Englewood Cliffs/New Jersey, 1962; J. H. Hildebrand, J. M. Prausnitz, and R. L. Scott: *Regular and Related Solutions*, Van Nostrand-Reinhold, Princeton/New Jersey, 1970. [99] A. F. M. Barton: *Handbook of Solubility Parameters and other Cohesion Parameters*, CRC Press, Boca Raton/Florida, 1983. [100] M. R. J. Dack, Aust. J. Chem. *28*, 1643 (1975).

[101] M. R. J. Dack: *The Influence of Solvent on Chemical Reactivity*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. VIII, Part II, p. 95ff., in A. Weissberger (ed.): *Techniques of Chemistry*, Wiley-Interscience, New York, 1976. [102] M. Bohle, W. Kollecker, and D. Martin, Z. Chem. *17*, 161 (1977). [103] A. D. Buckingham, E. Lippert, and S. Bratos: *Organic Liquids. Structure, Dynamics, and Chemical Properties*, Wiley-Interscience, Chichester, 1978. [104] D. Stoye: *Solvents*, in *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed., Vol. A 24, p. 437ff., VCH, Weinheim/Germany, 1993. [105] H. Gnamm and O. Fuchs: *Lösungsmittel und Weichmachungsmittel*, 8th ed., Vols. I and II, Wissenschaftliche Verlagsgesellschaft, Stuttgart, 1980. [106] M. Ash and I. Ash (eds.): *The Index of Solvents: An International Guide to more than 1700 Products by Trade Name, Chemical, Application and Manufacture*, Gower, Brookfield/VI, USA, 1996. [107] O. Popovych and R. P. T. Tomkins: *Nonaqueous Solution Chemistry*, Wiley-Interscience, New York, 1981. [108] N. H. March and M. P. Tosi: *Coulomb Liquids*, Academic Press, New York, London, 1984. [109] D. Demus, J. Goodby, G. W. Gray, H.-W. Spiess, and B. Vill (eds.): *Handbook of Liquid Crystals*, Vols. 1–4, Wiley-VCH, Weinheim/Germany, 1998. [110] D. Demus, J. Goodby, G. W. Gray, H.-W. Spiess, and B. Vill (eds.): *Physical Properties of Liquid Crystals*, Wiley-VCH, Weinheim/Germany, 1999.

[111] T. Nakano and H. Hirata, Bull. Chem. Soc. Jpn. *55*, 947 (1982). [112] D. Inman and D. G. Lovering (eds.): *Ionic Liquids*, Plenum Publ. Corp., New York, 1981. [113] R. M. Pagni: *Organic and Organometallic Reactions in Molten Salts and Related Melts*, in G. Mamantov, C. B. Mamantov, and J. Braunstein (eds.): *Advances in Molten Salt Chemistry*, Vol. 6, Elsevier, Amsterdam 1987. [114] C. L. Hussey: *Room-Temperature Molten Salt Systems*, in G. Mamantov (ed.): *Advances in Molten Salt Chemistry*, Vol. 5, p. 185ff., Elsevier, Amsterdam, 1983. [115] L. G. Wade, K. J. Acker, R. A. Earl, and R. A. Osteryoung, J. Org. Chem. *44*, 3724 (1979). [116] L. Gattermann, H. Wieland, Th. Wieland, and W. Sucrow: *Die Praxis des organischen Chemikers*, 43th ed., de Gruyter, Berlin, New York, 1982, p. 261. [117] L. K. Nash, J. Chem. Educ. *61*, 981 (1984). [118] L. R. Snyder, CHEMTECH *9*, 750 (1979); ibid. *10*, 188 (1980). [119] N. B. Godfrey, CHEMTECH *1972*, 359; *cf.* also I. Vavruch, Chemie für Labor und Betrieb *35*, 385 (1984). [120] American Society for Testing and Materials: *Annual Book of ASTM Standards 1982*, Part 29, p. 142–144 (ASTM D 1133–78), Philadelphia, 1982.

[121] M. Claessens, L. Palombini, M.-L. Stien, and J. Reisse, Nouv. J. Chim. *6*, 595 (1982); M.-L. Stien, M. Claessens, A. Lopez, and J. Reisse, J. Am. Chem. Soc. *104*, 5902 (1982); J. Reisse: *Intermolecular Interaction and Molecular Multipolarity*, Plenary lecture, ESOC IV, Aix-en-Provence, September 4, 1985. [122] M.-E. Colnay, A. Vasseur, and M. Guérin, J. Chem. Res. (S) *1983*, 220; J. Chem. Res. (M) *1983*, 2050. [123] H. Rau: *Asymmetric Photochemistry in Solution*, Chem. Rev. *83*, 535 (1983); *cf.* p. 545. [124] W. H. Pirkle and D. J. Hoover: *NMR Chiral Solvating Agents*, Topics in Stereochemistry *13*, 263 (1982). [125] W. A. König, S. Sievers, and U. Schulze, Angew. Chem. *92*, 935 (1980); Angew. Chem. Int. Ed. Engl. *19*, 910 (1980). [126] M. Bucciarelli, A. Forni, I. Moretti, and G. Torre, J. Org. Chem. *48*, 2640 (1983). [127] L. Jalander and R. Strandberg, Acta Chem. Scand., Part B *37*, 15 (1983). [128] D. Seebach and A. Hidber, Org. Synth. *61*, 42 (1983). [129] P. A. Giguère, J. Chem. Educ. *56*, 571 (1979); *cf.* also Chemiker Ztg. *104*, 172 (1980). [130] L. P. Hammett and A. J. Deyrup, J. Am. Chem. Soc. *54*, 2721 (1932).

[131] G. A. Olah, G. K. Surya Prakash, and J. Sommer: *Superacids*, Science *206*, 13 (1979); *Superacids*, Wiley, Chichester, 1985. [132] T. A. O'Donnell: *Super-Acids and Acidic Melts as Inorganic Chemical Reaction Media*, VCH, Weinheim/Germany, 1993. [133] B. J. Barker, J. Rosenfarb, and J. A. Caruso, Angew. Chem. *91*, 560 (1979); Angew. Chem. Int. Ed. Engl. *18*, 503 (1979). [134] T. Mukhopadhyay and D. Seebach, Helv. Chim. Acta *65*, 385 (1982); *cf.* also Chimia *39*, 147 (1985), and Chem. Br. *21*, 632 (1985). [135] M. Mashima, R. T. McIver, R. W. Taft, F. G. Bordwell, and W. N. Olmstead, J. Am. Chem. Soc. *106*, 2717 (1984). [136] M. C. R. Symons and V. K. Thomas, J. Chem. Soc., Faraday Trans. I *1981*, 1891. [137] L. B. Kier: *Quantitation of Solvent Polarity Based on Molecular Structure*, J. Pharm. Sci. *70*, 930 (1981); Chem. Abstr. *96*, 6069h (1982). [138] M. Chastrette, Tetrahedron *35*, 1441 (1979); M. Chastrette and J. Carretto, Tetrahedron *38*, 1615 (1982); M. Chastrette, M. Rajzmann, M. Chanon, and K. F. Purcell, J. Am. Chem. Soc. *107*, 1 (1985). [139] R. D. Cramer III, J. Am. Chem. Soc. *102*, 1837, 1849 (1980). [140] P. Svoboda, O. Pytela, and M. Večeřa, Collect. Czech. Chem. Commun. *48*, 3287 (1983); see also O. Pytela, ibid. *55*, 634, 644 (1990).

[141] J. Elguero and A. Fruchier, Anales de Quimica, Ser. C. *79*, 72 (1983); Chem. Abstr. *100*, 5743n (1984). [142] R. Carlson, T. Lundstedt, and C. Albano, Acta Chem. Scand., Part B, *39*, 79 (1985); see also R. Carlson: *Design and Optimization in Organic Synthesis*, Elsevier, Amsterdam, 1992, as well as R. Carlson and A. Nordahl, Top. Curr. Chem. *166*, 1 (1993). [143] P.-C. Maria, J.-F. Gal, J. de Franceschi, and E. Fargin, J. Am. Chem. Soc. *109*, 483 (1987). [144] (a) J. Shorter: *Correlation Analysis of Organic Reactivity: With Particular Reference to Multiple Regression*, Wiley, Chichester, 1982; (b) O. Exner: *Correlation Analysis of Chemical Data*, Plenum Press, New York, and SNTL, Prague, 1988. [145] (a) E. R. Malinowsky and D. G. Howery: *Factor Analysis in Chemistry*, Wiley, New York, 1980; (b) R. I. Zalewski: *Applications of PCA in Organic Chemistry*, Progr. Phys. Org. Chem. *18*, 77 (1990). [146] M. A. Sharaf, D. L. Illman, and B. R. Kowalski: *Chemometrics*, Wiley, Chichester, New York, 1986. [147] B. G. M. Vandeginste: *Chemometrics – General Introduction and Historical Development*, Top. Curr. Chem. *141*, 1 (1987). [148] M. F. Delaney: *Chemometrics*, Anal. Chem. *56*, 261R (1984), and references cited therein. [149] S. Wold and M. Sjöström: *LFE Relationships as Tools for Investigating Chemical Similarity. Theory and Practice*, in N. B. Chapman and J. Shorter (eds.): *Correlation Analysis in Chemistry, Recent Advances*, Plenum, New York, 1978; S. Wold and M. Sjöström: *Chemometrics and its Roots in Organic Chemistry*, Acta Chem. Scand. *52*, 517 (1998). [150] C. Reichardt, Angew. Chem. *91*, 119 (1979); Angew. Chem. Int. Ed. Engl. *18*, 98 (1979).

[151] S. Wold and M. Sjöström in B. R. Kowalski (ed.): *Chemometrics. Theory and Practice*, ACS Symposium Series No. 52, American Chemical Society, Washington D.C. 1977, Chapter 7. [152] A. Thielemans and D. Luc Massart, Chimia *39*, 236 (1985). [153] C. Hansch, Acc. Chem. Res. *2*, 232 (1969). [154] G. Allen, G. Gee, and G. J. Wilson, Polymer *1*, 456 (1960); *cf.* also reference [99], table 10 in chapter 7, p. 118. [155] W. L. Leigh, D. T. Frendo, and P. J. Klawunn, Can. J. Chem. *63*, 2131 (1985); ibid. *63*, 2736 (1985). [156] (a) P. Walden, Bull. Acad. Imper. Sci. (St. Petersburg) *8*, 405 (1914); Chem. Zentralbl. 1914/I, p. 1800; (b) S. Sugden and H. Wilkins, J. Chem. Soc. 1291 (1929). [157] D. Enders and R. W. Hoffmann: *Asymmetrische Synthese*, Chemie in unserer Zeit *19*, 177 (1985). [158] D. Seebach and W. Langer, Helv. Chim. Acta *62*, 1701 (1979). [159] C. H. Heathcock: *The Aldol Condensation in Stereoselective Organic Synthesis*, in H. Nozaki (ed.): *Current Trends in Organic Synthesis*, Pergamon Press, Oxford, 1983, p. 27ff.; C. H. Heathcock: *Stereoselective Aldol Condensations*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part B. p. 177ff., Elsevier, Amsterdam, 1984. [160] A. Streitwieser jr., J. I. Brauman, J. H. Hammons, and A. H. Pudjaatmaka, J. Am. Chem. Soc. *87*, 384 (1965); A. Streitwieser jr., E. Juaristi, and L. L. Nebenzahl: *Equilibrium Carbon Acidities in Solution*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part A, p. 323ff., Elsevier, Amsterdam, 1980; see also A. Streitwieser *et al.*, J. Org. Chem. *56*, 1074 (1991).

[161] F. G. Bordwell: *Equilibrium Acidities of Carbon Acids*, Pure Appl. Chem. *49*, 963 (1977); F. G. Bordwell *et al.*, Acc. Chem. Res. *21*, 456, 463 (1988); ibid. *26*, 510 (1993). [162] (a) J. S. Wilkes, J. A. Levisky, R. A. Wilson, and C. L. Hussey, Inorg. Chim. Acta *21*, 1263 (1982); (b) J. S. Wilkes and M. J. Zaworotko, J. Chem. Soc., Chem. Commun. 965 (1992). [163] V. Ramesh and R. G. Weiss, J. Org. Chem. *51*, 2535 (1986). [164] H. U. Borgstedt: *Chemical Reactions in Alkali Metals*,

Top. Curr. Chem. *134*, 125 (1986). [165] R. Carlson, T. Lundstedt, and R. Shabana, Acta Chem. Scand., Part B *40*, 694 (1986); ibid. *41*, 164 (1987). [166] J. Michl and E. W. Thulstrup: *Spectroscopy with Polarized Light – Solute Alignment by Photoselection, in Liquid Crystals, Polymers, and Membranes*, VCH Verlagsgesellschaft, Weinheim, 1986. [167] (a) K. R. Seddon, Kinetika i Kataliz *37*, 743 (1996); Kinetics and Catalysis *37*, 693 (1996); Chem. Abstr. *125*, 285927s (1996); (b) K. R. Seddon, J. Chem. Technol. Biotechnol. *68*, 351 (1997); Chem. Abstr. *126*, 306898w (1997). [168] L. Šafařik and Z. Stránský: *Titrimetric Analysis in Organic Solvents*, in G. Svehla (ed.): *Wilson and Wilson's Comprehensive Analytical Chemistry*, Vol. XXII, Chapters 1–3, Elsevier, Amsterdam, New York, 1986. [169] C. G. Swain, A. Ohno, D. K. Roe, R. Brown, and T. Maugh, J. Am. Chem. Soc. *89*, 2648 (1967). [170] R. A. Y. Jones: *Physical and Mechanistic Organic Chemistry*, 2nd ed., Cambridge University Press, Cambridge/U.K., 1984, p. 83ff.

[171] Y. Marcus, J. Phys. Chem. *91*, 4422 (1987); J. Chem. Soc., Perkin Trans. II 1751 (1994). [172] M. Răileanu: *Rolul Solventilor in Chimie*, Editura Academiei Republici Socialiste Romania, Bucureşti, 1988. [173] G. Gritzner, J. Mol. Liquids *73/74*, 487 (1997). [174] Yu. Ya. Fialkov: *Rastvoritel' kak Sredstvo Upravleniya Khimicheskim Protsessom (Solvents as an Agent for Controlling Chemical Processes)*, Publisher Khimiya, Leningrad/USSR, 1990. [175] Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998. [176] P. Knochel (ed.): *Modern Solvents in Organic Synthesis*, Top. Curr. Chem., Vol. 206, Springer, Berlin, 1999. [177] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, William Andrew Publishing, New York, 2001. [178] R. Breslow, Acc. Chem. Res. *24*, 150 (1991). [179] P. A. Grieco: *Organic Synthesis in Water*, Blackie Academic and Professional, Hampshire, 1998. [180] C.-J. Li and T.-H. Chan: *Organic Reactions in Aqueous Media*, Wiley, Chichester, 1997.

[181] A. Lubineau and J. Augé: *Water as Solvent in Organic Synthesis*, Top. Curr. Chem. *206*, 1 (1999). [182] A. Endres and G. Maas, Chem. Unserer Zeit *34*, 382 (2000). [183] A. Maciejewski, J. Photochem. Photobiol. A *51*, 87 (1990); Chem. Abstr. *112*, 226494g (1990). [184] B. Betzemeier and P. Knochel: *Perfluorinated Solvents – A Novel Medium in Organic Chemistry*, Top. Curr. Chem. *206*, 61 (1999). [185] T. Horváth, Acc. Chem. Res. *31*, 641 (1998). [186] R. H. Fish, Chem. Eur. J. *5*, 1677 (1999). [187] C. L. Hussey, Adv. Molten Salt Chem. *5*, 185 (1983); Pure Appl. Chem. *60*, 1763 (1988). [188] T. Welton, Chem. Rev. *99*, 2071 (1999); P. Wasserscheid and T. Welton: *Ionic Liquids in Synthesis*, Wiley-VCH, Weinheim, 2002. [189] P. Wasserscheid and W. Keim, Angew. Chem. *112*, 3926 (2000); Angew. Chem. Int. Ed. Engl. *39*, 3772 (2000). [190] D. W. Rooney and K. R. Seddon: *Ionic Liquids*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, William Andrew Publishing, New York, 2001; Chapter 21.2, p. 1459ff.

[191] A. J. Carmichael and K. R. Seddon, J. Phys. Org. Chem. *13*, 591 (2000). [192] S. N. V. K. Aki, J. F. Brennecke, and A. Samanta, J. Chem. Soc., Chem. Commun. 413 (2001). [193] (a) W. Schroth, J. Andersch, H.-D. Schädler, and R. Spitzner, Chemiker-Ztg. *113*, 261 (1989); (b) W. Schroth, H.-D. Schädler, and J. Andersch, Z. Chem. *29*, 56, 129 (1989). [194] (a) H. H. Rüttinger, J. Schiewe, and H. Matschiner, Spectrochim. Acta *37*, 759 (1992); Electrochim. Acta *38*, 553 (1993); (b) C.-P. Maschmeier and H. Matschiner: *Das System Dimethylamin/Kohlendioxid – ein neuer Elektrolyt*, in G. Kreysa and G. Sandstede (eds.): *Elektrochemische Stoffgewinnung: Grundlagen und Verfahrenstechnik*, DECHEMA Monographien, VCH, Weinheim/Germany, 1992, Vol. 125, p. 707ff. [195] (a) P. A. Grieco: *Organic Chemistry in Lithium Perchlorate/Diethyl Ether*, in M. V. Kisakürek (ed.): *Organic Chemistry: Its Language and Its State of the Art*, HVCA, Basel, and VCH, Weinheim, 1993, p. 133ff.; (b) P. A. Grieco, Aldrichimica Acta *24*(3), 59 (1991); Chem. Abstr. *116*, 99425r (1992). [196] (a) H. Waldmann, Angew. Chem. *103*, 1335 (1991); Angew. Chem. Int. Ed. Engl. *30*, 1306 (1991); (b) A. Flohr and H. Waldmann, J. Prakt. Chem. *337*, 609 (1995). [197] S. Sankararaman and J. E. Nesakumar, Eur. J. Org. Chem. 2003 (2000). [198] S. Winstein, S. Smith, and D. Darwish, J. Am. Chem. Soc. *81*, 5511 (1959). [199] F. Hensel and R. Winter: *Flüssige Metalle*, Chem. Unserer Zeit *22*, 185 (1988). [200] Y. Marcus, J. Sol. Chem. *21*, 1217 (1992); ibid. *25*, 455 (1996).

[201] H. Ohtaki, J. Sol. Chem. *21*, 39 (1992). [202] M. Handa, M. Kataoka, W. Wakaumi, and Y. Sasaki, Bull. Chem. Soc. Jpn. *70*, 315 (1997). [203] M. Dutkiewicz, J. Chem. Soc., Faraday Trans. *86*, 2237 (1990); ibid. *92*, 637 (1996). [204] U. Zilian, Chemiker-Ztg. *115*, 261 (1991). [205] F.

M. Menger and A. M. Sanchez, J. Chem. Soc., Chem. Commun. 199 (1997). [206] E. L. Eliel, S. H. Wilen, and M. P. Doyle: *Basic Organic Chemistry*, Wiley-Interscience, New York, 2001; Chapter 12-5, p. 585ff. [207] H. Huber, A. J. Dyson, and B. Kirchner, Chem. Soc. Rev. *28*, 121 (1999); Chimia *53*, 205 (1999). [208] A. R. Katritzky, L. Mu, V. S. Lobanov, and M. Karelson, J. Phys. Chem. *100*, 10400 (1996). [209] T. Clifford: *Fundamentals of Supercritical Fluids*, Oxford University Press, New York, 1999. [210] P. G. Debenedetti: *Metastable Liquids – Concepts and Principles*, Princeton University Press, Princeton/NJ, USA, 1996.

[211] G. Kaupp, Angew. Chem. *106*, 1519 (1994); Angew. Chem. Int. Ed. Engl. *33*, 1452 (1994). [212] P. E. Savage, S. Gopalan, T. I. Mizan, C. J. Martino, and E. E. Brock, AIChe J. *41*, 1723 (1995). [213] C. L. Phelps, N. G. Smart, and C. M. Wai, J. Chem. Educ. *73*, 1163 (1996). [214] W. Leitner, Acc. Chem. Res. *35*, 746 (2002). [215] R. Noyori (ed.): *Supercritical Fluids*, Chem. Rev. *99*, 353–633 (1999). [216] A. K. Sunol and S. G. Sunol: *Supercritical Fluids*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, William Andrew Publishing, New York, 2001; Chapter 21.1, p. 1419ff. [217] S. L. Wells and J. M. De Simone, Angew. Chem. *113*, 534 (2001); Angew. Chem. Int. Ed. Engl. *40*, 518 (2001). [218] C. Cagniard de la Tour, Ann. Chim. (Paris) *22*, 410 (1822). [219] J. B. Hannay and J. Hogarth, Proc. Roy. Soc. (London) A *29*, 324 (1879). [220] A. Baiker, Chem. Rev. *99*, 453 (1999).

[221] R. Eberhardt, S. Löbbecke, B. Neidhart, and C. Reichardt, Liebigs Ann./Recueil 1195 (1997), and references cited therein. [222] M. Maiwald and G. M. Schneider, Ber. Bunsenges. Phys. Chem. *102*, 960 (1998). [223] J. F. Kauffman, J. Phys. Chem. A *105*, 3433 (2001). [224] D. Bröll, C. Kaul, A. Krämer, P. Krammer, T. Richter, M. Jung, H. Vogel, and P. Zehner, Angew. Chem. *111*, 3180 (1999); Angew. Chem. Int. Ed. Engl. *38*, 299 (1999). [225] P. E. Savage, Chem. Rev. *99*, 603 (1999). [226] M. M. Hoffmann and M. S. Conradi, J. Am. Chem. Soc. *119*, 3811 (1997). [227] H. Ohtaki, T. Radnai, and Y. Yamaguchi, Chem. Soc. Rev. *26*, 41 (1997). [228] A. R. Katritzky, S. M. Allin, and M. Siskin, Acc. Chem. Res. *29*, 399 (1996). [229] M. Siskin and A. R. Katritzky, Chem. Rev. *101*, 825 (2001). [230] A. R. Katritzky, D. A. Nichols, M. Siskin, R. Muruyan, and M. Balasubramanian, Chem. Rev. *101*, 837 (2001).

[231] F. Franks: *Water – A Matrix of Life*, 2nd ed., Royal Society of Chemistry, Cambridge/U.K., 2000. [232] A. Bagno and G. Scorrano, J. Am. Chem. Soc. *110*, 4577 (1988). [233] J. L. Gázquez, J. Phys. Chem. A *101*, 4657 (1997). [234] P. K. Chattaraj, J. Phys. Chem. A *105*, 511 (2000). [235] G. Gritzner and S. Sperker, J. Solution Chem. *17*, 1133 (1988). [236] G. Gritzner, J. Chem. Soc., Faraday Trans. I *82*, 1955 (1986); ibid. *84*, 1047 (1988). [237] W. Mayrhofer and G. Gritzner, J. Chem. Soc., Faraday Trans. *86*, 823 (1990). [238] W. R. Fawcett, J. Phys. Chem. *97*, 9540 (1993). [239] T. Chen, G. Hefter, and Y. Marcus, J. Solution Chem. *29*, 201 (2000). [240] G. Gritzner *et al.*, Z. Phys. Chem. N. F. (München) *158*, 99 (1988); J. Chem. Soc., Faraday Trans. *89*, 3557 (1993).

[241] I. Persson, M. Sandström, and P. L. Goggin, Inorg. Chim. Acta *129*, 183 (1987). [242] R. I. Zalewski, H. Kokocińska, and C. Reichardt, J. Phys. Org. Chem. *2*, 232 (1989). [243] A. Henrion, R. Henrion, W. Abraham, and D. Kreysig, Z. Chem. *29*, 347 (1989); Z. Phys. Chem. (Leipzig) *271*, 29 (1990). [244] A. R. Katritzky, T. Tamm, Y. Wang, S. Sild, and M. Karelson, J. Chem. Inf. Comput. Sci. *39*, 684 (1999). [245] R. Katritzky, T. Tamm, Y. Wang, and M. Karelson, J. Chem. Inf. Comput. Sci. *39*, 692 (1999). [246] V. Palm and N. Palm, Org. React. (Tartu) *28*, 125 (1993); ibid. *31*, 141 (1997); Russ. J. Org. Chem. *36*, 1075 (2000); see also N. Palm, Ph.D. Thesis, University of Tartu/Estonia, 2001.

Chapter 4

[1] J. E. Leffler and E. Grunwald: *Rates and Equilibria of Organic Reactions*, Wiley, New York, London, 1963, p. 15ff. [2] E. A. Moelwyn-Hughes: *The Chemical Statics and Kinetics of Solutions*, Academic Press, London, New York, 1971, Chapter 3. [3] M. J. Blandamer: *Chemical Equilibria in Solution, Dependence of Rate and Equilibrium Constants on Temperature and Pressure*, Ellis Horwood and Prentice-Hall, London, 1992. [4] P. W. Atkins: *Physical Chemistry*, 6th ed., Oxford

University Press, Oxford/U.K., 1998, Chapter 9. [5] H. Martin: *Über Beziehungen zwischen Lösungs-und Gasphasenreaktionen*, Angew. Chem. *78*, 73 (1966); Angew. Chem. Int. Ed. Engl. *5*, 78 (1966). [6] S. W. Benson and G. D. Mendenhall: *Comparison of Equilibrium Reactions in the Gaseous and Liquid Phases*, J. Am. Chem. Soc. *98*, 2046 (1976). [7] A. Ben-Naim: *Solvent and Solute Effects on Chemical Equilibria*, J. Chem. Phys. *63*, 2064 (1975). [8] I. M. Kolthoff, M. K. Chantoni, A. I. Popov, J. A. Caruso, and J. Steigman: *Acid-Base Equilibria in Non-Aqueous Solvents*, in I. M. Kolthoff and P. J. Elving (eds.): *Treatise on Analytical Chemistry*, 2nd ed., Part I, Vol. 2, p. 237ff., Wiley, New York, 1979. [9] R. P. Bell: *The Proton in Chemistry*, Methuen, London, 1959, p. 36ff. [10] E. J. King: *Acid-Base Equilibria*, Pergamon Press, Oxford, 1965, Chapter 11, p. 280ff.

[11] G. Charlot and B. Trémillon: *Réactions Chimiques dans les Solvants et les Sels Fondus*, Gauthier-Villars, Paris, 1963; *Chemical Reactions in Solvents and Melts*, Pergamon Press, Oxford, 1969, Chapters 1 and 2, p. 10ff. and 45ff. [12] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 10. [13] See also references [44, 45, 49, 55, 56] in reference list to Chapter 3. [14] M. Born, Z. Physik *1*, 45 (1920); for modifications of the Born equation see (a) A. R. Rashin and B. Honig, J. Phys. Chem. *89*, 5588 (1985); (b) T. Abe, J. Phys. Chem. *90*, 713 (1986); (c) P. M. Qureshi and R. K. Varshney, J. Chem. Educ. *66*, 641 (1989); ibid. *68*, 109 (1991). [15] L. P. Hammett and A. J. Deyrup, J. Am. Chem. Soc. *54*, 2721 (1932); L. P. Hammett: *Physical Organic Chemistry*, McGraw-Hill, New York, 1970; *Physikalische Organische Chemie*, Verlag Chemie, Weinheim, 1973. [16] B. Gutbezahl and E. Grunwald, J. Am. Chem. Soc. *75*, 559, 565 (1953). [17] R. G. Bates: *Acidity Functions in Aqueous and Nonaqueous Media*, Bull. Soc. Chim. Belg. *84*, 1139 (1975). [18] E. M. Arnett: *Gas-Phase Proton Transfer – A Breakthrough for Solution Chemistry*, Acc. Chem. Res. *6*, 404 (1973). [19] C. Agami: *Acidités et Basicités en Phase Vapeur*, Bull. Soc. Chim. Fr. *1974*, 869. [20] O. A. Reutov, K. P. Butin, and I. P. Beletskaya: *Equilibrium Acidity of Carbon-Hydrogen Bonds in Organic Compounds*, Usp. Khim. *43*, 35 (1974); Russ. Chem. Rev. *43*, 17 (1974).

[21] R. W. Taft: *Gas-Phase Proton-Transfer Equilibria*, in E. F. Caldin and V. Gold (eds.): *Proton Transfer Reactions*, Chapman and Hall, London, 1975. [22] M. S. B. Munson, J. Am. Chem. Soc. *87*, 2332 (1965). [23] P. Kebarle: *Ions and Ion-Solvent Molecule Interactions in the Gas Phase*, in M. Szwarc (ed.): *Ions and Ion Pairs in Organic Reactions*, Wiley-Interscience, New York, 1972, Vol. I, p. 27ff. [24] H. Hartmann, K.-H. Lebert, and K.-P. Wanczek: *Ion Cyclotron Resonance Spectroscopy*, Fortschr. Chem. Forsch. *43*, 57 (1973). [25] R. C. Dunbar: *Ion Cyclotron Resonance and Fourier-Transform Mass Spectrometry*, in C. F. Bernasconi (ed.): *Investigation of Rates and Mechanisms of Reactions* (Volume VI of the Series *Techniques of Chemistry*), 4th ed., Wiley, New York, 1986, Part I, p. 903ff. [26] T. A. Lehman and M. M. Bursey: *Ion Cyclotron Resonance Spectrometry*, Wiley-Interscience, New York, 1976, p. 55ff. [27] H. C. Brown, H. Bartholomay, and M. D. Taylor, J. Am. Chem. Soc. *66*, 435 (1944). [28] J. I. Brauman and L. K. Blair, J. Am. Chem. Soc. *91*, 2126 (1969); J. I. Brauman, J. M. Riveros, and L. K. Blair, ibid. *93*, 3914 (1971). [29] E. M. Arnett, F. M. Jones, M. Taagepera, W. G. Henderson, J. L. Beauchamp, D. Holtz, and R. W. Taft, J. Am. Chem. Soc. *94*, 4724 (1972). [30] D. H. Aue, H. M. Webb, and M. T. Bowers, J. Am. Chem. Soc. *94*, 4726 (1972).

[31] A. F. Trotman-Dickenson, J. Chem. Soc. *1949*, 1293. [32] J. Long and B. Munson, J. Am. Chem. Soc. *95*, 2427 (1973). [33] M. Taagepera, W. G. Henderson, R. T. C. Brownlee, J. L. Beauchamp, D. Holtz, and R. W. Taft, J. Am. Chem. Soc. *94*, 1369 (1972). [34] J. I. Brauman and L. K. Blair, J. Am. Chem. Soc. *90*, 6561 (1968); ibid. *92*, 5986 (1970). [34a] R. Yamdagni and P. Kebarle, J. Am. Chem. Soc. *95*, 4050 (1973); K. Hiraoka, R. Yamdagni, and P. Kebarle, ibid. *95*, 6833 (1973); *cf.* also P. Haberfield and A. K. Rakshit, ibid. *98*, 4393 (1976). [35] J. Hine and M. Hine, J. Am. Chem. Soc. *74*, 5266 (1952). [36] *Cf.* references [11–19] in reference list to Chapter 1. [37] O. Dimroth, Liebigs Ann. Chem. *377*, 127 (1910); *399*, 91 (1913); *438*, 58 (1924). [38] G. Briegleb and W. Strohmeier, Angew. Chem. *64*, 409 (1952). [39] H. A. Staab: *Einführung in die theoretische organische Chemie*, 4th ed., Verlag Chemie, Weinheim, 1964, p. 642ff. [40] S. Forsén and M. Nilsson: *Enolization*, in J. Zabicky (ed.): *The Chemistry of the Carbonyl Group*, Wiley-Interscience, New York, 1970, Vol. 2, p. 157ff.

[41] A. I. Kol'tsov and G. M. Kheifets, Usp. Khim. *40*, 1646 (1971); Russ. Chem. Rev. *40*, 773 (1971). [42] Z. Rappoport (ed.): *The Chemistry of Enols (The Chemistry of Functional Groups)*, Wiley-Interscience, New York, 1990. [43] M. B. Smith and J. March: *March's Advanced Organic Chemistry*, 5th ed., Wiley, New York, 2001, pp. 73ff. and 773ff. [44] R. Matusch, Angew. Chem. *87*, 283 (1975); Angew. Chem. Int. Ed. Engl. *14*, 260 (1975). [45] R. S. Noy, V. A. Gindin, B. A. Ershov, A. I. Kol'tsov, and V. A. Zubkov, Org. Magn. Reson. *7*, 109 (1975). [46] B. Eistert and W. Riess, Chem. Ber. *87*, 92 (1954); B. Eistert and F. Geiss, Tetrahedron *7*, 1 (1959); B. Eistert and K. Schank, Tetrahedron Lett. 1964, 429. [47] M. T. Rogers and J. L. Burdett, J. Am. Chem. Soc. *86*, 2105 (1964); Can. J. Chem. *43*, 1516 (1965). [48] G. Allen and R. A. Dwek, J. Chem. Soc. *B 1966*, 161. [49] K. L. Lockwood, J. Chem. Educ. *42*, 481 (1965). [50] E. J. Drexler and K. W. Field, J. Chem. Educ. *53*, 392 (1976).

[51] G. Schwarzenbach and E. Felder, Helv. Chim. Acta *27*, 1044 (1944). [52] A. Yogev and Y. Mazur, J. Org. Chem. *32*, 2162 (1967). [53] P. B. Russell and J. Mentha, J. Am. Chem. Soc. *77*, 4245 (1955). [54] A new approach deriving the van't Hoff-Dimroth relationship has been given by S. Tanaka, Bull. Chem. Soc. Jpn. *39*, 84 (1966). [55] H. Mauser and B. Nickel, Chem. Ber. *97*, 1745, 1753 (1964). [56] J. G. Kirkwood, J. Chem. Phys. *2*, 351 (1934). [57] J. Powling and H. J. Bernstein, J. Am. Chem. Soc. *73*, 4353 (1951). [58] J. E. Dubois and J. Toullec, Tetrahedron *29*, 2859 (1973). [59] R. Hagen, E. Heilbronner, and P. A. Straub, Helv. Chim. Acta *50*, 2504 (1967). [59a] K. Hafner in R. N. Castle and M. Tišler (eds.): *Lectures in Heterocyclic Chemistry*, Vol. III, p. 33ff.; Supplementary Issue to J. Heterocycl. Chem. *13* (1976). [60] H. Baba and T. Takemura, Tetrahedron *24*, 4779 (1968).

[61] H. Sterk, Monatsh. Chem. *100*, 916 (1969). [62] A. I. Kol'tsov and G. M. Kheifets: *Study of Tautomerism by NMR Spectroscopy*, Usp. Khim. *41*, 877 (1972); Russ. Chem. Rev. *41*, 452 (1972). [63] A. R. Katritzky and J. M. Lagowski: *Prototropic Tautomerism of Heteroaromatic Compounds I–IV*, in A. R. Katritzky (ed.): Advances in Heterocyclic Chemistry, Academic Press, New York, London, 1963, Vol. 1, p. 311 and 339; Vol. 2, p. 1 and 27. [64] J. Elguero, C. Marzin, A. R. Katritzky, and P. Linda, *The Tautomerism of Heterocycles*, in A. R. Katritzky and A. J. Boulton (eds.): Advances in Heterocyclic Chemistry, Supplement 1, Academic Press, New York, 1976. [65a] P. Beak, Acc. Chem. Res. *10*, 186 (1977); P. Beak, J. B. Covington, and J. M. Zeigler, J. Org. Chem. *43*, 177 (1978). [65b] A. Maquestiau, Y. van Haverbeke, C. de Meyer, A. R. Katritzky, M. J. Cook, and A. D. Page, Can. J. Chem. *53*, 490 (1975); Bull. Soc. Chim. Belg. *84*, 465 (1975). [65c] M. J. Cook, S. El-Abbady, A. R. Katritzky, C. Guimon, and G. Pfister-Guillouzo, J. Chem. Soc., Perkin Trans. II 1652 (1977). [65d] R. D. Brown *et al.*, J. Phys. Chem. 97, 46 (1993). [65e] M. J. Nowak, L. Lapinski, J. Fulava, A. Les, and L. Adamowicz, J. Phys. Chem. *96*, 1562 (1992). [65f] S. Stoyanov, I. Petkov, L. Antonov, T. Stoyanova, P. Karagiannidis, and P. Aslanidis, Can. J. Chem. *68*, 1482 (1990). [66] G. Simchen, Chem. Ber. *103*, 398 (1970). [67] A. Gordon and A. R. Katritzky, Tetrahedron Lett. *1968*, 2767; J. Frank and A. R. Katritzky, J. Chem. Soc., Perkin Trans. II, 1428 (1976). [68] E. Daltrozzo, G. Hohlneicher, and G. Scheibe, Ber. Bunsenges. Phys. Chem. *69*, 190 (1965); G. Scheibe and E. Daltrozzo: *Diquinolylmethane and Its Analogs*, in A. R. Katritzky and A. J. Boulton (eds.): Advances in Heterocyclic Chemistry, Academic Press, New York, London, 1966, Vol. 7, p. 153ff. [69] H. Ahlbrecht, Tetrahedron Lett. *1968*, 4421; H. Ahlbrecht and S. Fischer, Tetrahedron *26*, 2837 (1970). [69a] R. Scheffold, J. Löliger, H. Blaser, and P. Geisser, Helv. Chim. Acta *58*, 49 (1975). [69b] M. Dreyfus, G. Dodin, O. Bensaude, and J. E. Dubois, J. Am. Chem. Soc. *98*, 6338 (1976); ibid. *99*, 7027 (1977). [70] M. Mousseron-Canet, J. P. Boca, and V. Tabacik, Spectrochim. Acta, Part A *23*, 717 (1967).

[71] I. Ya. Bershtein and O. F. Ginsburg: *Tautomerism of Aromatic Azo-Compounds*, Usp. Khim. *41*, 177 (1972); Russ. Chem. Rev. *41*, 97 (1972). [72] A. Burawoy and A. R. Thompson, J. Chem. Soc. *1953*, 1443. [73] B. L. Kaul, P. M. Nair, A. V. R. Rao, and K. Venkataraman, Tetrahedron Lett. *1966*, 3897. [74] R. Korewa and H. Urbanska, Rocz. Chem. *46*, 2007 (1972); Chem. Abstr. *78*, 147128s (1973). [74a] M. Misuishi, R. Kamimura, K. Shinohara, and N. Ishii, Sen'i Gakkaishi *32*, T 382 (1976); Chem. Abstr. *85*, 194065y (1976). [74b] R. Hempel, H. Viola, J. Morgenstern, and R. Mayer, J. Prakt. Chem. *318*, 983 (1976). [75] R. K. Norris and S. Sternhell, Aust. J. Chem. *19*, 841 (1966). [76] H. Uffmann, Z. Naturforsch. *22b*, 491 (1967). [76a] Ya. I. Shpinel' and Yu. I. Tarnopol'skii, Zh. Org. Khim. *13*, 1030 (1977); J. Org. Chem. USSR *13*, 948 (1977). [77] R. Escale and J.

Verducci: *Facteurs influencant la tautomérie cycle-chaîne*, Bull. Soc. Chim. Fr. *1974*, 1203. [78] H. Sterk, Monatsh. Chem. *99*, 1764 (1968). [79] J. Whiting and J. T. Edward, Can. J. Chem. *49*, 3799 (1971). [80] S. J. Angyal, Adv. Carbohydr. Chem. Biochem. *42*, 15 (1984). [80a] H. Möhrle, M. Lappenberg, and D. Wendisch, Monatsh. Chem. *108*, 273 (1977). [80b] H. B. Stegmann, R. Haller, and K. Scheffler, Chem. Ber. *110*, 3817 (1977).

[81] R. U. Lemieux, Pure Appl. Chem. *25*, 527 (1971); ibid. *27*, 527 (1971). [82] E. L. Eliel, S. H. Wilen, and L. N. Mander: *Stereochemistry of Organic Compounds*, Wiley-Interscience, New York, 1994, Chapters 9–11; see also E. L. Eliel, Angew. Chem. *84*, 779 (1972); Angew. Chem. Int. Ed. Engl. *11*, 739 (1972). [83] R. J. Abraham and E. Bretschneider: *Medium Effects on Rotational and Conformational Equilibria*, in W. J. Orville-Thomas (ed.): *Internal Rotation in Molecules*, Wiley-Interscience, New York, 1974, p. 481ff. [84] G. Maier: *Valenzisomerisierungen*, Verlag Chemie, Weinheim, 1972. [85] U. Mayer: *Ionic Equilibria in Donor Solvents*, Pure Appl. Chem. *41*, 291 (1975). [86] S. D. Christian and E. H. Lane: *Solvent Effects on Molecular Complex Equilibria*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Wiley-Interscience, New York, 1975; Vol. VIII, Part 1 of A. Weissberger (ed.): *Techniques of Chemistry*, p. 327ff. [87] G. J. Karabatsos and D. J. Fenoglio, J. Am. Chem. Soc. *91*, 1124 (1969). [88] R. J. Abraham and M. A. Cooper, J. Chem. Soc. *B 1967*, 202. [89] E. L. Eliel, Pure Appl. Chem. *25*, 509 (1971); E. L. Eliel and O. Hofer, J. Am. Chem. Soc. *95*, 8041 (1973). [90] *Cf.* Table 13.23, p. 555, in reference [83].

[91] P. Klaboe, J. J. Lothe, and K. Lunds, Acta Chem. Scand. *11*, 1677 (1957). [92] K. Kozima and K. Sabashita, Bull. Chem. Soc. Jpn. *31*, 796 (1958). [93] R. U. Lemieux and J. W. Loan, Can. J. Chem. *42*, 893 (1964). [94] O. D. Ul'yanova, M. K. Ostrovskii, and Yu. A. Pentin, Zh. Fiz. Khim. *44*, 1013 (1970); Russ. J. Phys. Chem. *44*, 562 (1970); and references cited therein. [94a] K. W. Baldry, M. H. Gordon, R. Hafter, and M. J. T. Robinson, Tetrahedron *32*, 2589 (1976). [94b] J. G. Vinter and H. M. R. Hoffmann, J. Am. Chem. Soc. *96*, 5466 (1974). [95] J. P. Snyder, Tetrahedron Lett. *1971*, 215. [96] C. J. Devlin and B. J. Walker, Tetrahedron Lett. *1971*, 4923. [97] R. E. Lutz and A. B. Turner, J. Org. Chem. *33*, 516 (1968). [98] R. Huisgen, W. Scheer, and H. Mäder, Angew. Chem. *81*, 619 (1969); Angew. Chem. Int. Ed. Engl. *8*, 602 (1969). [99] (a) J. B. Flannery, J. Am. Chem. Soc. *90*, 5660 (1968); (b) Y. Sueishi, M. Ohcho, and N. Nishimura, Bull. Chem. Soc. Jpn. *58*, 2608 (1985); (c) A. K. Chibisov and H. Görner, J. Phys. Chem. A *101*, 4305 (1997); (d) J. T. C. Wojtyk, A. Wasey, P. M. Kazmaier, S. Hoz, and E. Buncel, J. Phys. Chem. A *104*, 9046 (2000). [100] J. Elguero, R. Faure, J. P. Galy, and E. J. Vincent, Bull. Soc. Chim. Belg. *84*, 1189 (1975); L. A. Burke, J. Elguero, G. Leroy, and M. Sana, J. Am. Chem. Soc. *98*, 1985 (1976).

[101] E. Vogel, W. A. Böll, and H. Günther, Tetrahedron Lett. *1965*, 609; E. Vogel and H. Günther, Angew. Chem. *79*, 429 (1967); Angew. Chem. Int. Ed. Engl. *6*, 385 (1967). [102] E. Buncel and H. Wilson, Acc. Chem. Res. *12*, 42 (1979); J. Chem. Educ. *57*, 629 (1980). [103] M. H. Abraham, Progr. Phys. Org. Chem. *11*, 2 (1974). [104] R. Stewart: *The Proton: Applications to Organic Chemistry*, Academic Press, Orlando, 1985. [105] K. M. Dyumaev and B. A. Korolev: *The Influence of Solvation on the Acid-Base Properties of Organic Compounds in Various Media*, Usp. Khim. *49*, 2065 (1980); Russ. Chem. Rev. *49*, 1021 (1980). [106] M. I. Kabachnik: *Progress in the Theory of Acids and Bases*, Usp. Khim. *48*, 1523 (1979); Russ. Chem. Rev. *49*, 814 (1979). [107] H. L. Finston and A. C. Rychtman: *A New View of Current Acid-Base Theories*, Wiley, Chichester, 1982. [108] F. Strohbusch: *Neue Erkenntnisse der Säure-Basen-Theorie*, Chemie in unserer Zeit *16*, 103 (1982). [109] R. A. Cox and K. Yates: *Acidity Functions – An Update*, Can. J. Chem. *61*, 2225 (1983). [110] (a) E. P. Serjeant and B. Dempsey: *Ionisation Constants of Organic Acids in Aqueous Solution*, Pergamon Press, Oxford/U.K., 1979; (b) K. Izutsu: *Acid-Base Dissociation Constants in Dipolar Aprotic Solvents*, Blackwell Scientific Publishers, Oxford/U.K., 1990.

[111] A. Albert and E. P. Serjeant: *The Determination of Ionisation Constants – A Laboratory Manual*, 3rd ed., Chapman and Hall, London, New York, 1984. [112] D. D. Perrin, B. Dempsey, and E. P. Serjeant: *pK_a-Prediction for Organic Acids and Bases*, Chapman and Hall, London, 1981. [113] A. Streitwieser, E. Juaristi, and L. L. Nebenzahl: *Equilibrium Carbon Acidities in Solution*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part A, p. 323ff., Elsevier, Amsterdam, 1980. [114] J. H. Futrell: *Gaseous Ion Chemistry and Mass Spectrometry*, Wiley-Interscience, New York, 1986. [115] R. T. McIver: *Chemical Reactions without Solvation*, Scientific

American *243*, 148 (1980); *Chemische Reaktionen ohne Lösungsmittel*, Spektrum der Wissenschaft, January 1981, p. 27ff. [116] R. W. Taft: *Protonic Acidities and Basicities in the Gas Phase and in Solution: Substituent and Solvent Effects*, Progr. Phys. Org. Chem. *14*, 247 (1983). [117] C. R. Moylan and J. I. Brauman: *Gas-Phase Acid-Base Chemistry*, Annu. Rev. Phys. Chem. *34*, 187 (1983). [118] P. Kebarle: *Ion Thermochemistry and Solvation from Gas-Phase Ion Equilibrium*, Annu. Rev. Phys. Chem. *28*, 445 (1977). [119] D. K. Bohme, E. Lee-Ruff, and L. B. Young, J. Am. Chem. Soc. *94*, 5153 (1972); D. K. Bohme in P. Ausloos (ed.): *Interactions between Ions and Molecules*, Plenum, New York, 1974, p. 489ff. [120] M. J. Pellerite and J. I. Brauman: *Gas-Phase Acidities of Carbon Acids*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part A, p. 55ff., Elsevier, Amsterdam, 1980.

[121] J. E. Bartmess and R. T. McIver: *The Gas-Phase Acidity Scale*, in M. T. Bowers (ed.): *Gas-Phase Ion Chemistry*, Vol. 2, p. 87ff., Academic Press, New York, 1979. [122] T. B. McMahon and P. Kebarle, J. Am. Chem. Soc. *107*, 2612 (1985); G. Caldwell, R. Renneboog, and P. Kebarle, Can. J. Chem. *67*, 611 (1989). [123] O. A. Reutov, I. P. Beletskaya, and K. P. Butin: *CH-Acids*, Pergamon Press, Oxford, 1978. [124] F. G. Bordwell, Pure Appl. Chem. *49*, 963 (1977); Acc. Chem. Res. *21*, 456, 463 (1988). [125] J. E. Bartmess, J. A. Scott, and R. T. McIver, J. Am. Chem. Soc. *101*, 6046, 6056 (1979). [126] G. Boand, R. Houriet, and T. Gäumann, J. Am. Chem. Soc. *105*, 2203 (1983). [127] W. N. Olmstead, Z. Margolin, and F. G. Bordwell, J. Org. Chem. *45*, 3295 (1980). [128] G. Caldwell, M. D. Rozeboom, J. P. Kiplinger, and J. E. Bartmess, J. Am. Chem. Soc. *106*, 4660 (1984). [129] R. T. McIver, J. A. Scott, and J. M. Riveros, J. Am. Chem. Soc. *95*, 2706 (1973). [130] D. K. Bohme, A. B. Rakshit, and G. I. Mackay, J. Am. Chem. Soc. *104*, 1100 (1982).

[131] M. Mashima, R. T. McIver, R. W. Taft, F. G. Bordwell, and W. N. Olmstead, J. Am. Chem. Soc. *106*, 2717 (1984); R. W. Taft, I. A. Koppel, R. D. Topsom, and F. Anvia, ibid. *112*, 2047 (1990). [132] G. Caldwell, T. B. McMahon, P. Kebarle, J. E. Bartmess, and J. P. Kiplinger, J. Am. Chem. Soc. *107*, 80 (1985). [133] M. S. Gordon and J. H. Jensen, Acc. Chem. Res. *29*, 536 (1996). [134] (a) S. G. Mills and P. Beak, J. Org. Chem. *50*, 1216 (1985); (b) K. Almdal, H. Eggert, and O. Hammerich, Acta Chem. Scand., Part B, *40*, 230 (1986); (c) M. Moriyasu, A. Kato, and Y. Hashimoto, J. Chem. Soc., Perkin Trans. II 515 (1986); (d) J. Emsley and N. J. Freeman, J. Mol. Struct. *161*, 193 (1987). [135] J. N. Spencer, E. S. Holmboe, M. R. Kirshenbaum, D. W. Firth, and P. B. Pinto, Can. J. Chem. *60*, 1178 (1982). [136] K. D. Grande and S. M. Rosenfeld, J. Org. Chem. *45*, 1626 (1980). [137] H. Hart: *Simple Enols*, Chem. Rev. *79*, 515 (1979). [138] M. Melzig, S. Schneider, F. Dörr, and E. Daltrozzo, Ber. Bunsenges. Phys. Chem. *84*, 1108 (1980). [139] R. Roussel, M. Oteyza de Guerrero, P. Spegt, and J. C. Galin, J. Heterocycl. Chem. *19*, 785 (1982). [140] A. R. Katritzky, M. Karelson, and P. A. Harris, Heterocycles *32*, 329 (1991).

[141] P. Beak, J. B. Covington, and J. M. White, J. Org. Chem. *45*, 1347 (1980). [142] P. Beak, J. B. Covington, S. G. Smith, J. M. White, and J. M. Zeigler, J. Org. Chem. *45*, 1354 (1980). [143] M. Chevrier, O. Bensaude, J. Guillerez, and J. E. Dubois, Tetrahedron Lett. *21*, 3359 (1980). [144] M. J. Scanlan, I. H. Hillier, and R. H. Davies, J. Chem. Soc., Chem. Commun. *1982*, 685. [145] G. Pfister-Guillouzo, C. Guimon, J. Frank, J. Ellison, and A. R. Katritzky, Liebigs Ann. Chem. *1981*, 366. [146] A. I. Artemenko, E. K. Anufriev, I. V. Tikunova, and O. Exner, Zh. Prikl. Spektrosk. *33*, 131 (1980); Chem. Abstr. *94*, 46316b (1981). [147] H. Ahlbrecht and R.-D. Kalas, Liebigs Ann. Chem. *1979*, 102. [148] S. Ogawa and S. Shiraishi, J. Chem. Soc., Perkin Trans. I *1980*, 2527. [149] A. L. Weis, Tetrahedron Lett. *23*, 449 (1982). [150] P. Ball and C. H. Nicholls: *Azo-Hydrazone Tautomerism of Hydroxyazo Compounds – A Review*, Dyes Pigm. *3*, 5 (1982); Chem. Abstr. *96*, 124495s (1982).

[151] S. Kishimoto, S. Kitahara, O. Manabe, and H. Hiyama, J. Org. Chem. *43*, 3882 (1978). [152] E. Hofer, Chem. Ber. *112*, 2913 (1979). [153] G. V. Sheban, B. E. Zaitsev, and K. M. Dyumaev, Teor. Eksp. Khim. *16*, 249 (1980); Chem. Abstr. *93*, 113715a (1980). [154] J. O. Morley, J. Phys. Chem. *98*, 13177 (1994). [155] O. I. Kolodyazhnyi, Tetrahedron Lett. *23*, 499 (1982). [156] A. Munoz, Bull. Soc. Chim. Fr. *1977*, 728. [157] F. G. Kamaev, N. I. Baram, A. I. Ismailov, V. B. Leont'ev, and A. S. Sadykov, Izv. Akad. Nauk SSSR, Ser. Khim. *1979*, 1003; Chem. Abstr. *91*, 55761h (1979). [158] S. Chimichi, R. Nesi, M. Scotton, C. Mannucci, and G. Adembri, Gazz. Chim. Ital. *109*, 117 (1979); J. Chem. Soc., Perkin Trans. II *1980*, 1339. [159] F. Franks: *Physical*

Chemistry of Small Carbohydrates – Equilibrium Solution Properties, Pure Appl. Chem. *59*, 1189 (1987); and references cited therein. [160] L. A. Fedorov, D. N. Kravtsov, and A. S. Peregudov: *Metallotropic Tautomeric Transformations of the σ, σ-Type in Organometallic and Complex Compounds*, Usp. Khim. *50*, 1304 (1981); Russ. Chem. Rev. *50*, 682 (1981).

[161] I. F. Lutsenko, Yu. I. Baukov, and I. Yu. Belavin, J. Organomet. Chem. *24*, 359 (1970). [162] G. Boche, F. Heidenhain, and B. Staudigl, Angew. Chem. *91*, 228 (1979); Angew. Chem. Int. Ed. Engl. *18*, 218 (1979). [163] J. Malecki: *Solvent Effects on Molecular Complexes*, in H. Ratajczak and W. J. Orville-Thomas (eds.): *Molecular Interactions*, Vol. 3, Chapter 4, p. 183ff., Wiley-Interscience, New York, 1982. [164] E. A. Yerger and G. M. Barrow, J. Am. Chem. Soc. *76*, 5211 (1954); ibid. *77*, 4474, 6206 (1955); ibid. *78*, 5802 (1956). [165] (a) H. Baba, A. Matsuyama, and H. Kokubun, J. Chem. Phys. *41*, 895 (1964); Spectrochim. Acta, Part A, *25*, 1709 (1969); (b) Z. Pawelka and T. Kuc, Polish J. Chem. *75*, 845 (2001). [166] L. Elégant, J. Fidanza, J.-F. Gal, and S. N. Vinogradov, J. Chim. Phys. *75*, 914 (1978). [167] J. Fritsch and G. Zundel, J. Phys. Chem. *85*, 556 (1981); R. Krämer and G. Zundel, Z. Phys. Chem. (Frankfurt) N. F. *144*, 265 (1985). [168] Z. Dega-Szafran and E. Dulewicz, Org. Magn. Reson. *16*, 214 (1981). [169] A. Koll, M. Rospenk, and L. Sobczyk, J. Chem. Soc., Faraday Trans. I *1981*, 2309; A. Sucharda-Sobczyk and L. Sobczyk, J. Chem. Res. (S) *1985*, 208. [170] M. Rospenk, J. Fritsch, and G. Zundel, J. Phys. Chem. *88*, 321 (1984); M. Rospenk and T. Zeegers-Huyskens, Spectrochim. Acta, Part A *42*, 499 (1986).

[171] B.-L. Poh and H.-L. Siow, Aust. J. Chem. *33*, 491 (1980). [172] F. M. Menger and T. P. Singh, J. Org. Chem. *45*, 183 (1980). [173] H. Rosotti: *The Study of Ionic Equilibria. An Introduction*, Longman, London, 1978. [174] V. Gutmann: *The Donor-Acceptor Approach to Molecular Interactions*, Plenum Press, New York, 1978. [175] I. Rosenthal, P. Peretz, and K. A. Muszkat, J. Phys. Chem. *83*, 351 (1979). [176] D. Fompeydie and P. Levillain, Bull. Soc. Chim. Fr. *1980*. I-459. [177] M. Feigel, H. Kessler, and A. Walter, Chem. Ber. *111*, 2947 (1978). [178] M. Feigel, H. Kessler, D. Leibfritz, and A. Walter, J. Am. Chem. Soc. *101*, 1943 (1979); *cf.* also H. Kessler and M. Feigel, Acc. Chem. Res. *15*, 2 (1982). [179] E. M. Arnett and E. B. Troughton, Tetrahedron Lett. *24*, 3299 (1983). [180] E. B. Troughton, K. E. Molter, and E. M. Arnett, J. Am. Chem. Soc. *106*, 6726 (1984); E. M. Arnett and K. E. Molter, Acc. Chem. Res. *18*, 339 (1985); J. Phys. Chem. *90*, 383 (1986).

[181] V. N. Sheinker, A. D. Garnovskii, and O. A. Osipov: *Advances in the Study of s-cis-trans-Isomerism – Stereochemistry of Carbonyl Derivatives of Five-Membered Heterocycles*, Usp. Khim. *50*, 632 (1981); Russ. Chem. Rev. *50*, 336 (1981). [182] W. L. Jorgensen: *Theoretical Studies of Medium Effects on Conformational Equilibria*, J. Phys. Chem. *87*, 5304 (1983). [183] J. Fruwert, H. Böhlig, and G. Geiseler: *Gehemmte innere Rotation und Flexibilität kettenförmiger Moleküle*, Z. Chem. *25*, 41 (1985). [184] V. V. Samoshin and N. S. Zefirov: *Conformational Transformations of Organic Molecules in Solutions*, Zh. Vses. Khim. O-va. im. D. I. Mendeleeva (Moskva) *29*, 521 (1984); Chem. Abstr. *102*, 5205e (1985). [185] R. J. Abraham and T. M. Siverns, Tetrahedron *28*, 3015 (1972). [186] G. Alberghina, F. A. Bottino, S. Fisichella, and C. Arnone, J. Chem. Res. (S) *1985*, 108; J. Chem. Res. (M) *1985*, 1201. [187] C. Lopez-Mardomingo, R. Perez-Ossorio, and J. Plumet, J. Chem. Res. (S) *1983*, 150. [188] Z. Friedl, P. Fiedler, and O. Exner, Collect Czech. Chem. Commun. *45*, 1351 (1980); Z. Friedl, P. Fiedler, J. Biroš, V. Uchytilová, I. Tvaroška, S. Böhm, and O. Exner, ibid. *49*, 2050 (1984). [189] F. M. Menger and B. Boyer, J. Org. Chem. *49*, 1826 (1984). [190] L. Došen-Mićović and N. L. Allinger, Tetrahedron *34*, 3385 (1978).

[191] R. J. Abraham and L. Griffiths, Tetrahedron *37*, 575 (1981). [192] J. J. Moura-Ramos, L. Dumont, M. L. Stien, and J. Reisse, J. Am. Chem. Soc. *102*, 4150 (1980). [193] M. J. Cook, M. H. Abraham, L. E. Xodo, R. Cruz, Tetrahedron Lett. *22*, 2991 (1981). [194] M. H. Abraham, L. E. Xodo, R. J. Abraham, and M. J. Cook, Tetrahedron Lett. *22*, 5183 (1981). [195] M. Manoharan, E. L. Eliel, and F. I. Carroll, Tetrahedron Lett. *24*, 1855 (1983). [196] V. V. Samoshin and N. S. Zefirov, Zh. Org. Khim. *17*, 1319, 1771 (1981); J. Org. Chem. USSR *17*, 1170, 1585 (1981); Tetrahedron Lett. *22*, 2209 (1981). [197] L. Došen-Mićović and V. Žigman, J. Chem. Soc., Perkin Trans. II *1985*, 625. [198] F. Dietz, W. Förster, R. Thieme, and C. Weiss, Z. Chem. *22*, 144 (1982). [199] L. Onsager, J. Am. Chem. Soc. *58*, 1486 (1936). [200] L. Crombie: *Geometrical Isomerism about Carbon-Carbon Double Bonds*, Quart. Rev. (London) *6*, 101 (1952).

[201] R. E. Wood and D. P. Stevenson, J. Am. Chem. Soc. *63*, 1650 (1941). [202] M.-L. Stien, M. Claessens, A. Lopez, and J. Reisse, J. Am. Chem. Soc. *104*, 5902 (1982). [203] H. McNab, J. Chem. Soc., Perkin Trans. II *1981*, 1283. [204] M. V. Jovanovic, Heterocycles *20*, 1987 (1983). [205] Y. Inagaki, R. Okazaki, N. Inamoto, K. Yamada, and H. Kawazura, Bull. Chem. Soc. Jpn. *52*, 2008 (1979). [206] R. Thieme and C. Weiss, Stud. Biophys. *93*, 273 (1983); Chem. Abstr. *99*, 139087w (1983). [207] A. Maercker and J. D. Roberts, J. Am. Chem. Soc. *88*, 1742 (1966). [208] G. Boche and F. Heidenhain, J. Am. Chem. Soc. *101*, 738 (1979); G. Boche, F. Heidenhain, W. Thiel, and R. Eiben, Chem. Ber. *115*, 3167 (1982). [209] J. N. Spencer, C. L. Campanella, E. M. Harris, and W. S. Wolbach, J. Phys. Chem. *89*, 1888 (1985). [210] R. E. Valters and W. Flitsch: *Ring-Chain Tautomerism*, Plenum Publishing Corp., New York, 1985.

[211] O. Arjona, R. Pérez-Ossorio, A. Pérez-Rubalcaba, J. Plumet, and M. J. Santesmases, J. Org. Chem. *49*, 2624 (1984). [212] C. Reichardt, K.-Y. Yun, W. Massa, R. E. Schmidt, O. Exner, and E.-U. Würthwein, Liebigs Ann. Chem. *1985*, 1997. [213] E. M. Arnett: *Solvation Energies of Organic Ions*, J. Chem. Educ. *62*, 385 (1985). [214] G. Modena, C. Paradisi, and G. Scorrano: *Solvation Effects on Basicity and Nucleophilicity*, in F. Bernardi, I. G. Csizmadia, and A. Mangini (eds.): *Organic Sulfur Chemistry*, Elsevier, Amsterdam, 1985, Chapter 10, p. 568ff. [215] M. Mohammad, A. Y. Khan, M. Iqbal, R. Iqbal, and M. Razzaq, J. Am. Chem. Soc. *100*, 7658 (1978); J. Phys. Chem. *85*, 2816 (1981). [216] E. M. Kosower: *Stable Pyridinyl Radicals*, Top. Curr. Chem. *112*, 117 (1983). [217] (a) E. Juaristi and G. Cuevas: *The Anomeric Effect*, CRC Press, Boca Raton/FL, USA, 1994; (b) P. P. Graczyk and M. Mikolajczyk: *Anomeric Effect: Origin and Consequences*, Top. Stereochem. *21*, 159 (1994). [218] W. Schilf, L. Stefaniak, M. Witanowski, and G. A. Webb, Magn. Reson. Chem. *23*, 181 (1985), and references cited therein. [219] D. L. Hughes, J. J. Bergan, and E. J. J. Grabowski, J. Org. Chem. *51*, 2579 (1986). [220] N. Stahl and W. P. Jencks, J. Am. Chem. Soc. *108*, 4196 (1986).

[221] D. A. Hinckley and P. G. Seybold, Spectrochim. Acta, Part A, *44*, 1053 (1988); J. Chem. Educ. *64*, 362 (1987). [222] I. Lopez Arbeloa and K. K. Rohatgi-Mukherjee, Chem. Phys. Lett. *128*, 474 (1986). [223] H. Meier, W. Lauer, and V. Krause, Chem. Ber. *119*, 3382 (1986). [224] S. E. Biali and Z. Rappoport, J. Am. Chem. Soc. *106*, 5641 (1984). [225] J. F. Bunnett and F. P. Olsen, Can. J. Chem. *44*, 1899, 1917 (1966). [226] A. Bagno, G. Scorrano, and R. A. More O'Ferrall; *Stability and Solvation of Organic Cations*, Rev. Chem. Intermed. *7*, 313 (1987). [227] R. Wolfenden, Y.-L. Liang, M. Matthews, and R. Williams, J. Am. Chem. Soc. *109*, 463 (1987). [228] H. Mustroph: *Über die Azo-Hydrazon-Tautomerie*, Z. Chem. *27*, 281 (1987). [229] E. M. Arnett, K. E. Molter, E. C. Marchot, W. H. Donovan, and P. Smith, J. Am. Chem. Soc. *109*, 3788 (1987); see also E. M. Arnett *et al.*, Science *247*, 423 (1990) and Chem. Soc. Rev. *22*, 9 (1993). [230] J. K. Kochi, Angew. Chem. *100*, 1331 (1988); Angew. Chem. Int. Ed. Engl. *27*, 1227 (1988).

[231] C. J. Cramer and D. G. Truhlar, Chem. Rev. *99*, 2161 (1999); particularly Table 3 in Section 6. [232] E. D. Raczynska, M. Decouzon, J.-F. Gal, P.-C. Maria, K. Wozniak, R. Kurg, and S. N. Carins, Trends in Organic Chemistry *7*, 95 (1998). [233] I. A. Koppel *et al.*, J. Am. Chem. Soc. *116*, 3047 (1994), ibid. *122*, 5114 (2000). [234] I. A. Koppel, J. B. Koppel, and V. O. Pihl, Org. React. (Tartu) *25*, 77 (1988). [235] I. Koppel, J. Koppel, P.-C. Maria, J.-F. Gal, R. Notario, V. M. Vlasov, and R. W. Taft, Int. J. Mass Spectrom. and Ion Processes *175*, 61 (1998). [236] I. A. Koppel *et al.*, J. Org. Chem. *63*, 7868 (1998); ibid. *65*, 6202 (2000). [237] I. A. Koppel *et al.*, J. Chem. Soc., Perkin Trans. II 1125 (2000). [238] I. A. Topol *et al.*, J. Phys. Chem. A 104, 9619 (2000). [239] M. Kawata, S. Tenno, S. Kato, and F. Hirata, Chem. Phys. Lett. *240*, 199 (1995). [240] A. Bagno and G. Scorrano, J. Chem. Soc., Perkin Trans. II 1601 (1991).

[241] O. Exner and P. Naus, J. Phys. Org. Chem. *13*, 693 (2000). [242] S. J. Grabowski, T. M. Krygowski, and B. Stępień, J. Phys. Org. Chem. *13*, 740 (2000). [243] A. D. Headley, J. Org. Chem. *56*, 3688 (1991). [244] H. Yamataka, Mustamir, M. Mishima, J. Am. Chem. Soc. *121*, 10223 (1999). [245] W. Blokzijl, J. B. F. N. Engberts, and M. J. Blandamer, J. Chem. Soc., Perkin Trans. II 455 (1994). [246] K. Yamasaki and O. Kajimoto, Chem. Phys. Lett. *172*, 271 (1990). [247] J. J. Dannenberg and R. Rios, J. Phys. Chem. *98*, 6714 (1994). [248] M. Karelson, U. Maran, and A. R. Katritzky, Tetrahedron *52*, 11325 (1996). [249] A. Steigel, R. W. Veith, and M. Braun, Liebigs Ann. Chem. 99 (1989). [250] M. Holler, A. Burger, and J.-F. Biellmann, J. Am. Chem. Soc. *118*, 2153 (1996).

[251] M. W. Wong, K. B. Wiberg, and M. J. Frisch, J. Am. Chem. Soc. *114*, 1645 (1992). [252] O. G. Parchment, N. A. Burton, I. H. Hillier, and M. A. Vincent, J. Chem. Soc., Perkin Trans. II 861 (1993). [253] J. Gao and L. Shao, J. Phys. Chem. *98*, 13772 (1994). [254] D. G. Peck, A. J. Mehta, and K. P. Johnston, J. Phys. Chem. *93*, 4297 (1989). [255] S. Millefiori and A. Millefiori, Bull. Chem. Soc. Jpn. *63*, 2981 (1990). [256] R. Herzfeld and P. Nagy, Curr. Org. Chem. *5*, 373 (2001). [257] L. Antonov, W. M. F. Fabian, D. Nedeltcheva, and F. S. Kamounah, J. Chem. Soc., Perkin Trans. II 1173 (2000). [258] M. Cao, B. J. Teppen, D. M. Miller, J. Pranata, and L. Schäfer, J. Phys. Chem. *98*, 11353 (1994). [259] E. Kleinpeter and A. Koch, J. Phys. Org. Chem. *14*, 566 (2001). [260] E. D. Raczynska and R. Gawinecki, Trends in Organic Chemistry *7*, 85 (1998).

[261] M. W. Wong, K. B. Wiberg, and M. J. Frisch, J. Am. Chem. Soc. *114*, 523 (1992). [262] F. Diederich: *Cyclophanes (Monographs in Supermolecular Chemistry)*, Royal Society of Chemistry, London, 1991, Chapter 3, p. 106ff. [263] V. P. Solov'ev, N. N. Strakhova, O. A. Raevsky, V. Rüdiger, and H.-J. Schneider, J. Org. Chem. *61*, 5221 (1996). [264] K. Izutsu: *Acid-Base Dissociation Constants in Dipolar Aprotic Solvents*, Blackwell Scientific Publications, Oxford/U.K., 1990. [265] T. Bountis: *Proton-Transfer in Hydrogen-Bonded Systems*, Plenum, New York, 1992. [266] J. J. Timoneda and J. T. Hynes, J. Phys. Chem. *95*, 10431 (1991). [267] A. Bagno *et al.*, Chem. Eur. J. *5*, 523 (1999). [268] J.-F. Gal and P.-C. Maria, Progr. Phys. Org. Chem. *17*, 159 (1990). [269] Y. Sueishi, Y. Sugiyama, S. Yamamoto, and N. Nishimura, Bull. Chem. Soc. Jpn. *67*, 572 (1994). [270] H. S. Rzepa and G. A. Suner, J. Chem. Soc., Perkin Trans. II 1397 (1994).

[271] F. Dickert: *Chemosensoren für Gase und Lösungsmitteldämpfe*, Chem. Unserer Zeit *26*, 138 (1992); Fresenius' Z. Anal. Chem. *333*, 615 (1989). [272] Yu. G. Shtyrlin, N. I. Tyryshkin, G. G. Iskhakova, V. V. Gavrilov, V. D. Kiselev, and A. I. Konovalov, Zh. Obshch. Khim. *65*, 1101 (1995); Russ. J. Gen. Chem. *65*, 1004 (1995). [273] R. Makitra and Ya. N. Pirig, Zh. Obshch. Khim. *68*, 425 (1998); Russ. J. Gen. Chem. *68*, 395 (1998). [274] K. B. Wiberg, T. A. Keith, M. J. Frisch, and M. Murcko, J. Phys. Chem. *99*, 9072 (1995). [275] A. A. Stolov and A. B. Remizov, Spectrochim. Acta, Part A *51*, 1919 (1995). [276] W. L. Jorgensen, N. A. McDonald, M. Selmi, and P. R. Rablen, J. Am. Chem. Soc. *117*, 11809 (1995). [277] W. A. Herrebout and B. J. van der Veeken, J. Phys. Chem. *100*, 9671 (1996). [278] M. Kato, I. Abe, and Y. Taniguchi, J. Chem. Phys. *110*, 11982 (1999). [279] C. Cappelli, S. Corni, and J. Tomasi, J. Phys. Chem. A *105*, 10807 (2001). [280] K. B. Wiberg, P. R. Rablen, D. J. Rush, and T. A. Keith, J. Am. Chem. Soc. *117*, 4261 (1995).

[281] M. P. Freitas, R. Rittner, C. F. Tormena, and R. J. Abraham, J. Phys. Org. Chem. *14*, 317 (2001). [282] C. L. Perrin, K. B. Armstrong, and M. A. Fabian, J. Am. Chem. Soc. *116*, 715 (1994). [283] K. B. Wiberg and M. Marquez, J. Am. Chem. Soc. *116*, 2197 (1994). [284] S. A. Maw, R. A. Bryce, R. J. Hall, A. J. Masters, and I. H. Hillier, J. Phys. Chem. B *102*, 4089 (1998). [285] S. Ha, J. Gao, B. Tidor, J. W. Brady, and M. Karplus, J. Am. Chem. Soc. *113*, 1553 (1991); see also B. E. Thomas and K. N. Houk, Chemtracts – Organic Chemistry *4*, 210 (1991). [286] C. J. Cramer and D. G. Truhlar, J. Am. Chem. Soc. *115*, 5745 (1993). [287] J. Sandström, Top. Stereochem. *14*, 84 (1983). [288] R. R. Pappalardo, E. S. Marcos, M. F. Ruiz-López, D. Rinaldi, and J.-L. Rivail, J. Am. Chem. Soc. *115*, 3722 (1993). [289] N. E. Takenaka, R. Hamlin, and D. M. Lemal, J. Am. Chem. Soc. *112*, 6715 (1990). [290] J. M. L. Martin, D. Milstein *et al.*, J. Am. Chem. Soc. *122*, 8797 (2000).

[291] R. A. Ransdell and C. C. Wamser, J. Phys. Chem. *96*, 10572 (1992). [292] K. Komatsu *et al.*, J. Org. Chem. 55, 996 (1990); see also reference [292] in Chapter 2. [293] G. Gritzner, Rev. Inorg. Chem. *11*, 81 (1991). [294] D. J. Cram, Angew. Chem. *98*, 1041 (1986); Angew. Chem. Int. Ed. Engl. *25*, 1039 (1986). [295] (a) F. Vögtle: *Supramolecular Chemistry*, Wiley, Chichester, 1993; (b) H. R. Christen and F. Vögtle: *Organische Chemie – Von den Grundlagen zur Forschung*, Salle, Frankfurt, and Sauerländer, Aarau, 1994, Vol. III, Chapter 54, p. 206ff. (Supramolekulare Chemie). [296] J.-M. Lehn: *Supramolecular Chemistry – Concepts and Perspectives*, VCH, Weinheim, 1995. [297] J.-M. Lehn *et al.* (eds.): *Comprehensive Supramolecular Chemistry*, Vols. 1–11, Pergamon-Elsevier, New York, 1996. [298] F. Diederich *et al.*, J. Chem. Educ. *67*, 813 (1990); Pure Appl. Chem. *62*, 2227 (1990). [299] F. Diederich: *Cyclophanes (Monographs in Supramolecular Chemistry)*, Royal Society of Chemistry, London, 1991; Chapters 3 and 7, particularly Figure 7.1.

[300] V. P. Solov'ev, N. N. Strakhova, O. A. Raevsky, V. Rüdiger, and H.-J. Schneider, J. Org. Chem. *61*, 5221 (1996), and refs. cited therein.

[301] T. Burchard, B. G. Cox, P. Firman, and H. Schneider, Ber. Bunsenges. Phys. Chem. *98*, 1526, 1534 (1994). [302] T. Kuokkanen and S. Sarpola, Acta Chem. Scand. *48*, 258 (1994), and refs. cited therein. [303] K. A. Connors, M. J. Mulski, and A. Paulson, J. Org. Chem. *57*, 1794 (1992). [304] F. Vögtle: *Cyclophane Chemistry – Synthesis, Structures and Reactions*, Wiley, New York, 1993. [305] F. Diederich *et al.*, J. Am. Chem. Soc. *112*, 339 (1990); ibid. *113*, 5420 (1991); ibid. *118*, 6044 (1996). [306] A. Mirzoian and A. E. Kaifer, J. Org. Chem. *60*, 8093 (1995). [307] Y. Tokunaga, D. M. Rudkevich, J. Santamaria, G. Hilmersson, and J. Rebek, Chem. Eur. J. *4*, 1449 (1998), and refs. cited therein. [308] E. Fischer, Ber. Dtsch. Chem. Ges. *27*, 2992 (1894). [309] A. M. Klibanov, Acc. Chem. Res. *23*, 114 (1990). [310] M. N. Gupta, Eur. J. Biochem. *203*, 25 (1992).

[311] A. M. P. Koskinen and A. M. Klibanov (eds.): *Enzymatic Reactions in Organic Media*, Chapman and Hall, London, 1995. [312] G. Carrea and S. Riva, Angew. Chem. *112*, 2312 (2000); Angew. Chem. Int. Ed. Engl. *39*, 2226 (2000). [313] C. R. Wescott and A. M. Klibanov, J. Am. Chem. Soc. *115*, 1629, 10362 (1993); see also A. J. Russell *et al.*, J. Am. Chem. Soc. *118*, 12891 (1996). [314] E. Rubio, A. Fernandez-Mayorales, and A. M. Klibanov, J. Am. Chem. Soc. *113*, 695 (1991). [315] S. Tawaki and A. M. Klibanov, J. Am. Chem. Soc. *114*, 1882 (1992). [316] Y. Hirose, K. Kariya, I. Sasaki, Y. Kurono, H. Ebiike, and K. Achiwa, Tetrahedron Lett. *33*, 7157 (1992). [317] K. Griebenow and A. M. Klibanow, J. Am. Chem. Soc. *118*, 11965 (1996). [318] N. A. Izmailov, Zh. Fiz. Khim. 29, 1050 (1955); see also Yu. Ya. Fialkov: *Rastvoritel' Kak Sredstvo Upravleniya Khimicheskim Protsessom (Solvents as an Agent for Controlling Chemical Processes)*, Publisher Khimiya, Leningrad/USSR, 1990, p. 111 and 123.

Chapter 5

[1] *Cf.* references [7–10] in reference list to Chapter 1. [2] S. Glasstone, K. J. Laidler, and H. Eyring: *The Theory of Rate Processes*, McGraw Hill, New York, London, 1941, p. 400ff. [3] E. A. Moelwyn-Hughes; *Kinetics of Reactions in Solution*, 2nd ed., Oxford University Press, London, 1947. [4] K. A. Connors: *Chemical Kinetics, The Study of Reaction Rates in Solution*, VCH, Weinheim, 1990, Chapter 8, p. 385ff. [5] J. W. Moore and R. G. Pearson: *Kinetics and Mechanism – The Study of Homogeneous Chemical Reactions*, 3rd ed., Wiley, New York, 1981; *Kinetik und Mechanismen homogener chemischer Reaktionen* (translated into German by F. Helfferich and U. Schindewolf), Verlag Chemie, Weinheim, 1964, p. 114ff. [6] A. J. Parker, Quart. Rev. (London) *16*, 163 (1962); Usp. Khim. *32*, 1270 (1963); Adv. Org. Chem. *5*, 1 (1965); Adv. Phys. Org. Chem. *5*, 173 (1967); Usp. Khim. *40*, 2203 (1971); Chem. Rev. *69*, 1 (1969); Pure Appl. Chem. *53*, 1437 (1981). [7] J. E. Leffler and E. Grunwald: *Rates and Equilibria of Organic Reactions*, Wiley, New York, 1963, p. 263ff. [8] K. B. Wiberg: *Physical Organic Chemistry*, Wiley, New York, 1964, p. 374ff. [9] M. Ben-Nun and R. D. Levine, Int. Rev. Phys. Chem. *14*, 215 (1995). [10] B. Tchoubar, *Quelques aspects du rôle des solvants en chimie organique*, Bull. Soc. Chim. Fr. *1964*, 2069; Usp. Khim. *34*, 1227 (1965).

[11] K. J. Laidler: *Chemical Kinetics*, 3rd ed., Harper and Row, Hilversum, 1987. [12] E. S. Amis: *Solvent Effects on Reaction Rates and Mechanisms*, Academic Press, New York, London, 1966. [13] V. A. Pal'm: *Osnovy kolichestvennoi teorii organicheskikh reaktsii*, Izdatel'stvo Khimiya, Leningrad, 1967; *Grundlagen der quantitativen Theorie organischer Reaktionen* (translated into German by G. Heublein), Akademie-Verlag, Berlin, 1971. [14] A. Pross: *Theoretical and Physical Principles of Organic Reactivity*, Wiley, New York, 1995. [15] E. M. Kosower: *An Introduction to Physical Organic Chemistry*, Wiley, New York, 1968, p. 259ff. [16] C. K. Ingold: *Structure and Mechanism in Organic Chemistry*, 2nd ed., Cornell University Press, Ithaca/N.Y., and London, 1969, p. 457ff. and 680ff. [17] L. P. Hammett: *Physical Organic Chemistry – Reaction Rates, Equilibria, and Mechanisms*, 2nd ed., McGraw Hill, New York, 1970, Chapter 8; *Physikalische Organische Chemie – Reaktionsgeschwindigkeiten, Gleichgewichte, Mechanismen* (translated into German by P. Schmid), Verlag Chemie, Weinheim, 1973, Chapter 8, p. 221ff. [18] P. L. Houston: *Chemical*

Kinetics and Reaction Dynamics, McGraw-Hill, New York, 2001. [19] E. A. Moelwyn-Hughes: *The Chemical Statics and Kinetics of Solutions*, Academic Press, London, New York, 1971. [20] K. Schwetlick: *Kinetische Methoden zur Untersuchung von Reaktionsmechanismen*, Deutscher Verlag der Wissenschaften, Berlin, 1971, Chapter 4, p. 139ff.

[21] E. S. Amis and J. F. Hinton: *Solvent Effects on Chemical Phenomena*, Vol. I, Academic Press, New York, London, 1973. [22] J. Burgess: *Solvent Effects*, Inorg. React. Mech. *3*, 312 (1974); ibid. *4*, 236 (1976); ibid. *5*, 260 (1977); ibid. *6*, 278 (1979); ibid. *7*, 287 (1981). [23] M. H. Abraham: *Solvent Effects on Transition States and Reaction Rates*, in A. Streitwieser and R. W. Taft (eds.): Progr. Phys. Org. Chem. *11*, 1 (1974); M. H. Abraham *et al.*, Can. J. Chem. *66*, 2673 (1988). [24] S. H. Lin, K. P. Li, and H. Eyring: *Theory of Reaction Rates in Condensed Phases*, in H. Eyring, D. Henderson, and W. Jost (eds.): *Physical Chemistry – An Advanced Treatise*, Academic Press, New York, 1975, Vol. VII, p. 1ff. [25] C. A. Eckert: *Molecular Thermodynamics of Reactions in Solution*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. VIII, Part I, of A. Weissberger (ed.): *Techniques of Chemistry*, Wiley-Interscience, New York, 1975, p. 1ff. [26] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 13: *Effect of Solvent on Chemical Reactions and Reactivity*. [27] M. R. J. Dack: *The Influence of Solvent on Chemical Reactivity*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. VIII, Part II, of A. Weisberger (ed.): *Techniques of Chemistry*, Wiley-Interscience, New York, 1976, p. 95ff. [28] S. G. Entelis and R. P. Tiger: *Kinetika reaktsii v zhidkoi faze*, Izdatel'stvo Khimiya, Moscow, 1973; *Reaction Kinetics in the Liquid Phase* (translated into English by R. Kondor), Wiley, New York, and Israel Program for Scientific Translations, Jerusalem, 1976. [29] V. Gutmann: *Solvent Effects on the Reactivities of Organometallic Compounds*, Coord. Chem. Rev. *18*, 225 (1976). [30] N. Menschutkin, Z. Phys. Chem. *6*, 41 (1890); ibid. *34*, 157 (1900).

[31] D. J. Cram, B. Rickborn, C. A. Kingsbury, and P. Haberfield, J. Am. Chem. Soc. *83*, 3678 (1961). [32] H. Martin, *Über Beziehungen zwischen Lösungs- und Gasphasenreaktionen*, Angew. Chem. *78*, 73 (1966); Angew. Chem. Int. Ed. Engl. *5*, 78 (1966); Chimia *21*, 439 (1967). [33] J. B. Harkness, G. B. Kistiakowsky, and W. H. Mears, J. Chem. Phys. *5*, 682 (1937). [34] H. Kaufmann and A. Wassermann, J. Chem. Soc. *1939*, 870; A. Wassermann, Monatsh. Chem. *83*, 543 (1952). [35] G. Cöster and E. Pfeil, Chem. Ber. *101*, 4248 (1968); and references cited therein. [36] J. H. Raley, F. F. Rust, and W. E. Vaughan, J. Am. Chem. Soc. *70*, 88, 1336 (1948). [37] E. S. Huyser and R. M. VanScoy, J. Org. Chem. *33*, 3524 (1968). [38] A. Rembaum and M. Szwarc, J. Am. Chem. Soc. *76*, 5975 (1954); M. Levy, M. Steinberg, and M. Szwarc, ibid. *76*, 5978 (1954). [39] H. J. Shine, J. A. Waters, and D. M. Hofman, J. Am. Chem. Soc. *85*, 3613 (1963). [40] S. Winstein and A. H. Fainberg, J. Am. Chem. Soc. *78*, 2770 (1956); ibid. *79*, 5937 (1957).

[41] A. F. Moroni: *Die Abhängigkeit der thermodynamischen Größen des Übergangszustandes einer Reaktion vom Lösungsmittel*, Z. Phys. Chem. (Frankfurt) N. F. *59*, 1 (1968). [42] O. Dimroth, Liebigs Ann. Chem. *377*, 127 (1910); ibid. *399*, 91 (1913). [43] J. H. van't Hoff: *Vorlesungen über theoretische und physikalische Chemie*, Vieweg, Braunschweig, 1898, 1. Heft, p. 217. [44] E. D. Hughes and C. K. Ingold, J. Chem. Soc. *1935*, 244; Trans. Faraday Soc. *37*, 603, 657 (1941); K. A. Cooper, M. L. Dhar, E. D. Hughes, C. K. Ingold, B. J. MacNulty, and L. I. Woolf, J. Chem. Soc. *1948*, 2043. [45] C. A. Bunton: *Nucleophilic Substitution at a Saturated Carbon Atom*, Elsevier, Amsterdam, 1963, Chapter 4. [46] Y. Pocker: *Nucleophilic Substitutions at a Saturated Carbon Atom in Non-Hydroxylic Solvents*, in G. Porter and B. Stevens (eds.): Progress in Reaction Kinetics, Pergamon Press, Oxford, 1961, Vol. 1, p. 215. [47] M. H. Abraham, J. Chem. Soc., Perkin Trans. II, 1343 (1972). [48] D. J. Raber, R. C. Bingham, J. M. Harris, J. L. Fry, and P. v. R. Schleyer, J. Am. Chem. Soc. *92*, 5977 (1970). [49] H. Meerwein and K. van Emster, Ber. Dtsch. Chem. Ges. *55*, 2500 (1922). [50] H. von Halban, Z. Phys. Chem. *67*, 129 (1909).

[51] A. von Hemptinne and A. Bekaert, Z. Phys. Chem. *28*, 225 (1899). [52] H. von Halban, Z. Phys. Chem. *84*, 129 (1913). [53] J. F. Norris and S. W. Prentiss, J. Am. Chem. Soc. *50*, 3042 (1928). [54] A. G. Grimm, H. Ruf, and H. Wolff, Z. Phys. Chem. *B 13*, 301 (1931). [55] N. J. T. Pickles and C. N. Hinshelwood, J. Chem. Soc. *1936*, 1353. [56] E. Tommila and P. Kauranen, Acta Chem. Scand. *8*, 1152 (1954); *13*, 622 (1959). [57] J. D. Reinheimer, J. D. Harley, and W. W.

Meyers, J. Org. Chem. *28*, 1575 (1963). [58] H. Heydtmann, A. P. Schmidt, and H. Hartmann, Ber. Bunsenges. Phys. Chem. *70*, 444 (1966). [59] H. Hartmann and A. P. Schmidt, Z. Phys. Chem. (Frankfurt) N.F. *66*, 183 (1969). [60] C. Lassau and J.-C. Jungers, Bull. Soc. Chim. Fr. *1968*, 2678.

[61] Y. Drougard and D. Decroocq, Bull. Soc. Chim. Fr. *1969*, 2972; G. Berrebi and D. Decroocq, J. Chim. Phys., Phys. Chim. Biol. *71*, 673 (1974). [62] T. Matsui and N. Tokura, Bull. Chem. Soc. Jpn. *43*, 1751 (1970); Y. Kondo, M. Ohnishi, and N. Tokura, ibid. *45*, 3579 (1972). [63] P. Haberfield, A. Nudelman, A. Bloom, R. Romm, and H. Ginsberg, J. Org. Chem. *36*, 1792 (1971). [64] M. H. Abraham, J. Chem. Soc. *B 1971*, 229; M. H. Abraham and R. J. Abraham, J. Chem. Soc., Perkin Trans. II, *1975*, 1677; M. Abraham and P. C. Grellier, ibid. *1976*, 1735. [65] M. Auriel and E. de Hoffmann, J. Am. Chem. Soc. *97*, 7433 (1975); J. Chem. Soc., Perkin Trans. II *1979*, 325. [66] E. R. Swart and L. J. LeRoux, J. Chem. Soc. *1956*, 2110; ibid. *1957*, 406. [67] J. J. Delpuech, Tetrahedron Lett. *1965*, 2111. [68] P. Müller and B. Siegfried, Helv. Chim. Acta *55*, 2400 (1972). [69] E. D. Hughes and D. J. Whittingham, J. Chem. Soc. *1960*, 806; A. V. Eltsov, N. V. Pavlish, and V. A. Ketlinski, Zh. Org. Khim. *14*, 1751 (1978); J. Org. Chem. USSR *14*, 1630 (1978). [70] J. L. Gleave, E. D. Hughes, and C. K. Ingold, J. Chem. Soc. *1935*, 236.

[71] S. Oae and Y. H. Khim, Bull. Chem. Soc. Jpn. *42*, 3528 (1969). [72] D. J. Raber and J. M. Harris, J. Chem. Educ. *49*, 60 (1971). [73] D. V. Banthorpe: *Elimination Reactions*, Elsevier, Amsterdam, 1963, Chapter 2, p. 40. [74] W. H. Saunders and A. F. Cockerill: *Mechanisms of Elimination Reactions*, Wiley-Interscience, New York, 1973. [75] A. Loupy and J. Seyden-Penne, Bull. Soc. Chim. Fr. *1971*, 2306. [76] M. Cocivera and S. Winstein, J. Am. Chem. Soc. *85*, 1702 (1963). [77] P. B. D. de la Mare and R. Bolton: *Electrophilic Additions to Unsaturated Systems*, 2nd ed., Elsevier, Amsterdam, 1981; P. B. D. de la Mare: *Electrophilic Halogenation, Reaction Pathways Involving Attack by Electrophilic Halogens on Unsaturated Compounds*, Cambridge University Press, London, 1976. [78] F. Freeman: *Possible Criteria for Distinguishing between Cyclic and Acyclic Activated Complexes in Addition Reactions*, Chem. Rev. *75*, 439 (1975). [79] G. Heublein: *Neuere Aspekte der elektrophilen Halogenaddition an Olefine*, Z. Chem. *9*, 281 (1969). [80] R. S. Brown, Acc. Chem. Res. *30*, 131 (1997).

[81] M.-F. Ruasse, Adv. Phys. Org. Chem. *28*, 207 (1993). [81a] G. Modena, F. Rivetti, and U. Tonellato, J. Org. Chem. *43*, 1521 (1978). [82] D. S. Campbell and D. R. Hogg, J. Chem. Soc. *B 1967*, 889. [83] L. Rasteikiene, D. Greiciute, M. G. Lin'kova, and I. L. Knunyants: *The Addition of Sulfenyl Chlorides to Unsaturated Compounds*, Usp. Khim. *46*, 1041 (1977); Russ. Chem. Rev. *46*, 548 (1977). [84] T. Beier, H. G. Hauthal, and W. Pritzkow, J. Prakt. Chem. *26*, 304 (1964); G. Collin, U. Jahnke, G. Just, G. Lorenz, W. Pritzkow, M. Röllig, L. Winguth, P. Dietrich, C. E. Doring, H. G. Hauthal, and A. Wiedenhoft, ibid. *311*, 238 (1969). [85] P. P. Kadzyauskas and N. S. Zefirov: *Nitrosochlorination of Alkenes*, Usp. Khim. *37*, 1243 (1968); Russ. Chem. Rev. *37*, 543 (1968). [86] K. N. Houk, J. Liu, N. C. DeMello, and K. R. Condroski, J. Am. Chem. Soc. *119*, 10147 (1997). [86a] R. Renolen and J. Ugelstad, J. Chim. Phys., *57*, 634 (1960). [87] H. Kropf and M. R. Yazdanbachsch, Tetrahedron *30*, 3455 (1974). [88] B. Giese and R. Huisgen, Tetrahedron Lett., *1967*, 1889. [89] M. Neuenschwander and P. Bigler, Helv. Chim. Acta *56*, 959 (1973). [90] (a) R. B. Woodward and R. Hoffmann, Angew. Chem. *81*, 797 (1969); Angew. Chem. Int. Ed. Engl. *8*, 781 (1969); (b) E. A. Halevi: *Orbital Symmetry and Reaction Mechanism*, Springer, Berlin, 1992; (c) K. N. Houk, J. González, and Y. Li, *Pericyclic Reaction Transition States: Passions and Punctilios, 1935–1995*. Acc. Chem. Res. *28*, 81 (1995); (d) I. Fleming: *Pericyclic Reactions*, Oxford University Press, Oxford/U.K., 1999.

[91] R. Huisgen: *Cycloadditionen – Begriff, Einteilung und Kennzeichnung*, Angew. Chem. *80*, 329 (1968); Angew. Chem. Int. Ed. Engl. *7*, 321 (1968). [92] R. Gompper: *Cycloadditionen mit polaren Zwischenstufen*, Angew. Chem. *81*, 348 (1969); Angew. Chem. Int. Ed. Engl. *8*, 312 (1968). [93] P. D. Bartlett: *Mechanisms of Cycloadditons*, Quart. Rev. (London) *24*, 473 (1970). [94] G. Steiner and R. Huisgen, Tetrahedron Lett. *1973*, 3763, 3769; J. Am. Chem. Soc. *95*, 5054, 5055, 5056 (1973); R. Huisgen, R. Schug, and G. Steiner, Bull. Soc. Chim. Fr. *1976*, 1813; R. Huisgen, Acc. Chem. Res. *10*, 117, 199 (1977). [95] J. v. Jouanne, H. Kelm, and R. Huisgen, J. Am. Chem. Soc. *101*, 151 (1979). [96] W. J. Le Noble and R. Mukhtar, J. Am. Chem. Soc. *97*, 5938 (1975). [97] R. Huisgen, R. Schug, and G. Steiner, Angew. Chem. *86*, 47, 48 (1974); Angew. Chem. Int. Ed. Engl.

13, 80, 81 (1974). [98] I. Karle, J. Flippen, R. Huisgen, and R. Schug, J. Am. Chem. Soc. *97*, 5285 (1975); R. Huisgen, Chimia *31*, 13 (1976). [99] S. Proskow, H. E. Simmons, and T. L. Cairns, J. Am. Chem. Soc. *88*, 5254 (1966). [99a] M. Nakahara, Y. Tsuda, M. Sasaki, and J. Osugi, Chem. Lett. (Tokyo) *1976*, 731. [100] R. Huisgen, L. A. Feiler, and P. Otto, Tetrahedron Lett. *1968*, 4485; Chem. Ber. *102*, 3444 (1969); R. Huisgen, L. A. Feiler, and G. Binsch, Chem. Ber. *102*, 3460 (1969); G. Swieton, J. v. Jouanne, H. Kelm, and R. Huisgen, J. Chem. Soc., Perkin Trans. II *1983*, 37.

[101] R. Huisgen and P. Otto, J. Am. Chem. Soc. *90*, 5342 (1968). [102] R. Huisgen and P. Otto, J. Am. Chem. Soc. *91*, 5922 (1969). [103] R. Graf, Liebigs Ann. Chem. *661*, 111 (1963); Org. Synth. *46*, 51 (1966). [104] E. J. Moriconi and W. C. Meyer, J. Org. Chem. *36*, 2841 (1971). [105] J. K. Rasmussen and A. Hassner: *Recent Developments in the Synthetic Use of Chlorosulfonyl Isocyanates*, Chem. Rev. *76*, 389 (1976). [106] K. Clauß, Liebigs Ann. Chem. *722*, 110 (1969). [107] H. Bestian, Pure Appl. Chem. *27*, 611 (1971). [108] P. Bickart, F. W. Carson, J. Jacobus, E. G. Miller, and K. Mislow, J. Am. Chem. Soc. *90*, 4869 (1968); R. Tang and K. Mislow, ibid. *92*, 2100 (1970). [109] C. A. Grob and P. W. Schiess: *Die heterolytische Fragmentierung als Reaktionstypus in der organischen Chemie*, Angew. Chem. *79*, 1 (1967); Angew. Chem. Int. Ed. Engl. *6*, 1 (1967); C. A. Grob: *Mechanismus und Stereochemie der heterolytischen Fragmentierung*, Angew. Chem. *81*, 543 (1969); Angew. Chem. Int. Ed. Engl. *8*, 535 (1969). [110] R. E. Pincock, J. Am. Chem. Soc. *86*, 1820 (1964); ibid. *87*, 1274 (1965).

[111] W. H. Richardson and R. S. Smith, J. Am. Chem. Soc. *91*, 3610 (1969). [112] H. Kwart and P. A. Silver, J. Org. Chem. *40*, 3019 (1975); H. Kwart and T. H. Lilley, ibid. *43*, 2374 (1978). [113] R. B. Woodward and R. Hoffmann: *Die Erhaltung der Orbitalsymmetrie*, Verlag Chemie, Weinheim, 1970; *The Conservation of Orbital Symmetry*, Academic Press, New York, London, 1970. [114] J. Sauer and R. Sustmann: *Mechanistische Aspekte der Diels-Alder-Reaktion. Ein kritischer Überblick*, Angew. Chem. *92*, 773 (1980); Angew. Chem. Int. Ed. Engl. *19*, 779 (1980). [115] S. Seltzer: *The Mechanism of the Diels-Alder Reaction*, in H. Hart and G. J. Karabatsos (eds.): *Advances in Alicyclic Chemistry*, Academic Press, New York, London, 1968, Vol. 2, p. 1ff. (especially p. 13ff.). [116] G. Desimoni, G. Faita, P. P. Righetti, and L. Toma, Tetrahedron *46*, 7951 (1990). [117] R. Huisgen: *Kinetik und Mechanismus 1,3-dipolarer Cycloadditionen*, Angew. Chem. *75*, 604, 742 (1963); Angew. Chem. Int. Ed. Engl. *2*, 565, 633 (1963). [118] R. A. Firestone: *On the Mechanism of 1,3-Dipolar Cycloadditions*, J. Org. Chem. *33*, 2285 (1968); ibid. *37*, 2181 (1972); Tetrahedron *33*, 3009 (1977). [119] R. Huisgen: *The Concerted Nature of 1,3-Dipolar Cycloadditions and the Question of Diradical Intermediates*, J. Org. Chem. *33*, 2291 (1968); ibid. *41*, 403 (1976). [120] A. Padwa: *1,3-Dipolar Cycloaddition Chemistry*, Vols. 1–2, Wiley, New York, 1984.

[121] R. A. Fairclough and C. N. Hinshelwood, J. Chem. Soc. *1938*, 236 (Cyclopentadiene + *p*-benzoquinone). [122] Y. Yukawa and A. Isohisa, Mem. Inst. Sci. Ind. Research Osaka Univ. *10*, 191 (1953); Chem. Abstr. *48*, 7598 (1954) (Cyclohexa-1,3-diene + maleic anhydride). [123] L. J. Andrews and R. M. Keefer, J. Am. Chem. Soc. *77*, 6284 (1955) (9,10-Dimethylanthracene + maleic anhydride). [124] J. A. Berson, Z. Hamlet, and W. A. Mueller, J. Am. Chem. Soc. *84*, 297 (1962) (Cyclopentadiene + methyl acrylate). [125] P. Brown and R. C. Cookson, Tetrahedron *21*, 1977 (1965) (Anthracene + tetracyanoethene). [126] K. F. Wong and C. A. Eckert, Ind. Eng. Chem. Process Design Develop. *8*, 568 (1969); Chem. Abstr. *71*, 116937 (1969); Trans. Faraday Soc. *66*, 2313 (1970); R. A. Grieger and C. A. Eckert, J. Am. Chem. Soc. *92*, 7149 (1970); see also ref. [25] (Isoprene + maleic anhydride). [127] M. J. S. Dewar and R. S. Pyron, J. Am. Chem. Soc. *92*, 3098 (1970) (Isoprene + maleic anhydride). [128] P. Haberfield and A. K. Ray, J. Org. Chem. *37*, 3093 (1972). [129] B. Blankenburg, H. Fiedler, M. Hampel, H. G. Hauthal, G. Just, K. Kahlert, J. Korn, K.-H. Müller, W. Pritzkow, Y. Reinhold, M. Röllig, E. Sauer, D. Schnurpfeil, and G. Zimmermann, J. Prakt. Chem. *316*, 804 (1974). [130] M. E. Burrage, R. C. Cookson, S. S. Gupte, and I. D. R. Stevens, J. Chem. Soc., Perkin Trans. II *1975*, 1325.

[131] R. Huisgen, H. Stangl, H. J. Sturm, and H. Wagenhofer, Angew. Chem. *73*, 170 (1961). [132] R. Huisgen, H. Seidl, and I. Brüning, Chem. Ber. *102*, 1102 (1969). [133] M. K. Meilahn, B. Cox, and M. E. Munk, J. Org. Chem. *40*, 819 (1975). [134] P. K. Kadaba, J. Heterocycl. Chem. *6*, 587 (1969); Tetrahedron *25*, 3053 (1969); J. Org. Chem. *41*, 1073 (1976). [134a] A. Eckell, M. V.

George, R. Huisgen, and A. S. Kende, Chem. Ber. *110*, 578 (1977). [135] H. M. R. Hoffmann: *Die En-Reaktion*, Angew. Chem. *81*, 597 (1969); Angew. Chem. Int. Ed. Engl. *8*, 556 (1969). [136] R. Huisgen and H. Pohl, Chem. Ber. *93*, 527 (1960). [137] A. A. Frimer; *The Reaction of Singlet Oxygen with Olefins*, Chem. Rev. *79*, 359 (1979). [138] C. S. Foote, Pure Appl. Chem. *27*, 635 (1971); C. S. Foote and R. W. Denny, J. Am. Chem. Soc. *93*, 5168 (1971). [139] R. H. Young, K. Wehrly, and R. L. Martin, J. Am. Chem. Soc. *93*, 5774 (1971). [140] J. E. Baldwin and J. A. Kapecki, J. Am. Chem. Soc. *92*, 4868 (1970).

[141] E. Schaumann, Chem. Ber. *109*, 906 (1976). [142] G. Wittig: *From Diyls over Ylides to My Idyll*, Acc. Chem. Res. *7*, 6 (1974). [143] E. Vedejs and M. J. Peterson, Top. Stereochem. *21*, 1 (1994); Adv. Carbanion Chem. *2*, 1 (1996). [144] A. J. Speziale and D. E. Bissing, J. Am. Chem. Soc. *85*, 3878 (1963). [145] C. Rüchardt, P. Panse, and S. Eichler, Chem. Ber. *100*, 1144 (1967). [146] P. Frøyen, Acta Chem. Scand. *26*, 2163 (1972). [147] G. Aksnes and F. Y. Khalil, Phosphorus *2*, 105 (1972); ibid. *3*, 37, 79, 103 (1973). [148] E. Maccarone and G. Perrini, Gazz. Chim. Ital. *112*, 447 (1982). [149] F. Ramirez and S. Dershowitz, J. Org. Chem. *22*, 41 (1957). [150] D. I. Coomber and J. R. Partington, J. Chem. Soc. *1938*, 1444.

[151] (a) D. C. Wigfield and S. Feiner, Can. J. Chem. *48*, 855 (1970); (b) T. Mitsuhashi and G. Yamamoto, Bull. Chem. Soc. Jpn. *63*, 643 (1990). [152] D. C. Berndt, J. Chem. Eng. Data *14*, 112 (1969). [153] W. N. White and E. F. Wolfarth, J. Org. Chem. *35*, 2196, 3585 (1970). [154] S. J. Rhoads and N. R. Raulins: *The Claisen and Cope Rearrangements*, Org. React. *22*, 1 (1975). [155] R. Wehrli, D. Bellus, H.-J. Hansen, and H. Schmid, Chimia *30*, 416 (1976); Helv. Chim. Acta *60*, 1325 (1977). [156] S. G. Smith, J. Am. Chem. Soc. *83*, 4285 (1961). [157] R. Huisgen, A. Dahmen, and H. Huber, Tetrahedron Lett. *1969*, 1461, 1465. [158] H. Mayr and R. Huisgen, J. Chem. Soc., Chem. Commun. *1976*, 57. [158a] N. S. Isaacs and A. A. R. Laila, J. Chem. Soc., Perkin Trans. II, 1470 (1976). [158b] M. T. Reetz, Chem. Ber. *110*, 965 (1977); M. T. Reetz, N. Greif, and M. Kliment, ibid. *111*, 1095 (1978); M. T. Reetz, Angew. Chem. *91*, 185 (1979); Angew. Chem. Int. Ed. Engl. *18*, 173 (1979). [159] C. Walling: *Free Radicals in Solutions*, Wiley, New York, London, 1957. [160] E. S. Huyser: *Solvent Effects in Radical Reactions*, in G. H. Williams (ed.): *Advances in Free-Radical Chemistry*, Logos Press and Academic Press, New York, London, 1965, Vol. 1, p. 77ff.

[161] H. Sakurai and A. Hosomi: *Solvent Effects in Free-Radical Reactions*, Yuki Gosei Kagaku Kyokai Shi *25*, 1108 (1967); Chem. Abstr. *68*, 104, 254f (1968). [162] Yu. L. Spirin: *Reactivity of Radicals and Molecules in Radical Reactions*, Usp. Khim. *38*, 1201 (1969); Russ. Chem. Rev. *38*, 529 (1969). [163] J. C. Martin: *Solvation and Association*, in J. K. Kochi (ed.): *Free Radicals*, Wiley, New York, 1973, Vol. 1, Chapter 20, p. 493ff. [164] E. T. Denisov: *Konstanty Skorosti Gomoliticheskikh Zhidkofaznykh Reaktsii*, Nauka Press, Moscow, 1971; *Liquid-Phase Reaction Rate Constants* (translated into English by R. K. Johnston), IFI/Plenum Data Company, New York, 1974. [165] J. M. Tanko and N. K. Suleman: *Solvent Effects in the Reactions of Neutral Free Radicals*, in J. A. M. Simões, A. Greenberg, and J. F. Lieberman (eds.): *Energetics and Reactivities of Organic Free Radical Reactions*, Kluwer, Dordrecht/Netherlands, 1996, Vol. 4, p. 224; Chem. Abstr. *125*, 274936t (1996). [166] See also reference [28], chapter IX, concerning homolytic reactions, and references cited therein. [167] K. Ziegler, P. Orth, and K. Weber, Liebigs Ann. Chem. *504*, 131 (1933); K. Ziegler, A. Seib, K. Knoevenagel, P. Herte, and F. Andreas, ibid. *551*, 150 (1942). [168] H. Lankamp, W. T. Nauta, and C. MacLean, Tetrahedron Lett. *1968*, 249. [169] H. A. Staab, H. Brettschneider, and H. Brunner, Chem. Ber. *103*, 1101 (1970). [170] P. Jacobson, Ber. Dtsch. Chem. Ges. *38*, 196 (1905).

[171] B. S. Tanaseichuk, L. G. Tikhonova, and A. P. Dydykina, Zh. Org. Khim. *9*, 1273 (1973); J. Org. Chem. USSR *9*, 1301 (1973). [172] E. S. Huyser and R. M. VanScoy, J. Org. Chem. *33*, 3524 (1968). [173] See reference [164], table 7 on page 37, and the accompanying references. [174] S. D. Ross and M. A. Fineman, J. Am. Chem. Soc. *73*, 2176 (1951). [175] M. Levy, M. Steinberg, and M. Szwarc, J. Am. Chem. Soc. *76*, 5978 (1954). [176] H. J. Shine, J. A. Waters, and D. M. Hoffman, J. Am. Chem. Soc. *85*, 3613 (1963). [177] K. Nozaki and P. D. Bartlett, J. Am. Chem. Soc. *68*, 1686 (1946). [178] C. G. Swain, W. H. Stockmayer, and J. T. Clarke, J. Am. Chem. Soc. *72*, 5426 (1950). [179] W. R. Foster and G. H. Williams, J. Chem. Soc. *1962*, 2862. [180] R. C. Lamb, J. G. Pacifici, and P. W. Ayers, J. Am. Chem. Soc. *87*, 3928 (1965).

[181] C. Walling, H. P. Waits, J. Milanovic, and C. G. Pappiaonnou, J. Am. Chem. Soc. *92*, 4927 (1970). [182] R. C. Petersen, J. H. Markgraf, and S. D. Ross, J. Am. Chem. Soc. *83*, 3819 (1961). [183] See reference [164], table 12 on page 57, and the accompanying references. [184] A. F. Moroni, Makromol. Chem. *105*, 43 (1967). [185] O. Yamamoto, J. Yamashita, and H. Hashimoto, Kogyo Kagaku Zasshi *71*, 223 (1968); Chem. Abstr. *69*, 66644u (1968). [186] R. Huisgen and H. Nakaten, Liebigs Ann. Chem. *586*, 70 (1954). [187] M. G. Alder and J. E. Leffler, J. Am. Chem. Soc. *76*, 1425 (1954); J. E. Leffler and R. A. Hubbard, J. Org. Chem. *19*, 1089 (1954). [188] W. G. Bentrude and A. K. MacKnight, Tetrahedron Lett. *1966*, 3147. [189] R. Kerber, O. Nuyken, and R. Steinhausen, Makromol. Chem. *175*, 3225 (1975). [190] C. E. Dykstra and H. F. Schaefer, J. Am. Chem. Soc. *97*, 7210 (1975).

[191] W. Lobunez, J. R. Rittenhouse, and J. G. Miller, J. Am. Chem. Soc. *80*, 3505 (1958). [192] P. S. Engel, Chem. Rev. *80*, 99 (1980). [193] A. Schulz and C. Rüchardt, Tetrahedron Lett. *1976*, 3883; W. Duismann and C. Rüchardt, Chem. Ber. *111*, 596 (1978). [194] C. G. Overberger, J.-P. Anselme, and J. R. Hall, J. Am. Chem. Soc. *85*, 2752 (1963). [195] C. Rüchardt: *Nichtkatalysierte Perester-Zersetzungen*, Fortschr. Chem. Forsch. *6*, 251 (1996); Usp. Khim. *37*, 1402 (1968). [196] C. Rüchardt: *Zusammenhänge zwischen Struktur und Reaktivität in der Chemie freier Radikale*, Angew. Chem. *82*, 845 (1970); Angew. Chem. Int. Ed. Engl. *9*, 830 (1970). [197] P. D. Bartlett and R. R. Hiatt, J. Am. Chem. Soc. *80*, 1398 (1958). [198] C. Rüchardt and H. Böck, Chem. Ber. *104*, 577 (1971). [199] M. Trachtman and J. G. Miller, J. Am. Chem. Soc. *84*, 4828 (1962). [200] J. P. Engstrom and J. C. DuBose, J. Org. Chem. *38*, 3817 (1973).

[201] P. D. Bartlett and C. Rüchardt, J. Am. Chem. Soc. *82*, 1756 (1960). [202] C. Rüchardt and H. Schwarzer, Chem. Ber. *99*, 1861 (1966). [203] A. A. Turovskii, R. V. Kucher, A. M. Ustinova, and A. E. Batog, Zh. Obshch, Khim. *45*, 860 (1975); J. Gen. Chem. USSR *45*, 844 (1975). [204] C. Rüchardt and H. Schwarzer, Chem. Ber. *99*, 1871 (1966). [205] T. W. Koenig and J. C. Martin, J. Org. Chem. *29*, 1520 (1964); J. Am. Chem. Soc. *86*, 1771 (1964). [206] D. L. Tuleen, W. G. Bentrude, and J. C. Martin, Tetrahedron Lett. *1962*, 229; J. Am. Chem. Soc. *85*, 1938 (1963); P. Livant and J. C. Martin, ibid. *98*, 7851 (1976). [207] C. Rüchardt and H.-J. Quadbeck-Seeger, Chem. Ber. *102*, 3525 (1969). [208] R. Criegee and R. Kaspar, Liebigs Ann. Chem. *560*, 127 (1948); R. Criegee, Fortschr. Chem. Forsch. *1*, 508 (1950), especially p. 551. [209] P. D. Bartlett and B. T. Storey, J. Am. Chem. Soc. *80*, 4954 (1958). [210] T. Koenig and W. Brewer, J. Am. Chem. Soc. *86*, 2728 (1964).

[211] E. Hedaya, R. L. Hinman, L. M. Kibler, and S. Theodoropulos, J. Am. Chem. Soc. *86*, 2727 (1964). [212] C. Rüchardt and H. Schwarzer, Chem. Ber. *99*, 1878 (1966). [213] E. M. Kosower and M. Mohammad, J. Am. Chem. Soc. *93*, 2709 (1971). [214] E. M. Kosower and I. Schwager, J. Am. Chem. Soc. *86*, 4493, 5528 (1964). [215] E. M. Kosower and M. Mohammad, J. Am. Chem. Soc. *90*, 3271 (1968); ibid. *93*, 2709, 2713 (1971). [216] R. Kerber, Z. Elektrochem., Ber. Bunsenges. Phys. Chem. *63*, 296 (1959). [217] G. E. Zaikov and Z. K. Maizus, Dokl. Akad. Nauk SSSR *150*, 116 (1963); Chem. Abstr. *59*, 5831 (1963); G. E. Zaikov, Z. K. Maizus, and N. M. Emanuel, Kinetika i Kataliz *7*, 401 (1966); Chem. Abstr. *65*, 10459 (1966); G. E. Zaikov, A. A. Vichutinskii, Z. K. Maizus, and N. M. Emanuel, Dokl. Akad. Nauk SSSR *168*, 1096 (1966); Chem. Abstr. *65*, 15184 (1966). [218] D. G. Hendry and G. A. Russell, J. Am. Chem. Soc. *86*, 2368 (1964). [219] J. A. Howard and K. U. Ingold, Can. J. Chem. *42*, 1044, 1250 (1964); ibid. *44*, 1119 (1966). [220] E. Niki, Y. Kamiya, and N. Ohta, Bull. Chem. Soc. Jpn. *42*, 3224 (1969).

[221] G. A. Russell, J. Am. Chem. Soc. *79*, 2977 (1957); ibid. *80*, 4987, 4997, 5002 (1958). [222] C. Walling and B. B. Jacknow, J. Am. Chem. Soc. *82*, 6108, 6113 (1960). [223] C. Walling: *Some Aspects of the Chemistry of Alkoxy Radicals*, Pure Appl. Chem. *15*, 69 (1967). [224] C. Walling and P. Wagner, J. Am. Chem. Soc. *85*, 2333 (1963); ibid. *86*, 3368 (1964). [225] C. D. Cook and B. E. Norcross, J. Am. Chem. Soc. *81*, 1176 (1959). [226] R. G. Pearson, J. Chem. Phys. *20*, 1478 (1952). [227] C. Walling and D. Bristol, J. Org. Chem. *36*, 733 (1971). [228] J. H. Hildebrand and R. L. Scott; *The Solubility of Nonelectrolytes*, 3rd ed., Dover, New York, 1964. [229] J. H. Hildebrand, J. M. Prausnitz, and R. L. Scott: *Regular and Related Solutions*, Van Nostrand-Reinhold, Princeton, 1970. [230] G. Scatchard, Chem. Rev. *8*, 321 (1931); *10*, 229 (1932).

[231] A. F. M. Barton: *Handbook of Solubility Parameters and other Cohesion Parameters*, CRC Press, Boca Raton/Florida, 1983. [232] M. R. J. Dack: *The Importance of Solvent Internal Pressure*

and Cohesion to Solution Phenomena, Chem. Soc. Rev. (London) *4*, 211 (1975); M. R. J. Dack, Aust. J. Chem. *28*, 1643 (1975). [233] M. Richardson and F. G. Soper, J. Chem. Soc. *1929*, 1873. [234] S. Glasstone, J. Chem. Soc. *1936*, 723. [235] R. J. Ouellette and S. H. Williams, J. Am. Chem. Soc. *93*, 466 (1971). [236] H. F. Herbrandson and F. R. Neufeld, J. Org. Chem. *31*, 1140 (1966). [237] R. C. Neuman, J. Org. Chem. *37*, 495 (1972). [238] H. Burrell: *Solubility Parameter Values*, in J. Brandrup and E. H. Immergut (eds.): *Polymer Handbook*, 2nd ed., Wiley-Interscience, New York, 1975, p. IV-337ff. [239] M. R. J. Dack, J. Chem. Educ. *51*, 231 (1974). [240] J. G. Kirkwood: *Theory of Solutions of Molecules Containing Widely Separated Charges*, J. Chem. Phys. *2*, 351 (1934).

[241] J. G. Kirkwood and F. Westheimer, J. Chem. Phys. *6*, 506 (1938); C. Tanford and J. G. Kirkwood, J. Am. Chem. Soc. *79*, 5333, 5340, 5348 (1957). [242] K. J. Laidler and P. A. Landskroener, Trans. Faraday Soc. *52*, 200 (1956); K. J. Laidler, Suomen Kemistilehti *33*, A 44 (1960); Chem. Abstr. *56*, 2920 (1962). [243] K. Hiromi, Bull. Chem. Soc. Jpn. *33*, 1251, 1264 (1960). [244] E. S. Amis, J. Chem. Educ. *29*, 337 (1952); ibid. *30*, 351 (1953); Anal. Chem. *27*, 1672 (1955). [245] A. D. Stepukhovich, N. I. Lapshova, and T. D. Efimova, Zh. Fiz. Khim. *35*, 2532 (1961); Chem. Abstr. *59*, 3739 (1963). [246] M. Watanabe and R. M. Fuoss, J. Am. Chem. Soc. *78*, 527 (1965). [247] L. M. Litvinenko and V. A. Savelova, Reakts. Sposobnost Org. Soedin. *5*, 838 (1968); Chem. Abstr. *70*, 86878x (1969). [248] E. T. Caldin and J. Peacock, Trans. Faraday Soc. *51*, 1217 (1955). [249] J. H. Beard and P. H. Plesch, J. Chem. Soc. *1965*, 3682. [250] I. A. Koppel and V. A. Pal'm, Reakts. Sposobnost Org. Soedin. *4*, 862 (1967); Chem. Abstr. *69*, 99978j (1968); ibid. *4*, 892 (1967); Chem. Abstr. *69*, 99980d (1968).

[251] K. J. Laidler and H. Eyring, Ann. N. Y. Acad. Sci. *39*, 303 (1940); Chem. Abstr. *35*, 2777 (1941). [252] J. E. Quinlan and E. S. Amis, J. Am. Chem. Soc. *77*, 4187 (1955). [253] G. Scatchard, Chem. Rev. *10*, 229 (1932). [254] E. S. Amis and V. K. LaMer, J. Am. Chem. Soc. *61*, 905 (1939). [255] E. S. Amis and J. E. Price, J. Phys. Chem. *47*, 338 (1943). [256] J. N. Brønsted, Z. Phys. Chem. *102*, 169 (1922); *115*, 337 (1925). [257] N. Bjerrum, Z. Phys. Chem. *108*, 82 (1924); *118*, 251 (1925). [258] J. A. Christiansen, Z. Phys. Chem. *113*, 35 (1924). [259] P. Debye and E. Hückel, Phys. Z. *24*, 185, 305 (1923); E. Hückel, ibid. *26*, 93 (1925). [260] C. W. Davies; *Salt Effects in Solution Kinetics*, in G. Porter (ed.): *Progress in Reaction Kinetics*, Vol. 1, p. 161ff., Pergamon Press, Oxford, 1961.

[261] A. Streitwieser jr.: *Solvolytic Displacement Reactions*, McGraw Hill, New York, 1962; Chem. Reviews *56*, 571 (1956). [262] E. S. Gould: *Mechanism and Structure in Organic Chemistry*, Holt, New York, 1959; *Mechanismus und Struktur in der organischen Chemie* (translated by G. Koch), 2nd ed., Verlag Chemie, Weinheim, 1969. [263] H. G. O. Becker: *Einführung in die Elektronentheorie organisch-chemischer Reaktionen*, 3rd ed., Deutscher Verlag der Wissenschaften, Berlin, 1974. [264] C. G. Swain and R. W. Eddy, J. Am. Chem. Soc. *70*, 2989 (1948). [265] S. G. Smith, A. H. Fainberg, and S. Winstein, J. Am. Chem. Soc. *83*, 618 (1961). [266] J. A. Leary and M. Kahn, J. Am. Chem. Soc. *81*, 4173 (1959). [267] E. A. S. Cavell, J. Chem. Soc. *1958*, 4217; E. A. S. Cavell and J. A. Speed, ibid. *1960*, 1453. [268] S. Wideqvist, Arkiv Kemi *9*, 475 (1956); Chem. Abstr. *50*, 15187 (1956). [269] J. Murto, Suomen Kemistilehti *34*, 92 (1961); Chem. Abstr. *58*, 7803 (1963). [270] S. R. Palit, J. Org. Chem. *12*, 752 (1947).

[271] J. F. Bunnett: *Nucleophilic Reactivity*, in H. Eyring, C. J. Christensen, and H. S. Johnston (eds.): Ann. Rev. Phys. Chem. *14*, 271 (1963); Chem. Abstr. *59*, 14588 (1963). [272] C. G. Swain and C. B. Scott, J. Am. Chem. Soc. *75*, 141 (1953). [273] J. O. Edwards, J. Am. Chem. Soc. *76*, 1540 (1954); J. O. Edwards and R. G. Pearson, ibid. *84*, 16 (1962). [274] C. D. Ritchie and P. O. I. Virtanen, J. Am. Chem. Soc. *94*, 4966 (1972); ibid. *95*, 1882 (1973); ibid. *97*, 1170 (1975); ibid. *105*, 7313 (1983); Acc. Chem. Res. *5*, 348 (1972). [275] R. G. Pearson: *The Influence of the Reagent on Organic Reactivity*, in N. B. Chapman and J. Shorter (eds.): *Advances in Linear Free Energy Relationships*, Plenum Press, London, New York, 1972, Chapter 6, p. 281ff. [276] R. G. Pearson, H. Sobel, and J. Songstad, J. Am. Chem. Soc. *90*, 319 (1968). [277] R. G. Pearson and J. Songstad, J. Org. Chem. *32*, 2899 (1967). [278] W. M. Weaver and J. D. Hutchison, J. Am. Chem. Soc. *86*, 261 (1964). [279] S. Winstein, L. G. Savedoff, S. Smith, I. D. R. Stevens, and J. S. Gall, Tetrahedron Lett. No. 9, 24 (1960). [280] R. Fuchs and K. Mahendran, J. Org. Chem. *36*, 730 (1971).

[281] R. F. Rodewald, K. Mahendran, J. L. Bear, and R. Fuchs, J. Am. Chem. Soc. *90*, 6698 (1968); R. Fuchs, J. L. Bear, and R. F. Rodewald, ibid. *91*, 5797 (1969). [282] C. L. Liotta, E. E. Grisdale, and H. P. Hopkins, Tetrahedron Lett. *1975*, 4205. [283] W. T. Ford, R. J. Hauri, and S. G. Smith, J. Am. Chem. Soc. *96*, 4316 (1974). [284] J. E. Gordon and P. Varughese, J. Chem. Soc., Chem. Commun. *1971*, 1160. [285] J. I. Brauman, W. N. Olmstead, and C. A. Lieder, J. Am. Chem. Soc. *96*, 4030 (1974); W. N. Olmstead and J. I. Brauman, ibid. *99*, 4219 (1977). [286] M. S. Puar: *Nucleophilic Reactivities of the Halide Anions*, J. Chem. Educ. *47*, 473 (1970). [287] C. Minot and N. T. Anh: *Reversal of Nucleophilic Orders by Solvent Effects*, Tetrahedron Lett. *1975*, 3905. [288] G. Choux and R. L. Benoit, J. Am. Chem. Soc. *91*, 6221 (1969). [289] J. E. Gordon: *The Organic Chemistry of Electrolyte Solutions*, Wiley-Interscience, New York, 1975, p. 299ff. and 471ff. [290] D. K. Bohme, G. I. Mackay, and J. D. Payzant, J. Am. Chem. Soc. *96*, 4027 (1974); K. Tanaka, G. I. Mackay, J. D. Payzant, and D. K. Bohme, Can. J. Chem. *54*, 1643 (1976).

[291] R. Alexander, E. C. F. Ko, A. J. Parker, and T. J. Broxton, J. Am. Chem. Soc. *90*, 5049 (1968); see also B. G. Cox, G. R. Hedwig, A. J. Parker, and D. W. Watts, Aust. J. Chem. *27*, 477 (1974). [292] B. G. Cox and A. J. Parker, J. Am. Chem. Soc. *95*, 408 (1973). [292a] A. J. Parker, U. Mayer, R. Schmid, and V. Gutmann, J. Org. Chem. *43*, 1843 (1978). [293] H. Suhr, Ber. Bunsenges. Phys. Chem. *67*, 893 (1963); Chem. Ber. *97*, 3277 (1964); Liebigs Ann. Chem. *687*, 175 (1965). [294] F. Madaule-Aubry: *Le Rôle en Chimie de Certaines Solvants Dipolaires Aprotiques*, Bull. Soc. Chim. Fr. *1966*, 1456. [295] H. Liebig: *Präparative Chemie in aprotischen Lösungsmitteln*, Chemiker-Ztg. *95*, 301 (1971). [296] J. F. Coetzee: *Ionic Reactions in Acetonitrile*, Progr. Phys. Org. Chem. *4*, 45 (1967). [297] R. S. Kittila: *A Review of Catalytic and Synthetic Applications for DMF and DMAC*, Bulletin DuPont de Nemours and Co., Wilmington/USA, 1960 (and Supplement 1962). [298] S. S. Pizey: *Dimethylformamide*, in S. S. Pizey, *Synthetic Reagents*, Wiley, New York, London, 1974, Vol. I, P. 1ff. [299] D. Martin and H. G. Hauthal: *Dimethylsulfoxid*, Akademie-Verlag, Berlin, 1971; *Dimethyl sulphoxide*, Van Nostrand-Reinhold, London, 1975. [299a] E. Buncel and H. Wilson: *Physical Organic Chemistry of Reactions in Dimethyl sulphoxide*, Adv. Phys. Org. Chem. *14*, 133 (1977). [300] H. Normant: *Hexamethylphosphorsäuretriamid*, Angew. Chem. *79*, 1029 (1967); Angew. Chem. Int. Ed. Engl. *6*, 1046 (1967); Bull. Soc. Chim. Fr. *1968*, 791; Usp. Khim. *39*, 990 (1970); Russ. Chem. Rev. *39*, 457 (1970).

[301] P. E. Pfeffer, T. A. Foglia, P. A. Barr, I. Schmeltz, and L. S. Silbert, Tetrahedron Lett. *1972*, 4063. [302] J. E. Shaw, D. C. Kunerth, and J. J. Sherry, Tetrahedron Lett. *1973*, 689. [303] R. G. Smith, A. Vanterpool, and H. J. Kulak, Can. J. Chem. *47*, 2015 (1969). [304] D. J. Cram, B. Rickborn, and G. R. Knox, J. Am. Chem. Soc. *82*, 6412 (1960). [305] A. J. Parker, J. Chem. Soc. *1961*, 1328. [306] T. J. Wallace, J. E. Hoffmann, and A. Schriesheim, J. Am. Chem. Soc. *85*, 2739 (1963). [307] J. T. Maynard, J. Org. Chem. *28*, 112 (1963). [308] J. F. Normant and H. Deshayes, Bull. Soc. Chim. Fr. *1967*, 2455. [309] S. Bank, J. Org. Chem. *37*, 114 (1972). [310] A. I. Shatenshtein, E. A. Gvozdeva, and Yu. I. Ranneva, Zh. Obshch. Khim. *41*, 1818 (1971); J. Gen. Chem. USSR *41*, 1827 (1971).

[311] P. K. Kadaba, Synthesis *1973*, 71; J. Org. Chem. *41*, 1073 (1976). [312] E. Tommila and M.-L. Murto, Acta Chem. Scand. *17*, 1947 (1963). [313] R. Goitein and T. C. Bruice, J. Phys. Chem. *76*, 432 (1972). [314] H.-L. Pan and T. L. Fletcher, Chem. Ind. (London) *1969*, 240. [315] G. Bähr and G. Schleitzer, Chem. Ber. *88*, 1771 (1955). [316] O. Popovych: *Estimation of Medium Effects for Single Ions in Nonaqueous Solvents*, Critical Reviews in Analytical Chemistry *1*, 73 (1970); Chem. Abstr. *75*, 133680s (1971). [317] B. G. Cox: *Electrolyte Solutions in Dipolar Aprotic Solvents*, Annual Reports on the Progress of Chemistry *A 70*, 249 (1973), and references cited therein. [318] B. G. Cox and A. J. Parker, J. Am. Chem. Soc. *95*, 402 (1973). [319] D. W. Watts: *Reaction Kinetics and Mechanism*, in A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, Chapter 6, p. 681ff. [320] R. Alexander and A. J. Parker, J. Am. Chem. Soc. *90*, 3313 (1968); R. Alexander, A. J. Parker, J. H. Sharp, and W. E. Waghorne, ibid. *94*, 1148 (1972).

[321] D. J. Cram: *Fundamentals of Carbanion Chemistry*, Academic Press, New York, London, 1965, p. 32ff. [322] E. Buncel: *Carbanions: Mechanistic and Isotopic Aspects*, Elsevier, Amsterdam, 1975. [323] A. J. Hubert and H. Reimlinger: *The Isomerization of Olefins, Part I, Base-Catalysed*

Isomerization of Olefins, Synthesis *1969*, 97. [324] H. Pines: *Base-Catalysed Carbon-Carbon Addition of Hydrocarbons and of Related Compounds*, Acc. Chem. Res. *7*, 155 (1974). [325] C. C. Price and W. H. Snyder, J. Am. Chem. Soc. *83*, 1773 (1961); Tetrahedron Lett. No. 2, 69 (1962). [326] J. Sauer and H. Prahl, Chem. Ber. *102*, 1917 (1969). [327] R. Greenwald, M. Chaykovsky, and E. J. Corey, J. Org. Chem. *28*, 1128 (1963). [328] D. J. Cram, M. R. V. Sahyun, and G. R. Knox, J. Am. Chem. Soc. *84*, 1734 (1962). [329] H. H. Szmant: *Der Mechanismus der Wolff-Kishner-Reaktionen: Reduktion, Eliminierung und Isomerisierung*, Angew. Chem. *80*, 141 (1968); Angew. Chem. Int. Ed. Engl. *7*, 120 (1968). [330] H. H. Szmant and M. N. Roman, J. Am. Chem. Soc. *88*, 4034 (1966).

[331] M. R. V. Sahyun and D. J. Cram, J. Am. Chem. Soc. *85*, 1263 (1963). [332] S. F. Acree, Am. Chem. J. *48*, 352 (1912). [333] M. Szwarcz (ed.): *Ions and Ion Pairs in Organic Reactions*, Wiley-Interscience, New York, 1972/74, Vols. 1 and 2. [334] I. P. Beletskaya: *Ions and Ion-Pairs in Nucleophilic Aliphatic Substitutions*, Usp. Khim. *44*, 2205 (1975); Russ. Chem. Rev. *44*, 1067 (1975). [335] F. Guibe and G. Bram: *Réactivité S_N2 des formes dissociée et associée aux cations alcalins des nucléophiles anioniques*, Bull. Soc. Chim. Fr. *1975*, 933. [336] E. A. Kovrizhnykh and A. I. Shatenshtein: *Effect of Electron-Donor Solvents on the Reactivity of Lithium Alkyls*, Usp. Khim. *38*, 1836 (1969); Russ. Chem. Rev. *38*, 840 (1969). [337] V. S. Petrosyan and O. A. Reutov: *Effect of Solvent upon the Rates and Mechanisms of Organometallic Reactions, General Aspects*, J. Organomet. Chem. *52*, 307 (1973). [338] V. Gutmann: *Solvent Effects on the Reactivities of Organometallic Compounds*, Coord. Chem. Rev. *18*, 225 (1976). [339] *Cf.* also references [166–168] to Chapter 2. [340] J. Ugelstad, T. Ellingsen, and A. Berge, Acta Chem. Scand. *20*, 1593 (1966).

[341] P. Müller and B. Siegfried, Helv. Chim. Acta *55*, 2965 (1972). [342] U. Mayer, V. Gutmann, and A. Lodzinski, Monatsh. Chem. *104*, 1045 (1973). [343] C. J. Pedersen, J. Am. Chem. Soc. *89*, 2495, 7017 (1967); C. J. Pedersen and H. K. Frensdorff: *Makrocyclische Polyäther und ihre Komplexe*, Angew. Chem. *84*, 16 (1972); Angew. Chem. Int. Ed. Engl. *11*, 16 (1972). [344] B. Dietrich, J.-M. Lehn, and J.-P. Sauvage, Tetrahedron Lett. *1969*, 2885, 2889; Chemie in unserer Zeit *7*, 120 (1973); J.-M. Lehn: *Supramolekulare Chemie*, Angew. Chem. *100*, 92 (1988); Angew. Chem. Int. Ed. Engl. *27* (1988); J.-M. Lehn: *Cryptates: The Chemistry of Macropolycyclic Inclusion Complexes*, Acc. Chem. Res. *11*, 49 (1978). [345] F. Vögtle and E. Weber (eds.): *Host Guest Complex Chemistry – Macrocycles – Synthesis, Structures, Applications*, Springer, Berlin, 1985. [346] R. Izatt and J. J. Christensen (eds.): *Synthetic Multidentate Macrocyclic Compounds: Synthesis, Properties, and Uses*, Academic Press, New York, London, 1978. [347] G. W. Gokel and H. D. Durst: *Principles and Synthetic Applications in Crown Ether Chemistry*, Synthesis *1976*, 168. [348] F. Vögtle and E. Weber: *Progress in Crown Ether Chemistry*, Part IV A–IV E, Kontakte (Merck, Darmstadt) *1980* (2), 36ff.; *1981* (1), 24ff.; *1982* (1), 24ff.; *1983* (1), 38ff.; *1984* (1), 26ff. [349] E. Graf and J.-M. Lehn, J. Am. Chem. Soc. *97*, 5022 (1975); B. Metz, J. M. Rosalky, and R. Weiss, J. Chem. Soc., Chem. Commun. *1976*, 533. [350] H. E. Zaugg, B. W. Horrom, and S. Borgwardt, J. Am. Chem. Soc. *82*, 2895 (1960); H. E. Zaugg, ibid. *83*, 837 (1961).

[351] H. E. Zaugg, J. F. Ratajczak, J. E. Leonard, and A. D. Schaefer, J. Org. Chem. *37*, 2249 (1972). [352] H. D. Zook, T. J. Russo, E. F. Ferrand, and D. S. Stotz, J. Org. Chem. *33*, 2222 (1968). [353] H. D. Zook and T. J. Russo, J. Am. Chem. Soc. *82*, 1258 (1960); H. D. Zook and W. L. Gumby, ibid. *82*, 1386 (1960). [354] L. M. Thomasson, T. Ellingsen, and J. Ugelstad, Acta Chem. Scand. *25*, 3024 (1971). [355] C. L. Liotta and H. P. Harris, J. Am. Chem. Soc. *96*, 2250 (1974). [356] F. L. Cook, C. W. Bowers, and C. L. Liotta, J. Org. Chem. *39*, 3416 (1974). [357] D. J. Sam and H. E. Simmons, J. Am. Chem. Soc. *96*, 2252 (1974). [358] M. Cinquini, F. Montanori, and P. Tundo, J. Chem. Soc., Chem. Commun. *1975*, 393. [359] C. L. Liotta, H. P. Harris, M. McDermott, T. Gonzalez, and K. Smith, Tetrahedron Lett. *1974*, 2417. [360] H. D. Durst, Tetrahedron Lett. *1974*, 2421; H. D. Durst, M. Milano, E. J. Kikta, S. A. Connelly, and E. Grushka, Anal. Chem. *47*, 1797 (1975).

[361] J. W. Zubrick, B. I. Dunbar, and H. D. Durst, Tetrahedron Lett. *1975*, 71. [362] R. A. Bartsch: *Ionic Association in Base-Promoted β-Elimination Reactions*, Acc. Chem. Res. *8*, 239 (1975). [363] N. Kornblum, R. A. Smiley, R. K. Blackwood, and D. C. Iffland, J. Am. Chem. Soc. *77*, 6269 (1955). [364] R. Gompper: *Beziehungen zwischen Struktur und Reaktivität ambifunktioneller nucleophiler Verbindungen*, Angew. Chem. *76*, 412 (1964); Angew. Chem. Int. Ed. Engl. *3*,

560 (1964); Usp. Khim. *36*, 803 (1967). [365] W. J. Le Noble: *Conditions for the Alkylation of Ambident Anions*, Synthesis *1970*, 1. [366] S. A. Shevelev: *Dual Reactivity of Ambident Anions*, Usp. Khim. *39*, 1773 (1970); Russ. Chem. Rev. *39*, 844 (1970). [367] R. Gompper and H.-U. Wagner: *Das Allopolarisierungs-Prinzip. Substituenteneinflüsse auf Reaktionen ambifunktioneller Anionen*, Angew. Chem. *88*, 389 (1976); Angew. Chem. Int. Ed. Engl. *15*, 321 (1976). [367a] L. M. Jackman and B. C. Lange: *Structure and Reactivity of Alkali Metal Enolates*, Tetrahedron *33*, 2737 (1977). [367b] O. A. Reutov, I. P. Beletskaya, and A. L. Kurts: *Ambident Anions* (translated from Russian by J. P. Michael), Consultants Bureau, Plenum Publishing Corp., New York, 1983. [368] S. Hünig: *Die Reaktionsweise ambidenter Kationen*, Angew. Chem. *76*, 400 (1964); Angew. Chem. Int. Ed. Engl. *3*, 548 (1964); Usp. Khim. *36*, 693 (1967). [369] L. Claisen, Z. Angew. Chem. *36*, 478 (1923); L. Claisen, F. Kremers, F. Roth, and E. Tietze, Liebigs Ann. Chem. *442*, 210 (1925). [370] N. Kornblum, P. J. Berrigan, and W. J. Le Noble, J. Am. Chem. Soc. *82*, 1257 (1960); ibid. *85*, 1141 (1963).

[371] N. Kornblum, R. Seltzer, and P. Haberfield, J. Am. Chem. Soc. *85*, 1148 (1963). [372] A. L. Kurts, A. Macias, N. K. Genkina, I. P. Beletskaya, and O. A. Reutov, Dokl. Akad. Nauk SSSR *187*, 807 (1969); Chem. Abstr. *71*, 112364u (1969). [373] A. L. Kurts, P. I. Dem'yanov, A. Macias, I. P. Beletskaya, and O. A. Reutov, Tetrahedron Lett. *1968*, 3679; Tetrahedron *27*, 4759, 4769, 4777 (1971). [374] G. Brieger and W. M. Pelletier, Tetrahedron Lett. *1965*, 3555. [375] S. J. Rhoads and R. W. Holder, Tetrahedron *25*, 5443 (1969). [376] S. G. Smith and M. P. Hanson, J. Org. Chem. *36*, 1931 (1971). [377] A. L. Kurts, P. I. Dem'yanov, I. P. Beletskaya, and O. A. Reutov, Zh. Org. Khim. *9*, 1313 (1973); J. Org. Chem. USSR *9*, 1341 (1973). [378] W. J. Le Noble and S. K. Palit, Tetrahedron Lett. *1972*, 493. [379] G. Klopman, J. Am. Chem. Soc. *90*, 223 (1968); *Chemical Reactivity and Reaction Paths*, Wiley, New York, 1974. [380] A. Bertho, Liebigs Ann. Chem. *714*, 155 (1968).

[381] P. G. Duggan and W. S. Murphy, J. Chem. Soc., Perkin Trans. II, 1291 (1975). [382] O. A. Reutov: *Mechanism of Reactions of Electrophilic Substitution at a Saturated Carbon Atom*, Usp. Khim. *36*, 414 (1967); Russ. Chem. Rev. *36*, 163 (1967); Fortschr. Chem. Forsch. *8*, 61 (1967); J. Organomet. Chem. *100*, 219 (1975). [383] M. H. Abraham: *Constitutional Effects, Salt Effects, and Solvent Effects in Electrophilic Substitution at Saturated Carbon*, Compr. Chem. Kinet. *12*, 211 (1973); Chem. Abstr. *83*, 8562u (1975). [384] O. A. Reutov, V. I. Sokolov, I. P. Beletskaya, Yu. S. Ryabokobylko, and B. Prajsnar, Izvest. Akad. Nauk SSSR *1963*, 966, 970; Chem. Abstr. *59*, 5837 (1963). [385] E. D. Hughes, C. K. Ingold, and R. M. G. Roberts, J. Chem. Soc. *1964*, 3900. [386] G. Heublein, J. Prakt. Chem. *31*, 84 (1966). [387] J. H. Rolston and K. Yates, J. Am. Chem. Soc. *91*, 1477 (1969). [388] S. P. McManus and P. E. Peterson, Tetrahedron Lett. *1975*, 2753; S. P. McManus and S. D. Worley, ibid. *1977*, 555; and references cited therein. [389] K. Nakagawa, Y. Ishii, and M. Ogawa, Chem. Lett. (Tokyo) *1976*, 511; Tetrahedron *32*, 1427 (1976). [390] G. Mark, F. Mark, and O. E. Polansky, Liebigs Ann. Chem. *719*, 151 (1968).

[391] R. Steinmetz, W. Hartmann, and G. O. Schenck, Chem. Ber. *98*, 3854 (1965); I.-M. Hartmann, W. Hartmann, and G. O. Schenck, ibid. *100*, 3146 (1967). [392] F. D. Lewis and R. J. DeVoe, Tetrahedron *38*, 1069 (1982). [393] H. van Zwet and E. C. Kooyman, Recl. Trav. Chim. Pays-Bas *86*, 1143 (1967). [394] J. Brokken-Zijp and H. v. d. Bogaert, Tetrahedron *29*, 4169 (1973). [395] J. Sicher: *Der syn- und anti-koplanare Verlauf bimolekularer olefin-bildender Eliminierungen*, Angew. Chem. *84*, 177 (1972); Angew. Chem. Int. Ed. Engl. *11*, 200 (1972). [396] J. Avraamides and A. J. Parker, Tetrahedron Lett. *1971*, 4043. [397] H. Kessler: *Nachweis gehinderter Rotationen und Inversionen durch NMR-Spektroskopie*, Angew. Chem. *82*, 237 (1970); Angew. Chem. Int. Ed. Engl. *9*, 219 (1970). [398] U. Berns, G. Heinrich, and H. Güsten, Z. Naturforsch. *31b*, 953 (1976); and references cited therein. [399] *Cf.* also references [74–77, 80, 128, 131, 132, 220–224] in reference list to Chapter 2. [400] (a) C. A. Bunton and G. Savelli: *Organic Reactivity in Aqueous Micelles and Similar Assemblies*, Adv. Phys. Org. Chem. *22*, 213 (1986); (b) C. A. Bunton, F. Nome, F. H. Quina, and L. S. Romsted, Acc. Chem. Res. *24*, 357 (1991).

[401] M. Grätzel and K. Kalyanasundaram (eds.): *Kinetics and Catalysis in Microheterogeneous Systems*, Dekker, New York, 1991. [402] E. J. Fendler and J. H. Fendler: *Micellar Catalysis in Organic Reactions: Kinetic and Mechanistic Implications*, Adv. Phys. Org. Chem. *8*, 271 (1970).

[403] J. H. Fendler and E. J. Fendler: *Catalysis in Micellar and Macromolecular Systems*, Academic Press, New York, 1975; J. H. Fendler: *Membrane Mimetic Chemistry*, Wiley, New York, 1982. [404] H. Chaimovich, A. Blanco, L. Chayet, L. M. Costa, P. M. Monteiro, C. A. Bunton, and C. Paik, Tetrahedron *31*, 1139 (1975); C. A. Bunton, Pure Appl. Chem. *49*, 969 (1977). [405] C. A. Blyth and J. R. Knowles, J. Am. Chem. Soc. *93*, 3017 (1971). [406] D. Oakenfull, J. Chem. Soc., Perkin Trans. II *1973*, 1006. [407] J. H. Fendler, E. J. Fendler, and L. W. Simth, J. Chem. Soc., Perkin Trans. II *1972*, 2097. [408] R. M. Noyes: *Effects of Diffusion Rates on Chemical Kinetics*, in G. Porter und B. Stevens (eds.): *Progress in Chemical Kinetics*, Pergamon Press, Oxford, 1961, Vol. 1, p. 129ff. [409] A. M. North: *Diffusion-Controlled Reactions*, Quart. Rev. *20*, 421 (1966). [409a] S. A. Rice (ed.): *Diffusion-limited Reactions*, in C. H. Bamford, C. F. H. Tipper, and R. G. Compton (eds.): *Comprehensive Chemical Kinetics*, Vol. 25, Elsevier, Amsterdam, 1985. [410] K. U. Ingold: *Rate Constants for Free Radical Reactions in Solution*, in J. K. Kochi (ed.): *Free Radicals*, Wiley-Interscience, New York, 1973, Vol. 1, Chapter 2, p. 37ff.

[411] M. Eigen: *Protonenübertragung, Säure-Base-Katalyse und enzymatische Hydrolyse*, Angew. Chem. *75*, 489 (1963). [412] J. Frank and E. Rabinowitsch, Trans. Faraday Soc. *30*, 120 (1934); E. Rabinowitsch and W. C. Wood, ibid. *32*, 1381 (1936). [413] T. Koenig and H. Fischer: *Cage Effects*, in J. K. Kochi (ed.): *Free Radicals*, Wiley-Interscience, New York, 1973, Vol. 1, Chapter 4, p. 157ff. [414] M. J. Gibian and R. C. Corley: *Organic Radical-Radical Reactions, Disproportionation vs. Combination*, Chem. Rev. *73*, 441 (1973). [415] L. Herk, M. Feld, and M. Szwarc, J. Am. Chem. Soc. *83*, 2998 (1961); O. Dobis, J. M. Pearson, and M. Szwarc, ibid. *90*, 278 (1968). [416] R. K. Lyon and D. H. Levy, J. Am. Chem. Soc. *83*, 4290 (1961); R. K. Lyon, ibid. *86*, 1907 (1964). [417] R. E. Rebbert and P. Ausloos, J. Phys. Chem. *66*, 2253 (1962). [418] G. S. Hammond, J. N. Sen, and C. E. Boozer, J. Am. Chem. Soc. *77*, 3244 (1955). [419] G. Ayrey, Chem. Rev. *63*, 645 (1963). [420] J. W. Taylor and J. C. Martin, J. Am. Chem. Soc. *88*, 3650 (1966); ibid. *89*, 6904 (1967).

[421] C. Rüchardt and M. Grundmaier, Chem. Ber. *108*, 2448 (1975). [422] J. K. Kochi, J. Am. Chem. Soc. *92*, 4395 (1970). [423] N. Nodelman and J. C. Martin, J. Am. Chem. Soc. *98*, 6597 (1976). [424] R. Criegee: *Mechanismus der Ozonolyse*, Angew. Chem. *87*, 765 (1975); Angew. Chem. Int. Ed. Engl. *14*, 745 (1975). [425] L. D. Loan, R. W. Murray, and P. R. Story, J. Am. Chem. Soc. *87*, 737 (1965). [426] O. Lorenz and C. R. Parks, J. Org. Chem. *30*, 1976 (1965). [427] R. Criegee: *The Course of Ozonization of Unsaturated Compounds*, Record Chem. Progr. (Kresge-Hooker Sci. Lib.) *18*, 111 (1957); Chem. Abstr. *51*, 11982 (1957). [428] R. W. Murray: *The Mechanism of Ozonolysis*, Acc. Chem. Res. *1*, 313 (1968). [428a] V. Ramachandran and R. W. Murray, J. Am. Chem. Soc. *100*, 2197 (1978). [429] (a) W. J. le Noble, J. Chem. Educ. *44*, 729 (1967); (b) W. J. le Noble, Chem. Unserer Zeit *17*, 152 (1983); (c) W. J. le Noble (ed.): *Organic High Pressure Chemistry*, Elsevier, Amsterdam, 1988. [430] R. van Eldik, T. Asano, W. J. le Noble *et al.*, Chem. Rev. *78*, 407 (1978); ibid. *89*, 549 (1989); ibid. *98*, 2167 (1998).

[431] J. Jurczak and B. Baranovski (eds.): *High-Pressure Chemical Synthesis*, Elsevier, Amsterdam, 1989. [432] R. van Eldik and F.-G. Klärner: *High Pressure Chemistry*, Wiley-VCH, Weinheim, 2002. [433] G. Jenner, J. Phys. Org. Chem. *15*, 1 (2002). [434] M. G. Evans and M. Polanyi, Trans. Faraday Soc. *31*, 875 (1935); ibid. *32*, 1333 (1936). [435] R. C. Neuman: *Pressure Effects as Mechanistic Probes of Organic Radical Reactions*, Acc. Chem. Res. *5*, 381 (1972). [436] J. Buchanan and S. D. Hamann, Trans. Faraday Soc. *49*, 1425 (1953). [437] R. A. Grieger and C. A. Eckert, Trans. Faraday Soc. *66*, 2579 (1970). [438] R. J. Ouellette and S. H. Williams, J. Am. Chem. Soc. *93*, 466 (1971). [439] K. R. Brower, J. Am. Chem. Soc. *85*, 1401 (1963). [440] H. Hartmann, H. Kelm, and G. Rinck, Z. Phys. Chem. (Frankfurt) N. F. *44*, 335 (1965).

[441] H. Hartmann, H.-D. Brauer, H. Kelm, and G. Rinck, Z. Phys. Chem. (Frankfurt) N.F. *61*, 47, 53 (1968). [442] H.-D. Brauer and H. Kelm, Z. Phys. Chem. (Frankfurt) N.F. *76*, 98 (1971); ibid. *79*, 96 (1972). [443] Y. Kondo, M. Ohnishi, M. Uchida, and N. Tokura, Bull. Chem. Soc. Jpn. *41*, 992 (1968); ibid. *45*, 3579 (1972). [444] J. R. McCabe, R. A. Grieger, and C. A. Eckert, Ind. Eng. Chem., Fundam. *9*, 156 (1970); Chem. Abstr. *72*, 71368y (1970). [445] K. Tamura, Y. Ogo, and T. Imoto, Bull. Chem. Soc. Jpn. *46*, 2988 (1973); ibid. *48*, 369 (1975). [446] E. M. Arnett and D. R. McKelvey: *Solvent Isotope Effects on Thermodynamics of Nonreacting Solutes*, in J. F.

Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York, 1969, Vol. 1, p. 344ff. [447] P. M. Laughton and R. E. Robertson: *Solvent Isotope Effects for Equilibria and Reactions*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York 1969, Vol. 1, p. 399ff. [448] C. J. Collins and N. S. Bowman (eds.): *Isotope Effects in Chemical Reactions*, Van Nostrand Reinhold, New York, 1970, p. 246 and 266ff. [449] R. L. Schowen: *Mechanistic Deductions from Solvent Isotope Effects*, Progr. Phys. Org. Chem. *9*, 275 (1972). [450] C. G. Swain and R. F. W. Bader, Tetrahedron *10*, 182 (1960).

[451] D. R. Lide (ed.): *Handbook of Chemistry and Physics*, 82nd ed., CRC Press, Boca Raton/FL, USA, 2001. [452] E. F. Caldin: *Reaction Kinetics and Solvation in Non-Aqueous Solvents*, Pure Appl. Chem. *51*, 2067 (1979). [453] M. J. Blandamer and J. Burgess: *Initial State and Transition State Solvation in Inorganic Reactions*, Coord. Chem. Rev. *31*, 93 (1980); M. J. Blandamer, J. Burgess, and J. B. F. N. Engberts: *Activation Parameters for Chemical Reactions in Solution*, Chem. Soc. Rev. *14*, 237 (1985). [454] O. Popovych and R. P. T. Tompkins: *Nonaqueous Solution Chemistry*, Wiley, New York, 1981. [455] I. Bertini, L. Lunazzi, and A. Dei (eds.): *Advances in Solution Chemistry*, Plenum Press, New York, London, 1981. [456] C. Reichardt: *Der Lösungsmitteleinfluß auf chemische Reaktionen*, Chemie in unserer Zeit *15*, 139 (1981). [457] R. Schmid and V. N. Sapunov: *Non-Formal Kinetics*, Verlag Chemie, Weinheim, 1982. [458] K. Burger: *Solvation, Ionic and Complex Formation Reactions in Non-Aqueous Solvents*, Elsevier, Amsterdam, 1983. [459] R. A. Y. Jones: *Physical and Mechanistic Organic Chemistry*, 2nd edition, Cambridge University Press, Cambridge, 1984, Chapter 5, p. 94ff. [460] N. M. Emanuel, G. E. Zaikov, and Z. K. Maizus: *Oxidation of Organic Compounds, Medium Effects in Radical Reactions*, Pergamon Press, Oxford, 1984.

[461] J. Hayami: *Solvent Effects in the Organic Reactions*, Yuki Gosei Kagaku Kyokaishi *42*, 1107 (1984); Chem. Abstr. *102*, 148339 (1985). [462] A. L. Kurts: *Role of the Solvent in Organic Reactions*, Zh. Vses. Khim. Obshchestva im. D. I. Mendeleeva *29*, 530 (1984); Chem. Abstr. *102*, 5206f (1985). [463] (a) J. T. Hynes: *Chemical Reaction Dynamics in Solution*, Annu. Rev. Phys. Chem. 36, 573 (1985); (b) J. D. Simon: *Time-Resolved Studies of Solvation in Polar Media*, Acc. Chem. Res. *21*, 128 (1988); (c) B. Bagchi: *Dynamics of Solvation and Charge-Transfer Reactions in Dipolar Liquids*, Annu. Rev. Phys. Chem. *40*, 115 (1989). [464] K. J. Laidler: *A Lifetime of Transition-State Theory*, The Chemical Intelligencer *4*(3), 39 (1998). [465] (a) H. J. Kim and J. T. Hynes, J. Am. Chem. Soc. *114*, 10508, 10528 (1992); ibid. *115*, 8248 (1993); (b) J. R. Mathis and J. T. Hynes, J. Phys. Chem. *98*, 5445 (1994); (c) B. K. Carpenter, Angew. Chem. *110*, 3532 (1998); Angew. Chem. Int. Ed. Engl. *37*, 3340 (1998). [466] J. Bertrán and F. S. Burgos: *The Question of Equilibrium in Transition State Solvation*, J. Chem. Educ. *61*, 416 (1984). [467] E. Buncel and H. Wilson: *Initial-State and Transition-State Solvent Effects on Reaction Rates and the Use of Thermodynamic Transfer Functions*, Acc. Chem. Res. *12*, 42 (1979); *Solvent Effects on Rates and Equilibria*, J. Chem. Educ. *57*, 629 (1980). [468] C. Reichardt: *Solvent Effects on Chemical Reactivity*, Pure Appl. Chem. *54*, 1867 (1982). [469] R. T. McIver: *Chemical Reactions without Solvation*, Scientific American *243*, 148 (1980); Spektrum der Wissenschaft, Januar 1981, p. 27ff. [470] P. Ausloos (ed.): *Kinetics of Ion-Molecule Reactions*, (NATO Advanced Study Institute Series B, Vol. 40), Plenum Publishing Company, New York, 1979.

[471] M. T. Bowers (ed.): *Gas-Phase Ion Chemistry*, Vols. 1 and 2, Academic Press, New York, 1979. [472] J. H. Bowie: *Bimolecular Reactions of Nucleophiles in the Gas Phase*, Acc. Chem. Res. *13*, 76 (1980). [473] T. H. Morton: *Gas-Phase Analogues of Solvolysis Reactions*, Tetrahedron *38*, 3195 (1982). [474] W. N. Olmstead and J. I. Brauman: *Gas-Phase Nucleophilic Displacement Reactions*, J. Am. Chem. Soc. *99*, 4219 (1977); M. J. Pellerite and J. I. Brauman: *Intrinsic Barriers in Nucleophilic Displacements*, J. Am. Chem. Soc. *102*, 5993 (1980); J. I. Brauman *et al.*, Pure Appl. Chem. *56*, 1809 (1984); J. Phys. Chem. *90*, 471 (1986). [475] D. K. Bohme: *Gas-Phase Studies of the Influence of Solvation on Ion Reactivity*, in M. A. Almoster Ferreira (ed.): *Ionic Processes in the Gas Phase*, Reidel Publishing Company, Dordrecht, 1984, p. 111ff. [476] P. Kebarle, G. Caldwell, T. Magnera, and J. Sunner: *Ions-Gas Phase and Solution-Dipolar Aprotic Solvents*, Pure Appl. Chem. *57*, 339 (1985). [477] J. M. Riveros, S. M. José, and K. Takashima; *Gas-Phase Nucleophilic Displacement Reactions*, Adv. Phys. Org. Chem. *21*, 197 (1985). [478] E. M. Arnett: *Solvation Energies of Organic Ions*, J. Chem. Educ. *62*, 385 (1985). [479] H. A. Kramers: *Brownian Motion in a Field of Force and the Diffusion Model of Chemical Reactions*, Physica *7*, 284 (1940); Chem. Abstr. *34*, 4968

(1940). [480] K. Tanaka, G. I. Mackay, J. D. Payzant, and D. K. Bohme, Can. J. Chem. *54*, 1643 (1976).

[481] J. Chandrasekhar, S. F. Smith, and W. L. Jorgensen, J. Am. Chem. Soc. *106*, 3049 (1984); ibid. *107*, 154 (1985). [482] F. Carrion and M. J. S. Dewar, J. Am. Chem. Soc. *106*, 3531 (1984); M. J. S. Dewar and D. M. Storch, J. Chem. Soc., Chem. Commun. *1985*, 94. [483] S. S. Shaik, H. B. Schlegel, and S. Wolfe: *Theoretical Aspects of Physical Organic Chemistry – The S_N2 Mechanism*, Wiley, New York, 1992, Chapters 5 and 6. [484] D. K. Bohme and L. B. Young, J. Am. Chem. Soc. *92*, 7354 (1970). [485] D. K. Bohme and G. I. Mackay, J. Am. Chem. Soc. *103*, 978 (1981). [486] D. K. Bohme and A. B. Raksit, J. Am. Chem. Soc. *106*, 3447 (1984); Can. J. Chem. *63*, 3007 (1985). [487] M. Henchman, J. F. Paulson, and P. M. Hierl, J. Am. Chem. Soc. *105*, 5509 (1983); P. M. Hierl, A. F. Ahrens, M. Henchman, A. A. Viggiano, J. F. Paulson, and D. C. Clary, ibid. *108*, 3142 (1986). [488] G. Caldwell, M. D. Rozeboom, J. P. Kiplinger, and J. E. Bartmess, J. Am. Chem. Soc. *106*, 809 (1984). [489] K. Morokuma, J. Am. Chem. Soc. *104*, 3732 (1982). [490] D. N. Kevill and G. M. L. Liu, J. Am. Chem. Soc. *101*, 3916 (1979).

[491] F. Quemeneur and B. Bariou, J. Chem. Res. (S) *1979*, 187, 188; J. Chem. Res. (M) *1979*, 2344, 2357. [492] E. M. Arnett and R. Reich, J. Am. Chem. Soc. *102*, 5892 (1980). [493] D. N. Kevill, J. Chem. Soc., Chem. Commun. *1981*, 421. [494] M. H. Abraham and A. Nasehzadeh, J. Chem. Soc., Chem. Commun. *1981*, 905. [495] Y. Kondo, M. Ogasa, and S. Kusabayashi, J. Chem. Soc., Perkin Trans. II, 2093 (1984). [496] V. Bekárek, T. Nevěčná, and J. Vymětalová, Collect. Czech. Chem. Commun. *50*, 1928 (1985); ibid. *51*, 2071 (1986). [497] Y. Kondo, M. Ittoh, and S. Kusabayashi, J. Chem. Soc., Faraday Trans. I *78*, 2793 (1982). [498] K. C. Westaway, Can. J. Chem. *56*, 2691 (1978). [499] H. Maskill: *The Physical Basis of Organic Chemistry*, Oxford University Press, Oxford, New York, 1985, p. 409ff. [500] Y. Kondo, A. Zanka, and S. Kusabayashi, J. Chem. Soc., Perkin Trans. II, 827 (1985).

[501] C. F. Bernasconi in H. Zollinger (ed.): *MTP Rev. Sci., Org. Chem. Ser. One*, Vol. 3, p. 35ff., Butterworths, London, 1973; Acc. Chem. Res. *11*, 147 (1978). [502] L. Forlani, J. Chem. Res. (S) *1984*, 260; J. Chem. Res. (M) *1984*, 2379. [503] P. M. E. Mancini, R. D. Martinez, L. R. Vottero, and N. S. Nudelman, J. Chem. Soc., Perkin Trans. II, 1133 (1984), ibid. 1427 (1986), ibid. 951 (1987). [504] I. L. Bagal, S. A. Skvortsov, and A. V. El'tsov, Zh. Org. Khim. *14*, 361 (1978); J. Org. Chem. USSR *14*, 328 (1978). [505] Y. Hashida, F. Tanabe, and K. Matsui, Nippon Kagaku Kaishi *1980*, 865. [506] I. Szele and H. Zollinger: *Azo Coupling Reactions, Structures and Mechanisms*, Top. Curr. Chem. *112*, 1 (1983). [507] M. Gielen, Acc. Chem. Res. *6*, 198 (1973). [508] J. M. Fukuto and F. R. Jensen, Acc. Chem. Res. *16*, 177 (1983). [509] S. Fukuzumi and J. K. Kochi, J. Phys. Chem. *84*, 2254 (1980); J. Am. Chem. Soc. *102*, 2141 (1980). [510] V. S. Petrosyan, J. Organomet. Chem. *250*, 157 (1983); Advances in the USSR Organometallic Chemistry, Chapter 3, p. 73ff., MIR Publishers, Moscow, 1985.

[511] K. Ziegler *et al.*, Liebigs Ann. Chem. *473*, 1 (1929); ibid. *479*, 135 (1930). [512] G. Belluci, R. Bianchini, and R. Ambrosetti, J. Am. Chem. Soc. *107*, 2464 (1985). [513] M.-F. Ruasse and B.-L. Zhang, J. Org. Chem. *49*, 3207 (1984). [514] G. A. Jones, C. J. M. Stirling, and N. G. Bromby, J. Chem. Soc., Perkin Trans. II *1983*, 385. [515] W. A. Smit, N. S. Zefirov, I. V. Bodrikov, and M. Z. Krimer, Acc. Chem. Res. *12*, 282 (1979). [516] H. Mimoun, Angew. Chem. *94*, 750 (1982); Angew. Chem. Int. Ed. Engl. *21*, 734 (1982). [517] V. M. Vishnyakov, A. A. Bezdenezhnykh, Z. S. Zamchuk, and O. M. Kuznetsova, Zh. Org. Khim. *14*, 1238 (1978); J. Org. Chem. USSR *14*, 1146 (1978). [518] H. Graf and R. Huisgen, J. Org. Chem. *44*, 2594, 2595 (1979). [519] R. Huisgen and J. P. Ortega, Tetrahedron Lett. *1978*, 3975. [520] J. Drexler, R. Lindermeyer, M. A. Hassan, and J. Sauer, Tetrahedron Lett. *26*, 2559 (1985).

[521] H. K. Hale, L. C. Dunn, and A. B. Padias, J. Org. Chem. *45*, 835 (1980). [522] J. Mulzer and M. Zippel, Tetrahedron Lett. *21*, 751 (1980). [523] N. S. Isaacs and A. H. Laila, Tetrahedron Lett. *24*, 2897 (1983). [524] (a) R. Ocampo, W. R. Dolbier, M. D. Bartberger, and R. Paredes, J. Org. Chem. *62*, 109 (1997). (b) I. Morao, B. Lecea, A. Arrieta, and F. P. Cossio, J. Am. Chem. Soc. *119*, 816 (1997). [525] C. H. Heathcock; *Stereoselective Aldol Condensations*, in E. Buncel and T. Durst (eds.): *Comprehensive Carbanion Chemistry*, Part B, p. 177ff. (particularly p. 198), Elsevier, Amsterdam, 1984. [526] C. H. Heathcock and J. Lampe, J. Org. Chem. *48*, 4330 (1983). [527]

D.-M. Shin and D. G. Whitten, J. Am. Chem. Soc. *110*, 5206 (1988). [528] T. Asano and T. Okada, J. Org. Chem. *49*, 4387 (1984); ibid. *51*, 4454 (1986); Chem. Lett. (Tokyo) *1987*, 695. [529] N. Nishimura, T. Tanaka, M. Asano, and Y. Sueishi, J. Chem. Soc., Perkin Trans. II, 1839 (1986). [530] A. K. Dillow, J. S. Brown, C. L. Liotta, and C. A. Eckert, J. Phys. Chem. A *102*, 7609 (1998).

[531] G. Swieton and H. Kelm, J. Chem. Soc., Perkin Trans. II, 519 (1979). [532] A. I. Konovalov, I. P. Breus, I. A. Sharagin, and V. D. Kiselev, Zh. Org. Chem. *15*, 361 (1979); J. Org. Chem. USSR *15*, 315 (1979). [533] C. Rücker, D. Lang, J. Sauer, H. Friege, and R. Sustmann, Chem. Ber. *113*, 1663 (1980). [534] F. P. Ballistreri, E. Maccarone, G. Perrini, G. A. Tomaselli, and M. Torre, J. Chem. Soc., Perkin Trans. II, 273 (1982). [535] R. A. Firestone and S. G. Saffar, J. Org. Chem. *48*, 4783 (1983). [536] L. Pardo, V. Branchadell, A. Oliva, and J. Bertrán, J. Mol. Struct. *93*, 255 (1983). [537] G. Jenner, M. Papadopoulos, and J. Rimmelin, J. Org. Chem. *48*, 748 (1983). [538] M. Lotfi and R. M. G. Roberts, Tetrahedron *35*, 2137 (1979). [539] V. D. Kiselev, G. V. Mavrin, and A. I. Konovalov, Zh. Org. Khim. *16*, 1435 (1980); J. Org. Chem. USSR *16*, 1233 (1980). [540] O. B. Nagy, Can. J. Chem. *63*, 1382 (1985).

[541] R. Huisgen: *Cycloaddition Mechanism and the Solvent Dependence on Rate*, Pure Appl. Chem. *52*, 2283 (1980). [542] R. Huisgen: *1,3-Dipolar Cycloadditions – Introduction, Survey, Mechanism*, in A. Padwa (ed.): *1,3-Dipolar Cycloaddition Reactions*, Wiley-Interscience, New York, 1984, Vol. 1, p. 1ff. (particularly p. 76ff.). [543] M. S. Haque: *Concertedness of 1,3-Dipolar Cycloadditions*, J. Chem. Educ. *61*, 490 (1984). [544] R. Huisgen, H.-U. Reissig, and H. Huber, J. Am. Chem. Soc. *101*, 3647 (1979). [545] R. Huisgen, H.-U. Reissig, H. Huber, and S. Voss, Tetrahedron Lett. *1979*, 2987. [546] G. Bianchi, C. de Micheli, and R. Gandolfi: *1,3-Dipolar Cycloreversions*, Angew. Chem. *91*, 781 (1979); Angew. Chem. Int. Ed. Engl. *18*, 721 (1979). [547] R. Huisgen, R. Grashey, M. Seidel, G. Wallbillich, H. Knupfer, and R. Schmidt, Liebigs Ann. Chem. *653*, 105 (1962). [548] J.-M. Vandensavel, G. Smets, and G. L'abbé, J. Org. Chem. *38*, 675 (1973); G. L'abbé, Bull. Soc. Chim. Fr. *1975*, 1127. [549] J. P. Snyder and D. N. Horpp, J. Am. Chem. Soc. *98*, 7821 (1976); H. Olsen and J. P. Snyder, J. Am. Chem. Soc. *100*, 285 (1978). [550] L. M. Stephenson, M. J. Grdina, and M. Orfanopoulos, Acc. Chem. Res. *13*, 419 (1980).

[551] C. W. Jefford and S. Kohmoto, Helv. Chim. Acta *65*, 133 (1982). [552] A. Dondoni, A. Battaglia, F. Bernardi, and P. Giorgianni, J. Org. Chem. *45*, 3773 (1980). [553] M. Papadopoulos and G. Jenner, Nouv. J. Chim. *7*, 463 (1983). [554] H.-J. Bestmann, Pure Appl. Chem. *52*, 771 (1980); H.-J. Bestmann and O. Vostrowsky: *The Mechanism of the Wittig Reaction*, Top. Curr. Chem. *109*, 85 (1983). [555] N. S. Isaacs and O. H. Abed, Tetrahedron Lett. *27*, 995 (1986). [556] M. Schlosser and B. Schaub, J. Am. Chem. Soc. *104*, 5821 (1982); A. Piskala, A. H. Rehan, and M. Schlosser, Collect. Czech. Chem. Commun. *48*, 3539 (1983). [557] G. S. Bates and S. Ramaswamy, Can. J. Chem. *59*, 3120 (1981). [558] P. Schiess and H. Stalder, Tetrahedron Lett. *21*, 1413 (1980). [559] K. Harano and T. Taguchi, Chem. Pharm. Bull. (Tokyo) *23*, 467 (1975); Chem. Abstr. *83*, 96061h (1975). [560] R. Braun, F. Schuster, and J. Sauer, Tetrahedron Lett. *27*, 1285 (1986).

[561] B. Marcandalli, L. Pellicciari-Di Liddo, C. Di Fede, and I. R. Bellobono, J. Chem. Soc., Perkin Trans. II *1984*, 589. [562] M. G. Kulkarni, R. A. Mashelkar, and L. K. Doraiswamy, Chem. Eng. Sci. *35*, 823 (1980); Chem. Abstr. *93*, 94665r (1980). [563] C. Walling, R. R. W. Humphreys, J. P. Sloan, and T. Miller, J. Org. Chem. *46*, 5261 (1981). [564] R. Hiatt and P. M. Rahimi, Int. J. Chem. Kinet. *10*, 185 (1978); Chem. Abstr. *89*, 128858r (1978). [565] M. Schmittel, A. Schulz, C. Rüchardt, and E. Hädicke, Chem. Ber. *114*, 3533 (1981). [556] R. van Eldik, H. Kelm, M. Schmittel, and C. Rüchardt, J. Org. Chem. *50*, 2998 (1985). [567] R. C. Neuman and G. A. Binegar, J. Am. Chem. Soc. *105*, 134 (1983). [568] W. D. Hinsberg, P. G. Schultz, and P. B. Dervan, J. Am. Chem. Soc. *104*, 766 (1982). [569] W. Adam: *Oxygen Diradicals Derived from Cyclic Peroxides*, Acc. Chem. Res. *12*, 390 (1979). [570] E. M. Kosower: *Stable Pyridinyl Radicals*, Top. Curr. Chem. *112*, 117 (1983).

[571] J. P. Soumillion: *La Substitution Radicalaire des Composés Aliphatiques par le Chlore*, Ind. Chim. Belge *35*, 851, 1065 (1970). [572] N. J. Bunce, K. U. Ingold, J. P. Landers, J. Lusztyk, and J. C. Scaiano, J. Am. Chem. Soc. *107*, 5464 (1985). [573] A. Potter, J. M. Tedder, and J. C. Walton, J. Chem. Soc., Perkin Trans. II, 143 (1982). [574] A. Potter and J. M. Tedder, J. Chem. Soc., Perkin Trans. II, 1689 (1982). [575] S. Y. Atto, J. M. Tedder, and J. C. Walton, J. Chem. Soc., Perkin

Trans. II, 629 (1983). [576] O. Ito and M. Matsuda, J. Phys. Chem. *88*, 1002 (1984); J. Am. Chem. Soc. *104*, 568 (1982). [577] W. Offermann and F. Vögtle, Angew. Chem. *92*, 471 (1980); Angew. Chem. Int. Ed. Engl. *19*, 464 (1980). [578] O. Tapia: *Quantum Theories of Solvent-Effect Representation: An Overview of Methods and Results*, in H. Ratajczak and W. J. Orville-Thomas (eds.): *Molecular Interactions*, Vol. 3, Chapter 2, p. 47ff., Wiley, New York, 1982. [579] B. Ya. Simkin and I. I. Sheikhet; *Theoretical Methods of Calculating Solvation Effects*, J. Mol. Liquids *27*, 79 (1983). [580] B. M. Ladanyi and J. T. Hynes, J. Am. Chem. Soc. *108*, 585 (1986).

[581] L. M. Epshtein: *Hydrogen Bonds and the Reactivity of Organic Compounds in Proton Transfer and Nucleophilic Substitution Reactions*, Usp. Khim. *48*, 1600 (1979); Russ. Chem. Rev. *48*, 854 (1979). [582] K. Okamoto: *Solvent Molecules and Carbocation Intermediates in Solvolysis*, Pure Appl. Chem. *56*, 1797 (1984). [583] D. Landini, A. Maia, F. Montanari, and F. Rolla, J. Org. Chem. *48*, 3774 (1983). [584] M. Auriel and E. de Hoffmann, J. Chem. Soc., Perkin Trans. II, 325 (1979). [585] M. C. R. Symons, J. Chem. Soc., Chem. Commun. 418 (1978); M. J. Blandamer, J. Burgess, P. P. Duce, M. C. R. Symons, R. E. Robertson, and J. W. M. Scott, J. Chem. Res. (S) 130 (1982). [586] J. O. Edwards and R. G. Pearson, J. Am. Chem. Soc. *84*, 16 (1962). [587] C. H. Depuy, E. W. Della, J. Filley, J. J. Grabowski, and V. M. Bierbaum, J. Am. Chem. Soc. *105*, 2481 (1983). [588] S. Hoz and E. Buncel, Tetrahedron Lett. *25*, 3411 (1984); Israel J. Chem. *26*, 313 (1985). [589] D. S. Kemp and K. G. Paul, J. Am. Chem. Soc. *97*, 7305, 7312 (1975); D. S. Kemp, J. Reczek, and F. Vellaccio, Tetrahedron Lett. *1978*, 741. [590] M. Shirai and J. Smid, J. Am. Chem. Soc. *102*, 2863 (1980).

[591] J.-P. Kintzinger, J.-M. Lehn, E. Kauffmann, J. L. Dye, and A. I. Popov, J. Am. Chem. Soc. *105*, 7549 (1983). [592] J.-L. Pierre and P. Baret: *Complexes Moleculaires d'Anions*, Bull. Soc. Chim. Fr. *1983*, II-367. [593] T. Mitsuhashi, J. Am. Chem. Soc. *108*, 2400 (1986). [594] M. Mashima, R. R. McIver, R. W. Taft, F. G. Bordwell, and W. N. Olmstead, J. Am. Chem. Soc. *106*, 2717 (1984). [595] F. G. Bordwell and D. L. Hughes, J. Org. Chem. *45*, 3314 (1980). [596] T. F. Magnera, G. Caldwell, J. Sunner, S. Ikuta, and P. Kebarle, J. Am. Chem. Soc. *106*, 6140 (1984). [597] C. D. Ritchie, Can. J. Chem. *64*, 2239 (1986). [598] (a) H. Mayr and M. Patz, Angew. Chem. *106*, 990 (1994); Angew. Chem. Int. Ed. Engl. *33*, 938 (1994); (b) H. Mayr *et al.*, J. Am. Chem. Soc. *123*, 9500 (2001); (c) R. Lucius and H. Mayr, Angew. Chem. *112*, 2086 (2000); Angew. Chem. Int. Ed. Engl. *39*, 1995 (2000). [599] C. M. Sharts and W. A. Sheppard, Org. Reactions *21*, 125 (1974). [600] J. H. Clark, Chem. Rev. *80*, 429 (1980).

[601] R. O. Hutchins and I. M. Taffer, J. Org. Chem. *48*, 1360 (1983). [602] A. Lüttringhaus and H.-W. Dirksen; *Tetramethylharnstoff als Lösungsmittel und Reaktionspartner*, Angew. Chem. *75*, 1059 (1963). [603] B. J. Barker, J. Rosenfarb, and J. A. Caruso: *Harnstoffe als Lösungsmittel in der chemischen Forschung*, Angew. Chem. *91*, 560 (1979); Angew. Chem. Int. Ed. Engl. *18*, 503 (1979). [604] M. Bréant and G. Demange-Guerin: *Propriétés chimiques et électrochimiques dans le nitromethane*, Bull. Soc. Chim. Fr. *1975*, 163. [605] V. Gutmann and A. Scherhaufer: *Äthylensulfit als Lösungsmittel*, Monatsh. Chem. *99*, 1686 (1968). [606] R. S. Kittila: *A Review of Catalytic and Synthetic Applications for DMF and DMAC*, Bulletin DuPont de Nemours and Co., Wilmington/USA, 1960 (and Supplement 1962). [607] S. S. Pizey: *Dimethylformamide*, in S. S. Pizey (ed.): *Synthetic Reagents*, Wiley, New York, London, 1974, Vol. I, p. 1ff. [608] J. F. Coetzee (ed.): *Recommended Methods for Purification of Solvents and Tests for Impurities*, Pergamon Press, Oxford, 1982. [609] M. Bréant: *Propriétés chimiques et électrochimiques dans le N-méthylpyrrolidone*, Bull. Soc. Chim. Fr. *1971*, 725. [610] H. L. Huffman and P. G. Sears, J. Sol. Chem. *1*, 187 (1972).

[611] N. Saleh and J. A. Caruso, J. Sol. Chem. *8*, 197 (1979). [612] J. F. Coetzee: *Ionic Reactions in Acetonitrile*, Progr. Phys. Org. Chem. *4*, 45 (1967). [613] C. Agami: *Le Dimethylsulfoxyde en Chimie Organique*, Bull. Soc. Chim. Fr. *1965*, 1021. [614] D. Martin, A. Weise, and H.-J. Niclas: *Das Lösungsmittel Dimethylsulfoxid*, Angew. Chem. *79*, 340 (1967); Angew. Chem. Int. Ed. Engl. *6*, 318 (1967). [615] S. W. Jacob, E. E. Rosenbaum, and D. C. Wood (eds.): *Dimethyl Sulfoxide*, M. Dekker, New York, 1971, Vol. 1. [616] D. Martin and H. G. Hauthal: *Dimethylsulfoxid*, Akademie-Verlag, Berlin, 1971; *Dimethyl sulphoxide*, Van Nostrand-Reinhold, London, 1975. [617] T. Mukhopadhyay and D. Seebach: *Substitution of HMPT by the Cyclic Urea DMPU als a Cosolvent for Highly Reactive Nucleophiles and Bases*, Helv. Chim. Acta *65*, 385 (1982). [618] D.

Seebach, Chimia *39*, 147 (1985); Chemistry in Britain *21*, 632 (1985); *cf.* also Nachr. Chem. Techn. Labor *33*, 396 (1985). [619] R. Sowada: *Darstellung, Eigenschaften und Verwendung von Dimethylsulfon*, Z. Chem. *8*, 361 (1968). [620] N. Furukawa, F. Takahashi, T. Yoshimura, H. Morita, and S. Oae: *Enhanced Reactivities in Substitution and Elimination Reactions in Dimethyl Sulphoximide*, J. Chem. Soc., Perkin Trans. II, 432 (1981).

[621] H. G. Richey, R. D. Smith, B. A. King, T. C. Kester, and E. P. Squiller, J. Org. Chem. *46*, 2823 (1981); H. G. Richey and J. Farkas: *Sulfamides and Sulfonamides as Polar Aprotic Solvents*, J. Org. Chem. *52*, 479 (1987). [622] E. M. Arnett and C. F. Douty: *Sulfolane – A Weakly Basic Aprotonic Solvent of High Dielectric Constant*, J. Am. Chem. Soc. *86*, 409 (1964). [623] T. Yamamoto: *Sulfolane*, Yuki Gosei Kagaku Kyokai Shi *28*, 853 (1970); Chem. Abstr. *73*, 109587c (1970). [624] J. Martinma: *Sulfolane*, in J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvents*, Academic Press, New York, 1976, Vol. IV, p. 248ff. [625] W. H. Lee: *Cyclic Carbonates*, in J. J. Lagowski (ed.): *The Chemistry of Nonaqueous Solvents*, Academic Press, New York, 1976, Vol. IV, p. 167ff. [626] R. H. Wood and Q. D. Craft, J. Sol. Chem. *7*, 799 (1978). [627] Yu. I. Ranneva, I. S. Temnova, E. S. Petrov, A. I. Shatenshtein, E. N. Tsvetkov, and M. I. Kabachnik, Izv. Akad. Nauk SSSR, Ser. Khim. *1967*, 2129. [628] K. Weissermel, H.-J. Kleiner, M. Finke, and U.-H. Felcht, Angew. Chem. *93*, 256 (1981); Angew. Chem. Int. Ed. Engl. *20*, 223 (1981). [629] Jefferson Chemical Company, Inc.: *Propylene Carbonate*, Technical Bulletin, Houston/Texas, 1960. [630] J. Barthel and F. Feuerlein, J. Sol. Chem. *13*, 393 (1984).

[631] H. Normant: *Hexamethylphosphorsäuretriamid*, Angew. Chem. *79*, 1029 (1967); Angew. Chem. Int. Ed. Engl. *6*, 1046 (1967); Bull. Soc. Chim. Fr. *1968*, 791; Usp. Khim. *39*, 990 (1970); Russ. Chem. Rev. *39*, 457 (1970). [632] D. T. Sawyer, E. J. Nanni, and J. L. Roberts: *The Reaction Chemistry of Superoxide Ion in Aprotic Media*, in K. M. Kadish (ed.): Advances in Chemistry Series, No. 201, Electrochemical and Spectrochemical Studies of Biological Redox Components, American Chemical Society, Washington D.C., *1982*, p. 585ff. [633] A. F. Sowinski and G. M. Whitesides, J. Org. Chem. *44*, 2369 (1979). [634] Y. Marcus, Rev. Anal, Chem. *5*, 53 (1980); Pure Appl. Chem. *55*, 977 (1983); ibid. *57*, 1103 (1985). [635] (a) Y. Marcus: *Ion Solvation*, Wiley, Chichester, 1985; (b) Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998. [636] E. Buncel and E. A. Symons: *Initial State and Transition State Solvent Effects: Reactions in Protic and Dipolar Aprotic Media*, in I. Bertini, L. Lunazzi, and A. Dei (eds.): *Advances in Solution Chemistry*, Plenum Press, New York, 1981, p. 355ff. [637] M. H. Abraham: *Solvent Effects on Some Nucleophilic Substitutions*, in I. Bertini, L. Lunazzi, and A. Dei (eds.): *Advances in Solution Chemistry*, Plenum Press, New York, 1981, p. 341ff. [638] P. Haberfield and J. Pessin, J. Am. Chem. Soc. *105*, 526 (1983); P. Haberfield and D. Fortier, J. Org. Chem. *48*, 4554 (1983). [639] M. J. Blandamer *et al.*, J. Chem. Soc., Faraday Trans. I, 1471 (1986). [640] (a) I. M. Rozhkov and I. L. Knunyants, Dokl. Akad. Nauk SSR *199*, 614 (1971); Chem. Abstr. *76*, 7291 (1972); (b) K. O. Christe and W. W. Wilson, J. Fluorine Chem. *47*, 117 (1990).

[641] R. A. Bartsch, J. Org. Chem. *35*, 1023 (1970). [642] G. P. Schiemenz, J. Becker, and J. Stöckigt, Chem. Ber. *103*, 2077 (1970). [643] P. Viout: *Effects of Macrocyclic Cation Ligands and Quaternary Onium Salts on the Anionic Reactivity*, J. Mol. Catalysis *10*, 231 (1981). [644] D. J. Cram: *Cavitands: Organic Hosts with Enforced Cavities*, Science *219*, 1177 (1983); D. J. Cram: *Präorganisation – von Solventien zu Sphäranden*, Angew. Chem. *98*, 1041 (1986); Angew. Chem. Int. Ed. Engl. *25*, 1039 (1986). [645] F. De Jong and D. N. Reinhoudt: *Stability and Reactivity of Crown-Ether Complexes*, Adv. Phys. Org. Chem. *17*, 279 (1980). [646] F. Vögtle: *Supramolecular Chemistry – An Introduction*, Wiley, Chichester, 1993. [647] J.-M. Lehn et al. (eds.): *Comprehensive Supramolecular Chemistry*, Vols. 1–11, Pergamon-Elsevier, New York, 1996. [648] J.-M. Lehn: *Supramolecular Chemistry – Concepts and Perspectives*, VCH, Weinheim, 1995. [649] D. J. Cram, T. Kaneda, R. C. Helgeson, and G. M. Lein, J. Am. Chem. Soc. *101*, 6752 (1979). [650] B. G. Cox, J. Garcia-Rosas, and H. Schneier, J. Am. Chem. Soc. *103*, 1054 (1981); *cf.* also S. F. Lincoln, I. M. Brereton, and T. M. Spotswood, ibid. *108*, 8134 (1986).

[651] P. A. Mosier-Boss and A. I. Popov, J. Am. Chem. Soc. *107*, 6168 (1985); M. Shamsipur and A. I. Popov, J. Phys. Chem. *90*, 5997 (1986). [652] B. Dietrich and J. M. Lehn, Tetrahedron Lett. *1973*, 1225. [653] D. Landini, A. Maia, and F. Montanari, J. Am. Chem. Soc. *100*, 2796 (1978); J.

Chem. Soc., Perkin Trans. II, 46 (1980). [654] H. Handel and J. L. Pierre, Tetrahedron Lett. *1976*, 741. [655] J. L. Pierre, H. Handel, and R. Perraud, Tetrahedron Lett. *1977*, 2013. [656] G. W. Gokel and W. P. Weber, J. Chem. Educ. *55*, 350, 429 (1978). [657] C. M. Starks and C. Liotta: *Phase Transfer Catalysis: Principles and Techniques*, Academic Press, New York, 1978. [658] E. V. Dehmlow and S. S. Dehmlow: *Phase-Transfer Catalysis*, 3rd ed., VCH, Weinheim, 1993. [659] P. Sarthou, G. Bram, and F. Guibe, Can. J. Chem. *58*, 786 (1980). [660] S. Akabori and H. Tuji, Bull. Chem. Soc. Jpn. *51*, 1197 (1978).

[661] E. M. Arnett, S. G. Maroldo, G. W. Schriver, S. L. Schilling, and E. B. Troughton, J. Am. Chem. Soc. *107*, 2091 (1985). [662] C. Wesdemiotis and F. W. McLafferty, Tetrahedron *37*, 3111 (1981). [663] S. Hünig and G. Wehner, Chem. Ber. *113*, 302 (1980). [664] T. W. Bentley and P. v. R. Schleyer, J. Am. Chem. Soc. *98*, 7658 (1976); F. L. Schadt, T. W. Bentley, and P. v. R. Schleyer, ibid. *98*, 7667 (1976). [665] T. W. Bentley, C. T. Bowen, D. H. Morten, and P. v. R. Schleyer, J. Am. Chem. Soc. *103*, 5466 (1981); T. W. Bentley and G. E. Carter, ibid. *104*, 5741 (1982); T. W. Bentley, G. E. Carter, and H. C. Harris, J. Chem. Soc., Chem. Commun. 387 (1984). [666] D. Färcasiu, J. Jähme, and C. Rüchardt, J. Am. Chem. Soc. *107*, 5717 (1985). [667] A. D. Allen, V. M. Kanagasabapathy, and T. T. Tidwell, J. Am. Chem. Soc. *107*, 4513 (1985). [668] J. P. Richard and W. P. Jencks, J. Am. Chem. Soc. *106*, 1373, 1383, 1396 (1984). [669] G. A. Olah and G. K. S. Prakash, Chemistry in Britain *19*, 916 (1983). [670] A. P. Johnson and V. Vajs, J. Chem. Soc., Chem. Commun. 817 (1979).

[671] E. K. Fukuda and R. T. McIver, J. Am. Chem. Soc. *101*, 2498 (1979). [672] K. Takashima and J. M. Riveros, J. Am. Chem. Soc. *100*, 6128 (1978). [673] V. S. Petrosyan: *Solvent Effects on the Kinetics and Mechanism of Organomercury Reactions*, Advances in the USSR Organometallic Chemistry, MIR Publishers, Moscow, 1985, p. 73ff. [674] V. S. Petrosyan, J. Organomet. Chem. *250*, 157 (1983). [675] P. Kolsaker, H.-J. Storesund, T. Gulbrandsen, and G. Wøien, Acta Chem. Scand., Part B *37*, 187 (1983). [676] K. Müller and J. Sauer, Tetrahedron Lett. *25*, 2541 (1984). [677] W. Eberbach and J. C. Carré, Tetrahedron Lett. *21*, 1145 (1980). [678] N. Abe, T. Nishiwaki, and N. Komoto, Chem. Lett. (Tokyo) *1980*, 223. [679] R. Herges and I. Ugi, Angew. Chem. *97*, 596 (1985); Angew. Chem. Int. Ed. Engl. *24*, 594 (1985); Chem. Ber. *119*, 829 (1986). [680] S. Takamuku, T. Kuroda, and H. Sakurai, Chem. Lett. (Tokyo) *1982*, 377.

[681] A. A. Frimer: *The Reaction of Singlet Oxygen with Olefins: The Question of Mechanism*, Chem. Rev. *79*, 359 (1979). [682] K. Gollnick and G. O. Schenck: *Oxygen as Dienophile*, in J. Hamer (ed.): *1,4-Cycloaddition Reactions*, Academic Press, New York, London, 1967, Chapter 10, p. 255ff. [683] C. W. Jefford and S. Kohmoto, Helv. Chim. Acta *65*, 133 (1982). [684] C. W. Jefford, H. G. Grant, D. Jaggi, J. Boukouvalas, and S. Kohmoto, Helv. Chim. Acta *67*, 2210 (1984). [685] A. A. Frimer, P. D. Bartlett, A. F. Boschung, and J. G. Jewett, J. Am. Chem. Soc. *99*, 7977 (1977). [686] L. B. Harding and W. A. Goddard, J. Am. Chem. Soc. *102*, 439 (1980). [687] W. Adam and H. Rebollo, Israel J. Chem. *23*, 399 (1983). [688] M. L. Graziano, M. R. Iesce, S. Chiosi, and R. Scarpati, J. Chem. Soc., Perkin Trans. I, 2071 (1983). [689] E. L. Clennan and R. P. L'Esperance, J. Am. Chem. Soc. *107*, 5178 (1985); J. Org. Chem. *50*, 5424 (1985). [690] I. Szele and H. Zollinger, Helv. Chim. Acta *61*, 1721 (1978).

[691] H. Sawada, H. Hagii, K. Aoshima, M. Yoshida, and M. Kobayashi, Bull. Chem. Soc. Jpn. *58*, 3448 (1985). [692] Yu. I. Ovchinnikova, V. A. Fomin, A. I. D'yachkov, and V. S. Etlis, Zh. Obshch. Khim. *51*, 2355 (1981); J. Gen. Chem. USSR *51*, 2031 (1981). [693] R. A. Bartsch and J. Závada, Chem. Rev. *50*, 453 (1980). [694] M. Schlosser and T. D. An, Helv. Chim. Acta *62*, 1194 (1979). [695] S. Krishnamurthy, J. Org. Chem. *45*, 2550 (1980). [696] M. P. Hartshorn, R. S. Thompson, and J. Vaughan, Aust. J. Chem. *30*, 865 (1977); J. W. Blunt, M. P. Hartshorn, M. H. G. Munro, L. T. Soong, R. S. Thompson, and J. Vaughan, J. Chem. Soc., Chem. Commun. 820 (1980). [697] S. Kiyooka, F. Goto, and K. Suzuki, Chem. Lett. (Tokyo) *1981*, 1429. [698] H. O. House and T. V. Lee, J. Org. Chem. *43*, 4369 (1978). [699] J. B. Ousset, C. Mioskowski, and G. Solladié, Synth. Commun. *13*, 1193 (1984). [700] D. Enders and R. W. Hoffmann: *Asymmetrische Synthese*, Chemie in unserer Zeit *19*, 177 (1985).

[701] (a) E. L. Eliel, S. H. Wilen, and L. N. Mander: *Stereochemistry of Organic Compounds*, Wiley, New York, 1994, Chapter 12, p. 835ff.; (b) R. E. Gawley and J. Aubé: *Principles of Asym-*

metric Synthesis, Pergamon-Elsevier, Oxford/U.K., 1996. [702] R. Pérez-Ossorio, A. Pérez-Rubalcaba, M. L. Quiroga, and M. Lasperas, Tetrahedron Lett. *21*, 1565 (1980); O. Arjona, R. Pérez-Ossorio, A. Pérez-Rubalcaba, and M. L. Quiroga, J. Chem. Soc., Perkin Trans. II, 597 (1981). [703] J. Berlan, Y. Besace, D. Prat, and G. Pourcelot, J. Organomet. Chem. *264*, 399 (1984). [704] K. Soai, K. Komiya, Y. Shigematsu, H. Hasegawa, and A. Ookawa, J. Chem. Soc., Chem. Commun. 1282 (1982). [705] R. Schmierer, G. Grotemeier, G. Helmchen, and A. Selim, Angew. Chem. *93*, 209 (1981); Angew. Chem. Int. Ed. Engl. *20*, 207 (1981); G. Helmchen, A. Selim, D. Dorsch, and I. Taufer, Tetrahedron Lett. *24*, 3213 (1983). [706] R. E. Ireland, R. H. Mueller, and A. K. Willard, J. Am. Chem. Soc. *98*, 2868 (1976). [707] J. Sauer and J. Kredel, Tetrahedron Lett. *1966*, 6359. [708] T. Poll, G. Helmchen, and B. Bauer, Tetrahedron Lett. *25*, 2191 (1984). [709] K. Soai and A. Ookawa, J. Org. Chem. *51*, 4000 (1986). [710] A. R. Katritzky and B. Brycki, J. Am. Chem. Soc. *108*, 7295 (1986); and references cited therein.

[711] J. B. F. N. Engberts: *Organic Reactions in Highly Aqueous Binaries*, Pure Appl. Chem. *54*, 1797 (1982). [712] N. J. Turro, G. S. Cox, and M. A. Paczkowski: *Photochemistry in Micelles*, Top. Curr. Chem. *129*, 57 (1985). [713] (a) V. Ramamurthy, Tetrahedron *42*, 5753 (1986); (b) V. Ramamurthy (ed.): *Photochemistry in Organized and Constrained Media*, VCH, Weinheim, 1991. [714] (a) P. A. Grieco, Aldrichimica Acta *24*(3), 59 (1991); (b) P. A. Grieco: *Organic Syntheses in Water*, Blackie Academic and Professional, Hampshire/U.K., 1998. [715] S. Otto and J. B. F. N. Engberts, Pure Appl. Chem. *72*, 1365 (2000). [716] (a) A. Lubineau, J. Augé, and Y. Queneau, Synthesis 741 (1994); (b) A. Lubineau and J. Augé, Top. Curr. Chem. *206*, 1 (1999). [717] C.-J. Li and T.-H. Chan: *Organic Reactions in Aqueous Media*, Wiley, Chichester, 1997. [718] F. Fringuelli and A. Taticchi: *The Diels–Alder Reaction – Selected Practical Methods*, Wiley-VCH, Weinheim, 2002. [719] P. Laszlo and J. Lucchetti, Tetrahedron Lett. *25*, 2147 (1984); P. Laszlo, Acc. Chem. Res. *19*, 121 (1986). [720] Y. Nakamura, Y. Imakura, T. Kato, and Y. Morita, J. Chem. Soc., Chem. Commun. 887 (1977).

[721] H. Mayer, F. Schuster, and J. Sauer, Tetrahedron Lett. *27*, 1289 (1986). [722] K. A. Zachariasse, N. V. Phuc, and B. Kozankiewicz, J. Phys. Chem. *85*, 2676 (1981). [723] P. Plieninger and H. Baumgärtel, Ber. Bunsenges. Phys. Chem. *86*, 161 (1982). [724] (a) C. J. Drummond, F. Grieser, and T. W. Healy, Faraday Discuss. Chem. Soc. *81*, 95 (1986); (b) E. B. Tada, L. P. Novaki, and O. A. El Seoud, Langmuir *16*, 35 (2000); ibid. *17*, 652 (2001). [725] See references [17–22, 22a, 109, 110] to Chapter 3. [726] T. Svedberg: *Chemische Reaktionen in anisotropen Flüssigkeiten*, Kolloid Z. *18*, 54, 101 (1916); Chem. Zentralbl. *1916* I, 958; ibid. *1916* II, 211. [727] S. Ganapathy, R. G. Zimmermann, and R. G. Weiss, J. Org. Chem. *51*, 2529 (1986). [728] W. J. Leigh, D. T. Frendo, and P. J. Klawunn, Can. J. Chem. *63*, 2131 (1985). [729] J. P. Otruba and R. G. Weiss, J. Org. Chem. *48*, 3448 (1983). [730] D. A. Hrovat, J. H. Liu, N. J. Turro, and R. G. Weiss, J. Am. Chem. Soc. *106*, 7033 (1984); R. L. Treanor and R. G. Weiss, Tetrahedron *43*, 1371 (1987).

[731] V. Ramesh and R. G. Weiss, J. Org. Chem. *51*, 2535 (1986). [732] J. M. Nerbonne and R. G. Weiss, J. Am. Chem. Soc. *100*, 2571 (1978); ibid. *101*, 402 (1979). [733] T. Nakano and H. Hirata, Bull. Chem. Soc. Jpn. *55*, 947 (1982). [734] W. J. Leigh, Can. J. Chem. *63*, 2736 (1985). [735] M. Nakazaki, K. Yamamoto, and K. Fujiwara, Chem. Lett. (Tokyo) *1978*, 863. [736] A. Dondoni, A. Medici, S. Colonna, G. Gottarelli, and B. Samori, Mol. Cryst., Liq. Cryst. *55*, 47 (1979). [737] C. Eskenazi, J. F. Nicoud, and H. B. Kagan, J. Org. Chem. *44*, 995 (1979). [738] C. Cameron, P. P. S. Saluja, M. A. Floriano, and E. Whalley, J. Phys. Chem. *92*, 3417 (1988). [739] P. S. Bailey: *Ozonation in Organic Chemistry*, Vols. 1 and 2, Academic Press, New York, 1978 and 1982; *cf.* p. 89ff. in Vol. 1. [740] R. L. Kuczkowski: *Ozone and Carbonyl Oxides*, in A. Padwa (ed.): *1,3-Dipolar Cycloaddition Chemistry*, Wiley-Interscience, New York, 1984, Vol. 2, p. 197ff.; R. L. Kuczkowski: *Formation and Structure of Ozonides*, Acc. Chem. Res. *16*, 42 (1983).

[741] G. D. Fong and R. L. Kuczkowski, J. Am. Chem. Soc. *102*, 4763 (1980). [742] M. Miura, T. Fujisaka, M. Nojima, S. Kusabayashi, and K. J. McCullough, J. Org. Chem. *50*, 1504 (1985). [743] A. W. Castleman and R. G. Keesee: *Clusters – Bridging the Gas and Condensed Phases*, Acc. Chem. Res. *19*, 413 (1986). [744] A. Lubineau and Y. Queneau, Tetrahedron Lett. *26*, 2653 (1985). [745] H.-J. Schneider and N. K. Sangwan, J. Chem. Soc., Chem. Commun. 1787 (1986); Angew. Chem. *99*, 924 (1987); Angew. Chem. Int. Ed. Engl. *26*, 896 (1987). [746] A. Lubineau, J. Org.

Chem. *51*, 2144 (1986). [747] N. S. Isaacs, Tetrahedron *47*, 8463 (1991). [748] M. J. Blandamer, J. Burgess, R. E. Robertson, and J. M. W. Scott: *Dependence of Equilibrium and Rate Constants on Temperature and Pressure*, Chem. Rev. *82*, 259 (1982). [749] (a) K. Matsumoto, A. Sera, T. Uchida, Synthesis *1*, 999 (1985); (b) K. Matsumoto and R. M. Acheson (eds.): *Organic Synthesis at High Pressures*; (c) M. Ciobanu and K. Matsumoto, Liebigs Ann./Recueil 623 (1997). [750] R. van Eldik and C. D. Hubbard, Chem. Unserer Zeit *34*, 240, 306 (2000).

[751] B. Raistrick, R. H. Sapiro, and D. M. Newitt, J. Chem. Soc. *1939*, 1761; *cf.* also M. G. Gonikberg and L. F. Vereshchagin, Zh. Fiz. Khim. *23*, 1447 (1949); Chem. Abstr. *44*, 2832f (1950). [752] G. Swieton, J. v. Jouanne, H. Kelm, and R. Huisgen, J. Org. Chem. *48*, 1035 (1983). [753] Y. Yoshimura, J. Osugi, and M. Nakahara, Bull. Chem. Soc. Jpn. *56*, 680 (1983). [754] T. Asano, Pure Appl. Chem. *71*, 1691 (1999). [755] Y. Kondo, A. Zanka, and S. Kusabayashi, J. Chem. Soc., Perkin Trans. II *1985*, 827. [756] J. B. Hyne, H. S. Golinkin, and W. G. Laidlaw, J. Am. Chem. Soc. *88*, 2104 (1966). [757] W. Leitner: *Reactions in Supercritical Carbon Dioxide*, Top. Curr. Chem. *206*, 107 (1999). [758] S. L. Wells and J. M. DeSimone: *The CO_2 Technology*, Angew. Chem. *113*, 534 (2001); Angew. Chem. Int. Ed. Engl. *40*, 518 (2001). [759] M. E. Paulaitis and G. C. Alexander, Pure Appl. Chem. *59*, 61 (1987). [760] E. Buncel and C. C. Lee (eds.): *Secondary and Solvent Isotope Effects* (Vol. 7 of the series *Isotopes in Organic Chemistry*), Elsevier, Amsterdam, 1987.

[761] A. V. Willi: *Isotopieeffekte bei chemischen Reaktionen*, Thieme, Stuttgart, 1983. [762] R. A. More O'Ferrall, G. W. Koeppl, and A. J. Kresge, J. Am. Chem. Soc. *93*, 9 (1971). [763] C. A. Bunton and V. J. Shiner, J. Am. Chem. Soc. *83*, 42, 3207, 3214 (1961). [764] C. G. Mitton, M. Gressner, and R. L. Schowen, J. Am. Chem. Soc. *91*, 2045 (1963). [765] L. Verbit, T. R. Halbert, and R. B. Patterson, J. Org. Chem. *40*, 1649 (1975). [766] F. D. Saeva, P. E. Sharpe, and G. R. Olin, J. Am. Chem. Soc. *97*, 204 (1975). [767] R. M. Coates, B. D. Rogers, S. J. Hobbs, D. R. Peck, and D. P. Curran, J. Am. Chem. Soc. *109*, 1160 (1987). [768] H. Nöth, R. Rurländer, and P. Wolfgarth, Z. Naturforsch., Part B *35*, 31 (1981). [769] C. F. Bernasconi, R. D. Bunnell, D. A. Kliner, A. Mullin, P. Paschalis, and F. Terrier: *The Effect of Solvation on Intrinsic Rates of Proton-Transfer*, in M. Kobayashi (ed.): *Physical Organic Chemistry, 1986*, Elsevier, Amsterdam, 1987, p. 583ff.; C. F. Bernasconi, Acc. Chem. Res. *20*, 301 (1987). [770] W. J. Albery, Adv. Phys. Org. Chem. *28*, 139 (1993).

[771] D. G. Truhlar, B. C. Garrett, and S. J. Klippenstein: *Current Status of Transition-State Theory*, J. Phys. Chem. *100*, 12771 (1996). [772] C. Li, P. Ross, J. E. Szuleiko, and T. B. McMahon, J. Am. Chem. Soc. *118*, 9360 (1996). [773] T. Su, H. Wang, and W. L. Hase, J. Phys. Chem. A *102*, 9819 (1998). [774] H. Liu, F. Müller-Plathe, and W. F. van Gunsteren, Chem. Eur. J. *2*, 191 (1996). [775] B. Ensing, E. J. Meijer, P. E. Blöchl, and E. J. Baerends, J. Phys. Chem. A *105*, 3300 (2001). [776] M. N. Glukhovtsev, A. Pross, and L. Radom, J. Am. Chem. Soc. *118*, 6273, 11258 (1996). [777] A. A. Mohamed and F. Jensen, J. Phys. Chem. A *105*, 3259 (2001). [778] B. Safi, K. Choho, and P. Geerlings, J. Phys. Chem. A *105*, 591 (2001). [779] S. Parthiban, G. de Oliveira, and J. M. L. Martin, J. Phys. Chem. A *105*, 895 (2001). [780] N. M. M. Nibbering, Acc. Chem. Res. *23*, 279 (1990).

[781] J. Gao, Acc. Chem. Res. *29*, 298 (1996). [782] E. C. Ashby, Acc. Chem. Res. *21*, 414 (1988). [783] J.-M. Savéant, Adv. Phys. Org. Chem. *26*, 1 (1990). [784] A. R. Katritzky and B. Brycki, Chem. Soc. Rev. *19*, 83 (1990). [785] B. Speiser, Angew. Chem. *108*, 2623 (1996); Angew. Chem. Int. Ed. Engl. *35*, 2471 (1996). [786] J.-L. M. Abboud, R. Notario, J. Bertran, and M. Solá, Progr. Phys. Org. Chem. *19*, 1 (1993). [787] U. Maran, T. A. Pakkanen, and M. Karelson, J. Chem. Soc., Perkin Trans. II 2445 (1994). [788] S. S. Shaik, A. Ioffe, A. C. Reddy, and A. Pross, J. Am. Chem. Soc. *116*, 262 (1994). [789] C. Amovilli, B. Mennucci, and F. M. Floris, J. Phys. Chem. B *102*, 3023 (1998). [790] Yu. G. Shtyrlin, N. I. Tyryshkin, G. G. Iskhakova, V. V. Gavrilov, V. D. Kiselev, and A. I. Konovalov, Zh. Obshch. Khim. *65*, 1101 (1995); Russ. J. Gen. Chem. *65*, 1004 (1995).

[791] R. Makitra and Ya. N. Pirig, Zh. Obshch. Khim. *68*, 425 (1998); Russ. J. Gen. Chem. *68*, 395 (1998); [792] H. Mayr, M. Hartnagl, and K. Grimm, Liebigs Ann./Recueil 55 (1997). [793] S. I. Vdovenko, I. I. Gerus, and J. Wójcik, J. Phys. Org. Chem. *14*, 533 (2001). [794] F. P. Cossio, G. Roa, B. Lecea, and J. M. Ugalde, J. Am. Chem. Soc. *117*, 12306 (1995). [795] D. K. Jones-Hertzog and W. L. Jorgensen, J. Am. Chem. Soc. *117*, 9077 (1995). [796] (a) E. S. Lewis and C. E. Boozer,

J. Am. Chem. Soc. *74*, 308 (1952); (b) P. R. Schreiner, P. v. R. Schleyer, and R. K. Hill, J. Org. Chem. *58*, 2822 (1993). [797] W. L. Jorgensen, J. F. Blake, D. Lim, and D. L. Severance, J. Chem. Soc., Faraday Trans. *90*, 1727 (1994). [798] R. Sustmann and W. Sicking, J. Am. Chem. Soc. *118*, 12562 (1996). [799] C. Cativiela, J. I. Garcia, J. A. Mayoral, and L. Salvatella, Chem. Soc. Rev. *25*, 209 (1996). [800] R. Sustmann *et al.*, Chem. Ber. *124*, 2811 (1991).

[801] (a) D. C. Rideout and R. Breslow, J. Am. Chem. Soc. *102*, 7816 (1980); (b) R. Breslow, Acc. Chem. Res. *24*, 159 (1991); (c) R. Breslow, Green Chem. 225 (1998). [802] A. Kumar, Chem. Rev. *101*, 1 (2001). [803] R. Braun and J. Sauer, Chem. Ber. *119*, 1269 (1986). [804] (a) P. A. Grieco, J. J. Nunes, and M. Gaul, J. Am. Chem. Soc. *112*, 4595 (1990); (b) P. A. Grieco, Aldrichimica Acta *24*(3), 59 (1991); (c) P. A. Grieco, in M. V. Kisakürek (ed.): *Organic Chemistry: Its Language and Its State of Art*, HVCA, Basel, and VCH, Weinheim, 1993, p. 133ff. [805] (a) H. Waldmann, Angew. Chem. *103*, 1335 (1991); Angew. Chem. Int. Ed. Engl. *30*, 1306 (1991); (b) A. Flohr and H. Waldmann, J. Prakt. Chem. *337*, 609 (1995). [806] S. Sankararaman and J. E. Nesakumar, Eur. J. Org. Chem. 2003 (2000). [807] (a) M. E. Jung and J. Gervay, J. Am. Chem. Soc. *111*, 5469 (1989); see also Chemtracts-Organic Chemistry *3*, 284 (1990); (b) M. E. Jung, Synlett 186 (1990). [808] R. Sustmann *et al.*, Tetrahedron Lett. *33*, 8027 (1992); Tetrahedron *48*, 10293 (1992). [809] R. Huisgen, G. Mloston, and E. Langhals, J. Am. Chem. Soc. *108*, 6401 (1986); J. Org. Chem. *51*, 4085 (1986); Tetrahedron Lett. *30*, 5373 (1989). [810] G. Mloston and H. Heimgartner, Polish J. Chem. *74*, 1503 (2000).

[811] Y.-M. Chang, J. Sims, and K. N. Houk, Tetrahedron Lett. 4445 (1975). [812] (a) R. Huisgen, L. Fisera, H. Giera, and R. Sustmann, J. Am. Chem. Soc. *117*, 9671 (1995); (b) R. Huisgen *et al.*, Pure Appl. Chem. *68*, 789 (1996). [813] B. Lecea, A. Arrieta, G. Roa, J. M. Ugalde, and F. P. Cossio, J. Am. Chem. Soc. *116*, 9613 (1994). [814] (a) B. Ganem, Angew. Chem. *108*, 1014 (1996); Angew. Chem. Int. Ed. Engl. *35*, 936 (1996). (b) J. J. Gajewski, Acc. Chem. Res. *30*, 219 (1997). [815] H. Quast and M. Seefelder, Angew. Chem. *111*, 1132, 1136 (1999); Angew. Chem. Int. Ed. *38*, 1064, 1068 (1999). [816] E. Brandes, P. A. Grieco, and J. J. Gajewski, J. Org. Chem. *54*, 515 (1989). [817] G. Desimoni, G. Faita, S. G. Comini, P. P. Righetti, and G. Tacconi, Tetrahedron *49*, 2093 (1993). [818] F. L. Ansari, R. Qureshi, and M. L. Qureshi: *Electrocyclic Reactions – From Fundamentals to Research*, Wiley-VCH, Weinheim, 1998. [819] G. Desimoni, G. Faita, S. Guidetti, and P. P. Righetti, Eur. J. Org. Chem. 1921 (1999). [820] S. Celebi, M.-L. Tsao, and M. S. Platz, J. Phys. Chem. A *105*, 1158 (2001).

[821] P. Maslak, W. H. Chapman, T. M. Vallombroso, and B. A. Watson, J. Am. Chem. Soc. *117*, 12380 (1995). [822] D. B. Sclove, J. F. Pazos, R. L. Camp, and F. D. Greene, J. Am. Chem. Soc. *92*, 7488 (1970). [823] D. Lim, D. A. Hrovat, W. T. Borden, and W. L. Jorgensen, J. Am. Chem. Soc. *116*, 3494 (1994). [824] L. F. R. Cafferata, G. N. Eyler, E. L. Svartman, A. I. Cañizo, and E. Alvarez, J. Org. Chem. *56*, 411 (1991). [825] K. U. Ingold, J. Lusztyk, and K. D. Raner, Acc. Chem. Res. *23*, 219 (1990). [826] J. M. Tanko and N. K. Suleman, J. Am. Chem. Soc. *116*, 5162 (1994). [827] D. V. Avila, C. E. Brown, K. U. Ingold, and J. Lusztyk, J. Am. Chem. Soc. *115*, 466 (1993). [828] M. J. Blandamer, J. Burgess, J. B. F. N. Engberts, and W. Blokzijl, Annu. Rep. Progr. Chem. C 45 (1990). [829] (a) A. Loupy, B. Tchoubar, and D. Astruc, Chem. Rev. *92*, 1141 (1992); (b) A. Loupy and B. Tchoubar: *Salt Effects in Organic and Organometallic Chemistry*, VCH, Weinheim, 1992. [830] T. Mitsuhashi, H. Hirota, and G. Yamamoto, Bull. Chem. Soc. Jpn. *67*, 824, 831 (1994).

[831] D. Landini, A. Maia, and A. Rampoldi, J. Org. Chem. *54*, 328 (1989). [832] R. M. Takka and E. Buncel, J. Am. Chem. Soc. *117*, 1503 (1995). [833] H. Zipse, G. Apaydin, and K. N. Houk, J. Am. Chem. Soc. *117*, 8608 (1995). [834] J. W. Grate, R. A. McGill, and D. Hilvert, J. Am. Chem. Soc. *115*, 8577 (1993). [835] J. Catalán, C. Diaz, and F. Garcia-Blanco, J. Org. Chem. *65*, 3409 (2000). [836] G. M. Blackburn, A. Datta, H. Denham, and P. Wentworth, Adv. Phys. Org. Chem. *31*, 249 (1998); part. p. 273. [837] (a) A. Bianchi: *The Supramolecular Chemistry of Anions*, Wiley, Chichester, 1997; (b) F. P. Schmidtchen and M. Berger, Chem. Rev. *97*, 1609 (1997); (c) P. D. Beer and P. A. Gale, Angew. Chem. *113*, 503 (2001); Angew. Chem. Int. Ed. *40*, 486 (2001). [838] R. Singh and G. M. Whitesides, J. Am. Chem. Soc. *112*, 1190 (1990). [839] (a) G. T. Hefter, Pure Appl. Chem. *63*, 1749 (1991); (b) K. Seppelt, Angew. Chem. *104*, 299 (1992); Angew. Chem. Int.

Ed. Engl. *31*, 292 (1992). [840] X. Zhang, V. Groß, and K. Seppelt, Angew. Chem. *107*, 2019 (1995); Angew. Chem. Int. Ed. Engl. *34*, 1858 (1995).

[841] G. Aksnes and P. Stensland, Acta Chem. Scand. *43*, 893 (1989). [842] A. Maia, Pure Appl. Chem. *67*, 697 (1995). [843] A. Ostrowicki, E. Koepp, and F. Vögtle, Top. Curr. Chem. *161*, 37 (1991). [844] T. Flessner and S. Doye, J. Prakt. Chem. *341*, 186 (1999). [845] R. Hoss and F. Vögtle, Angew. Chem. *106*, 389 (1994); Angew. Chem. Int. Ed. Engl. *33*, 375 (1994). [846] P. W. Schultz, G. E. Leroi, and A. I. Popov, J. Am. Chem. Soc. *117*, 10735 (1995); ibid. *118*, 10617 (1996). [847] M. C. R. L. R. Lazana, M. L. T. M. B. Franco, and B. J. Herold, J. Chem. Soc., Perkin Trans. II 1399 (1987). [848] M. E. Jones, S. R. Kass, J. Filley, R. M. Barkley, and G. B. Ellison, J. Am. Chem. Soc. *107*, 109 (1985). [849] M. Zhong and J. I. Brauman, J. Am. Chem. Soc. *118*, 636 (1996). [850] Y. Okuno, J. Phys. Chem. A *103*, 190 (1999).

[851] H. Mestdagh and A. Puechberty, J. Chem. Educ. *68*, 515 (1991). [852] B. Weber and D. Seebach, Angew. Chem. *104*, 96 (1992); Angew. Chem. Int. Ed. Engl. *31*, 84 (1992). [853] G. Cainelli, P. Galletti, D. Giacomini, and P. Orioli, Angew. Chem. *112*, 533 (2000); Angew. Chem. Int. Ed. Engl. *39*, 523 (2000); J. Chem. Soc., Chem. Commun. 567 (1999). [854] F. M. Bickelhaupt, E. J. Baerends, and N. M. M. Nibbering, Chem. Eur. J. *2*, 196 (1996). [855] J. Crugeiras, J. R. Leis, and A. Rios, J. Chem. Educ. *78*, 1538 (2001). [856] F. M. Menger and Z. X. Fei, Angew. Chem. *106*, 329 (1994); Angew. Chem. Int. Ed. Engl. *33*, 346 (1994). [857] G. Jenner, J. Phys. Org. Chem. *12*, 619 (1999). [858] M. R. Gholami and A. H. Yangjeh, J. Chem. Res. (S) 226 (1999). [859] F. M. Menger, Angew. Chem. *103*, 1104 (1991); Angew. Chem. Int. Ed. Engl. *30*, 1086 (1991). [860] R. Schomäcker, Nachr. Chem. Tech. Lab. *40*, 1344 (1992).

[861] D. O. Shah (ed.): *Micelles, Microemulsions and Monolayers – Science and Technology*, Dekker, New York, 1998. [862] J. Klier, C. J. Tucker, T. H. Kalantar, and D. P. Green, Adv. Mater. *12*, 1751 (2000). [863] J. H. Schulman, W. Stoeckenius, and L. M. Prince, J. Phys. Chem. *63*, 1677 (1959). [864] F. M. Menger, J. A. Donohue, and R. F. Williams, J. Am. Chem. Soc. *95*, 286 (1973). [865] F. M. Menger and H. Park, Recl. Trav. Chim. Pays-Bas *113*, 176 (1994). [866] J. Scheffer (ed.), Tetrahedron *43*, 1197–1745 (1987). [867] R. G. Weiss, Tetrahedron *44*, 3413 (1988). [868] F. L. Dickert and K. D. Fritsch, J. Phys. Chem. *94*, 7722 (1990). [869] P. W. Atkins: *Physical Chemistry*, 6th ed., Oxford University Press, Oxford/U.K., 1998, Section 27.2, p. 825ff. [870] D. J. McAdoo and T. H. Morton, Acc. Chem. Res. *26*, 295 (1993).

[871] N. J. Turro, Pure Appl. Chem. *67*, 199 (1995). [872] R. Schmidt, J. Phys. Chem. A *102*, 9082 (1998); [873] (a) W. J. le Noble and T. Asano, J. Phys. Chem. A *105*, 3428 (2001); (b) R. A. Firestone and K. A. Swiss, J. Phys. Chem. A *105*, 3430 (2001). [874] V. Kiselev *et al.*, J. Phys. Org. Chem. *14*, 636 (2001). [875] Y. Marcus and G. T. Hefter, J. Mol. Liquids *73/74*, 61 (1997). [876] (a) P. Tundo and P. T. Anastas: *Green Chemistry: Challenging Perspectives*, Oxford University Press, Oxford/U.K., 2000; (b) P. Tundo *et al.*, Pure Appl. Chem. *72*, 1207 (2000). [877] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto; William Andrews Publishing, New York, 2001, Chapter 21, p. 1419ff. [878] J. Jarrouse, C. R. Acad. Sci., Ser. C. *232*, 1424 (1951); Chem. Abstr. *46*, 3973 (1952). [879] (a) M. Makosza and W. Wawrzyniewicz, Tetrahedron Lett. 4659 (1969); (b) M. Makosza and M. Fedorynski, Pol. J. Chem. *70*, 1093 (1996); Chem. Abstr. *125*, 300159p (1996); (c) M. Makosza, Pure Appl. Chem. *72*, 1399 (2000). [880] (a) C. M. Starks, J. Am. Chem. Soc. *93*, 195 (1971); (b) C. M. Starks, C. L. Liotta, and M. Halpern: *Phase-Transfer Catalysis, Fundamentals, Applications, and Industrial Perspectives*, Chapman and Hall, New York, 1994.

[881] Y. Sasson and R. Neumann (eds.): *Handbook of Phase-Transfer Catalysis*, Blackie Academic, London, 1997. [882] M.-L. Wang: *Effects of Organic Solvents on Phase-Transfer Catalysis*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Data Base on CD-ROM), ChemTec Publishing, Toronto; William Andrews Publishing, New York, 2001, Chapter 13.3, p. 798ff. [883] A. Nelson, Angew. Chem. *111*, 1685 (1999); Angew. Chem. Int. Ed. Engl. *38*, 1583 (1999). [884] M. Vogt, Ph.D. Thesis, RWTH Aachen/Germany, 1991 (superadvisor: W. Keim). [885] (a) I. T. Horváth and J. Rábai, Science *266*, 72 (1994); (b) I. T. Horváth, Acc. Chem. Res. *31*, 641 (1998); (c) I. T. Horváth *et al.*, J. Am. Chem. Soc. *120*, 3133 (1998). [886] D.-W. Zhu, Synthesis 953 (1993). [887] B. Cornils, Angew. Chem. *109*, 2147 (1997); Angew. Chem. Int. Ed. Engl. *36*, 2057 (1997). [888] D. P.

Curran, Angew. Chem. *110*, 1230 (1998); Angew. Chem. Int. Ed. Engl. *37*, 1174 (1998). [889] E. de Wolf, G. van Koten, and B.-J. Deelman, Chem. Soc. Rev. *28*, 37 (1999). [890] L. P. Barthel-Rosa and J. A. Gladysz, Coord. Chem. Rev. *190–192*, 587 (1999).

[891] B. Betzemeier and P. Knochel, Top. Curr. Chem. *206*, 61 (1999). [892] R. H. Fish, Chem. Eur. J. *5*, 1677 (1999). [893] A. Endres and G. Maas, Chem. Unserer Zeit *34*, 382 (2000). [894] I. Klement, H. Lütjens, and P. Knochel, Angew. Chem. *109*, 1606 (1997); Angew. Chem. Int. Ed. Engl. *36*, 1454 (1997). [895] B. Betzemeier, M. Cavazzini, S. Quici, and P. Knochel, Tetrahedron Lett. *41*, 4343 (2000). [896] B. Cornils, Top. Curr. Chem. *206*, 133 (1999). [897] M. J. Earle and K. R. Seddon, Pure Appl. Chem. *72*, 1391 (2000). [898] M. Freemantle, Chem. Eng. News (May 15), 37 (2000). [899] M. J. Earle, P. B. McCormac, and K. R. Seddon, J. Chem. Soc., Chem. Commun. 2245 (1998). [900] M. J. Earle, P. B. MacCormac, and K. R. Seddon, Green Chem. *1*, 23 (1999).

[901] T. Fischer, A. Sethi, T. Welton, and J. Woolf, Tetrahedron Lett. *40*, 793 (1999). [902] J. A. Boon, J. A. Lewisky, J. L. Pflug, and J. S. Wilkes, J. Org. Chem. *51*, 480 (1986). [903] C. J. Adams, M. J. Earle, G. Roberts, and K. R. Seddon, J. Chem. Soc., Chem. Commun. 2097 (1998). [904] V. P. W. Böhm and W. A. Herrmann, Chem. Eur. J. *6*, 1017 (2000). [905] Y. Chauvin, L. Mußmann, and H. Olivier, Angew. Chem. *107*, 2941 (1995); Angew. Chem. Int. Ed. Engl. *34*, 2941 (1995). [906] Y. Chauvin, H. Olivier, C. N. Wyrvalski, L. C. Simon, and R. F. De Souza, J. Catal. *165*, 275 (1997); Chem. Abstr. *126*, 118251f (1997). [907] B. Ellis, W. Keim, and P. Wasserscheid, J. Chem. Soc., Chem. Commun. 337 (1999). [908] J. F. Kauffman, J. Phys. Chem. A *105*, 3433 (2001). [909] (a) P. G. Jessop and W. Leitner (eds.): *Chemical Synthesis Using Supercritical Fluids*, Wiley-VCH, Weinheim, 1999; (b) W. Leitner, Acc. Chem. Res. *35*, 746 (2002). [910] U. E. O'Shea, K. M. Kirmse, M. A. Fox, and K. P. Johnston, J. Phys. Chem. *95*, 7863 (1991).

[911] K. P. Johnston and C. Haynes, AIChe J. *33*, 2017 (1987). [912] O. Kajimoto, Chem. Rev. *99*, 355 (1999). [913] J. F. Brennecke and J. E. Chateauneuf, Chem. Rev. *99*, 433 (1999). [914] F. Patat, Monatsh. Chem. *77*, 352 (1947). [915] C. M. Wai, F. Hunt, M. Ji, and X. Chen, J. Chem. Educ. *75*, 1614 (1998). [916] M. Buback, Pure Appl. Chem. *68*, 1501 (1996). [917] M. G. Hitzler and M. Poliakoff, J. Chem. Soc., Chem. Commun. 1667 (1997). [918] P. G. Jessop, T. Ikariya, and R. Noyori, Nature *368*, 231 (1994); Chem. Rev. *95*, 259 (1995). [919] M. T. Reetz, W. Könen, and T. Strack, Chimia *47*, 493 (1993). [920] J. M. Tanko and J. F. Blackert, Science *263*, 203 (1994).

[921] J. A. Hyatt, J. Org. Chem. *49*, 5097 (1984). [922] J. W. Tester *et al.*, J. Phys. Chem. *100*, 12337 (1996). [923] E. M. Glebov, L. G. Krishtopa, V. Stepanov, and L. N. Krasnoperov, J. Phys. Chem. A *105*, 9427 (2001). [924] Y. Ikushima *et al.*, Chem. Lett. (Tokyo) 109 (1993). [925] R. W. Shaw, T. B. Brill, A. A. Clifford, C. A. Eckert, and E. U. Franck, Chem. Eng. News (Dec. 23) 26 (1991).

Chapter 6

[1] A. E. Lutskii, V. V. Prezhdo, L. I. Degtereva, and V. G. Gordienko: *Spectroscopy of Intermolecular Field Interactions in Solutions*, Usp. Khim. *51*, 1398 (1982); Russ. Chem. Rev. *51*, 802 (1982). [2] T. E. Gough, D. E. Irish, and I. R. Lantzke: *Spectroscopic Measurements (electron absorption, infrared and Raman, ESR and NMR spectroscopy)*, in A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, Chapter 4, p. 405ff. [3] M. Jauquet and P. Laszlo: *Influence of Solvents on Spectroscopy*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. VIII, Part I, of A. Weissberger (ed.): *Techniques of Chemistry*, Wiley-Interscience, New York, 1975, p. 195ff. [4] C. N. R. Rao, S. Singh, and V. P. Senthilnathan: *Spectroscopic Studies of Solute-Solvent Interactions*, Chem. Soc. Rev. *5*, 297 (1976). [5] K. Dimroth: *Über den Einfluß von Lösungsmitteln auf die Farbe organischer Verbindungen*, Sitzungsberichte der Gesellschaft zur Beförderung der gesamten Naturwissenschaften zu Marburg *76*, Heft 3, p. 3ff. (1953); Chem. Zentralbl. *1954*, 9481; Chimia *15*, 80 (1961). [6] A. I. Kiprianov: *Effect of the Solvent on the Color of Dyes (Solvatochromism)*. Usp. Khim. *29*, 1336 (1960); Russ. Chem.

Rev. *29*, 618 (1960). [7] N. G. Bakhshiev (ed.): *Sol'vatokhromiya: Problemy i Metody (Solvatochromism: Problems and Methods)*, Izdatel'stvo Leningradskovo Universiteta, Leningrad/USSR, 1989; Chem. Abstr. *112*, 186279g (1990). [8] H. A. Staab: *Einführung in die theoretische organische Chemie*, 4th edition, Verlag Chemie, Weinheim, 1964, p. 336ff. and p. 385ff. [9] S. Hünig, G. Bernhard, W. Liptay, and W. Brenninger: *Zum Problem der Solvatochromie bei Merocyaninen*, Liebigs Ann. Chem. *690*, 9 (1965); and references cited therein. [10] C. Reichardt: *Solvatochromic Dyes as Solvent Polarity Indicators*, Chem. Rev. *94*, 2319 (1994).

[11] L. G. S. Brooker: *Sensitizing and Desensitizing Dyes*, in C. E. K. Mees and T. H. James (eds.): *The Theory of the Photographic Process*, 3rd ed., MacMillan, New York, London, 1966, Chapter II, p. 198ff. (especially p. 219ff.); Chimia *15*, 87 (1961); ibid. *20*, 327 (1966). [12] S. Nigam and S. Rutan: *Principles and Applications of Solvatochromism*, Appl. Spectrosc. *55*, 362A (2001). [13] W. Liptay: *Elektrochromie – Solvatochromie*, Angew. Chem. *81*, 195 (1969); Angew. Chem. Int. Ed. Engl. *8*, 177 (1969); Z. Naturforsch. *20a*, 1441 (1965); ibid. *21a*, 1605 (1966). [14] N. Mataga and T. Kubota: *Molecular Interactions and Electronic Spectra*, Dekker, New York, 1970. [15] (a) K. Sone and Y. Fukuda: *Solvatochromism of Transition Metal Complexes with Organic Ligands in Donor and Acceptor Solvents*, Rev. Inorg. Chem. *11*, 123 (1990); (b) U. El-Ayaan, F. Murata, and Y. Fukuda: *Thermochromism and Solvatochromism in Solution*, in W. Linert (ed.): *Highlights in Solute–Solvent Interactions*, Springer, Wien, New York, 2002, p. 43ff. [16] P. Suppan and N. Ghoneim: *Solvatochromism*, Royal Society of Chemistry, Cambridge/U.K., 1997. [17] (a) P. Chen and T. J. Meyer: *Medium Effects on Charge Transfer in Metal Complexes*, Chem. Rev. *98*, 1439 (1998); (b) N. S. Hush and J. R. Reimers: *Solvent Effects on the Electronic Spectra of Transition Metal Complexes*, Chem. Rev. *100*, 775 (2000). [18] S. Dähne: *Systematik und Begriffserweiterung der Polymethinfarbstoffe*, Z. Chem. *5*, 441 (1965); S. Dähne and D. Leupold: *Der Polymethinzustand*, Ber. Bunsenges. Phys. Chem. *70*, 618 (1966); S. Dähne and F. Moldenhauer, Progr. Phys. Org. Chem. *15*, 1 (1985). [19] J. Fabian and H. Hartmann: *π-Electron Structure of Polymethines*, J. Mol. Struct. *27*, 67 (1975). [20] S. Dähne, D. Leupold, H.-E. Nikolajewski, and R. Radeglia, Z. Naturforsch. *20b*, 1006 (1965); R. Radeglia and S. Dähne, J. Mol. Struct. *5*, 399 (1970).

[21] K. Lauer and R. Oda, Ber. Dtsch. Chem. Ges. *69*, 851 (1936). [22] N. S. Bayliss and L. Hulme, Aust. J. Chem. *6*, 257 (1953). [23] A. L. Le Rosen and C. E. Reid, J. Chem. Phys. *20*, 233 (1952). [24] F. Feichtmayr, E. Heilbronner, A. Nürrenbach, H. Pommer, and J. Schlag, Tetrahedron *25*, 5383 (1969). [25] H. Christen and P. A. Straub, Helv. Chim. Acta *56*, 739, 1752 (1973). [26] N. Tyutyulkov, J. Fabian, A. Mehlhorn, F. Dietz, and A. Tadjer: *Polymethine Dyes – Structure and Properties*, St. Kliment Ohridski University Press, Sofia/Bulgaria, 1991. [27] A. A. Ishchenko: *Stroenie i Spektral'no-Luminescentnye Svojstva Polimethinovych Krasitelei (Constitution and Spectral-Luminescent Properties of Polymethine Dyes)*, Naukova Dumka, Kiev/Ukraine, 1994. [28] G. Bach and S. Dähne: *Cyanine Dyes and Related Compounds*, in *Rodd's Chemistry of Carbon Compounds*, 2nd ed., 2nd suppl., Elsevier, Amsterdam, 1997. [29] K. Dimroth, C. Reichardt, T. Siepmann, and F. Bohlmann, Liebigs Ann. Chem. *661*, 1 (1963); K. Dimroth and C. Reichardt, ibid. *727*, 93 (1969); C. Reichardt, ibid. *752*, 64 (1971); C. Reichardt and E. Harbusch-Görnert, ibid. *1983*, 721. [30] K. Dimroth and C. Reichardt, Z. Anal. Chem. *215*, 344 (1966); Z. B. Maksimović, C. Reichardt, and A. Spirić, ibid. *270*, 100 (1974).

[31] J. Ortega, C. Ràfols, E. Bosch, and M. Rosés, J. Chem. Soc., Perkin Trans. II 1497 (1996). [32] N. G. Bakshiev, M. I. Knyazhanskii, V. I. Minkin, O. A. Osipov, and G. V. Saidov: *Experimental Determination of the Dipole Moments of Organic Molecules in Excited Electronic States*, Usp. Khim. *38*, 1644 (1969); Russ. Chem. Rev. *38*, 740 (1969). [33] W. Liptay: *Dipole Moments and Polarizabilities of Molecules in Excited Electronic States*, in E. C. Linn (ed.): *Excited States*, Academic Press, New York, London, 1974, Vol. I, p. 129ff. [34] L. G. S. Brooker, A. C. Craig, D. W. Heseltine, P. W. Jenkins, and L. L. Lincoln, J. Am. Chem. Soc. *87*, 2443 (1965). [35] M. J. Kamlet, E. G. Kayser, J. W. Eastes, and W. H. Gilligan, J. Am. Chem. Soc. *95*, 5210 (1973); M. J. Kamlet and R. W. Taft, ibid. *98*, 377, 2886, 3233 (1976). [36] P. Scheibe, S. Schneider, F. Dörr, and E. Daltrozzo, Ber. Bunsenges. Phys. Chem. *80*, 630 (1976); S. Schneider, ibid. *80*, 218 (1976). [37] S. Dähne, F. Schob, and K.-D. Nolte, Z. Chem. *13*, 471 (1973); S. Dähne, F. Schob, K.-D. Nolte, and R. Radeglia, Ukr. Khim. Zh. (Russ. Ed.) *41*, 1170 (1975); Chem. Abstr. *84*, 43086j (1976). [38]

D. J. Cowley, J. Chem. Soc., Perkin Trans. II, 287 (1975). [39] L. G. S. Brooker and R. H. Sprague, J. Am. Chem. Soc. *63*, 3214 (1941). [40] J. Figueras, J. Am. Chem. Soc. *93*, 3255 (1971).

[41] F. Terenziani and A. Painelli, J. Phys. Chem. A *104*, 11041, 11049 (2000). [42] E. M. Kosower, J. Am. Chem. Soc. *80*, 3261 (1958). [43] S. Dähne and M. Seebacher, Z. Chem. *12*, 141 (1972). [44] M. M. Davies and H. B. Hetzer, Anal. Chem. *38*, 451 (1966). [45] A. J. Carmichael and K. R. Seddon, J. Phys. Org. Chem. *13*, 591 (2000). [46] M. Klessinger and W. Lüttke, Chem. Ber. *99*, 2136 (1966); M. Klessinger, Tetrahedron *22*, 3355 (1966). [47] E. Lippert, Z. Elektrochem., Ber. Bunsenges. Phys. Chem. *61*, 962 (1957); Angew. Chem. *73*, 695 (1961). [47a] W. Baumann, H. Deckers, K.-D. Loosen, and F. Petzke, Ber. Bunsenges. Phys. Chem. *81*, 799 (1977). [48] L. G. S. Brooker, G. H. Keyes, and D. W. Heseltine, J. Am. Chem. Soc. *73*, 5350 (1951). [49] S. Hünig and O. Rosenthal, Liebigs Ann. Chem. *592*, 161 (1955). [50] H. G. Benson and J. N. Murrell, J. Chem. Soc., Faraday Trans. II *1972*, 137. [50a] M. J. Minch and S. S. Shah, J. Chem. Educ. *54*, 709 (1977).

[51] H. W. Gibson and F. C. Bailey, Tetrahedron *30*, 2043 (1974); Can. J. Chem. *53*, 2162 (1975); J. Chem. Soc., Perkin Trans. II *1976*, 1575. [52] L. G. S. Brooker, G. H. Keyes, R. H. Sprague, R. H. VanDyke, E. VanLare, G. VanZandt, F. L. White, H. W. J. Cressman, and S. G. Dent, J. Am. Chem. Soc. *73*, 5332 (1951). [53] N. S. Bayliss and E. G. McRae, J. Am. Chem. Soc. *74*, 5803 (1952). [54] T. Eichler and V. Schäfer, Tetrahedron *30*, 4025 (1974). [55] C. Reichardt and W. Mormann, Chem. Ber. *105*, 1815 (1972). [56] P. H. Vandewyer, J. Hoefnagels, and G. Smets, Tetrahedron *25*, 3251 (1969). [57] C. Reichardt, Tetrahedron Lett. *1965*, 429; Liebigs Ann. Chem. *715*, 74 (1968). [58] E. M. Kosower and B. G. Ramsay, J. Am. Chem. Soc. *81*, 856 (1959). [59] A. Schweig, Z. Naturforsch. *22a*, 724 (1967). [60] H. Prinzbach and E. Woischnik, Helv. Chim. Acta *52*, 2472 (1969).

[61] A. Schweig and C. Reichardt, Z. Naturforsch. *21a*, 1373 (1966). [62] W. Liptay, H.-J. Schlosser, B. Dumbacher, and S. Hünig, Z. Naturforsch. *23a*, 1613 (1968). [63] R. Foster: *Organic Charge-Transfer Complexes*, Academic Press, London, New York, 1969, p. 62ff. [64] S. Dupire, J. M. Mulindabyuma, J. B. Nagy, and O. B. Nagy, Tetrahedron *31*, 135 (1975). [65] E. M. Kosower, J. Am. Chem. Soc. *80*, 3253 (1958); J. Chim. Phys. *61*, 230 (1964); E. M. Kosower and M. Mohammad, J. Am. Chem. Soc. *90*, 3271 (1968); J. Phys. Chem. *74*, 1153 (1970). [66] E. M. Kosower: *Molecular Biochemistry*, McGraw Hill, New York, 1962, p. 180ff. [67] E. M. Kosower: *An Introduction to Physical Organic Chemistry*, Wiley, New York, 1968, p. 293ff. [68] M. J. Blandamer and M. F. Fox: *Theory and Applications of Charge-Transfer-to-Solvent Spectra*, Chem. Rev. *70*, 59 (1970). [69] N. S. Bayliss and E. G. McRae, J. Phys. Chem. *58*, 1002, 1006 (1954). [70] E. G. McRae, J. Phys. Chem. *61*, 562 (1957).

[71] O. E. Weigang and D. D. Wild, J. Chem. Phys. *37*, 1180 (1962). [72] Th. Förster, Z. Elektrochem., Angew. Phys. Chem. *45*, 548 (1939); especially p. 570. [73] G. Scheibe, W. Seiffert, G. Hohlneicher, C. Jutz, and H. J. Springer, Tetrahedron Lett. *1966*, 5053. [74] R. Radeglia, D. Leupold, and S. Dähne, Ber. Bunsenges. Phys. Chem. *70*, 745, 1167 (1966). [75] R. Radeglia, Z. Phys. Chem. (Leipzig) *235*, 335 (1967); Z. Chem. *15*, 355 (1975). [76] S. Dähne and R. Radeglia, Tetrahedron *27*, 3673 (1971). [77] R. Radeglia, G. Engelhardt, E. Lippmaa, T. Pehk, K.-D. Nolte, and S. Dähne, Org. Magn. Reson. *4*, 571 (1972). [78] M. Wähnert and S. Dähne, J. Prakt. Chem. *318*, 321 (1976). [78a] M. Wähnert and S. Dähne, J. Signal AM *4*, 403 (1976). [79] S. Dähne and K.-D. Nolte, J. Chem. Soc., Chem. Commun. 1056 (1972); J. Prakt. Chem. *318*, 643 (1976); Acta Chim. Acad. Sci. Hung. *97*, 147 (1978); Chem. Abstr. *90*, 38406g (1979). [80] L. Onsager. J. Am. Chem. Soc. *58*, 1486 (1936).

[81] W. Liptay, Ber. Bunsenges. Phys. Chem. *80*, 207 (1976). [82] H. Labhart, Ber. Bunsenges. Phys. Chem. *80*, 240 (1976). [83] A. Schweig, Mol. Phys. *15*, 1 (1968). [84] G. Scheibe, E. Daltrozzo, O. Wörz, and J. Heiss, Z. Phys. Chem. (Frankfurt) N. F. *64*, 97 (1969). [85] W. Liptay, Z. Naturforsch. *21a*, 1605 (1966). [86] Y. Oshika, J. Phys. Soc. Jpn. *9*, 594 (1954); Chem. Abstr. *49*, 15483c (1955). [87] N. G. Bakshiev, Optika i Spektroskopiya *10*, 717 (1961); Optics and Spectroscopy (Engl. Ed.) *10*, 379 (1961); Chem. Abstr. *58*, 4027h (1963). [88] L. Bilot and A. Kawski, Z. Naturforsch. *17a*, 621 (1962). [89] T. Abe, Bull. Chem. Soc. Jpn. *54*, 327 (1981). [90] W. Liptay in O. Sinanoğlu (ed.): *Modern Quantum Chemistry*, Academic Press, New York, London, 1965, Part II, p. 173ff.; Z. Naturforsch. *20a*, 1441 (1965).

[91] H. Kuhn and A. Schweig, Chem. Phys. Lett. *1*, 255 (1967). [92] M. F. Nicol, J. Swain, Y.-Y. Shum, R. Merin, and R. H. H. Chen, J. Chem. Phys. *48*, 3587 (1968). [93] P. Suppan, J. Chem. Soc. *A 1968*, 3125. [94] W. Liptay and G. Walz, Z. Naturforsch. *26a*, 2007 (1971). [95] F. J. Kampas, Chem. Phys. Lett. *26*, 334 (1974). [95a] H. A. Germer, Theor. Chim. Acta *34*, 145 (1974); ibid. *35*, 273 (1974). [95b] K. D. Nolte and S. Dähne, Advances in Molecular Relaxation and Interaction Processes *10*, 299 (1977); Chem. Abstr. *87*, 151529b (1977). [96] R. M. Hochstrasser and L. J. Noe, J. Mol. Spectrosc. *38*, 175 (1971). [97] H. McConnell, J. Chem. Phys. *20*, 700 (1952). [98] G. J. Brealey and M. Kasha, J. Am. Chem. Soc. *77*, 4462 (1955). [99] G. C. Pimentel, J. Am. Chem. Soc. *79*, 3323 (1957). [100] R. S. Becker, J. Mol. Spectrosc. *3*, 1 (1959).

[101] E. Lippert: *Der Einfluß von Wasserstoffbrücken auf Elektronenspektren*, in D. Hadži and H. W. Thompson (eds.): *Hydrogen Bonding*, Pergamon Press, London, 1959, p. 217ff. [102] M. Ito, K. Inuzuka, and S. Imanishi, J. Am. Chem. Soc. *82*, 1317 (1960); Bull. Chem. Soc. Jpn. *34*, 467 (1961). [103] J. E. Dubois, E. Goetz, and A. Bienvenüe, Spectrochim. Acta *20*, 1815 (1964). [103a] W. P. Hayes and C. J. Simmons, Spectrochim. Acta *21*, 529 (1965). [104] W. L. Dilling, J. Org. Chem. *31*, 1045 (1966). [105] J. E. Del Bene, J. Am. Chem. Soc. *95*, 6517 (1973); ibid. *96*, 5643 (1974). [106] S. Spange and D. Keutel, Liebigs Ann. Chem. 423 (1992). [107] O. Exner: *Dipole Moments in Organic Chemistry*, Thieme, Stuttgart, 1975, p. 122. [108] N. J. Turro: *Modern Molecular Photochemistry*, Benjamin, Menlo Park, 1978, p. 108. [109] G. Porter and P. Suppan, Trans. Faraday Soc. *61*, 1664 (1965); ibid. *62*, 3375 (1966). [110] J. H. Fendler, E. J. Fendler, G. A. Infante, P.-S. Shih, and L. K. Patterson, J. Am. Chem. Soc. *97*, 89 (1975).

[111] D. P. Stevenson, G. M. Coppinger, and J. W. Forbes, J. Am. Chem. Soc. *83*, 4350 (1961). [112] B. Valeur: *Molecular Fluorescence – Principles and Applications*, Wiley-VCH, Weinheim, 2002, Chapter 7, p. 200ff. [113] W. Schmidt: *Optische Spektroskopie – Eine Einführung*, Wiley-VCH, Weinheim, 2000, Chapter 5.2.4, p. 197ff. [114] A. K. Sharma and S. G. Schulman: *Introduction to Fluorescence Spectroscopy*, Wiley, Chichester, 1999. [115] O. S. Wolfbeis (ed.): *Fluorescence Spectroscopy – New Methods and Applications*, Springer, Berlin, 1992. [116] W. Rettig, Top. Curr. Chem. *169*, 253 (1994). [117] H. Heitele: *Dynamische Lösungsmitteleffekte bei Elektronentransferreaktionen*, Angew. Chem. *105*, 378 (1993); Angew. Chem. Int. Ed. Engl. *32*, 359 (1993). [118] R. A. Marcus: *Electron-Transfer Reactions in Solution*, Angew. Chem. *105*, 1161 (1993); Angew. Chem. Int. Ed. Engl. *32*, 1111 (1993). [119] E. M. Kosower, H. Dodiuk, K. Tanizawa, M. Ottolenghi, and N. Orbach, J. Am. Chem. Soc. *97*, 2167 (1975). [120] E. M. Kosower, Acc. Chem. Res. *15*, 259 (1982).

[121] A. Rodger and B. Nordén: *Circular Dichroism and Linear Dichroism*, Oxford University Press, Oxford/U.K., 1997. [122] G. Snatzke: *Anwendung von Polarimetrie, CD und ORD in der organischen Stereochemie*, in F. Korte (ed.): *Methodicum Chimicum*, Vol. I, Part 1, p. 426ff., Thieme, Stuttgart, and Academic Press, New York, 1973; G. Snatzke, Angew. Chem. *91*, 380 (1979); Angew. Chem. Int. Ed. Engl. *18*, 363 (1979). [123] (a) E. L. Eliel, S. H. Wilen, and L. N. Mander: *Stereochemistry of Organic Compounds*, Wiley, New York, 1994, Chapters 13-2 to 13-5, p. 992ff.; (b) E. L. Eliel, S. H. Wilen, and M. P. Doyle: *Basic Organic Chemistry*, Wiley-Interscience, New York, 2001, Chapter 12-5, p. 585ff. [124] N. Tokura: *Solvent Effects and Circular Dichroism*, Kagaku No Ryoiki *28*, 682 (1974); Chem. Abstr. *82*, 147175 (1975). [125] C. Coulombeau and A. Rassat, Bull. Soc. Chim. Fr. *1966*, 3752. [126] D. N. Kirk, W. Klyne, and S. R. Wallis, J. Chem. Soc. *C 1970*, 350. [127] R. Corriu and J. Masse, Tetrahedron *26*, 5123 (1970). [128] W. Klyne, D. N. Kirk, J. Tilley, and H. Suginome, Tetrahedron *36*, 543 (1980). [129] D. W. Miles, M. J. Robins, R. K. Robins, M. W. Winkley, and H. Eyring, J. Am. Chem. Soc. *91*, 824 (1969). [130] M. Cinquini, S. Colonna, I. Moretti, and G. Torre, Tetrahedron Lett. *1970*, 2773.

[131] B. Bosnich, J. Am. Chem. Soc. *89*, 6143 (1967). [132] N. Tokura, T. Nagai, S. Takenaka, and T. Oshima, J. Chem. Soc., Perkin Trans. II, 337 (1974). [133] Pham Van Huong and J. Lascombe, C. R. H. Acad. Sci., Ser *C 254*, 2543 (1962); Chem. Abstr. *57*, 297 (1962). [134] H. E. Hallam: *Hydrogen Bonding and Solvent Effects*, in M. Davis (ed.): *Infra-Red Spectroscopy and Molecular Structure*, Elsevier, Amsterdam, 1963, Chapter XII, p. 405ff. [135] M. D. Joesten and L. J. Schaad: *Hydrogen Bonding*, Dekker, New York, 1974. [136] L. J. Bellamy: *Solvent Effects on Infra-Red Group Frequencies*, in E. Thornton and H. W. Thompson (eds.): *Proceedings of the Conference on*

Molecular Spectroscopy, held in London, 1958, Pergamon Press, London, 1959, p. 216ff.; Spectrochim. Acta *14*, 192 (1959). [137] R. L. Williams: *Infrared Spectra and Molecular Interactions*, Ann. Rept. Progr. Chem. (Chem. Soc. London) *58*, 34 (1961); Chem. Abstr. *57*, 9219e (1962). [138] M.-L. Josien: *Infrared-Spectroscopy Study of Molecular Interactions in Liquids*, Pure Appl. Chem. *4*, 33 (1962); J. Chim. Phys. *61*, 245 (1964). [139] C. N. R. Rao: *Chemical Applications of Infrared Spectroscopy*, Academic Press, New York, London, 1963, p. 577ff. [140] M. Horák and J. Plíva: *Studies of Solute-Solvent Interactions. General Considerations*, Spectrochim. Acta *21*, 911 (1965).

[141] B. Schrader (ed.): *Infrared and Raman Spectroscopy*, Wiley-VCH, Weinheim, 1995. [142] N. Oi: *Infrared Solvents Shifts and Molecular Interactions*, Bunko Kenkyu *18*, 194 (1969); Chem. Abstr. *72*, 49158 (1970). [143] C. Agami: *Méthodes spectrographiques d'étude des solvants*, Bull. Soc. Chim. Fr. *1969*, 2183. [144] L. J. Bellamy: *The Infrared Spectra of Complex Molecules*, 3rd ed., Methuen, London, 1975; *Ultrarot-Spektrum und chemische Konstitution* (translated into German by W. Brügel), 2nd ed., Steinkopff, Darmstadt, 1966. [145] H. W. Thompson and D. J. Jewell, Spectrochim. Acta *13*, 254 (1958). [146] L. J. Bellamy and R. L. Williams, Trans. Faraday Soc. *55*, 14 (1959); Proc. Roy. Soc., London, Ser. A *255*, 22 (1960). [147] A. D. E. Pullin, Spectrochim. Acta *16*, 12 (1960). [148] C. Agami, Bull. Soc. Chim. Fr. *1968*, 2033. [149] A. G. Burden, G. Collier, and J. Shorter, J. Chem. Soc., Perkin Trans. II, 1627 (1976). [150] N. Oi and J. F. Coetzee, J. Am. Chem. Soc. *91*, 2473 (1969).

[151] L. J. Bellamy and P. E. Rogasch, Spectrochim. Acta *16*, 30 (1960). [152] Y. Ikegami, Bull. Chem. Soc. Jpn. *35*, 972 (1962). [153] T. Gramstad, Spectrochim. Acta *19*, 1363 (1963). [154] A. Allerhand and P. v. R. Schleyer, J. Am. Chem. Soc. *85*, 371 (1963); ibid. *86*, 5709 (1964). [155] T. Kagiya, Y. Sumida, and T. Inoue, Bull. Chem. Soc. Jpn. *41*, 767 (1968). [156] W. R. Fawcett, G. Liu, and T. E. Kessler, J. Phys. Chem. *97*, 9293 (1993). [157] J. S. Byrne, P. F. Jackson, and K. J. Morgan, J. Chem. Soc., Perkin Trans. II, 1291 (1972); ibid. *1976*, 1800. [158] M.-L. Josien and N. Fuson, J. Chem. Phys. *22*, 1169, 1264 (1954). [159] J. Devaure, Pham. Van Huong, and J. Lascombe, J. Chim. Phys. *65*, 1686 (1968). [160] W. Lesch and R. Ulbrich, Z. Naturforsch. *23a*, 1639 (1968).

[161] M. F. El Bermani, A. J. Woodward, and N. Jonathan, J. Am. Chem. Soc. *92*, 6750 (1970). [162] L. J. Bellamy, H. E. Hallam, and R. L. Williams, Trans. Faraday Soc. *54*, 1120 (1958). [163] H. E. Hallam, J. Mol. Struct. *3*, 43 (1969). [164] L. J. Bellamy and P. E. Rogasch, J. Chem. Soc. *1960*, 2218. [165] H. Götz, E. Heilbronner, A. R. Katritzky, and R. A. Jones, Helv. Chim. Acta *44*, 387 (1961). [166] J. G. Kirkwood, in W. West and R. T. Edwards, J. Chem. Phys. *5*, 14 (1937). [167] E. Bauer and M. Magat, J. Physique Radium *9*, 319 (1938); Chem. Abstr. *32*, 8856 (1938). [168] N. S. Bayliss, A. R. H. Cole, and L. H. Little, Aust. J. Chem. *8*, 26 (1955); Spectrochim. Acta *15*, 12 (1959). [169] A. D. E. Pullin, Spectrochim. Acta *13*, 125 (1958); ibid. *16*, 12 (1960). [170] A. D. Buckingham, Proc. Roy. Soc. London, Ser. A *248*, 169 (1957); ibid. A *255*, 32 (1960); Trans. Faraday Soc. *56*, 753 (1960).

[171] R. Ulbrich, Z. Naturforsch, *23a*, 1323 (1968). [172] J. Barriol, P. Bonnet, and J. Devaure, J. Chim. Phys. *71*, 107 (1974). [173] J. Fruwert and G. Geiseler: *Infrarotintensitäten*, Z. Chem. *20*, 157 (1980). [174] B. Lindner, J. Phys. Chem. *96*, 10708 (1992). [175] P. Mirone, Spectrochim. Acta *22*, 1897 (1966). [176] T. L. Brown, Spectrochim. Acta *10*, 149 (1957). [177] G. L. Caldow, D. C. Jones, and H. W. Thompson, Proc. Roy. Soc. London, Ser. A *254*, 17 (1960). [178] R. Heess and H. Kriegsmann, Spectrochim. Acta, Part A *24*, 2121 (1968). [179] J. Fruwert, G. Geiseler, and D. Luppa, Z. Chem. *8*, 238 (1968). [180] H. P. Figeys and V. Mahieu, Spectrochim. Acta, Part A *24*, 1553 (1968).

[181] M. Kakimoto and T. Fujiyama, Bull. Chem. Soc. Jpn. *48*, 2258 (1975). [182] C. M. Huggins and G. C. Pimentel, J. Chem. Phys. *23*, 896 (1955). [183] J. Gendell, J. H. Freed. and G. K. Fraenkel: *Solvent Effects in Electron Spin Resonance Spectra*, J. Chem. Phys. *37*, 2832 (1962). [184] N. M. Atherton: *Principles of Electron Spin Resonance*, Ellis Horwood/Prentice Hall, Hemel Hempstad, 1993. [185] M. C. R. Symons: *Application of ESR Spectroscopy to the Study of Solvation*, Pure Appl. Chem. *49*, 13 (1977). [186] (a) L. R. Dalton, A. Bain, and C. K. Westbrook: *Recent Advances in Electron Paramagnetic Resonance*, Annu. Rev. Phys. Chem. *41*, 389 (1990). (b) J. H. Freed: *New Techniques in ESR*, Annu. Rev. Phys. Chem. *51*, 655 (2000). [187] G. R. Luck-

hurst and L. E. Orgel, Mol. Phys. *8*, 117 (1964). [188] N. Hirota: *Metal Ketyls and Related Radical Ions*, in E. T. Kaiser and L. Kevan (eds.): *Radical Ions*, Wiley-Interscience, New York, 1968, Chapter 2, p. 35ff. [189] E. W. Stone and A. H. Maki, J. Chem. Phys. *36*, 1944 (1962); J. Am. Chem. Soc. *87*, 454 (1969). [190] E. A. C. Lucken, J. Chem. Soc. *1963*, 5123.

[191] P. J. Zandstra, J. Chem. Phys. *41*, 3655 (1964). [192] T. A. Claxton, J. Oakes, and M. C. R. Symons, Trans. Faraday Soc. *63*, 2125 (1967). [193] J. Oakes and M. C. R. Symons, Trans. Faraday Soc. *64*, 2579 (1968). [194] T. A. Claxton and D. McWilliams, Trans. Faraday Soc. *64*, 2593 (1968). [195] J. A. Pedersen and J. Spanget-Larsen, Chem. Phys. Lett. *35*, 41 (1975). [196] G. A. Russell: *Semidione Radical Anions*, in E. T. Kaiser and L. Kevan (eds.): *Radical Ions*, Wiley-Interscience, New York, 1968, Chapter 3, p. 87ff. [197] P. J. Zandstra and E. M. Evleth, J. Am. Chem. Soc. *86*, 2664 (1964). [198] K. Scheffler and H. B. Stegmann, Z. Phys. Chem. (Frankfurt) N. F. *44*, 353 (1965). [199] K. Mukai, H. Nishiguchi, K. Ishizu, Y. Deguchi, and H. Takaki, Bull. Chem. Soc. Jpn. *40*, 2731 (1967). [200] S. Aono and M. Suhara, Bull. Chem. Soc. Jpn. *41*, 2553 (1968).

[201] J. Q. Chambers, P. Ludwig, T. Layloff, and R. N. Adams, J. Phys. Chem. *68*, 661 (1964); J. Am. Chem. Soc. *86*, 4568 (1964). [202] J. Pannel, Mol. Phys. *7*, 317 (1964). [203] M. T. Jones: *Recent Advances in the Chemistry of Aromatic Anion Radicals*, in E. T. Kaiser and L. Kevan (eds.): *Radical Ions*, Wiley-Interscience, New York, 1968, p. 245ff. (especially p. 261ff.). [204] J. H. Sharp and M. C. R. Symons: *Electron Spin Resonance Studies of Ion Pairs*, in M. Szwarc (ed.): *Ions and Ion Pairs in Organic Reactions*, Wiley-Interscience, New York, 1972, Vol. 1, Chapter 5, p. 177ff. (especially fig. 4). [205] G. R. Stevenson and H. Hidalgo, J. Phys. Chem. *77*, 1027 (1973). [206] (a) H. G. Aurich and W. Weiss, Fortschr. Chem. Forsch. *59*, 65 (1975); (b) H. G. Aurich: *Nitroxides*, in S. Patai and Z. Rappoport (eds.): *Nitrones, Nitronates, and Nitroxides (Chemistry of Functional Groups)*, Wiley, Chichester, 1989, Chapter 4, p. 313ff. [207] Y. Deguchi, Bull. Chem. Soc. Jpn. *35*, 260 (1962); K. Mukai, H. Nishiguchi, K. Ishizu, Y. Deguchi, and H. Takaki, ibid. *40*, 2731 (1967). [208] A. V. Il'yasov, Zh. Strukt. Khim. *3*, 95 (1962); Chem. Abstr. *58*, 7523d (1963). [209] A. L. Buchachenko, Dokl. Akad. Nauk USSR *158*, 932 (1964); Chem. Abstr. *62*, 2390g (1965). [210] R. Briére, G. Chapelet-Letourneux, H. Lemaire, and A. Rassat, Tetrahedron Lett. *1964*, 1775; Bull. Soc. Chim. Fr. *1965*, 444, 3273.

[211] P. B. Ayscough and F. P. Sargent, J. Chem. Soc., Part B *1966*, 907. [212] T. Kawamura, S. Matsunami, and T. Yonezawa, Bull. Chem. Soc. Jpn. *40*, 1111 (1967). [213] Th. A. J. W. Wajer, A. Mackor, and Th. J. de Boer, Tetrahedron *25*, 175 (1969). [214] G. H. Dodd, M. D. Barratt, and L. Rayner, FEBS Letters *8*, 286 (1970). [215] E. F. Ullmann, L. Call, and J. H. Osiecki, J. Org. Chem. *35*, 3623 (1970); E. F. Ullmann and L. Call, J. Am. Chem. Soc. *92*, 7210 (1970). [216] H. Hayat and B. L. Silver, J. Phys. Chem. *77*, 72 (1973). [217] G. Stout and J. B. F. N. Engberts, J. Org. Chem. *39*, 3800 (1974). [218] B. R. Knauer and J. J. Napier, J. Am. Chem. Soc. *98*, 4395 (1976). [219] H. G. Aurich and J. Trösken, Liebigs Ann. Chem. *745*, 159 (1971). [220] H. G. Aurich, W. Dersch, and H. Forster, Chem. Ber. *106*, 2854 (1973).

[221] F. W. Heineken, M. Bruin, and F. Bruin, J. Chem. Phys. *37*, 452 (1962). [222] L. Grossi, F. Minisci, and G. F. Pedulli, J. Chem. Soc., Perkin Trans. II, 943 (1977); M. Guerra, F. Bernardi, and G. F. Pedulli, Chem. Phys. Lett. *48*, 311 (1977). [223] M. C. R. Symons, J. Phys. Chem. *71*, 172 (1967). [224] F. Gerson: *Hochauflösende ESR-Spektroskopie – dargestellt anhand aromatischer Radikal-Ionen*, Verlag Chemie, Weinheim 1967, p. 30ff., 128, and 173ff. [225] J. E. Gordon: *The Organic Chemistry of Electrolyte Solutions*, Wiley-Interscience, New York, 1975, p. 384ff. [226] A. A. Bothner-By and R. E. Glick, J. Chem. Phys. *26*, 1651 (1957). [227] L. W. Reeves and W. G. Schneider, Can. J. Chem. *35*, 251 (1957). [228] E. O. Bishop: *Chemical Shifts – Intermolecular Effects: Solvent Dependent Studies*, Annu. Rep. Progr. Chem. for 1961 (Chem. Soc. London) *58*, 71 (1962). [229] G. Mavel: *NMR and Molecular Interactions in the Liquid Phase*, J. Chim. Phys. *61*, 182 (1964). [230] I. Yamaguchi: *Solvent Effects on the High Resolution NMR Spectrum*, Kogyo Kagaku Zasshi *68*, 1328 (1965); Chem. Abstr. *64*, 2884g (1966); Kagaku (Kyoto) *20*, 328 (1965); Chem. Abstr. *64*, 10618g (1966).

[231] (a) H. Günther: *NMR Spectroscopy: Basic Principles, Concepts, and Applications in Chemistry*, 2nd ed., Wiley, New York, 1995; (b) R. S. Macomber: *A Complete Introduction to Modern*

NMR Spectroscopy, Wiley, New York, 1997; (c) C. J. Jameson: *Understanding NMR Chemical Shifts*, Annu. Rev. Phys. Chem. *47*, 135 (1996). [232] P. Laszlo: *Solvent Effects and Nuclear Magnetic Resonance*, in J. W. Emsley, F. Feeney, and L. H. Sutcliffe (eds.): *Progress in Nuclear Magnetic Resonance Spectroscopy*, Pergamon Press, Oxford, 1967, Vol. 3, Chapter 6, p. 231ff. [233] J. Ronayne and D. H. Williams: *Solvent Effects in Proton Magnetic Resonance Spectroscopy*, in E. F. Mooney (ed.): *Annual Review of NMR Spectroscopy*, Academic Press, London, New York, 1969, Vol. 2, p. 83ff. [234] M. I. Foreman: *Medium Effects in Nuclear Magnetic Resonance* (Specialist Periodical Report, Chem. Soc. London) *2*, 355 (1973); ibid. *5*, 292 (1976); Nucl. Magn. Reson. *6*, 233 (1977). [235] J. Homer: *Solvent Effects on Nuclear Magnetic Resonance Chemical Shifts*, Applied Spectroscopy Reviews *9*, 1 (1975). [236] S. L. Smith: *Solvent Effects and NMR Coupling Constants*, Fortschr. Chem. Forsch. *27*, 117 (1972). [237] M. Barfield and M. D. Johnston: *Solvent Dependence of Nuclear Spin-Spin Coupling Constants*, Chem. Rev. *73*, 53 (1973). [238] J. J. Dechter and J. I. Zink, J. Am. Chem. Soc. *97*, 2937 (1975). [239] J. F. Hinton and R. W. Briggs, J. Magn. Reson. *19*, 393 (1975); J. Sol. Chem. *6*, 827 (1977); J. F. Hinton, K. R. Metz, and R. W. Briggs, Progr. NMR Spectrosc. *13*, 211 (1982). [240] U. Mayer, V. Gutmann, and W. Gerger, Monatsh. Chem. *106*, 1235 (1975); ibid. *108*, 489 (1977).

[241] R. W. Taft, E. Price, I. R. Fox, I. C. Lewis, K. K. Andersen, and G. T. Davis, J. Am. Chem. Soc. *85*, 709, 3146 (1963). [242] M. R. Bacon and G. E. Maciel, J. Am. Chem. Soc. *95*, 2413 (1973). [243] D. Ziessow and M. Carroll, Ber. Bunsenges. Phys. Chem. *76*, 61 (1972). [244] Y. M. Cahen, P. R. Handy, E. T. Roach, and A. I. Popov, J. Phys. Chem. *79*, 80 (1975). [245] M. S. Greenberg, R. L. Bodner, and A. I. Popov, J. Phys. Chem. *77*, 2449 (1973). [246] W. J. DeWitte, R. C. Schoening, and A. I. Popov, Inorg. Nucl. Chem. Lett. *12*, 251 (1976); W. J. DeWitte, L. Liu, E. Mei, J. L. Dye, and A. I. Popov, J. Sol. Chem. *6*, 337 (1977). [247] See references [106–111] to Chapter 2. [248] W. F. Reynolds and U. R. Priller, Can. J. Chem. *46*, 2787 (1968). [249] R. G. Anderson and M. C. R. Symons, Trans. Faraday Soc. *65*, 2537 (1969). [250] T. G. Beaumont and K. M. C. Davis, J. Chem. Soc., Part B *1970*, 592.

[251] G. E. Maciel and R. V. James, J. Am. Chem. Soc. 86, 3893 (1964); G. E. Maciel and J. J. Natterstad, J. Chem. Phys. *42*, 2752 (1965). [252] G. L. Nelson, G. C. Levy, and J. D. Cargioli, J. Am. Chem. Soc. *94*, 3089 (1972). [253] S. Ueji and M. Nakamura, Tetrahedron Lett. *1976*, 2549. [254] H. A. Christ and P. Diehl, Helv. Phys. Acta *36*, 170 (1963); Chem. Abstr. *59*, 4702f (1963). [255] T. Schaefer and W. G. Schneider, J. Chem. Phys. *32*, 1218, 1224 (1960). [256] G. V. Rao, M. Balakrishnan, and N. Venkatasubramanian, Indian J. Chem. *13*, 1090 (1975). [257] R. L. Lichter and J. D. Roberts, J. Phys. Chem. *74*, 912 (1970). [258] F. A. L. Anet and I. Yavari, J. Org. Chem. *41*, 3589 (1976). [259] C. S. Giam and J. L. Lyle, J. Am. Chem. Soc. *95*, 3235 (1973). [260] E. Y. Lau and J. T. Gerig, J. Am. Chem. Soc. *118*, 1194 (1996).

[261] G. E. Maciel and R. V. James, Inorg. Chem. *3*, 1650 (1964). [262] A. D. Buckingham, T. Schaefer, and W. G. Schneider, J. Chem. Phys. *32*, 1227 (1960); ibid. *34*, 1064 (1961). [263] F. H. A. Rummens: *Van der Waals Forces in NMR – Intermolecular Shielding Effects*, in P. Diehl, E. Fluck, and R. Kosfield (eds.): *NMR – Basic Principles and Progress*, Springer, Berlin, 1975, Vol. 10, p. 1ff. [264] A. D. Buckingham, Can. J. Chem. *38*, 300 (1960). [265] I. D. Kuntz and M. D. Johnston, J. Am. Chem. Soc. *89*, 6008 (1967). [266] F. H. A. Rummens, J. Am. Chem. Soc. *92*, 3214 (1970). [267] H. J. Bernstein: *Solvent Effects on NMR Spectra of Gases and Liquids*, Pure Appl. Chem. *32*, 79 (1972). [268] J. T. Arnold and M. E. Packard, J. Chem. Phys. *19*, 1608 (1951). [269] U. Liddel and N. F. Ramsay, J. Chem. Phys. *19*, 1608 (1951). [270] E. Lippert: *Wasserstoffbrückenbindung und magnetische Protonenresonanz*, Ber. Bunsenges. Phys. Chem. *67*, 267 (1963).

[271] A. S. N. Murthy and C. N. R. Rao: *Spectroscopic Studies of the Hydrogen Bond*, in E. G. Brame (ed.): *Applied Spectroscopy Reviews*, Dekker, New York, London, 1969, Vol. 2, p. 69ff. (especially p. 106). [272] J. C. Davis and K. K. Deb: *Analysis of Hydrogen Bonding and Related Association Equilibria by Nuclear Magnetic Resonance*, in J. S. Waugh (ed.): *Advances in Magnetic Resonance*, Academic Press, New York, London, 1970, Vol. 4, p. 201ff. [273] J. V. Hatton and R. E. Richards, Trans. Faraday Soc. *57*, 28 (1961). [274] M. L. Martin, Annales de Physique *7*, 35 (1962); G. J. Martin and M. L. Martin, J. Chim. Phys. *61*, 1222 (1964). [275] C. Agami, M. Andrac-Taussig, and C. Prévost, Bull. Soc. Chim. Fr. *1968*, 952, 4467. [276] J. Reuben, J. Am.

Chem. Soc. *91*, 5725 (1969). [277] See also references [37–46] to Chapter 2. [278] P. Laszlo, Bull. Soc. Chim. Fr. *1964*, 2658. [279] E. M. Engler and P. Laszlo, J. Am. Chem. Soc. *93*, 1317 (1971). [280] See also references [17–22] to Chapter 3.

[281] G. R. Luckhurst: *Liquid Crystals as Solvents in NMR*, Quart. Rev. (London) *22*, 179 (1968). [282] J. Bulthuis: *NMR in Liquid-Crystalline Solvents*, Amsterdam, 1974; Chem. Abstr. *82*, 148359n (1975). [283] J. W. Emsley and J. C. Lindon: *NMR Spectroscopy Using Liquid-Crystal Solvents*, Pergamon Press, Oxford, 1975. [284] W. H. Pirkle and S. D. Beare, J. Am. Chem. Soc. *89*, 5485 (1967); W. H. Pirkle, S. D. Beare, and T. G. Burlingame, J. Org. Chem. *34*, 470 (1969). [285] D. F. Evans, J. Chem. Soc. *1963*, 5575. [286] P. Laszlo, Bull. Soc. Chim. Fr. *1966*, 558. [287] V. S. Watts and J. H. Goldstein, J. Phys. Chem. *70*, 3887 (1966). [288] S. L. Smith and R. H. Cox, J. Chem. Phys. *45*, 2848 (1966). [289] K. A. McLauchlan, L. W. Reeves, and T. Schaefer, Can. J. Chem. *44*, 1473 (1966). [290] R. L. Schmidt, R. S. Butler, and J. H. Goldstein, J. Phys. Chem. *73*, 1117 (1969).

[291] A. M. Ihrig and S. L. Smith, J. Am. Chem. Soc. *94*, 34 (1972). [292] A. Mishra, R. K. Behera, P. K. Behera, B. K. Mishra, and G. B. Behera, Chem. Rev. *100*, 1973 (2000). [293] D. J. Lougnot, P. Brunero, J. P. Fouassier, and J. Faure, J. Chim. Phys. Phys.-Chim. Biol. *79*, 343 (1982). [294] C. Reichardt: *Empirical Parameters of Solvent Polarity and Chemical Reactivity*, in H. Ratajczak and W. J. Orville-Thomas (eds.): *Molecular Interactions*, Wiley, Chichester, 1982, Vol. 3, Chapter 5, p. 241ff. [295] H. Langhals, Angew. Chem. *94*, 739 (1982); Angew. Chem. Int. Ed. Engl. *21*, 724 (1982). [296] H. Langhals: *Description of Properties of Binary Solvent Mixtures*, in R. I. Zalewski, T. M. Krygowski, and J. Shorter (eds.): *Similarity Models in Organic Chemistry, Biochemistry and Related Fields*, Elsevier, Amsterdam, 1991, Chapter 6, p. 283ff. [297] B. P. Johnson, B. Gabrielsen, M. Matulenko, J. G. Dorsey, and C. Reichardt: *Solvatochromic Solvent Polarity Measurements: Synthesis and Applications of ET-30*, Anal. Lett. *19*, 939 (1986). [298] P. Plieninger and H. Baumgärtel, Ber. Bunsenges. Phys. Chem. *86*, 161 (1982); Liebigs Ann. Chem. *1983*, 860. [299] K. A. Zachariasse, N. Van Phuc, and B. Kozankiewicz, J. Phys. Chem. *85*, 2676 (1981). [300] C. J. Drummond, F. Grieser, and T. W. Healy, Faraday Discuss. Chem. Soc. *81*, 95 (1986); Chem. Abstr. *106*, 126332k (1987).

[301] A. J. M. Van Beijnen, R. J. M. Nolte, and W. Drenth, Recl. Trav. Chim. Pays-Bas *105*, 255 (1986). [302] B. P. Johnson, M. G. Khaledi, and J. G. Dorsey, J. Chromatogr. *384*, 221 (1987); Anal. Chem. *58*, 2354 (1986); Chimia oggi (Milano) *1986* (11), 23. [303] For new methods for the determinatioon of excited-state dipole moments by solvatochromic effects, see: (a) J. J. Moura Ramos, M.-L. Stien, and J. Reisse, Recl. Trav. Chim. Pays-Bas *98*, 338 (1979); (b) P. Suppan, Chem. Phys. Lett. *94*, 272 (1983); (c) T. Abe and I. Iweibo, Bull. Chem. Soc. Jpn. *58*, 3415 (1985); (d) N. H. Ayachit, D. K. Deshpande, M. A. Shashidhar, and K. S. Rao, Spectrochim. Acta, Part A *42*, 585 (1986); (e) A. Samanta, T. P. Radhakrishnan *et al.*, J. Phys. Chem. *98*, 9133 (1994); J. Chem. Soc., Faraday Trans. *91*, 2739 (1995). [304] J. O. Morley, R. M. Morley, R. Docherty, and M. H. Charlton, J. Am. Chem. Soc. *119*, 10192 (1997). [305] K. Suzuki *et al.*, Anal. Chim. Acta *373*, 271 (1998); Anal. Chem. *70*, 1255 (1998). [306] I. Gruda and F. Bolduc, J. Org. Chem. *49*, 3300 (1984). [307] M. M. Habashy, F. El-Zawawi, M. S. Antonious, A. K. Sherif, and M. S. A. Abdel-Mottaleb, Indian J. Chem., Part A *24*, 908 (1985). [308] A. Botrel, A. Le Beuze, P. Jacques, and H. Straub, J. Chem. Soc., Faraday Trans. II *80*, 1235 (1984). [309] P. Jacques, J. Phys. Chem. *90*, 5535 (1986). [310] A. Le Beuze, A. Botrel, A. Samat, P. Appriou, and R. Guglielmetti, J. Chim. Phys. *75*, 255, 267 (1978).

[311] *Heterocyclic betaine dyes (mesoionic zwitterions)*: (a) Z. Pawelka and L. Sobczyk, J. Chem. Soc., Faraday Trans. I *76*, 43 (1980); (b) J.-H. Finkentey and H. W. Zimmermann, Liebigs Ann. Chem. *1981*, 1; (c) J. Bendig, D. Kreysig, E. Sauer, and M. Just, J. Prakt. Chem. *323*, 529 (1981); Z. Phys. Chem. (Leipzig) *263*, 442 (1982); (d) M. S. A. Abdel-Mottaleb, Z. Naturforsch., Part A *37*, 1353 (1982); Z. Phys. Chem. (Leipzig) *265*, 154 (1984); (e) S. Arai, M. Yamazaki, K. Nagakura, M. Ishikawa, and M. Hida, J. Chem. Soc., Chem. Commun. 1037 (1983); Sen-I Gakkaishi *42*, T-74 (1986); (f) V. Kampars, V. Kokars, Z. Bruvers, and O. Neilands, Zh. Obshch. Khim. *53*, 2299 (1983); Chem. Abstr. *100*, 50949v (1984); (g) P. de Mayo, A. Safarzadeh-Amiri, and S. K. Wong, Can. J. Chem. *62*, 1001 (1984); (h) H. Gotthardt and M. Oppermann, Chem. Ber. *119*, 2094 (1986). [312] *Substituted azobenzenes*: (a) H. Mustroph and J. Epperlein, J. Prakt. Chem. *322*, 305 (1980);

(b) B. Marcandalli, E. Dubini-Paglia, and L. Pellicciari-Di Liddo, Gazz. Chim. Ital. *111*, 187 (1981); (c) K. S. Schanze, T. F. Mattox, and D. G. Whitten, J. Org. Chem. *48*, 2808 (1983); (d) Y. Sueishi, M. Asano, S. Yamamoto, and N. Nishimura, Bull. Chem. Soc. Jpn. *58*, 2729 (1985). [313] *Quinone methides*: L. Pavličková, B. Koutek, V. Jehlička, and M. Souček, Collect. Czech. Chem. Commun. *48*, 2376 (1983). [314] *Thiolate ions and thiyl radicals*: (a) G. H. Morine and R. R. Kuntz, Chem. Phys. Lett. *67*, 552 (1979); (b) O. Ito and M. Matsuda, J. Phys. Chem. *88*, 1002 (1984); (c) T. J. Novak, S. G. Pleva, and J. Epstein, Anal. Chem. *52*, 1851 (1980). [315] J. W. Larsen, A. G. Edwards, and P. Dobi, J. Am. Chem. Soc. *102*, 6780 (1980). [316] T. R. Griffiths and R. H. Wijayanayake, Trans. Faraday Soc. *66*, 1563 (1970); J. Chem. Soc., Faraday Trans. I *69*, 1899 (1973). [317] J. L. Dye, Angew. Chem. *91*, 613 (1979); Angew. Chem. Int. Ed. Engl. *18*, 587 (1979). [318] (a) J. E. Brady and P. W. Carr: *An Analysis of Dielectric Models of Solvatochromism*, J. Phys. Chem. *89*, 5759 (1985); (b) B. S. Brunschwig, S. Ehrenson, and N. Sutin: *Solvatochromism and Intramolecular Electron-Transfer Barriers in Spheroidal Molecules*, J. Phys. Chem. *91*, 4714 (1987). [319] A. A. Ishchenko, N. A. Derevyanko, and A. I. Tolmachev, Dokl. Akad. Nauk SSSR *274*, 106 (1984); Chem. Abstr. *100*, 193516n (1984). [320] J. S. Ham, J. Chem. Phys. *21*, 756 (1953); G. Durocher and C. Sandorfy, J. Mol. Spectrosc. *20*, 410 (1966).

[321] A. B. Myers and R. R. Birge, J. Chem. Phys. *73*, 5314 (1980); ibid. *77*, 1075 (1982). [322] T. Abe, J. Chem. Phys. *77*, 1074 (1982); ibid. *83*, 1546 (1985); T. Abe and I. Iweibo, Bull. Chem. Soc. Jpn. *59*, 2381 (1986). [323] T. Shibuya, J. Chem. Phys. *78*, 5175 (1983); Bull. Chem. Soc. Jpn. *57*, 2991 (1984). [324] V. Bekárek *et al.*, Collect. Czech. Chem. Commun. *45*, 2063 (1980); ibid. *47*, 1060 (1982); ibid. *51*, 746 (1986). [325] Yu. T. Mazurenko, Opt. i Spektrosk. *55*, 471 (1983); Chem. Abstr. *100*, 42159x (1984). [326] W. Förster, F. Dietz, B. Mikutta, and C. Weiss, Z. Chem. *19*, 234 (1979). [327] A. Mehlhorn, J. Signal AM *6*, 211 (1978). [328] F. C. Grozema and P. Th. van Duijnen, J. Phys. Chem. A *102*, 7984 (1998). [329] J. Li, C. J. Cramer, and D. G. Truhlar, Int. J. Quantum Chem. *77*, 264 (2000). [330] A. F. Lagalante, R. J. Jacobson, and T. J. Bruno, J. Org. Chem. *61*, 6404 (1996).

[331] C. Haessner and H. Mustroph, Z. Chem. *26*, 170 (1986). [332] Y. Jinnouchi, M. Kohno, and A. Kuboyama, Bull. Chem. Soc. Jpn. *57*, 1147 (1984). [333] C. Benamou and L. Bellon, Bull. Soc. Chim. Fr. *1984*, I-321. [334] P. Mukerjee, C. Ramachandran, and R. A. Pyter, J. Phys. Chem. *86*, 3189 (1982). [335] M. C. R. Symons and A. S. Pena-Nuñez, J. Chem. Soc., Faraday Trans. I *81*, 2421 (1985). [336] A. Janowski, I. Turowska-Tyrk, and P. K. Wrona, J. Chem. Soc., Perkin Trans. II, 821 (1985). [337] W. D. Hinsberg, P. G. Schultz, and P. B. Dervan, J. Am. Chem. Soc. *104*, 766 (1982). [338] C. Ramachandran, R. A. Pyter, and P. Mukerjee, J. Phys. Chem. *86*, 3198 (1982). [339] G. van der Zwan and J. T. Hynes, J. Phys. Chem. *89*, 4181 (1985). [340] E. M. Kosower, J. Am. Chem. Soc. *107*, 1114 (1985); E. M. Kosower *et al.*, J. Phys. Chem. *87*, 2479 (1983); see also D. Huppert *et al.*, Chem. Phys. Lett. *144*, 15 (1988); ibid. *150*, 349 (1988).

[341] (a) G. Jones, W. R. Jackson, C. Choi, and W. R. Bergmark, J. Phys. Chem. *89*, 294 (1985); (b) G. Jones and M. A. Rahman, J. Phys. Chem. *98*, 13028 (1994). (c) L. Reynolds, J. A. Gardecki, S. J. V. Frankland, M. L. Horng, and M. Maroncelli, J. Phys. Chem. *100*, 10337 (1996); (d) A. Samanta and R. W. Fessenden, J. Phys. Chem. A *104*, 8577 (2000); (e) S. Nad and H. Pal, J. Phys. Chem. A *105*, 1097 (2001). [342] E. Lippert, W. Lüder, and H. Boos, Adv. Mol. Spectrosc., Proc. 4th Int. Meet., *1959* (1962), 443; E. Lippert *et al.*, J. Photochem. *17*, 237 (1981). [343] K. Rotkiewicz, K. H. Grellmann, and Z. R. Grabowski, Chem. Phys. Lett. *19*, 315 (1973); ibid. *21*, 212 (1973); Z. R. Grabowski *et al.*, Nouv. J. Chim. *3*, 443 (1979). [344] W. Rettig, Angew. Chem. *98*, 969 (1986); Angew. Chem. Int. Ed. Engl. *25*, 971 (1986). [345] J. W. Verhoeven *et al.*, J. Am. Chem. Soc. *106*, 6524 (1984); ibid. *112*, 1214 (1990); Recl. Trav. Chim. Pays-Bas *114*, 443 (1995); Eur. J. Org. Chem. 2373 (1998); ibid. 2391 (1999). [346] M. A. Kessler and O. S. Wolfbeis, Spectrochim. Acta, Part A, *47*, 187 (1991). [347] D. Banerjee, A. K. Laha, and S. Bagchi, J. Photochem. Photobiol. A *85*, 153 (1995). [348] M. Yasuda *et al.*, J. Photochem. Photobiol. A *121*, 149 (1999). [349] T. A. Fayed, J. Photochem. Photobiol. A *121*, 17 (1999). [350] H. Langhals, Z. Phys. Chem. (Frankfurt) N. F. *127*, 45 (1981); Tetrahedron *43*, 1771 (1987).

[351] P. Suppan, J. Chem. Soc., Faraday Trans. I *83*, 495 (1987). [352] T. Soujana, R. W. Fessenden, and A. Samanta, J. Phys. Chem. *100*, 3507 (1996). [353] D. E. Wetzler, C. Chesta, R.

Fernandez-Prini, and P. F. Aramendia, Pure Appl. Chem. *73*, 405 (2001). [354] G. Weber and F. J. Farris, Biochemistry *18*, 3075 (1979). [355] A. B. J. Parusel, W. Nowak, S. Grimme, and G. Köhler, J. Phys. Chem. A *102*, 7149 (1998). [356] A. Nakajima, Bull. Chem. Soc. Jpn. *44*, 3272 (1971); J. Mol. Spectrosc. *61*, 467 (1976). [357] D. C. Dong and M. A. Winnik, Photochem. Photobiol. *35*, 17 (1982); Chem. Abstr. *96*, 161966s (1982); Can. J. Chem. *62*, 2560 (1984). [358] K. W. Street and W. E. Acree, Analyst *111*, 1197 (1986). [359] D. S. Karpovich and G. J. Blanchard, J. Phys. Chem. *99*, 3951 (1995). [360] (a) K. Kalyanasundaram: *Photochemistry in Microheterogeneous Systems*, Academic Press, New York, 1987. (b) V. Ramamurthy (ed.): *Photochemistry in Organized and Constrained Media*, VCH, Weinheim, 1991.

[361] D. N. Kirk: *The Chiroptical Properties of Carbonyl Compounds*, Tetrahedron *42*, 777 (1986). [362] F. M. Menger and B. Boyer, J. Org. Chem. *49*, 1826 (1984). [363] S. Ohta *et al.*, Chem. Lett. (Tokyo) *1985*, 1331; Bull. Chem. Soc. Jpn. *59*, 1181 (1986). [364] M. Legrand: *Use of Solvent and Temperature Effects*, in F. Ciardelli and P. Salvadori (eds.): *Fundamental Aspects and Recent Developments in ORD and CD*, Heyden and Son, London, 1973, Chapter 4.2, p. 285ff.; and references cited therein. [365] B. Bosnich: *Induced Optical Activity*, in F. Ciardelli and P. Salvadori (eds.): *Fundamental Aspects and Recent Developments in ORD and CD*, Heyden and Son, London, 1973, Chapter 3.8, p. 254ff. [366] T. Hensel, J. Fruwert, M. Szombately, and H. Kriegsmann, Z. Phys. Chem. (Leipzig) *267*, 641, 650 (1986); T. Hensel, J. Fruwert, and K. Dathe, Collect. Czech. Chem. Commun. *52*, 22 (1987). [367] M. C. R. Symons, Chem. Soc. Rev. *12*, 1 (1983); Pure Appl. Chem. *58*, 1121 (1986). [368] M. C. R. Symons and G. Eaton, J. Chem. Soc., Faraday Trans. I *81*, 1963 (1985). [369] J. Manzur and G. Gonzalez, Z. Naturforsch., Part B *36*, 763 (1981). [370] (a) K. B. Patel, G. Eaton, and M. C. R. Symons, J. Chem. Soc., Faraday Trans. I *81*, 2775 (1985); (b) M. Meade, K. Hickey, Y. McCarthy, W. E. Waghorne, M. C. R. Symons, and P. P. Rastogi, J. Chem. Soc., Faraday Trans. *93*, 563 (1997).

[371] T. Yoshimura, Arch. Biochem. Biophys. *220*, 167 (1983). [372] M. C. R. Symons and G. Eaton, J. Chem. Soc., Faraday Trans. I *78*, 3033 (1982). [373] W. R. Fawcett and A. A. Kloss, J. Phys. Chem. *100*, 2019 (1996). [374] R. G. Makitra, Ya. N. Pirig, Ya. Tsikanchuk, and V. Ya. Zhukovskii, Ukr. Khim. Zh. *46*, 729 (1980); Chem. Abstr. *93*, 203859e (1980). [375] D. G. Cameron, S. C. Hsi, J. Umemura, and H. H. Mantsch, Can. J. Chem. *59*, 1357 (1981). [376] G. González and E. Clavijo, J. Chem. Soc., Perkin Trans. II, 1751 (1985). [377] R. G. Makitra, V. Ya. Zhukovskii, Ya. N. Pirig, and B. I. Chernyak, Zh. Prikl. Spektrosk. *36*, 962 (1982); Chem. Abstr. *97*, 82005v (1982). [378] C. Laurence, M. Berthelot, M. Lucon, M. Helbert, D. G. Morris, and J.-F. Gal, J. Chem. Soc., Perkin Trans. II, 705 (1984). [379] V. Bekárek and A. Mikulecká, Collect. Czech. Chem. Commun. *43*, 2879 (1978). [380] V. Bekárek, Acta Univ. Palacki. Olomuc., Fac. Rerum Nat. *61/65*, 87 (1980); Chem. Abstr. *94*, 147932e (1981).

[381] S. Kubota and Y. Ikegami, J. Phys. Chem. *82*, 2739 (1978). [382] D. M. Holton and D. Murphy, J. Chem. Soc., Faraday Trans. I *78*, 1223 (1982). [383] J. Spanget-Larsen, Theor. Chim. Acta *51*, 65 (1979). [384] H. G. Aurich, K. Hahn, K. Stork, and W. Weiss, Tetrahedron *33*, 969 (1977). [385] A. H. Reddoch and S. Konishi, J. Chem. Phys. *70*, 2121 (1979). [386] E. G. Janzen, G. A. Coulter, U. M. Oehler, and J. P. Bergsma, Can. J. Chem. *60*, 2725 (1982). [387] M. F. Ottaviani, G. Martini, and L. Nuti, Magn. Reson. Chem. *25*, 897 (1987). [388] R. Owenius, M. Engström, M. Lindgren, and M. Huber, J. Phys. Chem. A *105*, 10967 (2001). [389] O. W. Kolling, Anal. Chem. *55*, 143 (1983). [390] B. J. Tabner: *Organic Radicals in Solution*, in M. C. R. Symons (ed.): *Electron Spin Resonance*, Specialist Periodical Reports, Vol. 10A, Chapter 1, Royal Society of Chemistry, London, 1986; and previous volumes.

[391] K. Nakamura, Chem. Lett. (Tokyo) *1980*, 301; J. Am. Chem. Soc. *103*, 6973 (1981). [392] T. Abe, S. Tero-Kubota, and Y. Ikegami, J. Phys. Chem. *86*, 1358 (1982). [393] H. Block and S. M. Walker, Chem. Phys. Lett. *19*, 363 (1973). [394] P. Suppan, J. Chem. Soc., Faraday Trans. I *83*, 495 (1987). [395] P. Laszlo and A. Stockis, J. Am. Chem. Soc. *102*, 7818 (1980); B. Allard and E. Casadevall, Nouveau J. Chim. *9*, 565 (1985). [396] P. Laszlo: ^{23}*Na-NMR-Spektroskopie*. Nachr. Chem. Tech. Lab. *27*, 710 (1979). [397] B. Tiffon and J.-E. Dubois, Org. Magn. Reson. *11*, 295 (1978); B. Tiffon and B. Ancian, ibid. *16*, 247 (1981). [398] M. Jallali-Heravi and G. A. Webb, Org. Magn. Reson. *13*, 116 (1980). [399] W. Freitag and H.-J. Schneider, J. Chem. Soc., Perkin Trans. II, 1337 (1979). [400] V. Bekárek and V. Simánek, Collect. Czech. Chem. Commun. *46*, 1549 (1981).

[401] C. W. Fong and H. G. Grant, Z. Naturforsch., Part B *36*, 585 (1981). [402] R. Radeglia, R. Wolff, B. Bornowski, and S. Dähne, Z. Phys. Chem. (Leipzig) *261*, 502 (1980). [403] P. Plieninger and H. Baumgärtel, Liebigs Ann. Chem. *1983*, 860. [404] J. G. Dawber and R. A. Williams. J. Chem. Soc., Faraday Trans. I *82*, 3097 (1986). [405] J.-L. M. Abboud, A. Auhmani, H. Bitar, M. El Mouhtadi, J. Martin, and M. Rico, J. Am. Chem. Soc. *109*, 1332 (1987). [406] I. Ando and G. A. Webb, Org. Magn. Reson. *15*, 111 (1981). [407] M. Sugiura, N. Takao, and S. Ueji, Org. Magn. Reson. *18*, 128 (1982). [408] N. Kalyanam: *Application of Aromatic Solvent Induced Shifts in Organic Chemistry*, J. Chem. Educ. *60*, 635 (1983). [409] S. Ueji, M. Sugiura, and N. Takao, Tetrahedron Lett. *21*, 475 (1980). [410] J. D. Connolly and R. McCrindle, Chem. Ind. (London) *1965*, 379.

[411] D. H. Williams and N. S. Bhacca, Tetrahedron *21*, 2021 (1965). [412] K. Tori, I. Horibe, H. Sigemoto, and K. Umemoto, Tetrahedron Lett. *1975*, 2199. [413] K. Nikki and N. J. Nakagawa, Bull. Chem. Soc. Jpn. *51*, 3267 (1978); Org. Magn. Reson. *23*, 432 (1985). [414] M. Jutila, U. Edlund, D. Johnels, and E. Johansson, Acta Chem. Scand., Part B *38*, 131 (1984). [415] T. C. Morill (ed.): *Lanthanide Shifts Reagents in Stereochemical Analysis*, in A. P. Marchand (ed.): *Methods in Stereochemical Analysis*, Vol. 5, VCH Publishers, New York, 1986. [416] I. M. Dean and R. S. Varma, Tetrahedron *38*, 2069 (1982). [417] J. F. Bertrán and M. Rodríguez, Org. Magn. Reson. *21*, 1, 6 (1983). [418] J. C. Jochims: *Die magnetische Kernresonanz als Werkzeug des Stereochemikers*, in G. Hess (ed.): *Konstanzer Universitätsreden*, No. 56, Universitätsverlag, Konstanz, 1973, p. 26ff.; Chem. Abstr. *81*, 43982m (1974). [419] P. E. Hansen: *Carbon-Hydrogen Spin-Spin Coupling Constants*, Progr. NMR Spectrosc. *14*, 175 (1982). [420] I. Ando and S. Watanabe, Bull. Chem. Soc. Jpn. *53*, 1257 (1980); ibid. *54*, 1241 (1981).

[421] T. Schaefer, H. M. Hutton, and S. R. Salman, Can. J. Chem. *57*, 1877 (1979). [422] S. Watanabe and I. Ando, J. Mol. Struct. *84*, 77 (1982). [423] J. Burgess, J. G. Chambers, and R. I. Haines, Transition Met. Chem. *6*, 145 (1981). [424] W. Kaim, S. Ernst, and S. Kohlmann: *Farbige Komplexe: das Charge-Transfer-Phänomen*, Chemie in unserer Zeit *21*, 50 (1987). [425] D. M. Manuta and A. J. Lees, Inorg. Chem. *22*, 3825 (1983); ibid. *25*, 3212 (1986). [426] S. Ernst, Y. Kurth, and W. Kaim, J. Organomet. Chem. *302*, 211 (1986); W. Kaim and S. Kohlmann, Inorg. Chem. *25*, 3306 (1986). [427] R. E. Shepherd, M. F. Hoq, N. Hoblack, and C. R. Johnson, Inorg. Chem. *23*, 3249 (1984). [428] W. Kaim. S. Kohlmann, S. Ernst, B. Olbrich-Deussner, C. Bessenbacher, and A. Schulz, J. Organomet. Chem. *321*, 215 (1987). [429] J. Catalán, V. Lopez, P. Perez, R. Martin-Villamil, and J. G. Rodriguez, Liebigs Ann. 241 (1995). [430] G. L. Gaines, Angew. Chem. *99*, 346 (1987); Angew. Chem. Int. Ed. Engl. *26*, 341 (1987).

[431] A. M. Kjaer and J. Ulstrup, J. Am. Chem. Soc. *109*, 1934 (1987); ibid. *110*, 3874 (1988). [432] D. L. Richter-Egger, A. Tesfai, S. J. Flamm, and S. A. Tucker, J. Chem. Educ. *78*, 1375 (2001). [433] J.-J. La Clair, Angew. Chem. *110*, 339 (1998); Angew. Chem. Int. Ed. Engl. 37, 325 (1998). [434] M. V. Garcia and M. I. Redondo, Spectrochim. Acta, Part A *43*, 879 (1987). [435] (a) J. Tomasi and M. Persico, Chem. Rev. *94*, 2027 (1994); (b) J. Tomasi, B. Mennucci, and C. Cappelli: *Interactions in Solvents and Solutions*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 8, p. 419ff. [436] C. J. Cramer and D. G. Truhlar: *Implicit Solvation Models: Equilibria, Structure, Spectra, and Dynamics*, Chem. Rev. *99*, 2161 (1999). [437] M. Karelson: *Theoretical Treatment of Solvent Effects on Electronic and Vibrational Spectra of Compounds in Condensed Media*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 11.1, p. 639ff. [438] R. Kanski and C. J. Murray, Tetrahedron Lett. *34*, 2263 (1993). [439] M. C. Beard, G. M. Turner, and C. A. Schmuttenmaer, J. Phys. Chem. A *106*, 878 (2002). [440] D. A. Johnson, R. Shaw, and E. F. Silversmith, J. Chem. Educ. *71*, 517 (1994).

[441] M. F. Vitha, J. Chem. Educ. *78*, 370 (2001). [442] C. Machado and V. G. Machado, J. Chem. Educ. *78*, 649 (2001). [443] J. Kriwanek and R. Miller, Colloids Surfaces, Part A, *105*, 233 (1995). [444] N. O. Mchedlov-Petrossyan, G. P. Gorbenko, N. A. Vodolazkaya, V. I. Alekseeva, and L. P. Savvina, Functional Materials *7*, 138 (2000). [445] O. A. El Seoud *et al.*, Langmuir *16*, 35 (2000), ibid. *17*, 652 (2001). [446] R. A. McGill, M. S. Paley, and J. M. Harris, Macromolecules *25*, 3015

(1992). [447] Y. Imai and Y. Chujo, Macromolecules *33*, 3059 (2000). [448] C. Rottman, G. S. Grader, Y. DeHazan, and D. Avnir, Langmuir *12*, 5505 (1996). [449] D. J. Macquarrie *et al.*, New J. Chem. *23*, 725 (1999). [450] S. Spange *et al.*, Macromol. Rapid Commun. *21*, 643 (2000).

[451] K. Fischer and S. Spange, Macromol. Chem. Phys. *201*, 1922 (2000). [452] P. Blum, G. J. Mohr, K. Matern, J. Reichert, and U. E. Spichiger-Keller, Anal. Chim. Acta *432*, 269 (2001). [453] J. L. McHale, Acc. Chem. Res. *34*, 265 (2001). [454] S. Hogiu, J. Dreyer, M. Pfeiffer, K.-W. Brzezinka, and W. Werncke, J. Raman Spectrosc. *31*, 797 (2000). [455] P. F. Barbara *et al.*, J. Chem. Phys. *95*, 4188 (1991); J. Phys. Chem. *99*, 3554 (1995). [456] R. Scholz, M. Darwish, and M. Schreiber, J. Luminescence *76/77*, 404 (1998). [457] S. A. Kovalenko, N. Eilers-König, T. A. Senyushkina, and N. P. Ernsting, J. Phys. Chem. A *105*, 4834 (2001). [458] P. J. Rossky *et al.*, J. Phys. Chem. A *103*, 9432 (1999); ibid. *104*, 899 (2000); ibid. *105*, 558 (2001). [459] S. R. Mente and M. Maroncelli, J. Phys. Chem. B *103*, 7704 (1999). [460] J. Fabian, G. A. Rosquete, and L. A. Montero-Cabrera, J. Mol. Struct. (Theochem) *469*, 163 (1999).

[461] J. Lipinski and W. Bartkowiak, Chem. Phys. *245*, 263 (1999). [462] P. G. Jasien and L. L. Weber, J. Mol. Struct. (Theochem) *572*, 203 (2001). [463] S. H. Gallagher, R. S. Armstrong, P. A. Lay, and C. A. Reed, J. Phys. Chem. *99*, 5817 (1995). [464] J.-L. M. Abboud *et al.*, J. Phys. Org. Chem. *11*, 193 (1998). [465] F. Effenberger, F. Würthner, and F. Steybe, J. Org. Chem. *60*, 2082 (1995). [466] H. Hartmann, K. Eckert, and A. Schröder, Angew. Chem. *112*, 567 (2000); Angew. Chem. Int. Ed. Engl. *39*, 556 (2000). [467] A. A. Gorman, M. G. Hutchings, and P. D. Wood, J. Am. Chem. Soc. *118*, 8497 (1996). [468] W.-S. Sheu and P. J. Rossky, J. Am. Chem. Soc. *115*, 7729 (1993). [469] N. Ghoneim and P. Suppan, Spectrochim. Acta, Part A, *51*, 1043 (1995). [470] S. R. Mander *et al.*, Science *261*, 186 (1993); ibid. *265*, 632 (1994); J. Am. Chem. Soc. *116*, 10703 (1994).

[471] J. Gao and C. Alhambra, J. Am. Chem. Soc. *119*, 2962 (1997). [472] G. A. Rosquete and J. Fabian, J. Prakt. Chem. *341*, 631 (1999). [473] C. Machado, M. G. Nascimento, and M. C. Rezende, J. Chem. Soc., Perkin Trans. II, 2539 (1994). [474] L. da Silva, C. Machado, and M. C. Rezende, J. Chem. Soc., Perkin Trans. II, 483 (1995). [475] C. Aliaga, J. S. Galdames, and M. C. Rezende, J. Chem. Soc., Perkin Trans. II, 1055 (1997). [476] B. K. Mishra, M. Kuanar, A. Mishra, and G. B. Behera, Bull. Chem. Soc. Jpn. *69*, 2581 (1996). [477] E. Hammam and A. M. El-Nahas, J. Phys. Chem. A *102*, 9739 (1998). [478] P. Suppan: *Solvatochromic Shifts: The Influence of the Medium on the Energy of Electronic States*, J. Photochem. Photobiol. A *50*, 293 (1990). [479] H. Ågren and K. V. Mikkelsen: *Theory of Solvent Effects on Electronic Spectra*, J. Mol. Struct. (Theochem) *234*, 425 (1991). [480] A. Klamt, J. Phys. Chem. *100*, 3349 (1996).

[481] N. S. Bayliss and G. Wills-Johnson, Spectrochim. Acta, Part A, *24*, 551 (1968). [482] W. P. Hayes and C. J. Timmons, Spectrochim. Acta *21*, 529 (1965). [483] S. C. Beck and D. T. Cramb, J. Phys. Chem. B *104*, 2767 (2000). [484] L. Serrano-Andrés, M. Merchán, B. O. Roos, and R. Lindh, J. Am. Chem. Soc. *117*, 3189 (1995). [485] A. B. J. Parusel, W. Rettig, and W. Sudholt, J. Phys. Chem. A *106*, 804 (2002). [486] A. P. de Silva *et al.*, Chem. Rev. *97*, 1515 (1997). [487] T. Abe, Bull. Chem. Soc. Jpn. *61*, 3797 (1988). [488] E. Lippert, Z. Naturforsch., Part A, *10*, 541 (1955). [489] N. Mataga, Y. Kaifu, and M. Koizumi, Bull. Chem. Soc. Jpn. *29*, 465 (1956). [490] M. Kajtár, J. Kajtár, Zs. Majer, M. Zewdu, and M. Hollosi, Spectrochim. Acta, Part A, *48*, 87 (1992).

[491] J. Fidler, P. M. Rodger, and A. Rodger, J. Chem. Soc., Perkin Trans. II, 235 (1993). [492] C. Winther, Z. Phys. Chem. *60*, 590, 641 (1907); particularly Figs. 1 on pages 621 and 652. [493] H. G. Rule and A. McLean, J. Chem. Soc. 674 (1931). [494] O. E. Weigang, J. Chem. Phys. *41*, 1435 (1964). [495] G. Gottarelli, M. A. Osipov, and G. P. Spada, J. Phys. Chem. *95*, 3879 (1991). [496] J. B. F. N. Engberts, A. Perjéssy, and M. J. Blandamer, J. Chem. Soc., Faraday Trans. *89*, 4199 (1993). [497] M. M. Wohar, J. K. Seehra, and P. W. Jagodzinski, Spectrochim. Acta, Part A, *44*, 999 (1988). [498] F. Mao, D. R. Tyler, M. R. M. Bruce, A. E. Bruce, A. L. Rieger, and P. H. Rieger, J. Am. Chem. Soc. *114*, 6418 (1992). [499] O. W. Kolling, J. Phys. Chem. *100*, 16087 (1996). [500] Y. Sueishi, T. Isozaki, S. Yamamoto, and N. Nishimura, J. Phys. Org. Chem. *5*, 218 (1992).

[501] H. Takase, K. Morihashi, and O. Kikuchi, Bull. Chem. Soc. Jpn. *64*, 3511 (1991). [502] M. Witanowski, W. Sicinska, and G. A. Webb, Magn. Reson. Chem. *27*, 380 (1989). [503] M. Wita-

nowski, Z. Biedrzycka, W. Sicinska, and G. A. Webb, J. Chem. Soc., Perkin Trans. II, 533 (1997). [504] R. Streck and A. J. Barnes, Spectrochim. Acta, Part A *55*, 1049, 1059 (1999). [505] M. Luhmer and K. Bartik, J. Phys. Chem. A *101*, 5278 (1997). [506] V. Mäemets and I. Koppel, J. Chem. Res. (S) 480 (1994); J. Chem. Res. (M) 2864 (1994). [507] V. G. Malkin, O. L. Malkina, G. Steinebrunner, and H. Huber, Chem. Eur. J. *2*, 452 (1996). [508] B. Mennucci, J. M. Martinez, and J. Tomasi, J. Phys. Chem. A *105*, 7287 (2001). [509] A. D. Headley and N. M. Jackson, J. Phys. Org. Chem. *15*, 52 (2002). [510] M. H. Abraham, R. J. Abraham, P. Leonard, N. S. True, and C. Suarez, J. Chem. Soc., Perkin Trans. II, 463 (1991).

[511] C. Beeson, N. Pham, G. Shipps, and T. A. Dix, J. Am. Chem. Soc. *115*, 6803 (1993).

Chapter 7

[1] C. Reichardt: *Empirische Parameter der Lösungsmittelpolarität*, Angew. Chem. *77*, 30 (1965); Angew. Chem. Int. Ed. Engl. *4*, 29 (1965); C. Reichardt: *Empirical Parameters of Solvent Polarity as Linear Free-Energy Relationships*, Angew. Chem. *91*, 119 (1979); Angew. Chem. Int. Ed. Engl. *18*, 98 (1979). [2] C. Reichardt and K. Dimroth: *Lösungsmittel und empirische Parameter zur Charakterisierung ihrer Polarität*, Fortschr. Chem. Forsch. *11*, 1 (1968). [3] (a) C. Reichardt: *Lösungsmitteleffekte in der organischen Chemie*, 2nd ed., Verlag Chemie, Weinheim, 1973; (b) C. Reichardt: *Effets de solvant en chimie organique*, Flammarion, Paris, 1971 (translated into French by I. Tkatchenko); (c) C. Reichardt: *Youji Huaxue zhong de Rongji Jiaoying (Solvent Effects in Organic Chemistry)*, Huaxue Gongye Chubanshe (Publishing House of the Chemical Industry), Beijing/P.R. China, 1987 (translated into Chinese by P. Tang, Zh. Xie, Zh. Chen, and J. Lin); (d) C. Reichardt: *Rastvoriteli i Effekty Sredy v Organičeskoj Chimii (Solvent Effects in Organic Chemistry)*, Izdatel'stvo Mir, Moskva/Russia, 1991 (translated into Russian by A. A. Kiryuškin with V. S. Petrosyan as editor). [4] V. A. Palm: *Osnovy kolichestvennoi teorii organicheskikh reaktsii*, 2nd ed., Izdatel'stvo Khimiya, Leningrad, 1977; *Grundlagen der quantitativen Theorie organischer Reaktionen* (translated into German by G. Heublein), Akademie-Verlag, Berlin 1971. [5] E. M. Kosower: *An Introduction to Physical Organic Chemistry*, Wiley, New York, 1968, p. 293ff. [6] I. A. Koppel and V. A. Palm: *The Influence of the Solvent on Organic Reactivity*, in N. B. Chapman and J. Shorter (eds.): *Advances in Linear Free Energy Relationships*, Plenum Press, London, New York, 1972, Chapter 5, p. 203ff. [7] J. A. Hirsch: *Concepts in Theoretical Organic Chemistry*, Allyn and Bacon, Boston, 1974, Chapter 9, p. 207ff. [8] M. R. J. Dack: *The Influence of Solvent on Chemical Reactivity*, in M. R. J. Dack (ed.): *Solutions and Solubilities*, Vol. VIII, Part II, of A. Weissberger (ed.): *Techniques of Chemistry*, Wiley-Interscience, New York, 1976, p. 95ff. [9] V. Gutmann: *The Donor–Acceptor Approach to Molecular Interactions*, Plenum Publ. Corp., New York, 1978. [10] L. P. Hammett, J. Am. Chem. Soc. *59*, 96 (1937); Trans. Faraday Soc. *34*, 156 (1938); *cf.* also J. Shorter: *Die Hammett-Gleichung – und was daraus in fünfzig Jahren wurde*, Chemie in unserer Zeit *19*, 197 (1985).

[11] P. R. Wells: *Linear Free Energy Relationships*, Academic Press, London, New York, 1968. [12] N. B. Chapman and J. Shorter (eds.): *Advances in Linear Free Energy Relationships*, Plenum Press, London, New York, 1972. [13] N. B. Chapman and J. Shorter (eds.): *Correlation Analysis in Chemistry – Recent Advances*, Plenum Publ. Corp., New York, 1978. [14] J. Shorter: *Correlation Analysis in Organic Chemistry – An Introduction to Linear Free Energy Relationships*, Clarendon Press, Oxford, 1973. [15] J. Shorter: *Correlation Analysis of Organic Reactivity – With Particular Reference to Multiple Regression*, Research Studies Press, Chichester, 1982. [16] C. Reichardt and R. Müller, Liebigs Ann. Chem. *1976*, 1953. [17] J. N. Brønsted and K. J. Pedersen. Z. Phys. Chem. *108*, 185 (1924); Chem. Rev. *5*, 231 (1928). [18] J. E. Leffler and E. Grunwald: *Rates and Equilibria of Organic Reactions*, Wiley, New York, London, 1963. [19] A. R. Katritzky and R. D. Topsom: *Linear Free Energy Relationships and Optical Spectroscopy*, in N. B. Chapman and J. Shorter (eds.): *Advances in Linear Free Energy Relationships*, Plenum Press, London, New York, 1972, Chapter 3, p. 119ff. [20] M. T. Tribble and J. G. Traynham: *Linear Correlations of Substituent Effects in* ^{1}H, ^{19}F, *and* ^{13}C *Nuclear Magnetic Resonance Spectroscopy*, in N. B. Chapman and J.

Shorter (eds.): *Advances in Linear Free Energy Relationships*, Plenum Press, London, New York, 1972, Chapter 4, p. 143ff.

[21] E. M. Kosower, D. Hofmann, and K. Wallenfels, J. Am. Chem. Soc. *84*, 2755 (1962). [22] C. Reichardt and W. Grahn, Chem. Ber. *103*, 1072 (1970). [23] C. Reichardt and R. Müller, Liebigs Ann. Chem. *1976*, 1937. [24] K. H. Meyer, Ber. Dtsch. Chem. Ges. *47*, 826 (1914); ibid. *53*, 1410 (1920); ibid. *54*, 579 (1921). [25] E. L. Eliel, Pure Appl. Chem. *25*, 509 (1971); E. L. Eliel, Supplement to Pure Appl. Chem. 1971, p. 219ff.; E. L. Eliel and O. Hofer, J. Am. Chem. Soc. *95*, 8041 (1973). [26] V. Gutmann and E. Wychera, Inorg. Nucl. Chem. Lett. *2*, 257 (1966); V. Gutmann, Coord. Chem. Rev. *2*, 239 (1967); ibid. *18*, 225 (1976); V. Gutmann and A. Scherhaufer, Monatsh. Chem. *99*, 335 (1968); V. Gutmann, Chimia *23*, 285 (1969); V. Gutmann, Electrochimica Acta *21*, 661 (1976). [27] V. Gutmann: *Coordination Chemistry in Non-Aqueous Solvents*, Springer, Wien, New York, 1968; V. Gutmann: *Chemische Funktionslehre*, Springer, Wien, New York, 1971; V. Gutmann: *The Donor-Acceptor Approach to Molecular Interactions*, Plenum, New York, 1978. [28] G. Olofson and I. Olofson, J. Am. Chem. Soc. *95*, 7231 (1973). [29] M. S. Greenberg, R. L. Bodner, and A. I. Popov, J. Phys. Chem. *77*, 2449 (1973). [30] V. Gutmann, Pure Appl. Chem. *27*, 73 (1971); Fortschr. Chem. Forsch. *27*, 59 (1972); Structure and Bonding *15*, 141 (1973); Chimia *31*, 1 (1977); CHEMTECH *7*, 255 (1977); Pure Appl. Chem. *51*, 2197 (1979); A. J. Parker, U. Mayer, R. Schmid, and V. Gutmann, J. Org. Chem. *73*, 1843 (1978).

[31] (a) J. Catalán, J. Gómez, A. Couto, and J. Laynez, J. Am. Chem. Soc. *112*, 1678 (1990); (b) J. Catalán, A. Couto, J. Gómez, J. L. Saiz, and J. Laynez, J. Chem. Soc., Perkin Trans. II, 1181 (1992). [32] R. S. Drago and B. B. Wayland, J. Am. Chem. Soc. *87*, 3571 (1965); R. S. Drago, J. Chem. Educ. *51*, 300 (1974). [32a] J. Llor and M. Cortijo, J. Chem. Soc., Perkin Trans. II, 1111 (1977). [33] E. S. Gould: *Mechanism and Structure in Organic Chemistry*, Henry Holt and Co., New York, 1959, p. 299ff.; *Mechanismus und Struktur in der organischen Chemie* (translated into German by G. Koch), 2nd ed., Verlag Chemie, Weinheim, 1962, p. 353ff. [34] K. Schwetlick: *Kinetische Methoden zur Untersuchung von Reaktionsmechanismen*, Deutscher Verlag der Wissenschaften, Berlin, 1971, p. 162ff. [35] E. Grunwald and S. Winstein, J. Am. Chem. Soc. *70*, 846 (1948). [36] S. Winstein, E. Grunwald, and H. W. Jones, J. Am. Chem. Soc. *73*, 2700 (1951); A. H. Fainberg and S. Winstein, ibid. *78*, 2770 (1956); ibid. *79*, 1597, 1602, 1608 (1957); S. Winstein, A. H. Fainberg, and E. Grunwald, ibid. *79*, 4146, 5937 (1957). [37] S. G. Smith, A. H. Fainberg, and S. Winstein, J. Am. Chem. Soc. *83*, 618 (1961). [38] I. A. Koppel and V. A. Palm, Reakts. Sposobnost Org. Soedin. *6*, 504 (1969); Chem. Abstr. *72*, 6732y (1970). [39] D. J. Raber, R. C. Bingham, J. M. Harris, J. L. Fry, and P. v. R. Schleyer, J. Am. Chem. Soc. *92*, 5977 (1970). [40] D. N. Kevill, K. C. Kolwyck, and F. L. Weitl, J. Am. Chem. Soc. *92*, 7300 (1970).

[41] T. W. Bentley and P. v. R. Schleyer, J. Am. Chem. Soc. *98*, 7658 (1976); Adv. Phys. Org. Chem. *14*, 2 (1977). [42] S. Winstein, E. Grunwald, and H. W. Jones, J. Am. Chem. Soc. *73*, 2700 (1951); S. Winstein, A. H. Fainberg, and E. Grunwald, ibid. *79*, 4146 (1957). [43] P. E. Peterson and F. J. Waller, J. Am Chem. Soc. *94*, 991 (1972); P. E. Peterson, D. W. Vidrine, F. J. Waller, P. M. Henrichs, S. Magaha, and B. Stevens, ibid. *99*, 7968 (1977). [44] T. W. Bentley, F. L. Schadt, and P. v. R. Schleyer, J. Am. Chem. Soc. *94*, 992 (1972); ibid. *98*, 7667 (1976). [45] C. G. Swain, R. B. Mosely, and D. E. Bown, J. Am. Chem. Soc. *77*, 3731 (1955). [46] A. Streitwieser jr.: *Solvolytic Displacement Reactions*, McGraw-Hill, New York, 1962. [47] P. D. Bartlett, J. Am. Chem. Soc. *94*, 2161 (1972). [48] Y. Drougard and D. Decroocq, Bull. Soc. Chim. Fr. *1969*, 2972. [49] C. Lassau and J. C. Jungers, Bull. Soc. Chim. Fr. *1968*, 2678. [50] J. C. Jungers, L. Sajus, I. de Aguirre, and D. Decroocq: *L'Analyse cinétique de la transformation chimique*, Editions Technip, Paris, 1967/68, Vol. II, p. 627, 771, and 1220.

[51] M. Gielen and J. Nasielski, J. Organomet. Chem. *1*, 173 (1963); ibid. *7*, 273 (1967). [52] J. A. Berson, Z. Hamlet, and W. A. Mueller, J. Am. Chem. Soc. *84*, 297 (1962). [53] B. Blankenburg, H. Fiedler, M. Hampel, H. G. Hauthal, G. Just, K. Kahlert, J. Korn, K.-H. Müller, W. Pritzkow, Y. Reinhold, M. Röllig, E. Sauer, D. Schnurpfeil, and G. Zimmermann, J. Prakt. Chem. *316*, 804 (1974). [54] L. G. S. Brooker, G. H. Keyes, and D. W. Heseltine, J. Am. Chem. Soc. *73*, 5350 (1951). [55] E. M. Kosower, J. Am. Chem. Soc. *80*, 3253 (1958); J. Chim. Phys. *61*, 230 (1964); E. M. Kosower and M. Mohammad, J. Am. Chem. Soc. *90*, 3271 (1968); ibid. *93*, 2713 (1971); J.

Phys. Chem. *74*, 1153 (1970). [56] J. S. Gowland and G. H. Schmid, Can. J. Chem. *47*, 2953 (1969): *Z* values for water/dimethyl sulfoxide and water/ethanol mixtures. [57] M. Mohammad and E. M. Kosower, J. Phys. Chem. *74*, 1153 (1970). [58] J. H. Fendler and L.-J. Liu, J. Am. Chem. Soc. *97*, 999 (1975). [59] A. Ray, J. Am. Chem. Soc. *93*, 7146 (1971). [60] K. Tamura and T. Imoto, Chem. Lett. (Tokyo) *1973*, 1251.

[61] E. M. Kosower, M. Ito, and P.-K. Huang, Trans. Faraday Soc. *57*, 1662 (1961). [62] E. M. Kosower, J. Am. Chem. Soc. *80*, 3261, 3267 (1958). [63] E. M. Kosower, J. Org. Chem. *29*, 956 (1964). [64] N. S. Isaacs: *Experiments in Physical Organic Chemistry*, MacMillan Company, London, 1969, p. 161ff. [65] S. Brownstein, Can. J. Chem. *38*, 1590 (1960). [66] K. Dimroth, C. Reichardt, T. Siepmann, and F. Bohlmann, Liebigs Ann. Chem. *661*, 1 (1963); C. Reichardt, ibid. *752*, 64 (1971). [67] K. Dimroth and C. Reichardt, Liebigs Ann. Chem. *727*, 93 (1969). [68] K. Dimroth and C. Reichardt, Z. Anal. Chem. *215*, 344 (1966); Z. B. Maksimović, C. Reichardt, and A. Spirić, ibid. *270*, 100 (1974). [69] T. G. Beaumont and K. M. C. Davis, J. Chem. Soc., Part B *1968*, 1010 (dichloromethane/acetonitrile mixtures). [70] W. Koehler, P. Froelich, and R. Radeglia, Z. Phys. Chem. (Leipzig) *242*, 220 (1969) (acetone/water, 1,4-dioxane/water and tetrahydrofuran/water mixtures).

[71] K. Tamura, Y. Ogo, and T. Imoto, Bull. Chem. Soc. Jpn. *46*, 2988 (1973) (nitrobenzene/benzene mixtures). [72] E. M. Kosower, H. Dodiuk, K. Tanizawa, M. Ottolenghi, and N. Orbach, J. Am. Chem. Soc. *97*, 2167 (1975) (1,4-dioxane/water mixtures). [73] K. Dimroth, C. Reichardt, and A. Schweig, Liebigs Ann. Chem. *669*, 95 (1963). [74] K. Tamura, Y. Ogo, and T. Imoto, Chem. Lett. (Tokyo) *1973*, 625; K. Tamura and T. Imoto, Bull. Chem. Soc. Jpn. *48*, 369 (1975). [75] (a) R. Allmann, Z. Kristallogr. *128*, 115 (1969); (b) K. Stadnicka, P. Milart, A. Olech, P. K. Olszewski, J. Mol. Struct. *604*, 9 (2002). [76] H. C. Andersen: *Complete Fairy-Tales and Stories* (translated by E. Haugard), Gollantz, London, 1974. [77] L. G. S. Brooker, A. C. Craig, D. W. Heseltine, P. W. Jenkins, and L. L. Lincoln, J. Am. Chem. Soc. *87*, 2443 (1965). [78] S. Dähne, F. Schob, K.-D. Nolte, and R. Radeglia, Ukr. Khim. Zh. (Russ. Ed.) *41*, 1170 (1975); Chem. Abstr. *84*, 43086j (1976). [79] J.-E. Dubois, E. Goetz, and A. Bienvenüe, Spectrochim. Acta *20*, 1815 (1964); J.-E. Dubois and A. Bienvenüe, J. Chim. Phys. *65*, 1259 (1968). [80] I. A. Zhmyreva, V. V. Zelinskii, V. P. Kolobkov, and N. D. Krasnitskaya, Dokl. Akad. Nauk SSSR, Ser. Khim. *129*, 1089 (1959); Chem. Abstr. *55*, 26658e (1961).

[81] D. Walther, J. Prakt. Chem. *316*, 604 (1974). [82] D. M. Manuta and A. J. Lees, Inorg. Chem. *22*, 3825 (1983); ibid. *25*, 3212 (1986). [83] (a) W. Kaim, S. Kohlmann, S. Ernst, B. Olbrich-Deussner, C. Bessenbacher, and A. Schulz, J. Organomet. Chem. *321*, 215 (1987); (b) W. Kaim, B. Olbrich-Deussner, and T. Roth, Organometallics *10*, 410 (1991). [84] M. J. Kamlet and R. W. Taft, J. Am. Chem. Soc. *98*, 377, 2886 (1976). [84a] M. J. Kamlet, J. L. Abboud, and R. W. Taft, J. Am. Chem. Soc. *99*, 6027 (1977); ibid. *99*, 8325 (1977). [85] A. Allerhand and P. v. R. Schleyer, J. Am. Chem. Soc. *85*, 371 (1963); ibid. *86*, 5709 (1964). [86] T. Kagiya, Y. Sumida, and T. Inoue, Bull. Chem. Soc. Jpn. *41*, 767 (1968). [87] A. G. Burden, G. Collier, and J. Shorter, J. Chem. Soc., Perkin Trans. II *1976*, 1627. [88] I. A. Koppel and A. I. Paju, Reakts. Sposobnost. Org. Soedin. *11*, 121 (1974); Chem. Abstr. *82*, 42805q (1975). [89] B. R. Knauer and J. J. Napier, J. Am. Chem. Soc. *98*, 4395 (1976). [90] R. W. Taft, G. B. Klingensmith, E. Price, and I. R. Fox, Preprints of papers presented at the Symposium on LFE Correlations, Durham/N.C., October 1964, p. 265; R. T. C. Brownlee, S. K. Dayal, J. L. Lyle, and R. W. Taft, J. Am. Chem. Soc. *94*, 7208 (1972); and references cited therein.

[91] U. Mayer, V. Gutmann, and W. Gerger, Monatsh. Chem. *106*, 1235 (1975); ibid. *108*, 489, 757 (1977); V. Gutmann, Electrochimica Acta *21*, 661 (1976); V. Gutmann, CHEMTECH *7*, 255 (1977); Chem. Abstr. *86*, 178193d (1977); A. J. Parker, U. Mayer, R. Schmid, and V. Gutmann, J. Org. Chem. *43*, 1843 (1978). [92] C. D. Ritchie: *Interactions in Dipolar Aprotic Solvents*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York, London, 1969, Vol. 1, Table 4-13 on p. 281. [93] J. H. Hildebrand, J. M. Prausnitz, and R. L. Scott: *Regular and Related Solutions*, Van Nostrand-Reinhold, Princeton, 1970. [94] A. F. M. Barton: *Handbook of Solubility Parameters and other Cohesion Parameters*, CRC Press, Boca Raton/Florida, 1983. [95] H. Burrell: *Solubility Parameters*, in J. Brandrup and E. H. Immergut (eds.): *Polymer Handbook*,

2nd ed., Wiley-Interscience, New York, 1975, p. IV-337ff. [96] H. F. Herbrandson and F. R. Neufeld, J. Org. Chem. *31*, 1140 (1966). [97] J. E. Gordon, J. Phys. Chem. *70*, 2413 (1966). [98] L. Rohrschneider: *The Polarity of Stationary Liquid Phases in Gas-Chromatography*, in J. C. Giddings and R. A. Keller (eds.): *Advances in Chromatography*, Dekker, New York, 1967, Vol. 4, p. 333ff.; L. Rohrschneider: *Der Lösungsmitteleinfluß auf die gas-chromatographische Retention gelöster Stoffe*, Fortschr. Chem. Forsch. *11*, 146 (1968). [99] R. V. Golovnaya and Yu. N. Arsen'ev: *System of Retention Indices and Its Physicochemical Applications*, Usp. Khim. *42*, 2221 (1973); Russ. Chem. Rev. *42*, 1034 (1973). [100] E. Kováts, Z. Anal. Chem. *181*, 351 (1961); Adv. Chromatogr. *1*, 229 (1965).

[101] E. Kováts and P. B. Weiß, Ber. Bunsenges. Phys. Chem. *69*, 812 (1965). [102] L. Rohrschneider, J. Chromatogr. *22*, 6 (1966); ibid. *38*, 383 (1969); Z. Anal. Chem. *236*, 149 (1968). [103] W. O. McReynolds, J. Chromatogr. Sci. *8*, 685 (1970). [104] L. Rohrschneider, Anal. Chem. *45*, 1241 (1973). [105] L. R. Snyder: *The Role of the Mobile Phase in Liquid Chromatography*, in J. J. Kirkland (ed.): *Modern Practice of Liquid Chromatography*, Wiley-Interscience, New York, 1971, Chapter 4, p. 125. [106] L. R. Snyder and J. J. Kirkland; *Introduction to Modern Liquid Chromatography*, Wiley-Interscience, New York, 1974, p. 215ff. and p. 444ff. [107] L. R. Snyder, J. Chromatogr. Sci. *16*, 223 (1978). [108] P. H. Vandewyer, J. Hoefnagels, and G. Smets, Tetrahedron *25*, 3251 (1969). [109] P. Bickart, F. W. Carson, J. Jacobus, E. G. Miller, and K. Mislow, J. Am. Chem. Soc. *90*, 4869 (1968); R. Tang and K. Mislow, ibid. *92*, 2100 (1970). [110] M. H. Abraham and P. L. Grellier, J. Chem. Soc., Perkin Trans. II, 1735 (1976).

[111] F. W. Fowler, A. R. Katritzky, and R. J. D. Rutherford, J. Chem. Soc., Part B *1971*, 460. [112] I. A. Koppel and V. A. Palm, Reakts. Sposobnost Org. Soedin. *8*, 291 (1971); Organic Reactivity (Tartu) *8*, 296 (1971); Chem. Abstr. *76*, 28418k (1972). [113] T. M. Krygowski and W. R. Fawcett, J. Am. Chem. Soc. *97*, 2143 (1975); Aust. J. Chem. *28*, 2115 (1975); Can. J. Chem. *54*, 3283 (1976). [114] R. C. Dougherty, Tetrahedron Lett. *1975*, 385. [115] I. A. Koppel and A. Paju, Reakts. Sposobnost Org. Soedin. *11*, 139 (1974); Organic Reactivity (Tartu) *11*, 137 (1974); Chem. Abstr. *82*, 42806r (1975). [116] A. G. Burden, N. B. Chapman, H. F. Duggua, and J. Shorter, J. Chem. Soc., Perkin Trans II, 296 (1978); M. H. Aslam, G. Collier, and J. Shorter, ibid. 1572 (1981). [117] N. S. Isaacs and E. Rannala, J. Chem. Soc., Perkin Trans. II, 902 (1974). [118] W. G. Jackson, G. A. Lawrence, P. A. Lay, and A. M. Sargeson, Aust. J. Chem. *35*, 1561 (1982). [119] S. Glikberg and Y. Marcus, J. Sol. Chem. *12*, 255 (1983). [120] Z. Zhong-yuan, S. Cheng-E, and H. Te-kang, J. Chem. Soc., Perkin Trans. II, 929 (1985).

[121] M. Claessens, L. Palombini, M.-L. Stien, and J. Reisse, Nouv. J. Chim. *6*, 595 (1982). [122] T. R. Griffiths and D. C. Pugh, Coord. Chem. Rev. *29*, 129 (1979). [123] J.-L. M. Abboud, M. J. Kamlet, and R. W. Taft, Progr. Phys. Org. Chem. *13*, 485 (1981). [124] C. Reichardt: *Empirical Parameters of Solvent Polarity and Chemical Reactivity*, in H. Ratajczak and W. J. Orville-Thomas (eds.): *Molecular Interactions*, Vol. 3, p. 241ff., Wiley, Chichester, 1982. [125] (a) O. Exner: *Correlation Analysis of Chemical Data*, Plenum Press, New York, SNTL Publishers, Prague, 1988; (b) O. Exner: *Physicochemical Preconditions of Linear Free-Energy Relationships*, Progr. Phys. Org. Chem. *18*, 129 (1990). [126] M. Sjöström and S. Wold: *Linear Free Energy Relationships. Local Empirical Rules or Fundamental Laws of Chemistry*? Acta Chem. Scand., Part B *35*, 537 (1981); ibid. *40*, 270 (1986). [127] M. J. Kamlet and R. W. Taft: *Linear Solvation Energy Relationships. Local Empirical Rules – or Fundamental Laws of Chemistry? A Reply to the Chemometricians*, Acta Chem. Scand., Part B *39*, 611 (1985); ibid. *40*, 619 (1986); M. J. Kamlet, R. M. Doherty, G. R. Famini, and R. W. Taft, ibid. *41*, 589 (1987). [128] J.-C. Bollinger, G. Yvernault, and T. Yvernault, Thermochim. Acta *60*, 137 (1983). [129] H. D. Schädler, M. Riemer, and W. Schroth, Z. Chem. *24*, 407 (1984). [130] M. C. Day, J. H. Medley, and N. Ahmad, Can. J. Chem. *61*, 1719 (1983).

[131] S. Hahn, W. M. Miller, R. N. Lichtenthaler, and J. M. Prausnitz, J. Sol. Chem. *14*, 129 (1985). [132] R. W. Soukop, Chemie in unserer Zeit *17*, 129, 163 (1983); R. W. Soukop and R. Schmid, J. Chem. Educ. *62*, 459 (1985). [133] Y. Marcus, J. Sol. Chem. *13*, 599 (1984); *cf*. also Y. Marcus: *Ion Solvation*, Chapter 6, Wiley, Chichester, 1985. [134] R. Schmid, J. Sol. Chem. *12*, 135 (1983). [135] U. Mayer and V. Gutmann, Structure and Bonding *12*, 113 (1972); *cf*. p. 125. [136] A. I. Popov, Pure Appl. Chem. *41*, 275 (1975); and references cited therein. [137] T. Ogata, T.

Fujisawa, N. Tanaka, and H. Yokoi, Bull. Chem. Soc. Jpn. *49*, 2759 (1976). [138] P.-C. Maria and J.-F. Gal, J. Phys. Chem. *89*, 1296 (1985); P.-C. Maria, J.-F. Gal, J. de Franceschi, and E. Fargin, J. Am. Chem. Soc. *109*, 483 (1987). [139] A. Sabatino, G. La Manna, and L. Paoloni, J. Phys. Chem. *84*, 2641 (1980). [140] K. Hiraoka, Bull. Chem. Soc. Jpn. *59*, 2571 (1986).

[141] R. S. Drago, Coord. Chem. Rev. *33*, 251 (1980); Pure Appl. Chem. *52*, 2261 (1980). [142] L. Elégant, G. Fratini, J.-F. Gal, and P.-C. Maria, J. Calorim. Anal. Therm. *11*, 3/21/1–3/21/9 (1980); Chem. Abstr. *95*, 121811k (1981). [143] M. Berthelot, J.-F. Gal, M. Helbert, C. Laurence, and P.-C. Maria, J. Chim. Phys. *82*, 427 (1985). [144] I. Persson, Pure Appl. Chem. *58*, 1153 (1986). [145] (a) A. Leo, C. Hansch, and D. Elkins, Chem. Rev. *71*, 525 (1971); (b) C. Hansch, Acc. Chem. Res. *26*, 147 (1993). [146] C. Hansch, A. Leo, and D. Hoekman: *Exploring QSAR: (a) Fundamentals and Applications in Chemistry and Biology; (b) Hydrophobic, Electronic, and Steric Constants*, American Chemical Society, Washington D.C., 1995. [147] H. Kubinyi: *QSAR: Hansch Analysis and Related Approaches*, VCH, Weinheim, 1993. [148] A. J. Dallas and P. W. Carr, J. Chem. Soc., Perkin Trans. II, 2155 (1992). [149] R. W. Taft, M. H. Abraham, G. F. Famini, R. M. Doherty, J.-L. M. Abboud, and M. J. Kamlet, J. Pharm. Sci. *74*, 807 (1985). [150] F. M. Menger and U. V. Venkataram, J. Am. Chem. Soc. *108*, 2980 (1986).

[151] D. Fărcasiu, J. Jähme, and C. Rüchardt, J. Am. Chem. Soc. *107*, 5717 (1985). [152] M. H. Abraham, R. W. Taft, and M. J. Kamlet, J. Org. Chem. *46*, 3053 (1981). [153] T. W. Bentley and G. E. Carter, J. Am. Chem. Soc. *104*, 5741 (1982), and references cited therein. [154] T. W. Bentley, C. T. Bowen, D. H. Morten, and P. v. R. Schleyer, J. Am. Chem. Soc. *103*, 5466 (1981). [155] D. N. Kevill, W. A. Kamil, and S. W. Anderson, Tetrahedron Lett. *1982*, 4635. [156] C. A. Bunton, M. M. Mhala, and J. R. Moffatt, J. Org. Chem. *49*, 3639 (1984). [157] S. P. McManus and S. E. Zutaut, Tetrahedron Lett. *25*, 2859 (1984). [158] T. W. Bentley and G. E. Carter, J. Org. Chem. *48*, 579 (1983). [159] T. W. Bentley, S. Jurczyk, K. Roberts, and D. J. Williams, J. Chem. Soc., Perkin Trans. II, 293 (1987). [160] D. N. Kevill and G. M. L. Lin, J. Am. Chem. Soc. *101*, 3916 (1979); D. N. Kevill *et al.*: *Studies of Displacement Reactions Using* $R—X^{\oplus}$ *Substrates*, in M. Kobayashi (ed.): *Physical Organic Chemistry 1986*, Elsevier, Amsterdam, 1987, p. 311ff.

[161] T. W. Bentley, G. E. Carter, and K. Roberts, J. Org. Chem. *49*, 5183 (1984). [162] D. N. Kevill, M. S. Bahari, and S. W. Anderson, J. Am. Chem. Soc. *106*, 2895 (1984). [163] D. N. Kevill and S. W. Anderson, J. Am. Chem. Soc. *108*, 1579 (1986). [164] D. N. Kevill and S. W. Anderson, J. Org. Chem. *50*, 3330 (1985). [165] X. Creary and S. R. McDonald, J. Org. Chem. *50*, 474 (1985). [166] T. W. Bentley and K. Roberts, J. Org. Chem. *50*, 4821 (1985). [167] T. W. Bentley and K. Roberts, J. Org. Chem. *50*, 5852 (1985). [168] F. L. Schadt, C. J. Lancelot, and P. v. R. Schleyer, J. Am. Chem. Soc. *100*, 228 (1978). [169] T. Oshima, S. Arikata, and T. Nagai, J. Chem. Res. *1981* (S) 204; (M) 2518. [170] T. Oshima and T. Nagai, Bull. Chem. Soc. Jpn. *55*, 555 (1982); Tetrahedron Lett. *26*, 4785 (1985).

[171] J. W. Larsen, A. G. Edwards, and P. Dobi, J. Am. Chem. Soc. *102*, 6780 (1980). [172] T. R. Griffiths and D. C. Pugh, J. Sol. Chem. *8*, 247 (1979); Coord. Chem. Rev. *29*, 129 (1979). [173] P. Štrop, F. Mikeš, and J. Kálal, J. Phys. Chem. *80*, 694 (1976); F. Mikeš, J. Labský, P. Štrop, and D. Vyprachticky: *On the Microenvironment of Soluble and Crosslinked Polymers*, Sb. Vys. Sk. Chem.-Technol. Praze [Oddil] S *1984*, S11, 83–144; Chem. Abstr. *102*, 167386s (1985). [174] C. Reichardt and E. Harbusch-Görnert, Liebigs Ann. Chem. *1983*, 721. [175] C. Laurence, P. Nicolet, and C. Reichardt, Bull. Soc. Chim. Fr. *1987*, 125. [176] C. Laurence, P. Nicolet, M. Lucon, and C. Reichardt, Bull. Soc. Chim. Fr. *1987*, 1001. [177] Y. Marcus: *Ion Solvation*, Chapter 7, Table 7.3 on page 188, Wiley, New York, 1985. [178] T. M. Krygowski, P. K. Wrona, U. Zielkowska, and C. Reichardt, Tetrahedron *41*, 4519 (1985). [179] M. De Vijlder, Bull. Soc. Chim. Belg. *91*, 947 (1982). [180] J. R. Haak and J. B. F. N. Engberts, Recl. Trav. Chim. Pays-Bas *105*, 307 (1986).

[181] B. P. Johnson, M. G. Khaledi, and J. G. Dorsey, J. Chromatogr. *384*, 221 (1987). [182] W. S. Hammack, D. N. Hendrickson, and H. G. Drickamer, J. Phys. Chem. *93*, 3483 (1989). [183] S. Balakrishnan and A. J. Easteal, Aust. J. Chem. *34*, 933, 943 (1981). [184] H. Langhals, Nouv. J. Chim. *5*, 97, 511 (1981); ibid. *6*, 265 (1982). [185] I. A. Koppel and J. B. Koppel, Organic Reactivity (Tartu) *20*, 547 (1983); Chem. Abstr. *101*, 110274d (1984). [186] H. Elias, G. Gumbel, S. Neitzel, and H. Volz, Z. Anal. Chem. *306*, 240 (1981). [187] I. A. Koppel and J. B. Koppel,

Organic Reactivity (Tartu) *20*, 523 (1983); Chem. Abstr. *101*, 110180v (1984). [188] V. Bekárek and T. Nevěčná, Collect. Czech. Chem. Commun. *50*, 1928 (1985); ibid. *51*, 1942, 2071 (1986). [189] J. Hicks, M. Vandersall, Z. Babrogic, and K. B. Eisenthal, Chem. Phys. Lett. *116*, 18 (1985). [190] P. Nagy and R. Herzfeld, Acta Universitatis Szegediensis, Acta Physica et Chimica *31*, 736 (1985); Chem. Abstr. *107*, 22905d (1987).

[191] H. Langhals, Tetrahedron Lett. *27*, 339 (1986); Z. Phys. Chem. (Leipzig) *286*, 91 (1987). [192] (a) H. Langhals: *Polarität binärer Flüssigkeitsgemische*, Angew. Chem. *94*, 739 (1982); Angew. Chem. Int. Ed. Engl. *21*, 724 (1982); (b) H. Langhals: *Description of Properties of Binary Solvent Mixtures*, in R. I. Zalewski, T. M. Krygowski, and J. Shorter (eds.): *Similarity Models in Organic Chemistry, Biochemistry, and Related Fields*, Elsevier, Amsterdam, 1991, Chapter 6, p. 283ff. [193] M. Chastrette and J. Carretto, Tetrahedron *38*, 1615 (1982); Can. J. Chem. *63*, 3492 (1985). [194] V. Bekárek and J. Juřina, Collect. Czech. Chem. Commun. *47*, 1060 (1982). [195] H. Langhals, Angew. Chem. *94*, 452 (1982); Angew. Chem. Int. Ed. Engl. *21*, 432 (1982); Angew. Chem. Suppl. *1982*, 1138. [196] H. Langhals, Chem. Ber. *114*, 2907 (1981). [197] M. C. Rezende, Tetrahedron *44*, 3513 (1988). [198] S. Kumoi, K. Oyama, T. Yano, H. Kobayashi, and K. Ueno, Talanta *17*, 319 (1970). [199] H. Langhals, Z. Anal. Chem. *305*, 26 (1981); ibid. *308*, 441 (1981). [200] C. Reichardt: *Solvent Scales and Chemical Reactivity*, in A. D. Buckingham, E. Lippert, and S. Bratos (eds.): *Organic Liquids: Structure, Dynamics, and Chemical Properties*, Chapter 16, p. 269ff., Wiley, Chichester, 1978.

[201] C. Reichardt: *Solvent Effects on Chemical Reactivity*, Pure Appl. Chem. *54*, 1867 (1982). [202] C. Han: E_T *Empirical Parameters of Solvent Polarity and Their Applications*, Huaxue Tongbao [Chemistry, Beijing] *1985*, 40; Chem. Abstr. *104*, 147904x (1986). [203] B. P. Johnson, B. Gabrielsen, M. Matulenko, J. G. Dorsey, and C. Reichardt: *Solvatochromic Solvent Polarity Measurements in Analytical Chemistry: Synthesis and Applications of ET-30*, Anal. Lett. *19*, 939 (1986). [204] M. S. Tunuli, M. A. Rauf, and Farhataziz, J. Photochem. *24*, 411 (1984). [205] F. Ibáñez, J. Photochem. *30*, 245 (1985). [206] Y. Marcus, E. Pross, and J. Hormadaly, J. Phys. Chem. *84*, 2708 (1980). [207] R. Schmid and V. N. Sapunov: *Non-Formal Kinetics in Search for Chemical Reaction Pathways*, Verlag Chemie, Weinheim, 1982; R. Schmid, J. Sol. Chem. *12*, 135 (1983). [208] C. Reichardt, E. Harbusch, and R. Müller: *Pyridinium N-phenoxide Betaine Dyes as Solvent Polarity Indicators, Some New Findings*, in I. Bertini, L. Lunazzi, and A. Dei (eds.): *Advances in Solution Chemistry*, Plenum Press, New York, 1981, p. 275ff. [209] I. A. Koppel, J. B. Koppel *et al.*, Org. React. (Tartu) *21*, 98, 144 (1984); Chem. Abstr. *103*, 12232k, 37044p (1985). [210] (a) V. Gageiro, M. Aillon, and M. C. Rezende, J. Chem. Soc., Faraday Trans. *88*, 201 (1992); (b) V. G. Machado, C. Machado, M. de Gracia Nascimento, and M. C. Rezende, Quimica Nova *19*, 523 (1996); Chem. Abstr. *126*, 226509e (1997).

[211] G. Hollmann and F. Vögtle, Chem. Ber. *117*, 1355 (1984). [212] Y. Pocker and J. C. Ciula, J. Am. Chem. Soc. *111*, 4728 (1989). [213] H. Langhals, Tetrahedron *43*, 1771 (1987). [214] A. v. Baeyer and V. Villiger, Ber. Dtsch. Chem. Ges. *35*, 1189 (1902). [215] P. Plieninger and H. Baumgärtel, Liebigs Ann. Chem. *1983*, 860. [216] J. J. Dawber and R. A. Williams, J. Chem. Soc., Faraday Trans. I *82*, 3097 (1986). [217] P. de Mayo, A. Safarzadeh-Amiri, and S. K. Wong, Can. J. Chem. *62*, 1001 (1984). [218] P. Mukerjee, C. Ramachandran, and R. A. Pyter, J. Phys. Chem. *86*, 3189 (1982). [219] A. Janowski, I. Turowska-Tyrk, and P. K. Wrona, J. Chem. Soc., Perkin Trans. II, 821 (1985). [220] W. Walter and O. H. Bauer, Liebigs Ann. Chem. *1977*, 421.

[221] R. W. Soukup, Chemie in unserer Zeit *17*, 129, 163 (1983); R. W. Soukup and R. Schmid, J. Chem. Educ. *62*, 459 (1985). [222] D. C. Dong and M. A. Winnik, Photochem. Photobiol. *35*, 17 (1982); Can. J. Chem. *62*, 2560 (1984). [223] K. W. Street and W. E. Acree, Analyst *111*, 1197 (1986). [224] J.-L. M. Abboud, M. J. Kamlet, and R. W. Taft, Progr. Phys. Org. Chem. *13*, 485 (1981); R. W. Taft, J.-L. M. Abboud, M. J. Kamlet, and M. H. Abraham, J. Sol. Chem. *14*, 153 (1985). [225] H. C. Brown, G. K. Barbaras, H. L. Berneis, W. H. Bonner, R. B. Johannesen, M. Grayson, and K. L. Nelson, J. Am. Chem. Soc. *75*, 1 (1953). [226] M. J. Kamlet, J.-L. M. Abboud, M. H. Abraham, and R. W. Taft, J. Org. Chem. *48*, 2877 (1983). [227] C. Laurence, P. Nicolet, M. T. Dalati, J.-L. M. Abboud, and R. Notario, J. Phys. Chem. *98*, 5807 (1994). [228] D. V. Matyushov, R. Schmid, and B. M. Ladanyi, J. Phys. Chem. B *101*, 1035 (1997). [229] M. J.

Kamlet and R. W. Taft, J. Chem. Soc., Perkin Trans. II, 337 (1979). [230] M. J. Kamlet, R. M. Doherty, M. H. Abraham, Y. Marcus, and R. W. Taft, J. Phys. Chem. *92*, 5244 (1988).

[231] T. M. Krygowski, E. Milczarek, and P. K. Wrona, J. Chem. Soc., Perkin Trans. II 1563 (1980). [232] T. M. Krygowski *et al*., Tetrahedron *37*, 119 (1981); J. Chem. Res. (S) 116 (1983). [233] O. W. Kolling, Anal. Chem. *53*, 54 (1981); ibid. *56*, 430 (1984). [234] V. Bekárek, J. Phys. Chem. *85*, 722 (1981); J. Chem. Soc., Perkin Trans. II 1293 (1983); ibid. 1425 (1986); V. Bekárek and M. Stolařová, Collect. Czech. Chem. Commun. *48*, 1237 (1983); ibid. *49*, 2332 (1984). [235] J.-L. M. Abboud, R. W. Taft, and M. J. Kamlet, J. Chem. Soc., Perkin Trans. II, 815 (1985). [236] M. Sjöström and S. Wold, J. Chem. Soc., Perkin Trans. II, 104 (1981). [237] P. Nicolet and C. Laurence, J. Chem. Soc., Perkin Trans. II, 1071 (1986). [238] C. Laurence, P. Nicolet, and M. Helbert, J. Chem. Soc., Perkin Trans. II, 1081 (1986). [239] P. Nicolet, C. Laurence, and M. Lucon, J. Chem. Soc., Perkin Trans. II, 483 (1987). [240] C. Somolinos, I. Rodriguez, M. I. Redondo, and M. V. Garcia, J. Mol. Struct. *143*, 301 (1986).

[241] M. I. Redondo, C. Somolinos, and M. V. Garcia, Spectrochim. Acta, Part A *42*, 677 (1986). [242] R. G. Makitra, Ya. M. Vasyutin, and V. Ya. Zhukowskii, Org. React. (Tartu) *21*, 190 (1984); Chem. Abstr. *103*, 122883f (1985). [243] R. G. Makitra and Ya. N. Pirig, Org. React. (Tartu) *16*, 103 (1979); Chem. Abstr. *92*, 41142r (1980); Org. React. (Tartu) *17*, 180 (1980); Chem. Abstr. *95*, 79388a (1981). [244] O. W. Kolling, Anal. Chem. *49*, 591 (1977); ibid. *55*, 143 (1983). [245] A. H. Reddoch and S. Konishi, J. Chem. Phys. *70*, 2121 (1979). [246] M. C. R. Symons and A. S. Pena-Nuñez, J. Chem. Soc., Faraday Trans. I *81*, 2421 (1985). [247] Y. Liu and Y. Wang, Huaxue Xuebo *43*, 232 (1985); Chem. Abstr. *103*, 177687a (1985). [248] Z. Kecki, B. Łyczkowski, and W. Kołodziejski, J. Sol. Chem. *15*, 413 (1986). [249] B. Chawla, S. K. Pollack, C. B. Lebrilla, M. J. Kamlet, and R. W. Taft, J. Am. Chem. Soc. *103*, 6924 (1981). [250] H. Elias, M. Dreher, S. Neitzel, and H. Volz, Z. Naturforsch., Part B *37*, 684 (1982).

[251] U. Mayer: *NMR-Spectroscopic Studies on Solute-Solvent and Solute-Solute Interactions*, in N. Tanaka, H. Ohtaki, and R. Tamamushi (eds.): *Ions and Molecules in Solution*, Elsevier, Amsterdam, 1983, p. 219ff. [252] M. C. R. Symons and G. Eaton, J. Chem. Soc., Faraday Trans. I *78*, 3033 (1982). [253] P. Bacelon, J. Corset, and C. de Loze, J. Sol. Chem. *12*, 23 (1983). [254] B. Allard and E. Casadevall, Nouv. J. Chim. *9*, 565 (1985). [255] M. J. Kamlet, P. W. Carr, R. W. Taft, and M. H. Abraham, J. Am. Chem. Soc. *103*, 6062 (1981). [256] R. W. Taft, M. H. Abraham, R. M. Doherty, and M. J. Kamlet, J. Am. Chem. Soc. *107*, 3105 (1985). [257] M. H. Abraham, R. M. Doherty, M. J. Kamlet, J. M. Harris, and R. W. Taft, J. Chem. Soc., Perkin Trans. II, 913, 1097 (1987). [258] R. G. Makitra and Ya. N. Pirig, Zh. Obshch. Khim. *56*, 657 (1986); J. Gen. Chem. USSR *56*, 584 (1986). [259] F. Vernon: *The Characterization of Solute-Solvent Interactions in Gas-Liquid Chromatography*, in C. E. H. Knapman (ed.): *Developments in Chromatography*, Applied Science Publ., London, 1978, Vol. 1, Chapter 1, p. 1ff.; Chem. Abstr. *92*, 65125p (1980). [260] R. V. Golovnaya and T. A. Misharina: *Thermodynamic Treatment of the Polarity and Selectivity of Solvents in Gas Chromatography*, Usp. Khim. *49*, 171 (1980); Russ. Chem. Rev. *49*, 88 (1980).

[261] G. Tarján, Á. Kiss, G. Kocsis, S. Mészáros, and J. M. Takács, J. Chromatogr. *119*, 327 (1976). [262] S. Wold and K. Andersson, J. Chromatogr. *80*, 43 (1973). [263] J.-L. M. Abboud, R. W. Taft, and M. J. Kamlet, J. Chem. Res. (S) 98 (1984). [264] J. E. Brady and P. W. Carr, J. Phys. Chem. *88*, 5796 (1984). [265] C. G. Swain, M. S. Swain, A. L. Powell, and S. Alunni, J. Am. Chem. Soc. *105*, 502 (1983). [266] U. Mayer, Monatsh. Chem. *109*, 421, 775 (1978). [267] M. J. Kamlet, J.-F. Gal, P.-C. Maria, and R. W. Taft, J. Chem. Soc., Perkin Trans. II, 1583 (1985). [268] G. Collier, I. A. Pickering, and J. Shorter, unpublished work; *cf*. I. A. Pickering, Thesis, University of Hull 1985, and reference [15] , p. 164, equation (5.22). [269] J. E. Brady and P. W. Carr, J. Phys. Chem. *89*, 1813 (1985). [270] B. Eliasson, D. Johnels, S. Wold, U. Edlund, and M. Sjöström, Acta Chem. Scand., Part B *36*, 155 (1982); ibid. *41*, 291 (1987).

[271] M. J. Kamlet and R. W. Taft, Acta Chem. Scand., Part B *40*, 619 (1986). [272] M. J. Kamlet, M. H. Abraham, R. M. Doherty, and R. W. Taft, J. Am. Chem. Soc. *106*, 464 (1984); Nature (London) *313*, 384 (1985). [273] M. H. Abraham, M. J. Kamlet, R. W. Taft, and P. K. Weathersby, J. Am. Chem. Soc. *105*, 6797 (1983). [274] M. J. Kamlet, R. M. Doherty, M. H. Abraham, and R. W. Taft, Quant. Struct.-Act. Relat. *7*, 71 (1988); Chem. Abstr. *109*, 144105c (1988). [275] R.

W. Taft, J.-L. M. Abboud, and M. J. Kamlet, J. Org. Chem. *49*, 2001 (1984). [276] C. G. Swain, J. Org. Chem. *49*, 2005 (1984). [277] S. Glikberg and Y. Marcus, J. Sol. Chem. *12*, 255 (1983). [278] G. Heublein, P. Hallpap, R. Wondraczek, and P. Adler, Z. Chem. *20*, 11 (1980). [279] G. Heublein, J. Macromol. Sci., Part A *22*, 1277 (1985). [280] M. Kupfer, A. Henrion, and W. Abraham, Z. Phys. Chem. (Leipzig) *267*, 705 (1986).

[281] E. Fogassy, A. Lopata, F. Faigl, F. Darvas, M. Ács, and L. Töke, Tetrahedron Lett. *21*, 647 (1980); A. Lopata, F. Faigl, E. Fogassy, and F. Darvas, J. Chem. Res. (S) 322, 324 (1984). [282] M. H. Abraham, P. L. Grellier, and R. A. McGill, J. Chem. Soc., Perkin Trans. II, 339 (1988). [283] H.-J. Schneider and N. K. Sangwan, J. Chem. Soc., Chem. Commun. 1787 (1986); Angew. Chem. *99*, 924 (1987); Angew. Chem. Int. Ed. Engl. *26*, 896 (1987). [284] K. Popper: *Unended Quest, An Intellectual Autobiography*, Revised edition, Open Court, La Salle/Illinois, 1982, p. 28. [285] Y. Marcus, J. Phys. Chem. *91*, 4422 (1987). [286] M. J. Kamlet, R. M. Doherty, J.-L. M. Abboud, M. H. Abraham, and R. W. Taft: *Solubility – A New Look*, CHEMTECH *16*, 566 (1986). [287] I. Persson, M. Sandström, and P. L. Goggin, Inorg. Chim. Acta *129*, 183 (1987). [288] (a) M. H. Abraham, R. M. Doherty, M. J. Kamlet, J. M. Harris, and R. W. Taft, J. Chem. Soc., Perkin Trans. II, 913, 1097 (1987); (b) M. H. Abraham, P. L. Grellier, A. Nasehzadeh, R. A. C. Walker, J. Chem. Soc., Perkin Trans. II, 1717 (1988). [289] O. Pytela, Collect. Czech. Chem. Commun. *53*, 1333 (1988). [290] G. Gritzner, Z. Phys. Chem. (München) N. F. *158*, 99 (1988).

[291] P. Müller: *Glossary of Terms used in Physical Organic Chemistry – IUPAC Recommendations 1994*, Pure Appl. Chem. *66*, 1077 (1994), particularly p. 1151. [292] M. Sandström, I. Persson, and P. Persson, Acta Chem. Scand. *44*, 653 (1990). [293] C. Reichardt: *Solvatochromic Dyes as Solvent Polarity Indicators*, Chem. Rev. *94*, 2319 (1994). [294] (a) Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998, Chapter 4, p. 131ff. (b) Y. Marcus, Chem. Soc. Rev. *22*, 409 (1993). [295] J.-L. M. Abboud and R. Notario: *Critical Compilation of Scales of Solvent Polarities, Part I, Pure non-HBD Solvents*, Pure Appl. Chem. *71*, 645 (1999). [296] J. Catalán: *Solvent Effects based on Pure Solvent Scales*, in G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, William Andrew Publishing, New York, 2001, Chapter 10.3, p. 583ff. [297] C. Cativiela, J. I. Garcia, J. Gil, R. M. Martinez, J. A. Mayoral, L. Salvatella, J. S. Urieta, A. M. Mainar, and M. H. Abraham, J. Chem. Soc., Perkin Trans. II, 653 (1997). [298] V. S. Shmidt, E. A. Mezhov, and S. S. Novikova, Radiokhimiya *9*, 700 (1967); Soviet Radiochemistry *9*, 657 (1967); Chem. Abstr. *68*, 99202f (1968). [299] V. S. Shmidt: *Linear Relations of Free Energies as a Device for the Generalization and Analysis of Data on Extraction*, Usp. Khim. *47*, 1730 (1978); Chem. Abstr. *89*, 221660e (1978). [300] (a) V. S. Shmidt, N. L. Khananashvili, and E. A. Mezhov, Radiokhimiya *29*, 519 (1987); Soviet Radiochemistry *29*, 503 (1987); Chem. Abstr. *107*, 223988g (1987); (b) V. S. Shmidt, E. A. Mezhov, and N. L. Khananashvili, Zh. Fiz. Khim. *62*, 330 (1988); Chem. Abstr. *108*, 193653v (1988).

[301] E. A. Mezhov, N. L. Khananashvili, and V. S. Shmidt, Radiokhimiya *31*, 60 (1989); Chem. Abstr. *111*, 181942 (1989). [302] (a) R. G. Makitra: *Linear Free-Energy Relationships in Extraction*, Org. React. (Tartu) *31*, 79 (1997); Chem. Abstr. *128*, 53777b (1998); (b) R. G. Makitra, Ya. N. Pirig, and Ya. M. Vasyutyn, Radiochemistry (Transl. of Radiokhimiya) *38*, 485 (1996); Chem. Abstr. *127*, 225734h (1997). [303] G. Gritzner, J. Mol. Liquids *73/74*, 487 (1997). [304] T. Chen, G. Hefter, and Y. Marcus, J. Sol. Chem. *29*, 201 (2000). [305] D. N. Kevill: *Development and Uses of Scales of Solvent Nucleophilicity*, in M. Charton (ed.): *Advances in Quantitative Structure–Property Relationships*, JAI Press, Greenwich/CT, 1996, Vol. 1, p. 81ff. [306] T. W. Bentley and G. Llewellyn: Y_X *Scales of Solvent Ionizing Power*, Progr. Phys. Org. Chem. *17*, 121 (1990). [307] T. W. Bentley: *Nucleophilicities of Aqueous, Alcoholic, and Acidic Media*, Adv. Chem. Ser. No. 215 (Nucleophilicity), American Chemical Society, Washington D.C., 1987, p. 255ff. [308] D. N. Kevill and S. W. Anderson, J. Org. Chem. *56*, 1845 (1991). [309] D. N. Kevill and M. J. D'Souza, J. Chem. Res. (S) 174 (1993). [310] D. N. Kevill and M. J. D'Souza, J. Phys. Org. Chem. *5*, 287 (1992).

[311] K.-T. Liu and H.-C. Shen, J. Org. Chem. *56*, 3021 (1991). [312] T. W. Bentley, J.-P. Dau-Schmidt, G. Llewellyn, and H. Mayr, J. Org. Chem. *57*, 2387 (1992). [313] M. Fujio *et al.*, Bull. Chem. Soc. Jpn. *68*, 673, 2619 (1995). [314] K. Takeuchi *et al.*, J. Phys. Org. Chem. *14*, 229 (2001).

[315] H. Langhals and K. Hadizamani, Chem. Ber. *123*, 407 (1990). [316] E. Buncel and S. Rajagopal: *Solvatochromism and Solvent Polarity Scales*, Acc. Chem. Res. *23*, 226 (1990). [317] S. Bagchi *et al.*, J. Sol. Chem. *19*, 271 (1990); J. Chem. Soc., Perkin Trans. II, 343 (1992). [318] D. A. Johnson, R. Shaw, and E. F. Silversmith, J. Chem. Educ. *71*, 517 (1994). [319] B. R. Osterby and R. D. McKelvey, J. Chem. Educ. *73*, 260, 737 (1996). [320] T. Deng and W. E. Acree, J. Chem. Educ. *76*, 1555 (1999).

[321] M. F. Vitha, J. Chem. Educ. *78*, 370 (2001). [322] C. Machado and V. G. Machado, J. Chem. Educ. *78*, 649 (2001). [323] M. A. Kessler and O. S. Wolfbeis, Chem. Phys. Lipids *50*, 51 (1989); Spectrochim. Acta, Part A, *47*, 187 (1991). [324] C. Reichardt, M. Eschner, and G. Schäfer, J. Phys. Org. Chem. *14*, 737 (2001). [325] C. Reichardt, D. Che, G. Heckenkemper, and G. Schäfer, Eur. J. Org. Chem. 2343 (2001). [326] Y. Marcus, J. Chem. Soc., Perkin Trans. II, 1015, 1751 (1994). [327] R. D. Skwierczynski and K. A. Connors, J. Chem. Soc., Perkin Trans. II, 467 (1994); see also W. A. Acree, J. R. Powell, and S. A. Tucker, ibid. 529 (1995). [328] (a) E. Bosch, M. Rosés *et al.*, J. Chem. Soc., Perkin Trans. II, 1607 (1995); ibid. 1497, 2177 (1996); ibid. 243, 1341 (1997); (b) E. Bosch, M. Rosés *et al.*, J. Phys. Org. Chem. *9*, 403 (1996); ibid. *11*, 193 (1998); ibid. *12*, 109 (1999). [329] S. Bagchi *et al.*, J. Chem. Soc., Faraday Trans. *91*, 631 (1995); ibid. *92*, 1499 (1996). [330] P. M. E. Mancini *et al.*, J. Phys. Org. Chem. *8*, 617 (1995); ibid. *10*, 849 (1997); ibid. *12*, 207, 430, 713 (1999); ibid. *13*, 221 (2000).

[331] O. A. El Seoud *et al.*, Ber. Bunsenges. Phys. Chem. *101*, 105, 902 (1997); J. Phys. Org. Chem. *13*, 679 (2000). [332] C. Reichardt: *Solvatochromism, Thermochromism, Piezochromism, Halochromism, and Chiro-Solvatochromism of Pyridinium N-Phenoxide Betaine Dyes*, Chem. Soc. Rev. *21*, 147 (1992). [333] E. Buncel and S. Rajagopal, Acc. Chem. Res. *23*, 226 (1990); J. Org. Chem. *54*, 798 (1989); Dyes Pigm. *17*, 303 (1991). [334] M. Podsiadla, J. Rzeszotarska, and M. K. Kalinowski, Monatsh. Chem. *125*, 827 (1994). [335] J. Catalán, V. López, P. Pérez, R. Martin-Villamil, and J. G. Rodriguez, Liebigs Ann. 241, 793 (1995). [336] J. Catalán, C. Diaz, V. López, P. Pérez, J.-L. de Paz, and J. G. Rodriguez, Liebigs Ann. 1785 (1996). [337] (a) J. Catalán and C. Diaz, Liebigs Ann. 1941 (1997); (b) J. Catalán and C. Diaz, Eur. J. Org. Chem. 885 (1999). [338] T. Abe, Bull. Chem. Soc. Jpn. *63*, 2328 (1990). [339] C. Laurence: *Similarity Models in IR and UV Spectroscopy*, in R. I. Zalewski, T. M. Krygowski, and J. Shorter (eds.): *Similarity Models in Organic Chemistry, Biochemistry, and Related Fields*, Elsevier, Amsterdam, 1991, Chapter 5, p. 231ff. [340] J. Catalán, J. Org. Chem. *60*, 8315 (1995).

[341] J. Catalán, C. Diaz, and F. Garcia-Blanco, J. Org. Chem. *64*, 6512 (1999). [342] J. Catalán, C. Diaz, and F. Garcia-Blanco, J. Org. Chem. *65*, 3409 (2000). [343] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, William Andrew Publishing, New York, 2001, Chapter 4, p. 101ff., and Chapter 5, p. 243ff. [344] (a) C. F. Poole and S. K. Poole, Chem. Rev. *89*, 377 (1989); (b) C. F. Poole, T. O. Kollie, and S. K. Poole, Chromatographia *34*, 281 (1992). [345] R. G. Makitra, Ya. N. Pirig, and R. B. Kivelyuk, Zh. Obshch. Khim. *60*, 2209 (1990); Russ. J. Gen. Chem. *60*, 1973 (1990); Chem. Abstr. *114*, 246504e (1991). [346] A. R. Katritzky, T. Tamm, Y. Wang, S. Sild, and M. Karelson, J. Chem. Inf. Comput. Sci. *39*, 684 (1999). [347] A. R. Katritzky, T. Tamm, Y. Wang, and M. Karelson, J. Chem. Inf. Comput. Sci. *39*, 692 (1999). [348] V. Palm and N. Palm, Org. React. (Tartu) *28*, 125 (1993); ibid. *31*, 141 (1997); Russ. J. Org. Chem. *36*, 1075 (2000); see also N. Palm, Ph.D. Thesis, University of Tartu/Estonia, 2001. [349] O. Pytela, Collect. Czech. Chem. Commun. *54*, 136 (1989). [350] (a) L. Y. Wilson and G. R. Famini, J. Med. Chem. *34*, 1668 (1991); (b) G. R. Famini *et al.*, Quant. Struct.-Act. Relat. *10*, 344 (1991); ibid. *11*, 162 (1992); (c) G. R. Famini, C. A. Penski, and L. Y. Wilson, J. Phys. Org. Chem. *5*, 395 (1992); (d) C. J. Cramer, G. R. Famini, and A. H. Lowry, Acc. Chem. Res. *26*, 599 (1993); (e) G. R. Famini and L. Y. Wilson, Org. React. (Tartu) *29*, 117 (1995); Chem. Abstr. *124*, 145303q (1996).

[351] (a) R. S. Drago, J. Chem. Soc., Perkin Trans. II, 1827 (1992); (b) R. S. Drago, M. S. Hirsch, D. C. Ferris, and C. W. Chronister, J. Chem. Soc., Perkin Trans. II, 219 (1994). [352] O. Pytela, Collect. Czech. Chem. Commun. *53*, 1333 (1988). [353] G. A. Marschalok, Yu. I. Oglashennyi, R. G. Makitra, Ya. N. Pirig, and I. I. Yatchishin, Zh. Org. Khim. *34*, 1471 (1998); Chem. Abstr. *131*, 129619m (1999). [354] R. G. Makitra, Ya. N. Pirig, and I. I. Yatchishin, Zh. Org. Khim. *35*, 1073

(1999); Chem. Abstr. *132*, 122224r (2000). [355] R. G. Makitra, Ya. N. Pirig, and Ya. M. Vasyutyn, Zh. Obshch. Khim. *70*, 722 (2000); Russ. J. Gen. Chem. *70*, 672 (2000); Chem. Abstr. *134*, 106493p (2001). [356] R. Flunt, R. G. Makitra, O. Makohon, and Ya. N. Pirig, Org. React. (Tartu) *31*, 21 (1997); Chem. Abstr. *128*, 53813k (1998). [357] P. W. Carr, Microchem. J. *48*, 4 (1993); Chem. Abstr. *119*, 189467x (1993). [358] R. Fuchs, M. H. Abraham, M. J. Kamlet, and R. W. Taft, J. Phys. Org. Chem. *2*, 559 (1989). [359] M. J. Kamlet, M. H. Abraham, P. W. Carr, R. M. Doherty, and R. W. Taft, J. Chem. Soc., Perkin Trans. II, 2087 (1988). Because of the untimely death of Dr. M. J. Kamlet in 1988, not all papers of this and the preceding series have been published. His last paper, entitled "Linear Solvation Energy Relationships: An Improved Equation for Correlation and Prediction of Aqueous Solubilities of Aromatic Solutes Including Polycyclic Aromatic Hydrocarbons and Polychlorinated Biphenyls", was published posthumously in Progr. Phys. Org. Chem. *19*, 295 (1993). [360] (a) M. H. Abraham, Chem. Soc. Rev. *22*, 73 (1993); (b) M. H. Abraham, Pure Appl. Chem. *65*, 2503 (1993).

[361] M. H. Abraham, J. Chromatogr. *644*, 95 (1993). [362] M. H. Abraham *et al.*, Pesticide Science *55*, 78 (1999). [363] M. H. Abraham, C. E. Green, and W. E. Acree, J. Chem. Soc., Perkin Trans. II, 281 (2000). [364] G. R. Famini and L. Y. Wilson, J. Phys. Org. Chem. *12*, 645 (1999). [365] A. D. Headley, S. D. Starnes, L. Y. Wilson, and G. R. Famini, J. Org. Chem. *59*, 8040 (1994). [366] A. D. Headley and J. Nam, J. Phys. Org. Chem. *15*, 62 (2002). [367] G. R. Famini and L. Y. Wilson, J. Chem. Soc., Perkin Trans. II, 1641 (1994). [368] J. B. F. N. Engberts, G. R. Famini, A. Perjéssy, and L. Y. Wilson, J. Phys. Org. Chem. *11*, 261 (1998). [369] G. R. Famini, D. Benyamin, C. Kim, R. Veerawat, and L. Y. Wilson, Collect. Czech. Chem. Commun. *64*, 1727 (1999). [370] Y. Marcus, A. L. Smith, M. V. Korobov, A. L. Mirakyan, N. V. Avramenko, and E. B. Stukalin, J. Phys. Chem. B *105*, 2499 (2001).

[371] R. M. C. Gonçalves and L. M. P. C. Albuquerque, J. Phys. Org. Chem. *14*, 731 (2001).

Appendix

[1] A. Riddick, W. B. Bunger, and T. Sakano: *Organic Solvents, Physical Properties and Methods of Purification*, 4th ed., in A. Weissberger (ed.): *Techniques of Chemistry*, Vol. II, Wiley-Interscience, New York, 1986. [2] W. L. Archer: *Industrial Solvents Handbook*, Dekker, New York, 1996. [3] B. P. Whim and P. G. Johnson: *Directory of Solvents*, Blackie Academic and Professional, London, 1996. [4] A. J. Gordon and R. A. Ford: *The Chemist's Companion – A Handbook of Practical Data, Techniques, and References*, Wiley-Interscience, New York, 1972. [5] G. P. Nilles and R. D. Schuetz, J. Chem. Educ. *50*, 267 (1973). [6] R. L. Schneider: *Physical Properties of Some Organic Solvents*, Eastman Organic Chemical Bulletin *47*, No. 1 (1975). [7] D. R. Lide (ed.): *Handbook of Chemistry and Physics*, 82th ed., CRC Press, Boca Raton/FL, USA, 2001. [8] Y. Marcus: *The Properties of Solvents*, Wiley, Chichester, 1998. [9] M. Raban and K. Mislow, Top. Stereochem. *2*, 199 (1967). [10] H. Pracejus, Fortschr. Chem. Forsch. *8*, 493 (1967).

[11] T. D. Inch, Synthesis 466 (1970). [12] J. D. Morrison and H. S. Mosher; *Asymmetric Organic Reactions*, Prentice Hall, Englewood Cliffs/New Jersey 1971, pp. 411. [13] W. H. Pirkle and D. J. Hoover: *NMR Chiral Solvating Agents*, Topics in Stereochemistry *13*, 263 (1982). [14] V. Sedivec and J. Flek: *Handbook of Analysis of Organic Solvents*, Ellis Horwood, Chichester, and Wiley, New York, 1976. [15] M. Spiteller and G. Spiteller: *Massenspektrensammlung von Lösungsmitteln, Verunreinigungen, Säulenbelegmaterialien und einfachen aliphatischen Verbindungen*, Springer-Verlag, Wien, New York, 1973. [16] B. Hampel and K. Maas, Chemiker-Ztg. *95*, 316 (1971). [17] W. Bunge in Houben-Weyl-Müller, *Methoden der organischen Chemie*, 4th ed., Thieme, Stuttgart, 1959, Vol. I/2, pp. 765–868. [18] H. Rickert and H. Schwarz in Houben-Weyl-Müller, *Methoden der organischen Chemie*, 4th ed., Thieme, Stuttgart, 1959, Vol. I/2, pp. 869–885. [19] G. Wohlleben, Angew. Chem. *67*, 741 (1955); ibid. *68*, 752 (1956). [20] B. D. Pearson and J. E. Ollerenshaw, Chem. Ind. (London) *1966*, 370.

[21] G. Hesse and H. Schildknecht, Angew. Chem. *67*, 737 (1955); G. Hesse, B. P. Engelbrecht, H. Engelhardt, and S. Nitsch, Z. Anal. Chem. *241*, 91 (1968); G. Hesse: *Chromatographisches*

Praktikum, Akademische Verlagsgesellschaft, Frankfurt (Main), 1968. [22] ICN Biomedicals GmbH (formerly Woelm Pharma GmbH): *Lösungsmittel–Reinigung mit ICN Adsorbentien*, D-37269 Eschwege/Germany. [23] H. Engelhardt: *Purification of Organic Solvents by Frontal-Analysis Chromatography*, in M. Zief and R. M. Speights (eds.): *Ultrapurity-Methods and Techniques*, Dekker, New York, 1972, p. 71f. [24] B. P. Engelbrecht, GIT Fachz. Lab. *23*, 681 (1979); Chem. Abstr. *91*, 133523y (1979). [25] M. Pestemer, Angew. Chem. *63*, 118 (1951); ibid. *67*, 740 (1955); M. Pestemer: *Anleitung zum Messen von Absorptionsspektren im Ultravioletten und Sichtbaren*, Thieme, Stuttgart, 1964. [26] J. Schurz and H. Stübchen, Z. Elektrochem., Ber. Bunsenges. Physik. Chem. *61*, 754 (1957). [27] D. D. Tunnicliff, Talanta *2*, 341 (1959). [28] W. Esselborn and B. Hampel, Photoelectric Spectrometry Group Bulletin Nr. 17, 498 (1967); Chem. Abstr. *69*, 35055y (1968). [29] J. Ratcliffe: *Practical Hints on Absorption Spectrometry*, Plenum Publishing Corporation, New York, 1967. [30] N. Fisher and R. M. Cooper, Chem. Ind. (London) *1968*, 619.

[31] G. Belanger, P. Sauvageau, and C. Sandorfy, Chem. Phys. Lett. *3*, 649 (1969). [32] H. R. Dickinson and W. C. Johnson, Appl. Opt. *10*, 681 (1971); Chem. Abstr. *74*, 117764w (1971). [33] M. F. Fox and E. Hayon, J. Phys. Chem. *76*, 2703 (1972). [34] B. Hampel in: *DMS-UV-Atlas organischer Verbindungen*, Butterworths, London, and Verlag Chemie, Weinheim, 1966, Hauptgruppe M. [35] N. L. Alpert, W. E. Keiser, and H. A. Szymanski; *IR-Theory and Practice of Infrared Spectroscopy*, 2nd ed., Plenum Press, New York, 1973. [36] *DMS-Working Atlas of Infrared Spectroscopy*, Butterworths, London, and Verlag Chemie, Weinheim, 1972. [37] R. F. Goddu and D. A. Delker, Anal. Chem. *32*, 140 (1960). [38] A. Visapää, Kemian Teollisuus *22*, 487 (1965), Chem. Abstr. *63*, 15732 (1965); Tek. Tutkimuslaitos, Tiedotus, Sarja IV *76* (1965), Chem. Abstr. *64*, 18701 (1966). [39] F. F. Bentley, E. F. Wolfarth, N. E. Srp, and W. R. Powell, Spectrochim. Acta *13*, 1 (1958). [40] H. R. Wyss, R. D. Werder, and H. H. Günthard, Spectrochim. Acta *20*, 573 (1964).

[41] N. L. McNiven and R. Court: *IR-Spectra of Deuterated Solvents*, in L. May (ed.): *Spectroscopic Tricks*. Vol. 3, Plenum Press, New York, London, 1974, p. 148. [42] D. W. Henty and S. Vary, Chem. Ind. (London) 1782 (1967). [43] R. A. Fletton and J. E. Page, The Analyst *96*, 370 (1971). [44] H.-O. Kalinowski, S. Berger, and S. Braun: *Carbon–13 NMR Spectroscopy*, Wiley, Chichester, 1988. [45] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001; particularly Chapters 2.3 (p. 42ff.) and 14 (p. 847ff.). [46] A. Lüttringhaus in Houben-Weyl-Müller, *Methoden der organischen Chemie*, 4th ed., G. Thieme, Stuttgart, 1958, Vol. I/1, p. 341 (especially p. 369). [47] R. S. Tipson in A. Weissberger (ed.): *Technique of Organic Chemistry*, 2nd ed., Interscience, New York, London, 1956, Vol. III, Part I, p. 395 (especially p. 549). [48] B. Cornils and J. Falbe, in F. Korte (ed.): *Methodicum Chimicum*, G. Thieme, Stuttgart, and Academic Press, New York, London, 1973, Vol. I/1, pp. 39–45. [49] K. Schwetlick et al.: *Organikum – Organisch-chemisches Grundpraktikum*, 21st ed., Wiley-VCH, Weinheim/Germany, 2000, p. 36. [50] F. A. von Metzsch, Angew. Chem. *65*, 586 (1953).

[51] E. Hecker, Chimia *8*, 229 (1954); E. Hecker: *Verteilungsverfahren in Laboratorien*, Verlag Chemie, Weinheim, 1955, p. 92 and 139. [52] L. R. Snyder: *Solvent Selection for Separation Processes*, in E. S. Perry and A. Weissberger (eds.): *Techniques of Chemistry*, 3rd ed., Vol. XII, p. 25ff., Wiley-Interscience, New York, 1978. [53] T. C. Lo, M. H. J. Baird, and C. Hanson (eds.): *Handbook of Solvent Extraction*, Wiley, Chichester, 1983. [54] M. Hampe: *Flüssig/Flüssig-Extraktion: Einsatzgebiete und Lösungsmittel-Auswahl*, Chem.-Ing.-Tech. *50*, 647 (1978); Chem. Abstr. *90*, 8134a (1979). [55] F. J. Wolf: *Separation Methods in Organic and Biochemistry*, Academic Press, New York, London, 1969. [56] H. Pauschmann and E. Bayer, in F. Korte (ed.): *Methodicum Chimicum*. G. Thieme, Stuttgart, and Academic Press, New York, London, 1973, Vol. I/1, p. 148 (espicially p. 156) and references cited in this review. [57] G. Wohlleben, Fortschr. Chem. Forsch. *6*, 640 (1966). [58] C. F. Poole and S. K. Poole: *Chromatography*, Elsevier, Amsterdam, 1991. [59] L. R. Snyder: *Principles of Adsorption Chromatography*, Dekker, New York, 1968. [60] L. R. Snyder and J. J. Kirkland: *Introduction to Modern Liquid Chromatography*, 2nd ed., Wiley-Interscience, New York, 1979.

[61] D. F. G. Pusey, Chem. Brit. *5*, 408 (1969). [62] W. Huber: *Titrationen in nichtwäßrigen Lösungsmitteln*, Akademische Verlagsgesellschaft, Frankfurt (Main), 1964; *Titrations in Nonaqueous*

Solvents, Academic Press, New York, London, 1967. [63] J. Kucharský and L. Safařik: *Titrations in Non-Aqueous Solvents*, Elsevier, Amsterdam, 1965. [64] L. N. Bykova, A. P. Kreshkov, N. A. Kazaryan, and I. D. Pevzner, Usp. Khim. *37*, 677 (1968); Russ. Chem. Rev. *37*, 278 (1968). [65] L. Šafařik and Z. Stránský: *Titrimetric Analysis in Organic Solvents*, in G. Svehla (ed.): *Wilson and Wilson's Comprehensive Analytical Chemistry*, Vol. XXII, Elsevier, Amsterdam, New York, 1986. [66] G. Charlot and B. Trémillon: *Réactions Chimiques dans les Solvants et les Sels Fondus*, Gauthier-Villars, Paris, 1963; *Chemical Reactions in Solvents and Melts*, Pergamon Press, Oxford, 1969. [67] C. A. Streuli: *Acid-Base Titration in Nonaqueous Solvents*, in I. M. Kolthoff, P. J. Elving, and E. B. Sandell (eds.): *Treatise on Analytical Chemistry*, Wiley-Interscience, New York, 1975, Part I, Vol. 11, p. 7035–7116. [68] I. Gyenes: *Titrationen in nichtwäßrigen Medien*, 3rd ed., Akadémiai Kiadó, Budapest, and F. Enke, Stuttgart, 1970; *Titrations in Non-Aqueous Media*, Van Nostrand, New York, 1967. [69] B. Kratochvil, Anal. Chem. *54*, 105 R (1982). [70] L. N. Bykova and S. I. Petrov, Usp. Khim. *41*, 2065 (1972); Russ. Chem. Rev. *41*, 975 (1972).

[71] R. G. Bates: *Medium Effects and pH in Nonaqueous Solvents*, in J. F. Coetzee and C. D. Ritchie (eds.): *Solute-Solvent Interactions*, Dekker, New York, London, 1969, Vol. 1, p. 50ff. [72] H. B. van der Heijde, Anal. Chim. Acta *17*, 512 (1957). [73] K. Izutsu: *Electrochemistry in Nonaqueous Solutions*, Wiley-VCH, Weinheim, 2002. [74] G. J. Janz and R. P. T. Tomkins (eds.): *Nonaqueous Electrolytes Handbook*, Academic Press, London, New York, 1972/73, Vols. 1 and 2. [75] D. Orbakh (ed.): *Nonaqueous Electrochemistry*, Dekker, New York, 1999. [76] M. Salomon: *Electrochemical Measurements*, in A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, Chapter 2, Part 2, p. 137ff. [77] D. T. Sawyer and J. L. Roberts: *Experimental Electrochemistry for Chemists*, Wiley, New York, 1974, p. 167ff. [78] B. Kratochvil and H. L. Yeager: *Conductance of Electrolytes in Organic Solvents*, Fortschr. Chem. Forsch. *27*, 1 (1972). [79] R. Fernández-Prini: *Conductance*, in A. K. Covington and T. Dickinson (eds.): *Physical Chemistry of Organic Solvent Systems*, Plenum Press, London, New York, 1973, Chapter 5, Part 1, p. 525ff. [80] H. Strehlow: *Electrode Potentials in Non-Aqueous Solvents*, in J. J. Lagowski (ed.): *The Chemistry of Non-Aqueous Solvents*, Academic Press, New York, London, 1966, Vol. I, p. 129ff.

[81] J. Badoz-Lambling and G. Cauquis: *Voltammetry in Non-Aqueous Solvents and Melts*, in H. W. Nürnberg (ed.): *Electroanalytical Chemistry*, Vol. 10 in the series: *Advances in Analytical Chemistry and Instrumentation* (eds. C. N. Reilley and R. W. Murray). Wiley-Interscience, London, 1974, p. 335–419 (especially p. 397ff.). [82] K. Blumrich, H. Schwarz, and A. Wingler, in Houben-Weyl-Müller. *Methoden der organischen Chemie*, 4th ed., Thieme, Stuttgart, 1959, Vol. I/2, p. 887–942. [83] D. Stoye: *Solvents*, in: *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed., Vol. A 24, p. 437ff., VCH, Weinheim/Germany, 1993. [84] E. Browning: *Toxicity and Metabolism of Industrial Solvents*, 2nd ed., Elsevier, Amsterdam, 1965. [85] V. Riihimäki and U. Ulfvarson (eds.): *Safety and Health Aspects of Organic Solvents*, Alan R. Riss, New York, 1986. [86] Berufsgenossenschaft der chemischen Industrie: *Lösemittel*, Merkblatt M 017 (Ausgabe 5/92), Jedermann-Verlag, Heidelberg/Germany, 1992. [87] W. Dekant and S. Vamvakas: *Toxicology*, in: *Ullmann's Encyclopedia of Indiustrial Chemistry*, 5th ed., Vol. B 7, p. 155ff., VCH, Weinheim/Germany, 1995. [88] S. G. Luxon (ed.): *Hazards in the Chemical Laboratory*, 5th ed., Royal Society of Chemistry, Cambridge/U.K., 1992. [89] G. Duve, O. Fuchs, and H. Overbeck: *Lösemittel Hoechst*, 6th ed., Hoechst AG, Frankfurt (Main) 1976, p. 107ff. [90] American Conference of Governmental Industrial Hygienists (ACGIH): *2002 TLV's and BEI's – Threshold Limit Values for Chemical Substances and Physical Agents Biological Exposure Indices*, ACGIH, Cincinnati/Ohio, USA, 2002.

[91] Deutsche Forschungsgemeinschaft, Senatskommission zur Prüfung gesundheitsschädlicher Arbeitsstoffe: *List of MAK and BAT Values 2002*, *Maximum Concentrations and Biological Tolerance Values at the Workplace*, Report No. 38, Wiley-VCH, Weinheim/Germany, 2002. [92] H. Greim (ed.): *Gesundheitsschädliche Arbeitsstoffe – Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten*. 1.–33. Lieferung, 1972–2001, Wiley-VCH, Weinheim/Germany, 2001. [93] B. Meyer: *Low Temperature Spectroscopy*, Elsevier, New York, 1971, p. 191ff. [94] H. E. Hallam (ed.): *Vibrational Spectroscopy of Trapped Species*, Wiley-Interscience, London, 1973, p. 45. [95] S. Cradock and A. J. Hinchcliffe: *Matrix Isolation – A Technique for the Study of Reactive Inorganic*

Species, Cambridge University Press, Cambridge, 1975. [96] M. Moskovits and G. A. Ozin (eds.): *Cryochemistry*, Wiley-Interscience, New York, 1976. [97] D. Stoye (ed.): *Paints, Coatings and Solvents*, VCH, Weinheim/Germany, 1993. [98] H. Gnamm and O. Fuchs: *Lösungsmittel und Weichmachungsmittel*, 8th ed., Vol. I and II, Wissenschaftliche Verlagsgesellschaft, Stuttgart, 1980. [99] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001. [100] A. F. M. Barton: *Handbook of Solubility Parameters and Other Cohesion Parameters*, CRC Press, Boca Raton/Florida, 1985.

[101] J. H. Hildebrand and R. L. Scott: *The Solubility of Nonelectrolytes*, 3rd ed., Dover, New York, 1964. [102] L. R. Snyder: *Solutions to Solution Problems*, CHEMTECH *9*, 750 (1979); ibid. *10*, 188 (1980). [103] N. B. Godfrey: *Solvent Selection via Miscibility Numbers*, CHEMTECH *1972*, 359; *cf.* also I. Vavruch, Chemie für Labor und Betrieb *35*, 385 (1984). [104] W. L. F. Armarego and D. D. Perrin: *Purification of Laboratory Chemicals*, 4th ed., Butterworth-Heinemann, Oxford/U.K., 1996. [105] J. F. Coetzee (ed.): *Recommended Methods for Purification of Solvents and Tests for Impurities*, Pergamon Press, Oxford, 1982; *cf.* also Pure Appl. Chem. *57*, 634, 855, 860 (1985). [106] D. R. Burfield, K.-H. Lee, and R. H. Smithers, J. Org. Chem. *42*, 3060 (1977) (acetonitrile, benzene, 1,4-dioxane). [107] D. R. Burfield, G.-H. Gan, and R. H. Smithers, J. Appl. Chem. Biotechnol. *28*, 23 (1978); Chem. Abstr. *89*, 12551f (1978) (molecular sieves as desiccants). [108] D. R. Burfield and R. H. Smithers, J. Org. Chem. *43*, 3966 (1978) (dipolar aprotic solvents). [109] D. R. Burfield and R. H. Smithers, J. Chem. Technol. Biotechnol. *30*, 491 (1980); Chem. Abstr. *94*, 66822s (1981) (cation exchange resins as desiccants). [110] D. R. Burfield, R. H. Smithers, and A. S. C. Tan, J. Org. Chem. *46*, 629 (1981) (amines).

[111] D. R. Burfield and R. H. Smithers, J. Chem. Educ. *59*, 703 (1982) (diethyl ether). [112] D. R. Burfield and R. H. Smithers, J. Org. Chem. *48*, 2420 (1983) (alcohols). [113] D. R. Burfield, G. T. Hefter, and D. S. P. Koh, J. Chem. Technol. Biotechnol., Chem. Technol. Part A *34*, 187 (1984); Chem. Abstr. *101*, 133117u (1984) (ethanol). [114] D. R. Burfield, J. Org. Chem. *49*, 3852 (1984) (calcium sulfate as desiccant). [115] D. R. Burfield, J. Chem. Educ. *56*, 486 (1979); D. R. Burfield, E. H. Goh, E. H. Ong, and R. H. Smithers, Gazz. Chim. Ital. *113*, 841 (1983) (chloroform). [116] D. R. Burfield and R. H. Smithers, Chem. Ind. (London) 240 (1980) (chloroform, diethyl ether). [117] D. R. Burfield, J. Org. Chem. *47*, 3821 (1982) (deperoxidation of ethers). [118] Sadtler Research Laboratories: *Infrared Spectra Handbook of Common Organic Solvents*, Philadelphia/USA, 1983. [119] J. B. Baumann: *Solvent Selection for Recrystallization*, J. Chem. Educ. *56*, 64 (1979). [120] R. J. Davey: *Solvent Effects in Crystallization Processes*, Curr. Top. Mater. Sci. *8*, 429 (1982); Chem. Abstr. *97*, 154023z (1982).

[121] D. Bierne, S. Smith, and B. E. Hoogenboom, J. Chem. Educ. *51*, 602 (1974). [122] G. J. Eppert: *Flüssigchromatographie – HPLC – Theorie und Praxis*, 3rd ed., Vieweg, Braunschweig, Wiesbaden/Germany, 1997, Chapter 5, p. 83ff. (Elutionsmittel). [123] P. C. Sadek (ed.): *The HPLC Solvent Guide*, 2nd ed., Wiley, New York, 2002. [124] H. Engelhardt: *Hochdruck-Flüssigkeits-Chromatographie*. 2nd ed., Springer, Berlin, 1977; *High Performance Liquid Chromatography* (translated from German by G. Gutnikov), Springer, Berlin, 1979. [125] V. R. Meyer: *Practical High-Performance Liquid Chromatography*, 3rd ed., Wiley, Chichester, 1998. [126] B. Fried and J. Sherma: *Thin Layer Chromatography*, 4th ed., Dekker, New York, 1999. [127] J. C. Touchstone: *Practice of Thin-Layer Chromatography*, 2nd ed., Wiley, New York, 1983; J. C. Touchstone and J. Sherma (eds.): *Techniques and Applications of Thin-Layer Chromatography*, Wiley, New York, 1985. [128] G. Gritzner and J. Kuta: *Recommendations on Reporting Electrode Potentials in Nonaqueous Solvents*, Pure Appl. Chem. *56*, 461 (1984). [129] E. Pungor, K. Tóth, P. G. Klatsmányi, and K. Izutsu (eds.): *Applications of Ion-Selective Electrodes in Nonaqueous and Mixed Solvents*, Pure Appl. Chem. *55*, 2029 (1983). [130] A. J. Collings and S. G. Luxon (eds.): *Safe Use of Solvents*. Academic Press, London, New York, 1982.

[131] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and Willam Andrews Publishing, New York, 2001; Chapter 20 (p. 1315ff.): *Toxic Effects of Solvent Exposure*. [132] D. J. De Renzo (ed.): *Solvents Safety Handbook*, Noyes, Park Ridge/USA, 1986. [133] R. V. Golovnaya and T. A. Misharina: *Thermodynamic Treatment of the Polarity and the Selectivity of Sorbents in Gas Chromatography*, Usp. Khim. *49*, 171 (1980); Russ.

Chem. Rev. *49*, 88 (1980). [134] D. Parker: *NMR Determination of Enantiomeric Purity*, Chem. Rev. *91*, 1441 (1991). [135] H.-U. Blaser: *The Chiral Pool as a Source of Enantioselective Catalysts and Auxiliaries*, Chem. Rev. *92*, 935 (1992). [136] R. Herrmann: *Survey of Chiral Auxiliaries, Solvents, Reagents, and Catalysts*, in: *Methods of Organic Chemistry (Houben-Weyl)*, Vol. E 21f (Stereoselective Syntheses), Thieme, Stuttgart, 1996. [137] A. B. Pangborn, M. A. Girardello, R. H. Grubbs, R. K. Rosen, and F. J. Timmers, Organometallics *15*, 1518 (1996). [138] P. J. Alaimo, D. W. Peters, J. Arnold, R. G. Bergman, J. Chem. Educ. *78*, 64 (2001). [139] K. Fischer, Angew. Chem. *48*, 394 (1935). [140] E. Scholz: *Karl-Fischer-Titration*, Springer, Berlin, 1983.

[141] K. Schöffski, Chem. Unserer Zeit *34*, 170 (2000); GIT Fachz. Lab. *42*, 681 (1998). [142] H. Langhals, Fresenius' Z. Anal. Chem. *305*, 26 (1981); Anal. Lett. *23*, 2243 (1990); GIT Fachz. Lab. *35*, 766 (1991). [143] B. P. Johnson, B. Gabrielsen, M. Matulenko, J. G. Dorsey, and C. Reichardt, Anal. Lett. *19*, 939 (1986). [144] M. F. Vitha, J. Chem. Educ. *78*, 370 (2001). [145] C. Machado and V. G. Machado, J. Chem. Educ. *78*, 649 (2001). [146] P. Blum, G. J. Mohr, K. Matern, J. Reichert, and U. E. Spichiger-Keller, Anal. Chim. Acta *432*, 269 (2001). [147] E. Pretsch, P. Bühlmann, and C. Affolter: *Structure Determination of Organic Compounds – Tables of Spectral Data*, 3rd ed., Springer, Berlin, 2000; p. 157ff. (^{13}C NMR), p. 240ff. (^{1}H NMR), p. 310ff. (IR), and p. 404 (UV/Vis). [148] D. L. Hansen (ed.): *The Sprouse Collection of Infrared Spectra*, Book IV (Common Solvents), Elsevier, Amsterdam, New York, 1990. [149] I. T. Horváth, Acc. Chem. Res. *31*, 641 (1998). [150] D. P. Curran, Angew. Chem. *110*, 1230 (1998); Angew. Chem. Int. Ed. Engl. *37*, 1174 (1998).

[151] R. H. Fish, Chem. Eur. J. *5*, 1677 (1999). [152] B. Betzemeier and P. Knochel, Top. Curr. Chem. *206*, 61 (1999). [153] E. de Wolf, G. van Koten, and B.-J. Deelman, Chem. Soc. Rev. *28*, 37 (1999). [154] A. Endres and G. Maas, Chem. Unserer Zeit *34*, 382 (2000). [155] T. Welton, Chem. Rev. *99*, 2071 (1999). [156] P. Wasserscheid and W. Keim, Angew. Chem. *112*, 3926 (2000); Angew. Chem. Int. Ed. Engl. *39*, 3772 (2000). [157] R. E. R. Craig, J. Chem. Educ. *66*, 88 (1989). [158] K. Zosel: *Praktische Anwendung der Stofftrennung mit überkritischen Gasen*, Angew. Chem. *90*, 748 (1978); Angew. Chem. Int. Ed. Engl. *17*, 702 (1978). [159] S. B. Hawthorne, Anal. Chem. *62*, 633A (1990). [160] B. Wenclawiak and T. Paschke, Nachr. Chem. Tech. Lab. *41*, 806 (1993).

[161] M. E. P. McNally, Anal. Chem. *67*, 308A (1995). [162] C. L. Phelps, N. G. Smart, and C. M. Wai, J. Chem. Educ. *73*, 1163 (1996). [163] J. M. Levy: *Practical Supercritical Fluid Extraction*, Hüthig, Heidelberg, 1996. [164] A. K. Sunol and S. G. Sunol: *Supercritical Solvents*, in: G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrew Publishing, New York, 2001, Chapter 21.1, p. 1419ff. [165] H. Rizvi (ed.): *Supercritical Fluid Processing of Food and Biomaterials*, Blackie Academic and Professional, London, 1994. [166] L. R. Snyder, J. Chromatogr. *92*, 223 (1974); J. Chromatogr. Sci. *16*, 223 (1978). [167] S. C. Rutan and L. R. Snyder, J. Chromatogr. *463*, 21 (1989). [168] E. Klesper, *Chromatographie mit überkritischen fluiden Phasen*, Angew. Chem. *90*, 785 (1978); Angew. Chem. Int. Ed. Engl. *17*, 738 (1978). [169] M. Caude and D. Thiébaut: *Practical Supercritical Fluid Chromatography and Extraction*, Harwood Academic Publishers, Amsterdam, 2000. [170] A. Abbott: *Electrochemistry in Media of Low Dielectric Constants*, Chem. Soc. Rev. *22*, 435 (1993).

[171] D. W. Rooney and K. R. Seddon: *Ionic Liquids*, in: G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001; Chapter 21.2.4 on p. 1472ff. [172] G. Gritzner, Rev. Inorg. Chem. *11*, 81 (1991). [173] G. Gritzner, Pure Appl. Chem. *62*, 1839 (1990). [174] G. Wypych (ed.): *Handbook of Solvents* (+Solvent Database on CD-ROM), ChemTec Publishing, Toronto, and William Andrews Publishing, New York, 2001; Chapter 17 (p. 1149ff.): *Environmental Impact of Solvents*; Chapter 15 (p. 1053ff.): *Methods of Solvent Detection and Testing*; Chapter 22 (p. 1507ff.): *Solvent Recycling, Removal, and Degradation*; Chapter 21 (p. 1419ff.): *Substitution of Solvents by Safer Products and Processes*; Chapter 19 (p. 1267ff.): *Regulations*. [175] (a) N. Kob and G. Altnau, Umwelt-Technologie International (UTI) 1/2000, p. 10–13; GIT Verlag, Darmstadt/Germany; (b) G. Altnau, Eur. Coatings J. 9/2001, 39–40, 42, 44; Chem. Abstr. *136*, 57924 (2002). [176] Ausschuss für Gefahrstoffe: Technische Regeln für Gefahrstoffe (TRGS), TRGS 420, Anhang 1; Bundesarbeitsblatt 9/1999, p. 55ff.

Figure and Table Credits

Fig. 2-1 is reproduced from R. A. Horne, *Survey of Progress in Chemistry 4*, 1 (1968), by permission of the copyright owner, Academic Press, London, New York.

Fig. 2-2 is reproduced from G. Duve, O. Fuchs, and H. Overbeck, *Lösemittel Hoechst*. 6th edition, Hoechst Aktiengesellschaft, Frankfurt (Main), 1976. Used with permission of Hoechst AG, Federal Republic of Germany.

Fig. 2-7 is reproduced from K. L. Wolf, *Theoretische Chemie*. 4th edition, Johann Ambrosius Barth Verlag, Leipzig, 1959, by permission of the copyright owner.

Fig. 3-2 is reproduced from Chemical Reviews *99*, 455 (1999), by permission of the American Chemical Society, Washington D. C., and the author, Prof. A. Baiker, Zürich/Switzerland.

Fig. 3-3 is reproduced from H. H. Sisler, *Chemistry in Non-aqueous Solvents*. Reinhold Publishing Corporation, New York, and Chapman and Hall, London, 1961, with permission of the copyright owners, Litton Educational Publishing Corporation, New York, and the author.

Fig. 4-3 is reproduced from E. J. Drexler and K. W. Field, Journal of Chemical Education *53*, 392 (1976), by permission of the Division of Chemical Education of the American Chemical Society, Tucson/Arizona, and the authors.

Fig. 5-16 is reproduced from J. A. Leary and M. Kahn, Journal of the American Chemical Society *81*, 4173 (1959), by permission of the American Chemical Society, Washington D. C., and the authors.

Fig. 6-3 is reproduced from E. M. Kosower, *Molecular Biochemistry*. McGraw-Hill Book Co., New York, 1962, by permission of the copyright owner and the author.

Fig. 6-5 is reproduced from G. J. Brealey and M. Kasha, Journal of the American Chemical Society *77*, 4462 (1955), by permission of the American Chemical Society, Washington D. C., and the authors.

Fig. 6-6 is reproduced from N. J. Turro, *Molecular Photochemistry*. Benjamin, New York, 1965, by permission of the publishers, Addison-Wesley and W. A. Benjamin Inc., Reading/Massachusetts, and the author.

Table 5-25 is reproduced from M. R. J. Dack, Journal of Chemical Education *51*, 231 (1974), by permission of the Division of Chemical Education of the American Chemical Society, Tucson/Arizona, and the author.

Table A-5 is reprinted from N. L. Alpert, W. E. Keiser, and H. A. Szymanski, *IR – Theory and Practice of Infrared Spectroscopy*. 2nd edition, 1973, by courtesy of Plenum Press Publishing Corporation, New York.

The *MAK-values* of Table A-13 are reproduced with permission of the 'Senatskommission zur Prüfung gesundheitsschädlicher Arbeitsstoffe' of the 'Deutsche Forschungsgemeinschaft', Bonn-Bad Godesberg, Federal Republic of Germany.

The *TL-values* of Table A-13 are reprinted with permission of the 'American Conference of Governmental Industrial Hygienists (ACGIH)', Cincinnati/Ohio, U.S.A.

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt
Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim
ISBN: 3-527-30618-8

Subject Index

a

$a(^{14}N)$ values 414, 415, 437
α-effect of nucleophiles 242
α values of solvent HBD acidity 431–433, 447, 456–461
Acceptor number *AN* 25, 26, 49, 80, 438, 439, 465
Acenaphthylene, photo-dimerization of 296, 301
Acetone 4, 82, 252, 348
Acetonitrile 82, 252
Acetylacetone, keto/enol tautomerism of 107, 108
Acetylchloride/aluminium trichloride adduct, ionization of 51
Acidity constants 99–104
Acidity scales 99–104
Acity of solvents 462–464
Acrylonitrile, as 1,3-dipolarophile 191
Activated complexes 149–153, 309, 310
— dipolar 162, 173–187
— free-radical 163, 199–215
— isopolar 162, 163, 187–199
— solvent activity coefficients 254–258
Activity coefficients 94, 98, 99
1- and 2-Adamantyl tosylate, solvolysis of 274, 403, 405–407
Addition reactions, solvent effects on 176–178, 254, 278, 279
Alcohols, acidity and basicity of 102, 103
Aldol addition reaction, solvent effects on 183, 184
Alkahest 1
Alkene isomerization reaction, solvent effects on 253
t-Alkoxy radicals, decomposition of 212, 213
Alkyl amines, basicity anomaly of 104–106
O/*C* Alkylation ration, solvent effects on 269–273
Alkyl chlorosulfites, thermolysis of 187
Alkylidenephosphoranes, Wittig reaction of 194, 195, 260, 262
Allopolarization principle 273
Allyl silylmethyl ethers, dyotropic rearrangement of 199
Allyl thionbenzoates, sigmatropic rearrangement of 197
Allyl 4-tolyl ether, Claisen rearrangement of 196, 197
Allyl 4-tolyl sulfoxide, thermal racemization of 184, 449, 450
Ambident anions 269–273
Ambident cations 269, 273
Ambifunctional anions 269–273
α-Amino acids, intramolecular proton-transfer equilibrium 103, 104
4-Amino-5-methylacridine, proton-transfer equilibrium of 122, 123
Amphiphiles 42, 292–298
Amphiprotic solvents 18, 75, 96
Anomeric effect, solvent-dependence of 131
Anthracene, Diels–Alder cycloaddition reaction of 188
9-Anthrone, keto/enol tautomerism of 112, 113
Antimony pentachloride/adduct formation with EPD solvents 22–24
Apolar non-HBD solvents 82, 83, 88, 89
Apolar solvents 66, 67, 88, 89, 425
Aprotic solvents 18, 75, 82, 83, 88, 89
Arabinose, ring/chain tautomerism of 119
Aromatic compounds, 330
— π-electronic structure 330, 331
— UV/Vis spectra 331
Aromatic solvent-induced shift (ASIS),
— camphor 385
— 2-methylcyclohexanone 384
— pulegone 384, 385
— with hexafluorobenzene 386
Arrhenius theory of reaction kinetics 149
1-Arylethyl tosylates, solvolysis of 275
Atom-transfer reactions 163, 208–213, 223, 224
Autoprotolysis constants 74, 75, 496–499
Autoxidation reaction of hydrocarbons 210
7-Azidocycloheptatriene/tropylium azide, ionization equilibrium 51, 125
3-Azidopyrazine-1-oxide, valence isomerization of 135
2-Azidothiazole, valence isomerization of 135
Azoalkanes, thermolysis of 201, 203, 204, 305, 306
Azobisisobutyronitrile, thermolysis of 200, 201, 210, 305, 306
Azo coupling reactions, solvent effects on 174, 175
Azomethane, photolysis of 305
[1,1′]Azonorbornane, *cis*/*trans*-isomerization of 203

b

B values of Lewis basicity 453–455
B_{MeOD} values 436, 454, 455
B_{PhOH} values 436
β values of solvent HBA basicity 431–434, 456–460

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt

Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

ISBN: 3-527-30618-8

Basicity constants 98
Basicity of nucleophiles 244
Basicity scales 99–106
Basity of solvents 462–464
Basketane, valence isomerization of 61
Beckman–Chapman rearrangement of ketoximes, solvent effects on 411
Bellamy–Hallam–Williams (BHW) plot 367
Benzaldehyde, enantioselective additions to 69, 70, 289
Benzenediazo(aryl)sulfanes, thermolysis of 284
Benzonitrile 82
Benzophenone, UV/Vis spectrum 349, 350
1,4-Benzoquinone/1,4-hydroquinone, electron-transfer reaction 137, 138
1,4-Benzoquinone monoxime, tautomerism of 118
1,3,2-Benzoxazaphospholine, ring/chain tautomerism of 119
2-Benzoylbenzoic acid, induced CD 362
3-Benzoyl camphor, keto/enol tautomerism of 110
Bilayers 46
Binary solvent mixtures, 38–42
— $E_T(30)$ values 425–428
— equi- or iso-solvation point 40, 41
— preferential or selective solvation 38–42, 426, 427
— for recrystallization 488, 490, 491
— synergistic behaviour 42, 426
1,2-Bis(methoxycarbonyl)-3,4-bis(2-naphthyl)cyclobutane, cycloreversion of 282
1,2-Bis(trifluoromethyl)-1,2-dicyanoethene, cycloaddition of 180
Boiling points of solvents 60, 63, 472–477, 491, 504
Born equation 96, 235
Born–Haber cycle for solvation 32
Boron trifluoride, adduct formation with EPD solvents 24, 398
Brønsted acid/base interactions 73–79, 95–99
Bromine addition to alkenes, solvent effects on 176, 278, 279
1-Bromoadamantane, solvolysis of 165, 166, 403
1-Bromobutane, S_N2 reaction of 168, 234, 235, 249, 263, 267, 323
(2-Bromoethyl)benzene, Menschutkin reaction of 229, 230
Bromomethane, S_N2 reaction of 405
2-Bromo-2-methylpropane, acetolysis of 240
Bromonium ion as intermediate 176, 278
N-Bromosuccinimide, radical chain bromination with 212, 326
1-Bromo-*n*-undecane, chemoselective reduction of 287
Buckingham equation 368
Butenone, Diels–Alder cycloaddition reaction of 189
3-*t*-Butylaminopropenal, *cis*/*trans* equilibrium of 133
n-Butyl 4-bromobenzene sulfonate, S_N2 reaction of 263
t-Butyl chloride, solvolysis of 165, 215, 216, 231, 232, 313, 314, 402–406, 458, 459
t-Butyl cycloheptatrieneperoxycarboxylate, thermolysis of 205, 207
t-Butyl heptafluorobutyrate, solvolysis of 274, 403
t-Butyl heptafluoroperoxybutyrate, thermolysis of 286
t-Butyl hypochlorite, chlorination with 69, 212
t-Butyl peroxyformate, thermolysis of 185
trans-3-*t*-Butyl-4-phenyloxetan-2-one, decarboxylation of 182
t-Butyl 2-phenylthioperbenzoate, thermolysis of 206
t-Butyl 2-picolyl ketone, keto/enol tautomerism of 113
n-Butyl stearate as smectic solvent 299–302
n-Butyl vinyl ether, cycloaddition of 179–181
t-Butyl vinyl thioether, cycloaddition of 180

c

Cage ion pairs 53
Cage reactions 303–308
Carbon dioxide as supercritical-fluid solvent 70–73, 186, 315, 324–327, 492, 494, 504, 505
Carbon disulfide,
— index of refraction 14
— photochlorination reaction in 211, 212
— reaction with tri(*n*-butyl)phosphane 125, 168
Cation solvators 262–269
Cavitands as cation solvators 266–268
Cesium effect 264
n-Cetyl trimethylammonium bromide (CTAB), micellar catalysis by 44, 292, 293, 297
χ_B values 414, 415, 429
χ_R values 414, 415, 429
C—H acids, gas-phase acidity of 101, 102
Cheletropic reactions, solvent effects on 198, 199
Charge-transfer absorption 19, 20, 330, 338–340
Charge-transfer complexes 19–27
Charge-transfer-to-solvent (CTTS) absorption 340

Chemometrics 84, 395
Chemoselectivity, solvent effects on 287, 288
Chiral solvents 69, 70, 289, 361, 362, 386, 387, 476, 477
Chloroacetaldehyde, rotational equilibrium of 126, 127
α-Chlorobenzyl methyl ether, thermolysis of 187, 325
2-*exo*-Chlorobornane, Wagner–Meerwein rearrangement to 166
2-Chlorocyclohexanone, conformational equilibrium of 129
1-Chloro-2,4-dinitrobenzene, S_NAr reaction of 173, 174
Chloro-(2,4-dinitrophenyl)sulfane, electrophilic addition to cyclohexene 176, 177
Chloroethene, preparation of 61
(+)-*trans*-2-Chloro-5-methylcyclohexa-none, conformational equilibrium of 129
6-Chloro-2-hydroxy-4-methylpyridine, lactam/lactim tautomerism of 114
2-Chloro-2-methylpropane, solvolysis of 165, 215, 216, 231, 232, 313, 314, 402–406, 458, 459
Chloro-oxalato-tripyridine-rhodium(III) complex, selective solvation of 41, 42
Chlorosulfonyl isocyanate, cycloaddition to alkenes 181, 182
3-Chloro-2,3,3-trimethylnorbornane, Wagner–Meerwein rearrangement of 166
Chloro-triphenylmethane, ionization and dissociation of 49, 50
Cholesteryl benzoates as nematic solvents 302, 303
Cholesteryl tosylate, nucleophilic substitution of 274
Chromophore, definition 329
Circular dichroism (CD), solvent effects on, 359–362
— benzil 361, 362
— benzophenone 361, 362
— 2-benzoylbenzoic acid 362
— (*R*)-5-methylpyrrolidine-2-thione 360
Cis/*trans*-equilibria, solvent effects on,
— 3-*t*-butylaminopropenal 133
— 2,3-dibenzoyl-1-benzylaziridine 134
— 1,2-dichloroethene 67, 132, 133
— (ethoxycarbonyl)methylene triphenyl-phosphorane 127, 128
— formylbenzylidene triphenyl-phosphorane 128
ortho-Claisen rearrangement, solvent effects on 196, 197, 303
CODESSA program 69, 90, 91
Cohesion parameter 63, 64
Cohesive energy density 63, 64, 219, 220
Cohesive pressure 63, 64, 219, 220, 443, 444, 457
Conformational equilibria, solvent effects on, 126–132
— chloroacetaldehyde 126, 127
— 2-chlorocyclohexanone 129
— (+)-*trans*-2-chloro-5-methylcyclo-hexanone 129
— *trans*-1,2-dihalocyclohexanes 129
— (ethoxycarbonyl)methylene triphenyl-phosphorane 127, 128
— 2-, 3-, and 4-fluorobenzoates 128
— furfural 128
— 2- and 4-halocyclohexanones 129
— 2-isopropyl-5-methoxy-1,3-dioxane 130, 131
— 4-methoxycyclohexanone 129
— phencyclidine 129, 130
— 1-phenyl-1-(1-piperidinyl)cyclohexane 129, 130
Contact ion pairs 52–55
Coordinating solvents 22
Coordination number 35
Cope elimination reactions, solvent effects on 261
Cope rearrangements, solvent effects on,
— 2-allyl-2-(1-ethyl-1-propenyl)-malononitrile 196
— arylazo(α,α-dimethylallyl)-malononitrile 196
— 2,6-dicyano-4,8-diphenylbarbaralane 196
— diethyl allylisopropenylmalonate 196
— 1,5-hexadienes 163
Coronands as cation solvators 266–268, 271
Cosphere 37
Cotton effect, solvent effects on 359–362
Coulomb interaction forces 10, 11, 47, 233–235
Critical micelle concentration (*cmc*) 43, 44, 292
Crown ethers as cation solvators 266–268, 271
Cryptands as cation solvators 266–268, 271
Cuprate, lithium dimethyl-, ion-pair equilibrium of 56
Cut-off points of spectroscopic solvents 479, 482
Cyanine dyes, definition 330, 331
Cybotactic region 37
Cycloaddition reactions, solvent effects on 178–182, 280–282, 294–296
Cyclohexene,
— autoxidation reaction of 210
— electrophilic addition to 176, 177

Cyclopentadiene, Diels–Alder cycloaddition reaction of 155, 156, 189, 190, 223, 280, 281, 291, 295, 327, 410
1,3-Cyclopentadienyl-trimethylstannane, metallotropic tautomerism of 120
2-Cyclopenten-1-one, photodimerization of 281, 282
Cyclopropanone, ring-opening reaction 201
Cycloreversion reactions, solvent effects on 182, 183, 280, 282

d

trans-9-Decalyl peroxybenzoate, thermolysis of 205, 207
trans-9-Decalyl peroxyphenylacetate, thermolysis of 205, 207, 285, 286
all-cis-Deca-2,4,6,8-tetraene, electrocyclic reaction of 198
Dediazoniation of arenediazonium salts, solvent effects on 285
Desmotropic constant 396
Desoxybenzoin-phenylimine, imine/enamine tautomerism of 116
Diacetyl peroxide, thermolysis of 155, 201, 306
Diacyl peroxides, thermolysis of 155, 201, 202, 306
Dialkyl peroxides, thermolysis of 155, 201, 202, 213, 215, 216, 286
Diastereoselectivity, solvent effects on 288–291
1,4-Diazabicyclo[2.2.2]octane, Menschutkin reaction of 229, 230, 241
Diazo-diphenylmethane,
— 1,3-dipolar cycloaddition reaction of 191, 312, 410, 411
— reaction with carboxylic acids 455, 458
Diazomethane, 1,3-dipolar cycloaddition reaction of 191
2,3-Dibenzoyl-1-benzylaziridine, *cis*/*trans* equilibrium of 134
Dibenzoyl peroxide, thermolysis of 201, 202
Dibenzyl peroxydicarbonate, thermolysis of 286
Dibromomethane, atom-transfer reaction of 208, 209
erythro-1,2-Dibromo-1-(4-nitrophenyl)-2-phenylethane, debromination/dehydrobromination of 287
2,6-Di-*t*-butyl-4-*i*-propylphenoxy radical, disproportionation of 214
Di-*t*-butyl peroxide, thermolysis of 155, 201, 215, 216
1,3-Dicarbonyl compounds, keto/enol tautomerism of 3, 4, 106–111, 396
1,2-Dichloroethene, *cis*/*trans*-equilibrium of 67, 132, 133
Dielectric constant (relative permittivity) 3, 47, 48, 67, 68, 226, 227, 454, 472–475
Diels–Alder cycloaddition reactions, solvent effects on 155, 156, 163, 187–191, 223, 280, 281, 291, 294–296, 302, 311, 312, 315, 410
Diethyl azodicarboxylate as enophile 192
Diethyl sodio-*n*-buylmalonate, alkylation of 264, 265
Differentiating solvents 76
Diffusion-controlled reactions 304
3,3-Difluoro-4,4-dimethyloxetan-2-one, decarboxylation of 183
8,8-Diformylheptafulvene, valence isomerization of 136
trans-1,2-Dihalocyclohexanes, conformational equilibrium of 129
2,3-Dihydro-4-methyl-4*H*-pyrane, photooxygenation of 283
Di-*i*-butyryl peroxide, thermolysis of 201, 202
Dimcarb 60, 62
Di-*i*-propyl peroxide, thermolysis of 202, 203
2,2-Dimesitylethenol, keto/enol tautomerism of 109
N,*N*-Dimethylacetamide 82, 252
Dimethyl acetylenedicarboxylate as 1,3-dipolarophile 191, 312
3-(Dimethylamino)propenal, solvent effects on,
— IR spectrum 343
— ^{1}H NMR spectrum 343, 378, 379
— ^{13}C NMR spectrum 343, 378, 379
— ^{15}N NMR spectrum 343, 378, 379
4-(Dimethylamino)benzenethiyl radical, addition to α-methylstyrene 213, 214
4-(Dimethylamino)-4′-nitroazobenzene, *cis*/*trans* isomerization of 185, 186
3-(Dimethylamino)propynal, nucleophilic addition to 178
cis-2,2′-Dimethyl-[2,2′]azopropane, thermolysis of 203
2,6-Dimethyl-1,4-benzoquinone, Diels–Alder cycloaddition reaction of 295
2,3-Dimethylbutane, photochlorination of 210–212
5,5-Dimethylcyclohexan-1,3-dione, keto/enol tautomerism of 108, 109
2,5-Dimethyl-3,4-diphenylcyclopenta-dienone, Diels–Alder cycloaddition reaction of 269, 270
N,*N*′-Dimethylethylene urea (DMEU) 252, 504
N,*N*-Dimethylformamide (DMF) 82, 252
1,3-Dimethylimidazolidin-2-one (DMEU) 252, 504

Dimethyl fumarate as 1,3-dipolarophile 191
Dimethylketene, cycloaddition reaction of 181, 280
2,2-Dimethylpropenal, enantioselective addition of phenylmagnesium bromide to 69
N,*N*′-Dimethylpropylene urea (DMPU) 82, 252, 504
Dimethyl sulfone 82, 252
Dimethyl sulfoxide (DMSO) 82, 252
S,*S*-Dimethylsulfoximine (DMSOI) 253
2,4-Dinitrophenyl sulfate, aminolysis of 297
1,3,2-Dioxathiolane-2-oxide 252
1,3-Dioxolan-2-one 253
γ,γ-Diphenylallylcarbinyl lithium, valence isomerization of 136, 137
Diphenylcyclopropenylcarbinyl lithium, valence isomerization of 136, 137
Diphenylketene, cycloaddition of 180, 181, 193
Dipolar aprotic solvents 18, 82, 83, 88, 89, 240, 244, 245, 252, 253, 425
1,3-Dipolar cycloaddition reactions, solvent effects on 191, 192, 296, 312, 410, 411
Dipolar molecules, definition 11, 66, 67, 225
Dipolar non-HBD solvents 18, 82, 83, 88, 89, 239, 240, 245, 246, 252, 253, 425
Dipolar solvents 11, 66, 67, 75, 82, 83, 252, 253, 425
Dipolar transition-state reactions 162, 173–187, 225–232, 233, 234
Dipole moment, induced 13, 14
Dipole moment, permanent 11–13, 66–68, 252, 253, 472–475
Di-(2-quinolyl)methane, tautomerism of 115, 116
Dispersion forces 13, 14
Dissociating solvents 47, 52
Dissociation,
— definition 46, 52
— dissociating solvents 47, 52
— equilibria 46–56
— ionogens 46
— ionophores 46
— radical anions 55
— solvent effects on 46–56
Dissociation of ions, solvent effects on 41–56
N,*N*′-Distearoylindigo, *cis*/*trans* isomerization of 299
Donor number *DN* 22–24, 49, 80, 397, 398, 462, 465
Drago equation of Lewis acidity/basicity 25, 27
Drude–Nernst equation 309, 310
Dyotropic reactions, solvent effects on 199

e
E values of Lewis acidity 453–455
E_B^N values 351, 430
E_K values 430
E^*_{MLCT} values 430
$E_T(30)$ values 416–429, 446–451, 454, 462
E_T^N values 418–425, 427, 447, 448, 467, 472–475, 491
E_T^{SO} values 430
Electrochromism 342
Electrocyclic reactions, solvent effects on 197, 198
Electron affinity 20, 21
π-Electron systems, classification of 330, 331
Electron-transfer reactions, solvent effects on,
— bis(dimethylaminophenyl)-phenylcarbenium tetrafluoroborate/sodium tris-(4-nitrophenyl)methide 138
— 1-ethyl-4-(methoxycarbonyl)pyridinyl radical 137, 209
— 1,4-benzoquinone/1,4-hydroquinone 137, 138
— radicals 208–210
— S_N2 reactions 164
— time-dependent fluorescence 356
Electrophilic solvent assistance 274, 275
Electrostatic factor *EF* 67, 68
Electrostriction 309, 310
β-Elimination reactions, solvent effects on 169–172, 253, 261, 262, 279, 280, 286, 287
Eluotropic solvent series 494, 495
Empirical parameters of solvent polarity,
— $a(^{14}N)$ values 414, 415, 437
— α values of solvent HBD acidity 431–433, 447, 456–461
— acceptor number *AN* 25, 26, 49, 80, 438, 439, 465
— *B* values of Lewis basicity 453–455
— B_{MeOD} values of Lewis basicity 436, 454, 455
— B_{PhOH} values of Lewís basicity 436
— β values of solvent HBA basicity 431–434, 456, 460
— χ_B values 414, 415, 429
— χ_R values 414, 415, 429
— D_π values 411
— D_S and D_H values 81
— desmotropic constant 396
— donor number *DN* 22–24, 49, 80, 397, 398, 462, 465
— *E* values of Lewis acidity 453–455
— E_B^N values 351, 430
— E_K values 430
— E^*_{MLCT} values 430
— $E_T(30)$ values 416–429, 446–451, 454, 462

Empirical parameters of solvent polarity (cont.)
— E_T^N values 418–425, 427, 447, 448, 467, 472–475, 491
— E_T^{SO} values 430
— Φ values 414, 415, 429, 430
— *G* values 414, 415, 435
— ΔH°_{D-BF3} values 24, 398
— $\Delta I_{50}^{Cl/Br}$ values 444
— lg k_{ion} values 404, 408, 409
— μ values 81, 401, 402
— *N* values 405–407
— Ω values 404, 410
— *P* values 437
— *Py* values 358, 430
— π^* values 431–434, 447, 456–461
— π^*_{azo} values 432
— *RPM* values 429
— *S* values 413–416
— *SA* values 441–443, 461
— *SB* values 439–441, 461
— *SI* values 401
— *SPP* values 439–441, 461
— *Sp* values 295, 400, 401
— *SP* values 81, 402
— $\mathscr{S}$ values 404, 409
— "*W*" values 409
— *X* values 404, 409, 410
— ζ values 457, 458
— *Y* values 402–408, 453, 454
— Y_X values 406, 407
— *Z* values 412–416, 446, 447
Enantioselectivity, solvent effects on 69, 70, 288, 289
Ene reactions, solvent effects on 192, 193, 282, 283
Enolate ions, *O*/*C* alkylation of 269–273
Enol constant 396
α-Enones,
— ambident reactivity 273
— chemoselective reactivity 288
Enzyme-catalyzed reactions 143–145, 327
Eosin, intramolecular ionization of 124
EPA solvents 21–27, 48, 80, 81, 88, 89
EPD solvents 21–27, 48, 80, 81, 88, 89, 264, 265
EPD/EPA complexes 19–27, 79, 80, 338, 397, 398
Epoxidation of alkenes, solvent effects on 177
Equilibria, solvent effects on,
— amino/imino tautomeric 116, 117
— association 46–56, 121, 123
— azo/hydrazone tautomeric 3, 117, 118, 325
— Brønsted acid/base 73–79, 95–99, 121–123
— *cis*/*trans* isomerization 132–135
— conformational 126–132
— dissociation 46–56, 121, 123
— electron-transfer 137–139
— gas-phase acid/base 99–106
— hoest/guest complexation 139–145
— imine/enamine tautomeric 116
— ionization 46–56, 95, 121–123
— keto/enol tautomeric 3, 4, 106–113
— lactam/lactim tautomeric 113–115
— Lewis acid/base 79–81, 121–123
— metallotropic 120
— nitro/isonitro tautomeric 3
— nitrone/hydroxylamine tautomeric 117
— nitroso/oxime tautomeric 118
— ring/chain tautomeric 118, 119
— rotational 126–128
— tautomeric 106–120
— valence isomerization 135–137
Equilibrium constants, thermodynamic 98
Equisolvation point 40, 41
E_1 and E_2 reactions, solvent effects on 169–172, 253, 279, 280
ESR spectra, solvent effects on, 369–375
— 4-acetyl-1-methyl-1,4-dihydropyridinyl 373, 374, 437
— acyl aminyloxides 370
— azomethine aminyloxides 370
— 1,2- and 1,4-benzoquinone anion radicals 370, 372
— dialkyl and diaryl aminyloxides 370, 414, 415, 437
— 2-[4-(dimethylamino)phenyl]indane-1,3-dionyl 374
— ketyls 370, 372, 373
— 4-(methoxycarbonyl)-1-methylpyridinyl 369, 370, 373
— nitroaromatic anion radicals 370
— phenoxyl radicals 370
— semidiones 370
— sodium 9-fluorenone ketyl 372, 373
— sodium naphthalenides 55, 375
— 2,2,6,6-tetramethylpiperidine-1-oxyl 437
— 2,2,6,6-tetramethyl-4-piperidone-1-oxyl 370, 371
— thioindigo radicals 370
Esterification, solvent effects on 2, 3
(α-Ethoxycarbonyl)benzylmercuric bromide, isotope exchange reaction of 276, 277
Ethyl acetoacetate, keto/enol tautomerism of 3, 106–108, 396
Ethyl acrylate as 1,3-dipolarophile 191, 192
Ethylene carbonate 253
Ethylene sulfite 252
1-Ethyl-4-(methoxycarbonyl)pyridinium iodide,
— solvatochromism of 334, 338, 339, 411–415

— solvent polarity indicator 339, 411–415
— UV/Vis spectrum 338
1-Ethyl-4-(methoxycarbonyl)pyridinyl radical,
— atom-transfer reaction 208–210
— electron-transfer reaction 137
1-Ethyl-3-methylimidazolium (EMIM) salts 60, 322, 323, 387
Ethyl-phenylmalonic acid, decarboxylation of 303
Eutectics 60, 61
Evaporation number 63
Excited states, electronic
— carbonyl group 348–351
— dipole moments 335–337, 354
— solvation of 342, 353
External ion pair 53
External pressure 308–315
Extrathermodynamic relationships 33, 255
Eyring equation 149–152

f
Φ values 404, 410
Factor analysis (FA) 84, 85
Finkelstein halide exchange reaction, solvent effects on 168, 240
Fluorescein, intramolecular ionization of 124
Fluorescence spectra, solvent effects on, 352–359
— 4-aminophthalimide 357, 430
— 6,2-(arylamino)naphthalenesulfonates 355
— 2,5-bis[3-(*N*-methyl-*N*-phenylamino)-propen-1-yliden]cyclopentanone 357
— 7-dialkylamino-4-trifluoromethyl-coumarins 354
— 4-(*N*,*N*-dimethylamino)benzonitrile 354, 355
— 2-(dimethylamino)-7-nitrofluorene 357
— 2-(dimethylamino)-6-propanoyl-naphthalene (PRODAN) 357
— 2-[4-(dimethylamino)styryl]benzoxazol 357
— 2-(2-naphthoxy)ethyl 4-cyanobenzoate 357
— 8-phenylamino-1-naphthalenesulfonates 355
— 1-phenyl-4-(4-cyano-1-naphthyl-methylene)piperidine 356
— pyrene 358, 430
2-, 3-, and 4-Fluorobenzoates, conformational equilibrium of 128
1-Fluoro-2,4-dinitrobenzene, S_NAr reaction of 293
4-Fluoro-nitrobenzenes, S_NAr reaction of 248, 249
Fluorous biphase catalysis (FBC) 320–322
Fluorous biphasic systems (FBS) 4, 57, 58, 320–322
Fluorous solvents 4, 57, 58, 319–322, 504
Fragmentation reactions, solvent effects on 162, 185, 204, 205, 284, 285
Franck–Condon principle 341, 342, 348, 353
Free-radical transition state reactions 163, 199–215
Free radicals, generation of 200–215
Friedel–Crafts acylation reactions 61, 323
Fullerene C_{60}, solubility of 45, 468
Furfural, rotational equilibrium 128
2-Furfuryl methyl fumarate, intramolecular Diels–Alder cycloaddition of 189, 190

g
G values 414, 415, 435
Gas-phase acidities and basicities 99–106
Gas-phase proton affinities 101
Gas-phase reactions 99–106, 147, 148, 155–162, 245, 272, 276, 277
Gibbs energy diagrams 94, 104, 150, 153, 157, 160, 179, 183
Gibbs energy of hydration 32
Gibbs energy of solvation 31–33, 218, 254
Gibbs energy of transfer 33, 34, 93, 94, 254, 256, 468
Green chemistry 317, 325
Ground states, electronic,
— carbonyl group 348–351
— dipole moments 11, 335–337, 341, 342
— solvation of 342, 353
Grunwald–Winstein equation 402–407

h
Halide nucleophilicities 158, 159, 244–247, 263
Haloacetic acids, acidity of 103
Halochromism 428, 429
2- and 4-Halocyclohexanones, conformational equilibrium of 129
Halogen addition reactions, solvent effects on 176, 278
4-(Halomethyl)-nitrobenzenes, electron-transfer reactions of 209
1-Halo-1-phenylethanes, phenolysis of 240
Halo-solvatochromism 428, 429
Hammett acidity function H_0 77
Hammett equation 85, 390, 392–394
Hard and soft acids and bases 79–81, 244, 272, 273, 401, 402
HBA solvents 18, 88, 89
HBD solvents 18, 88, 89, 238–244, 425
Heavy water, properties of 316
n-Heptadecyl phenyl ketone, photolysis of 300
Hexamethyl phosphoric acid triamide (HMPT) 11, 82, 253, 504

"Hexaphenylethane", dissociation of 200, 201, 222, 224
Homofuran, Diels–Alder cycloaddition reaction of 281
Homomorphic molecules 431
Host/guest complexes 139–145,
— [18]crown-6/potassium thiocyanate 140
— α-cyclodextrin/methylorange 141
— cyclophanes/pyrene 141, 142
— 1,10-diaza-[18]crown-6/silver ion 140
— influence on reaction selectivities 142–144
— polyethers/benzenediazonium salts 140, 141
HSAB concept 79–81, 244, 272, 273
Hughes–Ingold rules 163–173, 215–217
Hydration,
— definition 30
— negative 37, 38
— positive 37
Hydrochloric acid, interaction forces of liquid 14
Hydrogen bonding, 15–19
— anion solvation by 18, 80
— antidromic 16
— circular 16
— dissociation enthalpy 16
— ESR spectra, influence on 371
— heterodromic 16
— hetero-intermolecular 15
— homodromic 16
— homo-intermolecular 15
— intermolecular/intramolecular 15
— IR spectra, influence on 364–367
— nature of 15–19
— NMR spectra, influence on 78, 381–384
— protomerism 17, 18
— solvation by 18
— type A/type B 18
— UV/Vis spectra, influence on 348–351
Hydrogen halides, acidity of 102
Hydrophobic effect 29–31, 189
Hydrophobic hydration 29–31
Hydrophobic interactions 29–31, 292–296, 400
Hydrophobicity substituent parameters (π_X, *MH*) 30, 400
6-Hydroxy-4,8-dimethylazulene, keto/enol tautomerism of 112
2-Hydroxy-7-isopropyl-1,4-dimethylazulene, keto/enol tautomerism of 112
9-(Hydroxymethyl)anthracene, Diels–Alder cycloaddition of 294, 295
2-Hydroxy-2-methyltetrahydrofuran, ring/chain tautomerism of 119
5-Hydroxypentan-2-one, ring/chain tautomerism of 118, 119
(2-Hydroxyphenylimino)phosphorane, ring/chain tautomerism of 119
5-Hydroxy-1-phenyl-1,2,3-triazole-4-carboxylic esters, rearrangement of 154
2-Hydroxypyridine, lactam/lactim tautomerism of 113, 114

i
$\Delta I_{50}^{Cl/Br}$ values 444
1-Indanone, synthesis of 61
Index of refraction 4, 68, 154
Infrared comparison method 436, 437
Interaction forces, intermolecular, 10–30
— and solvent polarity 8, 68
— Coulomb 10, 11, 47, 225, 235
— dipole-dipole 11, 12, 225
— dipole-induced dipole 13
— dispersion 13, 14
— EPD/EPA 19–27, 48
— hydrogen bonding 15–19
— hydrophobic 27, 28, 43, 292–296, 400
— instantaneous dipole-induced dipole 13, 14
— ion-dipole 10, 11, 233
— ion-ion 47, 234–237
— solvophobic 27–30, 292–296, 400
Internal ion pair 53
Internal pressure 63–65, 220–224
Intimate ion pair 53
2-Iodobutane, dehydrohalogenation of 261, 262
Iodoethane, S_N2 reactions of 167, 186, 227, 228, 312, 313, 451
Iodomethane, S_N2 reactions of 166–168, 241, 249, 257, 313, 314, 392, 409
(Iodomethyl)benzene, S_N2 reactions of 240, 241
Ionic liquids 4, 58–62, 322–324, 387, 504
Ionic strength 236
Ionization, 46–56
— haloalkanes 48
— definition 46
— equilibria 46–56, 100, 123–126
— ionogens 46, 121
— ionophores 46
— solvent effects on 46–54
Ionization energy 14, 121
Ionization of covalent bonds, solvent effects on 46–54, 123–126
Ionizing power of solvents 48, 52, 53, 402
Ionizing solvents 48, 52, 53
Ionogens 46, 121
Ionophores 46
Ion pairs,
— anion-nucleophilicity of 262, 263
— contact 53–56, 262, 286, 387

— definition 46
— ligand-separated 263, 266, 267
— penetrated 52
— product-separated 286
— solvent-separated 52–55, 262
— solvent-shared 52, 53
— supramolecular 55, 56
— tight and loose 53
IR spectra, solvent effects on, 363–369
— absorption intensities 369
— acetone 364, 365
— acetonitrile 365
— acetophenone 365
— 1-alkynes 365
— benzophenone 365, 435
— *t*-butyl hydroperoxide 365
— camphor 366
— carbonyl compounds 364–367
— 1-chloropropane 364
— *N*-cyclohexylpyrrolidin-2-one 365
— deuterio-trichloromethane 369
— deuteriomethanol 364, 367, 436
— 1,1-dichloroethene 363
— 3-(dimethylamino)propenal 343
— *N*,*N*-dimethylformamide 365, 435
— dimethyl sulfoxide 365, 435
— haloalkanes 365
— *N*-methylacetamide 365
— *n*-octane 365
— phenol 367, 436
— phenylacetylene 78
— 4-pyridones 365
— pyrrol 365
— silanes 365
— trichloroacetic acid 436, 437
— trichloromethane 78, 365
— trialkylphosphane oxides 365
— tropolone ands tropone 365
— urea 365
N-Isobutenylpyrrolidine, cycloaddition reaction of 181, 280
Isodielectric solvent mixtures 40
cis/*trans*-Isomerization reaction of
— alkenes 132, 133
— aziridines 134
— azoalkanes 203, 204
— azobenzenes 185, 186
— imines 291, 292
Isopolar transition-state reactions 162, 163, 187–199, 219–225
Isoprene,
— alkenylation reaction of 260
— Diels–Alder cycloaddition reaction of 188, 312, 315
2-Isopropyl-5-methoxy-1,3-dioxane, conformational equilibrium of 130, 396
4-Isopropylpyridine, base-catalyzed reaction with isoprene 260
Iso-solvation point 40, 41
7-Isothiocyanatocycloheptatriene/tropylium/isothiocyanate ionization equilibrium 125

k
Karl–Fischer titration 478
Kauri–Butanol (*KB*) numbers 66
α-Ketoamides, diastereoselectice reduction of 290
Kinetic solvent isotope effects (KSIE) 315–317
Kirkwood–Bauer–Magat (KBM) equation 367
Kirkwood equation 111, 225, 226
Kundt's rule 4

l
Levelling solvents 76
Lewis acid/base interactions 21–27, 79–81, 123–126, 397, 398
lg k_{ion} values 404, 408, 409
Ligand-to-metal charge-transfer (LMCT) absorption, solvatochromism of 339, 340
"Like dissolves like" rule 1, 6, 57, 80, 490, 492
Linear Gibbs energy (LGE) relationships 85, 391–396, 443, 452
Linear solvation energy relationships (LSER) 432, 456–461
Lippert–Mataga equation 359
Liquid crystals as solvents 58, 59, 298–303, 306, 386
Liquid metals as solvents 58, 62
Lithium dimethylcuprate, ion-pair equilibrium of 56
Lithium perchlorate in diethyl ether (LPDE) 62, 189
Lithium 10-phenylnonafulvene-10-oxide, valence isomerization of 55, 137
Lithium triphenylmethide, dissociation of 55
Lyate ion 74, 497, 498
Lyonium ion 74, 497, 498
Lyotropic liquid-crystalline solvents 58

m
μ values 81, 401, 402
MAK values 500–503
Maleic anhydride, Diels–Alder cycloaddition reaction of 188, 312, 315
Matrix isolation, solvents for 488
Melting points of solvents 60, 472–477
Menschutkin reaction, solvent effects on 2, 3, 121, 166–168, 215, 228–230, 312, 313, 409, 451

Menstruum universale 1
2- and 4-Mercaptopyridine, lactam/lactim tautomerism of 114
Merocyanine dyes,
— definition 331
— π-electronic structure 330, 331, 342–344
— solvatochromism of 331, 335–337, 342, 343
Meropolymethines,
— definition 330, 331
— π-electronic structure 330, 331, 342–344
— solvatochromism of 331, 335–337, 342, 343
Mesomorphic solvents 58, 59, 298–303, 306, 386
Metal-to-ligand charge-transfer (MLCT) absorption, solvatochromism of 339, 340, 430
Metastable liquids 73
N-(4-Methoxybenzylidene)-4-*n*-butylaniline (MBBA), nematic solvent 59
(Methoxycarbonyl)acetylene, nucleophilic addition to 178
4-Methoxycyclohexanone, conformational equilibrium of 129
2-(Methoxymethylidene)adamantane, photo-oxygenation of 284
4-Methoxyneophyl tosylate, S_N1 solvolysis of 239, 404, 408, 409
4-Methoxyphenylazo-2-methylpropanedinitrile, thermolysis of 200, 201
1-(4-Methoxyphenyl)-1-methyl-2,2-diphenyloxirane, Wagner–Meerwein rearrangement of 69
N-Methylacetamide 82
Methyl acrylate, Diels–Alder cycloaddition reaction of 190, 191, 280, 281, 327, 410
γ-Methylallyl 4-tolyl ether, *ortho*-Claisen rearrangement of 303
Methyl amines, basicity anomaly of 104–106
3-Methylcytosine, amino/imino tautomerism of 116, 117
2-Methyl-2,3-dihydro-4*H*-naphth[1,2-*e*] 1,3-oxazine-3-ol, ring/chain tautomerism of 119
3-Methyl-2,5-dihydrothiophene 1,1-dioxide, cheletropic reaction of 198, 199
2-Methyl-4,4-diphenylcyclobutenone, electrocyclic reaction of 198
1-Methyl-hexahydroazepin-2-one 252
Methyl 3-hydroxy-2,3-dimesityl-2-propenoate, keto/enol tautomerism of 109
2-Methyl-*N*-hydroxyindole, nitrone/hydroxylamine tautomerism of 117
2-Methylindolenine-*N*-oxide, nitrone/hydroxylamine tautomerism of 117
3-Methyloxazolidin-2-one 252
2-Methyl-2-pentene, addition of singlet oxygen to 193
N-Methyl-*C*-phenylnitrone, 1,3-dipolar cycloaddition reactions of 191, 192
2-Methyl-3-phenylpropionitrile, base-catalyzed H/D exchange reaction 260
Methylphosphonic acid bis(dimethylamide) 253
Methyl propionate, alkaline hydrolysis of 234, 235
1-Methylpyrrolidin-2-one 82, 252
Methyl tosylate, S_N2 reactions of 168, 258, 405, 406
Micellar catalysis 42–46, 292–298, 306
Micelles 43–46, 292, 306
Michaelis–Menten mechanism 143
Microemulsions 298
Microscopic hydrophobicity (*MH*) substituent parameter 400
Miscibility (*M*) numbers 65, 66, 471
Miscibility of organic solvents 9, 65, 66, 471, 493
Mixotropic solvent series 471, 493
Molten salts as solvents 58–62, 245, 322–324, 504
Montmorillonite as catalyst in Diels–Alder cycloaddition reactions 295
Multiple linear regression analysis (MRA) 84, 85, 452
Multipolar solvents 67

n

"Naked" ions 245, 246, 248, 251, 268
1,4-Naphthoquinone-monophenylhydrazone, azo/hydrazone tautomerism of 3, 117, 118
Neoteric solvents 4, 62, 317–328
Neutral solvents 75
Nitrobenzene 82
4-Nitrobenzenediazonium tetrafluoroborate, coupling reaction of 174, 175
6-Nitrobenzisoxazol-3-carboxylate, decarboxylation of 242, 243
Nitromethane 82, 252
2-Nitrophenol, inter/intramolecular H-bonding 16
4-Nitrophenyl decanoate, aminolysis of 293, 294
4-Nitrosophenol, tautomerism of 118
Nitrosyl chloride, electrophilic addition to alkenes 177
NMR spectra, solvent effects on, 376–388
— acetone 378
— aromatic hydrocarbons 378
— camphor 385
— 1-chloro-1-cyanoethene 388

— (−)-cocaine 387
— 1,4-diethylpyridinium salts 376
— 3-(dimethylamino)propenal 342, 343, 378, 379, 388
— *N*,*N*-dimethylbenzamide 378
— 4-fluoro-nitrosobenzene 377
— fluoropyridines 378
— meropolymethine dyes 342, 343, 378, 379
— metal cations 36, 376, 377
— methane 381
— 1-methylpyridinium salts 376
— 1-methylquinolinium salts 376
— phenylacetylene 382
— pulegone 385
— pyridine-1-oxide 378
— pyridinium *N*-phenolate betaine dye 378
— spin-spin coupling constants 387, 388
— tetra-*n*-butylammonium hexacyano-cobaltate(III) 383
— tetramethylsilane 376, 377
— thallous ion 36, 376, 377
— tribromomethane 388
— trichloromethane 78, 378, 387
— triethylphosphane oxide 25, 26, 377, 378, 438, 439
— trifluoroacetic acid/2,4,6-trimethylpyridine 383
— trifluoromethane 388
— triphenylphosphane oxide 378
— tropylium salts 376
— xenon 381
n-Nonane, solubilization of 45
non-HBD solvents 18, 82, 88, 425
Nonpolar solvents 66, 67, 425
(2-Norbornyl)formyl 3-chlorobenzoyl peroxide, thermolysis of 202
Nucleophilicity of nucleophiles 159–161, 243–254, 263
Nucleophilicity parameters N_+ and N 247, 248, 405–407
Nucleophilicity scales 157, 158, 243–251, 263
Nucleophilic solvent assistance 275, 455

o
Ω values 327, 404, 410
n-Octadecyl *trans*-cinnamate, photodimerization of 301, 302
1-Octanol/water partition coefficient $P_{O/W}$ 399, 400, 460, 461
Oligoethyleneglykol dialkyl ethers as cation solvators 266–268
Onsager's reaction field 38, 132, 341, 342, 367
Optical rotations, solvent effects on 362
— (2*R*, 3*R*)-(+)-diethyl tartrate 362
— (*S*)-(−)-nicotine 362
Optical rotatory dispersion (ORD), solvent effects on, 359–362
— (+)-*trans*-2-chloro-5-methylcyclohexanone 361
— (*S*)-5-hydroxy-1,7-diphenylheptan-3-one 361
Organometallics, ionization and dissociation of 54, 55
7-Oxabicyclo[2.2.1]hept-5-ene, retro-Diels–Alder reaction of 188
Oxepin/benzene oxide, valence isomerization of 135, 136
3-Oxo-2,3-diphenylpropionitrile, keto/enol tautomerism of 109
Oxonole dyes, definition 331
Ozonolysis of alkenes, solvent effects on 307, 308

p
P values 437
π^* values 431–434, 447, 456–461
π^*_{azo} values 432
Parker's solvent classification 82–84, 244
Penetrated ion pairs 52
1-Pentanol, solubilization of 45
1-Pentene, bromine addition to 176
Perfluoroalkanes 4, 57, 58, 319–322, 504
Pericyclic reactions, solvent effects on 163, 187–199
Peroxyacetic acid, epoxidation of trimethylethene with 177
Peroxybenzoic acid, epoxidation of cyclohexene with 177
Peroxycarboxylic esters, thermolysis of 204–207, 285
Phase-transfer catalysis (PTC) 318, 319
Phencyclidine, conformational equilibrium of 129, 130
Phenol,
— as HBD solvent 240, 241
— *O*/*C* alkylation of 269–271
Phenyl acetate, alkaline hydrolysis of 276, 277
Phenylazide, 1,3-dipolar cycloaddition reaction of 191
4-Phenylazo-1-naphthol, azo/hydrazone tautomerism of 3, 117, 118, 325
(±)-3-Phenylbutanone, diastereoselective Grignard reaction of 289, 290
10-Phenylnonafulvene-10-oxide, dissociation equilibrium 55, 56
1-Phenyl-1-(piperidin-1-yl)cyclohexane, conformational equilibrium of 129, 130
3-Phenyl-1-(4-tolyl)-1-propene, ene reaction of 192
Photochlorination of hydrocarbons 210–212

Photo-oxygenation reactions, solvent effects on 282–284
Phthaldehydic acid, ring/chain tautomerism 118
Phthalide, ring/chain tautomerism 118
Piezo-solvatochromism 428
Piperidine, nucleophilic addition to alkines 178
Podands as cation solvators 266–268
Polarity of solvents,
— definition 8, 68, 389, 390
— empirical parameters 68, 69, 389–445
Polarizability of molecules 13, 14
Polar solvents 66–68
Polyenes,
— π-electronic structure 330, 331
— UV/Vis spectra 329, 331, 340
Polyynes,
— π-electronic structure 330, 331
— UV/Vis spectra 331
Polymerization reaction, multiparameter treatment of 468
Polymethines,
— π-electronic structure 330, 331
— UV/Vis spectra 331, 332
"Polywater" 6
Potassium phenolate, alkylation with 1-bromobutane 267, 268
Preferential solvation 38–42, 426, 427
Princess and pea 429
Principal component analysis (PCA) 84–90
Propylene carbonate 82
Protic solvents 18, 82–84, 88, 89, 238–244, 425
Protogenic solvents 75
Proton abstraction reaction, solvent effects on 253, 259–262
Proton affinity 101
Proton-transfer reactions 95–99, 121–123, 259–262, 317
Protophilic solvents 75
Protophobic solvents 75
Purification of solvents 471, 478, 480–481
Py values 358, 430
Pyridazine, UV/Vis spectrum of 349
Pyridinium *N*-phenolate betaines,
— alkylation reaction 392
— as solvent polarity indicators 332–334, 416–429
— solvatochromism of 332–334, 336, 416–428
— UV/Vis absorption 332, 333, 393, 394
— X-ray structure analysis 425
3-Pyridinyl 2-picolyl ketone, keto/enol tautomerism of 113
2-Pyridone, lactam/lactim tautomerism of 113, 114

q
Quantitative structure-property relation-ships (QSPR) 69, 90, 91

r
Racemization reaction of allyl sulfoxides, solvent effects on 184, 185
Radical addition reactions, solvent effects on 208, 213, 214
Radical combination reactions, solvent effects on 208
Radical disproportionation reactions, solvent effects on 208, 214
Reaction field 38, 132, 341, 342, 367
Rearrangement reactions, solvent effects on 163, 166, 184, 195–199
Recrystallization, solvents for 488, 490, 491
Refractive index 14, 68, 154, 448, 453, 454, 472–475
Relative permittivity 3, 47, 48, 67, 68, 454, 472–475
Retention index 444, 445
Rhodamine B, intramolecular ionisation of 123, 124
RPM values 429
Ruggli–Ziegler's dilution principle 6

s
S values 413–416
$\mathscr{S}$ values 404, 409
Salt effects,
— primary 236, 237
— secondary 237
Salting-out effect 38
Selective solvation 38–42, 426, 427
S_E reactions, solvent effects on 174, 175, 277, 278
Sigmatropic reactions, solvent effects on 195–197
Single-electron transfer (SET) reactions 137, 138, 164, 208–210
Smenogenic solvents 47
Smenolytic solvents 47
SMIRC program 91
S_NAr reactions, solvent effects on 173, 174, 249, 276, 277
S_N1 reactions,
— ionization step in 54, 164
— solvent effects on 164–166, 215, 216, 238–240, 274–276, 313, 314
S_N2 reactions,
— solvation rule for 172, 173

— solvent effects on 156–162, 164–169, 238–242, 243–251, 262–265, 274–276, 313, 314
— substituent effects on 392, 393
S_N2(intermediate) mechanism 274–276, 405
S_N2(one-stage)mechanism 275, 405
Soaps, selective solvation of 42
Sodio-butyrophenone, alkylation of 265
Sodio-4-(3-hydroxyphenyl)butyl tosylate, intramolecular alkylation of 276
Sodium fluorenide, dissociation of 54, 55
Sodium naphthalenide, dissociation of 55
Sodium 2-naphtholate, *O*/*C* alkylation of 270, 271, 323
Sodium phenolate, *O*/*C* alkylation of 269–271
Softness of solvents 80, 81
Solubility of hydrocarbons 28–30
Solubility parameter δ 9, 10, 65, 66, 220–224, 443, 444, 457, 460, 471
Solubilizate 45
Solubilization 45
Solubilizer 45
Solutions,
— definition 5
— dilute 5
— ideal 5
— regular 220
Solvation,
— chemical 35
— coordination number 35
— definition 30
— equisolvation point 40, 41
— Gibbs energy of 30–33
— heteroselective 39, 40
— homoselective 30, 39
— internal 279
— ions by dipolar molecules 31
— iso-solvation point 40, 41
— micellar 42–46
— multizone models 37
— NMR spectroscopic investigation 36
— physical 35
— preferential 38–42, 426, 427
— primary 35
— quantum-chemical calculations 38
— secondary 35
— selective 38–42, 426, 427
— shell, models of 37
— solvation number 35
— specific 82–84
Solvation number 35
Solvation rule for S_N2 reactions 172, 173
Solvatochromic comparison method 431–435
Solvatochromic dyes 333, 335–337, 411–435
Solvatochromism,
— definition 4, 330
— McRae–Bayliss model of 340, 341, 345, 346
— microstructural model of 347
— qualitative results 330–340
— quantitative calculations 340–347
Solvatochromism of
— benzene 331, 340, 345
— benzophenone 348–351
— 2,2′-bipyridine tetracarbonyltungsten(0) 339
— 4,4′-bis(dimethylamino)benzophenone (Michler's ketone) 350, 351
— bis(pentacarbonyltungsten)pyrazine 339
— 4-(*t*-butoxycarbonyl)-1-ethylpyridinium iodide 413
— *o*-*t*-butylstilbazolium betaine 440, 442
— carotinoids 331
— 4-(diacetylmethylene)-1,4-dihydro-1-methyl-2,3,5,6-tetraphenylpyridine 337
— *o*,*o*′-di-*t*-butylstilbazolium betaine 440, 442
— 1,1-dicyano-10-(dimethylamino)deca-1,3,5,7,9-pentaene 335
— 2,3-dicyano-5,6-diphenylcalicene 337
— *N*,*N*-diethyl-4-nitroaniline 335, 431
— 5-(dimethylamino)-5′-nitro-2,2′-bithiophene 334
— 3-(dimethylamino)propenal 331, 342, 343
— 2,5-bis(dimethylamino)-1,4-benzo-quinone 335
— 2-dimethylamino-7-nitrofluorene 439–441
— 4-dimethylamino-4′-nitrostilbene 336
— 5-(dimethylamino)penta-2,4-dien-1-al 335, 347, 429
— 1-(4-dimethylaminophenyl)-2-nitroethene 335
— 5-{5,5-dimethyl-3-[2-(2,3,6,7-tetrahydro-1*H*,5*H*-benzo[ij]quinolizin-9-yl]-2-cyclohexen-1-ylidene}-1,3-diethyl-2-thiobarbituric acid 335
— *N*,*N*-(dimethyl)thiobenzamide-*S*-oxide 430
— 2,6-diphenyl-4-(2,4,6-triphenylpyridinium-1-yl)phenolate 332–334, 336, 416–428
— disodium imidazole pentacyano-ferrate(III) 339
— 1-ethyl-2-(3-formyl-3-nitroallylidene)-1,2-dihydroquinoline 337
—1-ethyl-4-(methoxycarbonyl)pyridinium iodide 334, 338, 339, 411–415
— 4-[5-(5-ethyl-1,2,3,5-tetrahydro-3-methylpyrido[1,2-*a*]benzimidazol-4-yl)penta-2,4-dienylidene]-2,2-dimethyl-cyclobutane-1,3-dione 336
— 2-fluoro-7-nitrofluorene 439–441
— indigo 336
— iodide ion 340

Solvatochromism of (cont.)
— lycopene 331
— merocyanines 331, 342, 343
—1-methyl-4-(3,3-diformylallylidene)-1,4-dihydroquinoline 337
— 1-methyl-5-nitroindole 439–441
— 4-methylpent-3-en-2-one 335, 413
— 4-[2-(1-methyl-4-pyridinium-1-yl)-ethenyl]phenolate 336, 344, 345
— 4-[4-(1-methyl-4(1*H*)-pyridylidene)-2-butenylidene]-3-phenyl-2-isoxazolin-5-one 336
— nil blue A oxazone 336
— 4-nitroaniline 431, 433
— 5-nitroindole 439–441
— phenanthrene 340
— phenol blue 335, 413
— polyacetylenes 331
— pyridazine 349
— pyridine-1-oxide 413
— pyridinium cyclopentadienide 337
— pyridinium *N*-phenolate betaine 332–334, 336, 416–428
— tetracarbonyl-[*N*-(2-pyridylmethylene)-benzylamino]molybdenum(0) 430
— 2,2,6,6-tetramethylpiperidine-1-oxide 351
— tropylium iodide 413
Solvatomers 40
Solvency 65
Solvent activity coefficient 33, 34, 254–259
Solvent cage effects 303–308
Solvent isotope effects 315–317
Solvent polarity,
— definition 8, 68, 389, 390
— empirical parameters 68, 69, 389–445
Solvents,
— acid/base behaviour 73–79
— amphiprotic 18, 75, 96
— apolar 66, 67, 88, 89, 425
— apolar non-HBD 82, 83, 88, 89
— aprotic 18, 75, 82, 83, 88, 89
— autoprotolysis constants 74, 75, 496–499
— boiling points 60, 63, 472–477, 491, 504
— chemical constitution 57, 58
— chiral 69, 70, 386, 387, 476, 477, 504
— cohesive pressure 62, 64, 219, 220, 443, 444, 457
— coordinating 22
— desmotropic constants 396
— differentiating 76
— dipolar 11, 66, 67, 75, 82, 83, 252, 253
— dipolar non-HBD 18, 82, 83, 88, 89, 239, 240, 245, 246, 252, 253, 425
— dissociating 47, 52
— electrostatic factor *EF* 67, 68
— EPA solvents 21–27, 48, 80, 81, 88, 89
— EPD solvents 21–27, 48, 80, 81, 88, 89, 263–265, 410
— evaporation numbers 63
— fluorous solvents 4, 57, 58, 319–322, 504
— HBA basicity 18, 82
— HBD acidity 18, 82
— internal pressure 63–65, 220–224
— ionizing 48, 52, 53
— ionizing power 48, 52, 53, 402
— levelling 76
— liquid crystals 58, 59, 298–303, 306, 307
— liquid metals 58, 62
— melting points 60, 472–477
— molten salts 58–62, 245, 322–324, 504
— neutral 75
— non-HBD 18, 82, 88, 425
— nonpolar 66, 67
— polar 66–68
— polarity 8, 68, 389, 390
— protic 18, 82–84, 88, 89, 238–244, 425
— protogenic 75
— protophilic 75
— protophobic 75
— purification 471, 478, 480–481
— smenogenic 47
— smenolytic 47
— soft and hard 80, 81, 244, 272, 273, 401, 402
— spectroscopic 479, 482–488
— supercritical 4, 70–72, 315, 324–328, 492, 494, 504
— Trouton constants 63
— viscosity 63, 307
Solvent-separated ion pairs 52–55
Solvent-shared ion pairs 52, 53
Solvent-transfer activity coefficients 33, 34, 254–259
Solvophobic interactions 27–30, 292–298, 400
Solvophobicity parameter S_p 295, 400, 401
Spectroscopic solvents 479, 482–488
Spherands as cation solvators 266, 267
Spiropyran, intramolecular ionization of 124, 125
Stern layer 43
Stilbene, bromine addition to 278, 279
Stokes shift 352, 353
Styrene,
— autoxidation reaction of 210
— cycloaddition reaction of 180, 193
— electrophilic addition to 177
Sulfamic acid, intramolecular proton-transfer equilibrium 104
Sulfolane 82, 253
Sulfur dioxide 49, 50
Super acid/base systems 77, 81

Supercritical-fluid (SCF) solvents 4, 70–72, 186, 315, 324–328, 492, 494, 504
Supramolecular ion pairs 56
Surfactants 44
Sydnones, dipole moments of 11
Synergetic effects 42, 426

t
(*R*,*R*)-TADDOL as chiral additive 289
Tautomeric equilibria, solvent effects on,
— amino/imino 116, 117
— azo/hydrazone 3, 117, 118, 325
— 1,3-dicarbonyl compounds 3, 4, 106–111
— imine/esamine 116, 117
— keto/enol 3, 4, 106–111
— lactam/lactim 113–115
— metallotropic 120
— nitro/isonitro 3
— nitrone/hydroxylamine 117
— nitroso/oxime 118
— ring/chain 118, 119
2,3,4,5-Tetrachloro-6-[(diethylamino)-methyl]phenol, proton-transfer equilibrium of 122
Tetracyanoethene, cycloaddition reactions of 179, 180, 281, 410
Tetraethylsulfamide (TES) 253, 504
3,4,5,6-Tetrahydro-1,3-dimethyl-2(1*H*)-pyrimidinone (DMPU) 82, 252, 504
Tetrahydrothiophene 1,1-dioxide 82, 253
Tetralin, autoxidation of 210
4,4,5,5-Tetramethyl-1,2-dioxolan-3-one, decarboxylation of 206
N-(2,2,5,5-Tetramethylpyrrolidyl)nitrene, thermolysis of 204
Tetramethyltin, S_E reaction of 175, 409
1,1,3,3-Tetramethylurea (TEMUR) 82, 252
Theoretical linear solvation energy relationships (TLSER) 466–468
Thermo-solvatochromic comparison method 434, 435, 459
Thermo-solvatochromism 428
Thermotropic liquid-crystalline solvents 58
Thiazolo[2,3-*e*]tetrazole, valence isomerization of 135
Thiocarbonyl ylides, 1,3-dipolar cycloaddition of 191
Threshold limit values (TLV) 500–503
Tight and loose ion-pairs 53
4-Toluoyl chloride/antimony pentachloride adduct, ionization of 50, 51
Transition-state theory 149–154, 218, 219, 308, 309, 314
Triarylsulfonium halides, alkaline hydrolysis of 169
Tri(*n*-butyl)phosphane, reaction with carbon disulfide 125, 168
Tri(*n*-butyl)phosphonium dithiocarboxylate, formation of 125
2,4,6-Tri-*t*-butyl-*N*-thiosulfinylaniline, valence isomerization of 135
Triethylamine, S_N2 alkylation reaction of 167, 227, 228, 313, 451
Triethyloxonium hexafluorophosphate, solvolysis of 407
Triethylphosphane oxide, ^{31}P-NMR spectrum of 25, 26, 376, 377, 438, 439
Triethylsulfonium bromide, thermolysis of 166
Trimethylamine, S_N reactions of 169, 170
2,4,6-Trimethylbenzenecarbonitrile oxide, 1,3-dipolar cycloaddition of 191
1′,3′,3′-Trimethyl-6-nitrospiro[2*H*-1-benzopyran-2,2′-indoline], intramolecular ionization of 124, 125
Trimethyl-phenacyltin, metallotropic tautomerism of 120
Trimethylsulfonium ion, S_N2 reactions of 169
(Triphenylcyclopropen-1-yl)-(4-nitrophenyl)malononitrile/triphenylcyclopropenium dicyano-(4-nitrophenyl)methide, ionization equilibrium of 51
Triphenylmethyl radical, generation of 200, 201, 222, 224
Triphenylphosphane, S_N2 reaction of 167
1,1,1-Tris(4-methoxyphenyl)-2,2-dicyano-2-(4-nitrophenyl)ethane, ionization equilibrium of 126
Tri-*n*-propylamine, Menschutkin reaction of 166, 409
Tropolone/triethylamine, proton-transfer equilibrium of 122
Trouton constant 63

u
UV/Vis absorption, $\pi \rightarrow \pi^*$, solvent effects on 330–340
UV/Vis absorption, specific solvent effects on, 348–351
— $\pi \rightarrow \pi^*$ absorption 348–351
— $n - \sigma^*$ absorption 351

v
Valence isomerization equilibria, solvent effects on, 135–137
— 3-azidopyrazine-1-oxide 135
— 2-azidothiazole/thiazolo[2,3-*e*]tetrazole 135
— 8,8-diformylheptafulvene/8a*H*-cyclohepta[*b*]furan-3-carbaldehyde 136
— γ,γ-diphenylallylcarbinyl lithium/diphenylcyclopropylcarbinyl lithium 136, 137

Valence isomerization equilibria, solvent effects on (cont.)
— lithium 10-phenylnonafulvene-10-oxide 55, 137
— oxepin/benzene oxide 135, 136
— 2,4,6-tri-*t*-butyl-*N*-thiosulfinylaniline 135
Van der Waals forces 10, 220, 380
Van't Hoff–Dimroth relationship 110, 154
Vapour hazard ratio (VHR) 503, 507
Vapour pressure of solvents 63, 501, 502
Vesicles 46
Vinyl chloride, preparation of 61
Viscosity of solvents 63, 307, 314
Volume of activation 221–223, 308–315

w
"*W*"-values 409
Wagner–Meerwein rearrangement, solvent effects on 69, 166
Water,
— acid/base behaviour 72, 75, 79
— flickering cluster model 6, 37
— heavy 316
— metastable 73
— "polywater" 6
— structure 6, 7, 37
— supercooled 73
— supercritical 71, 72, 327
— superheated 72, 73, 328
Willgerodt–Kindler reaction 87
Williamson ether synthesis, solvent effects on 251
Winstein solvolysis scheme 54, 238
Wittig reaction, solvent effects on 194, 195, 260, 262
Wolff–Kishner reduction, solvent effects on 260, 261

x
X values 404, 409, 410
ζ values 457, 458

y
Y values 402–408, 453, 454

z
Z values 412–416, 446, 447

Author Index

a
Aakeröy, C. B. 15, 17
Abbott, A. 47, 52, 500
Abboud, J.-L. M. 19, 27, 166, 168, 230, 331, 340, 379, 390, 393, 411, 429, 431–434, 436, 442, 443, 447, 448, 450, 453, 456–458, 460, 461, 464, 465
Abdel-Mottaleb, M. S. A. 334, 336
Abdoul-Carime, H. 39
Abe, I. 127, 131
Abe, T. 96, 337, 345, 354, 358, 373, 432, 440
Abed, O. H. 195
Abley, P. 477
Abraham, M. H. 19, 29, 32, 94, 129, 132, 147, 154, 165, 166, 215, 227, 228, 230–232, 250, 257, 259, 276, 278, 388, 396, 399–401, 403, 408, 429, 431, 432, 434, 436, 442, 443, 447, 449, 453, 456–458, 460, 461, 463, 464
Abraham, R. J. 121, 126, 128, 129, 131, 132, 166, 227, 228, 230, 257, 259, 388
Abraham, W. 87, 468
Accascina, F. 46, 47, 52
Acheson, R. M. 308, 309, 314, 315
Achiwa, K. 144
Acker, K. J. 61
Acree, S. F. 262
Acree jr., W. A. 42
Acree, W. E. 358, 416, 426, 430, 463
Ács, M. 468
Adam, C. 42
Adam, W. 206, 284
Adamowicz, L. 113, 114
Adams, C. J. 323
Adams, R. N. 370
Addison, C. C. 49, 57
Adembri, G. 119
Adler, P. 468
Afanas'ev, V. N. 39
Affolter, C. 479, 485, 487
Agami, C. 27, 78, 79, 82, 84, 99, 252, 364, 367, 383, 475
Ågren, H. 345
Aguirre, de, I. 404, 409
Ahlbrecht, H. 116
Ahmad, N. 397
Ahrens, A. F. 159–161
Aillon, M. 428
Akabori, S. 271, 272
Aki, S. N. V. K. 62
Akiya, N. 72, 328
Aksnes, G. 194, 259
Alaimo, P. J. 478
Albano, C. 84, 86, 87
Alberghina, G. 128
Albert, A. 99
Albuquerque, L. M. P. C. 468
Aldarova, N. Sh. 499
Alder, M. G. 201
Alei, M. 36
Alekseeva, V. I. 333
Alexander, G. C. 315
Alexander, R. 156, 249, 250, 254, 255, 257–259
Alhambra, C. 343
Aliaga, C. 345
Alkorta, I. 15
Allard, B. 376, 383
Allen, A. D. 275
Allen, G. 64, 65, 107
Allerhand, A. 365, 369, 414, 415, 435
Allin, S. M. 72, 73
Allinger, N. L. 129, 131, 132
Allmann, R. 425
Almdal, K. 106–109, 112, 113
Alpert, N. L. 479, 483
Altnau, G. 503
Alunni, S. 398, 408, 453, 462–465
Alvarez, E. 203
Ambrose, D. 72
Ambrose, J. 60
Ambrosetti, R. 176
Amey, R. L. 12
Amis, E. S. 30, 35, 36, 38, 40, 84, 147, 215, 219, 226, 234–236, 238
Ammon, R. von 69
Amovilli, C. 166, 168
Anastas, P. T. 317, 325
Andersch, J. 60, 62
Andersen, H. C. 429
Andersen, K. K. 376, 377
Anderson, J. E. 475
Anderson, R. G. 376
Anderson, S. W. 403, 407
Ando, I. 381, 388
Andrac-Taussig, M. 383
Andreas, F. 200, 201, 222, 224, 317
Andrews, L. J. 188
Anet, F. A. L. 378
Angyal, S. J. 119, 128
Anh, N. T. 244
Ansari, F. L. 197
Anselme, J.-P. 203
Antonious, M. S. 336
Antonov, L. 113–115
Anufriev, E. K. 115
Anvia, F. 103
Aono, S. 370
Aonuma, S. 51, 52
Aoshima, K. 286
Apaydin, G. 243
Appriou, P. 337
Arai, S. 334
Aramendia, P. F. 357, 358
Archer, W. L. 57, 471
Argaman, R. 36
Arikata, S. 410, 411
Arjona, O. 116, 289, 290
Armarego, W. L. F. 478
Armstrong, K. B. 131
Armstrong, R. S. 331
Arnett, E. M. 16, 24, 29, 30, 51, 52, 82, 99, 104, 106, 126, 138, 148, 166, 253, 315–317, 272
Arnold, J. 478
Arnold, J. T. 381, 382
Arnone, C. 128
Arrhenius, S. 2, 149
Arrieta, A. 183, 194
Arsenev, Yu. N. 444, 445
Artemenko, A. I. 115
Asano, M. 185, 186, 292, 300, 334
Asano, T. 185, 186, 300, 308–310, 314, 315
Ash, I. 57
Ash, M. 57
Ashby, E. C. 164
Aslam, M. H. 454
Aslanidis, P. 113, 114
Astle, M. J. 475
Astruc, D. 237, 263

Solvents and Solvent Effects in Organic Chemistry, Third Edition. Christian Reichardt

Copyright © 2003 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

ISBN: 3-527-30618-8

Atherton, N. M. 55, 369
Atkins, P. W. 10, 12, 13, 93, 304
Atto, S. Y. 212
Aubé, J. 288
Audrieth, L. F. 57
Aue, D. H. 104
Augé, J. 57, 294, 505
Auhmani, A. 379
Aurich, H. G. 370, 372
Auriel, M. 65, 229, 238, 241, 242
Ausloos, P. 100, 147, 305
Avila, D. V. 213
Avnir, D. 333
Avraamides, J. 287
Avramenko, N. V. 468
Axelrod, E. 477
Ayachit, N. H. 337, 354
Ayers, P. W. 201, 202
Ayrey, G. 306
Ayscough, P. B. 370

b

Baaz, M. 50
Baba, H. 112, 121
Babrogic, Z. 417, 425
Bacelon, P. 438
Bach, G. 331
Bacon, M. R. 376, 377
Bader, R. F. W. 316
Badoz-Lambling, J. 500
Baerends, E. J. 162, 172
Baeyer, A. von 428
Bagal, I. L. 174, 175
Bagchi, B. 36, 147–149, 151, 217, 218
Bagchi, S. 42, 357, 413, 426
Bagno, A. 40, 78, 79, 99, 103, 123
Bahari, M. S. 407
Bähr, G. 254
Baiker, A. 70, 71
Bailey, P. S. 307, 308, 336
Bain, A. 369
Baird, M. H. J. 490
Baker, B. R. 28, 30
Bakshiev, N. G. 330, 337, 345, 348, 354
Balakrishnan, M. 378
Balakrishnan, S. 417, 425, 426
Balasubramanian, M. 72, 73
Baldry, K. W. 129
Baldwin, J. E. 193
Ball, P. 6, 117
Ballistreri, F. P. 188
Bamford, C. H. 304
Banarjee, D. 357
Banarjee, S. 9
Bank, S. 253, 260
Banthorpe, D. V. 170, 253
Baram, N. I. 119
Baranovski, B. 308, 317
Barbara, P. F. 334
Barbaras, G. K. 431
Baret, P. 243
Barfield, M. 376, 378, 387, 388
Bariou, B. 166
Barker, B. J. 82, 252, 475
Barker, J. A. 5
Barkley, R. M. 272
Barnes, A. J. 378
Barr, P. A. 251
Barrat, M. D. 370
Barriol, J. 368
Barrow, G. M. 121
Bartberger, M. D. 183
Barth, G. 477
Barthel, J. 46, 47, 253
Barthel, J. M. G. 30
Barthel-Rosa, L. P. 320, 321
Bartholomay, H. 104
Bartik, K. 381
Bartkowiak, W. 334
Bartlett, P. D. 178, 188, 201, 204, 205, 207, 282, 283, 285, 409
Bartmess, J. E. 102, 103, 159, 162
Barton, A. F. M. 9, 10, 63–66, 68, 220, 222, 260, 308, 443, 471
Bartsch, R. A. 261, 268, 286
Basu, R. 21
Basu, S. 31
Bates, G. S. 195
Bates, R. G. 73, 75, 77, 79, 99, 496, 499
Bathgate, R. H. 156, 159
Batog, A. E. 205
Battaglia, A. 193
Battino, R. 9
Bauer, B. 291
Bauer, E. 367
Bauer, O. H. 429, 430
Bauer, S. H. 150
Bauer, W. 55
Bauernfeind, D. 50
Baukov, Yu. I. 120
Baumann, J. B. 490, 491
Baumann, W. 336, 354
Baumgärtel, H. 297, 333, 378, 429
Bayer, E. 492
Bayer, R. E. 499
Bayliss, N. S. 331, 336, 340, 342, 345, 349, 367, 368
Beak, P. 106–109, 111–115
Bear, J. L. 244–247
Beard, J. H. 229
Beard, M. C. 333
Beare, S. D. 386, 477
Beauchamp, J. L. 103, 104
Beaumont, T. G. 376, 417, 425
Beck, S. C. 353
Becker, H. G. O. 238
Becker, J. 262
Becker, R. S. 348
Beckett, M. A. 42
Beckmann, E. 47
Beckmann, O. 4
Beer, P. D. 243
Beeson, C. 388
Beguin, C. 498
Behera, G. B. 331, 340, 345
Behera, P. K. 331, 340
Behera, R. K. 331, 340
Beier, T. 177
Beither, U. 69
Bekaert, A. 166
Bekárek, V. 166, 345, 368, 378, 417, 425, 433, 434, 447, 448
Belanger, G. 479
Belavin, I. Yu. 120
Beletskaya, I. P. 52, 54, 99, 101, 263, 269, 271, 272, 276, 277
Bell, R. P. 73, 79, 95, 121
Bellamy, L. J. 364, 365, 367, 383
Bellobono, I. R. 185, 186, 292, 300
Bellon, L. 351
Belluci, G. 176
Bellus, D. 196
Benamou, C. 351
Bender, C. J. 19
Bendig, J. 334
Ben-Naim, A. 27, 29, 30, 36, 42, 93
Ben-Nun, M. 147
Benoit, R. L. 247
Bensasson, R. V. 45
Bensaude, O. 113–116

Benson, H. G. 336, 344, 378
Benson, S. W. 93
Bent, H. A. 19
Bentley, F. F. 479
Bentley, T. W. 274, 275, 403, 405–408, 432, 459
Bentrude, W. G. 201, 205, 206
Benyamin, D. 453, 467
Bergan, J. J. 104
Berge, A. 264
Berger, M. 243
Berger, S. 479, 485, 487
Bergman, R. G. 478
Bergmark, W. R. 354
Bergsma, J. P. 370
Berlan, J. 290
Bernardi, F. 103, 193, 370, 374
Bernasconi, C. F. 100, 173, 259
Berndt, D. C. 195, 196
Berneis, H. L. 431
Bernhard, G. 330
Berns, U. 292
Bernstein, H. J. 111, 381
Berrebi, G. 166, 227, 229
Berrigan, P. J. 270
Bershtein, I. Ya. 117
Berson, J. A. 188, 190, 191, 280, 291, 295, 404, 410, 446
Bertagnolli, H. 6
Berthelot, M. 2, 19, 398
Bertho, A. 274
Bertini, I. 147
Bertrán, J. 38, 148, 149, 151, 166, 168, 188, 217, 218, 230
Bertrán, J. F. 386
Besace, Y. 290
Bessenbacher, C. 340, 429, 430
Bestian, H. 182, 193
Bestmann, H.-J. 194, 195
Betzemeier, B. 57, 320, 321, 488, 505
Bezdenezhnykh, A. A. 178
Bhacca, N. S. 384
Bhattacharyya, K. 36
Biali, S. E. 109
Bianchi, A. 243
Bianchi, G. 192
Bianchini, R. 176
Bickart, P. 184, 185, 449
Bickelhaupt, F. M. 172
Biedrzycka, Z. 378
Biellmann, J.-F. 113
Bienvenüe, A. 348, 414, 415, 429
Bienvenue, E. 45
Bierbaum, V. M. 242
Bierne, D. 490
Bigler, P. 178
Bilot, L. 345
Bindewald, H. 3
Binegar, G. A. 203
Bingham, R. C. 166, 403
Binnewies, M. 4, 506
Binsch, G. 180, 181, 193
Birge, R. R. 345
Biroš, J. 128, 132
Bishop, E. O. 376, 378
Bissing, D. E. 194
Bitar, H. 379
Bjerrum, N. J. 2, 47, 237
Blackburn, G. M. 243
Blackert, J. F. 326, 327
Blackwood, R. K. 269
Blair, L. K. 102, 104
Blake, J. F. 188, 189, 197, 201
Blanchard, G. J. 358
Blanco, A. 292
Blandamer, M. J. 30, 93, 108, 147, 219, 237, 238, 242, 259, 308, 314, 340, 365
Blankenburg, B. 188, 410
Blaser, H. 116
Blaser, H.-U. 471
Blinov, L. M. 58
Blöchl, P. E. 162
Block, H. 373
Blokzijl, W. 27–29, 108, 237
Bloom, A. 166
Bloom, H. 59
Blum, P. 333, 478
Blumrich, K. 500
Blunt, J. W. 287
Blyth, C. A. 293, 294
Boand, G. 102
Boca, J. P. 117
Boche, G. 52, 55, 56, 120, 137
Böck, H. 205
Bockman, T. M. 52
Bodner, R. L. 376, 397
Bodrikov, I. V. 177
Boerhaave, H. 1
Bogaert, H. v. d. 284
Bohle, M. 84, 86
Böhlig, H. 126
Bohlmann, F. 332, 333, 416, 417, 449–451
Böhm, S. 128, 132
Böhm, V. P. W. 323, 506
Bohme, D. K. 100, 103, 147, 156, 158–160, 162, 245, 246
Bolduc, F. 336
Böll, W. A. 136
Bollinger, J.-C. 22, 397
Bolm, C. 4
Bolton, R. 176, 177
Bonner, W. H. 431
Bonnet, P. 368
Boon, J. A. 323
Boos, H. 354
Boozer, C. E. 187, 306
Borden, W. T. 201
Bordwell, F. G. 18, 77, 78, 82, 88, 101–103, 244
Borgstedt, H. U. 62
Borgwardt, S. 264, 265
Born, M. 51, 96
Bornowski, B. 343, 378, 379
Borst, J. P. 55
Bortiatynski, J. 55
Bosch, E. 42, 331, 426, 427
Boschung, A. F. 282, 283
Bosnich, B. 362
Bothner-By, A. A. 376
Botrel, A. 336, 337, 344, 345
Bottino, F. A. 128
Boubel, J. C. 36
Boukouvalas, J. 282, 283
Boulton, A. J. 113, 115, 116
Bountis, T. 121
Bowen, C. T. 274, 275, 406
Bowers, M. T. 104, 147, 268
Bowie, J. H. 147
Bowman, N. S. 315, 317
Bown, D. R. 408
Boyd, R. H. 77, 79
Boyer, B. 129, 361
Brackmant, J. C. 42
Brady, J. E. 340, 345, 346, 452, 459
Brady, J. W. 131
Bram, G. 263, 271
Branchadell, V. 188
Brandes, E. 197
Bratos, S. 57
Brauer, H.-D. 312, 313
Brauman, J. I. 77, 99, 101, 102, 104, 147, 156–159, 160, 162, 244–246, 272
Braun, M. 108
Braun, R. 189, 191, 268, 404
Braun, S. 479, 485, 487
Braunstein, J. 59

Brealey, G. J. 348–350
Bréant, M. 82, 252, 475, 498
Brennecke, J. F. 62, 325
Brenninger, W. 330
Brereton, I. M. 267
Breslow, R. 57, 189, 295, 505
Bretschneider, E. 121, 126, 128, 131, 132
Brettschneider, H. 200
Breus, I. P. 188
Brewer, W. 205, 207
Brewster, P. W. 498, 499
Brieger, G. 271
Briegleb, G. 19, 21, 106, 107, 111
Brière, R. 370
Briggs, R. W. 36, 376, 377
Brill, T. B. 327, 328
Bristol, D. 215, 216,
Brock, E. E. 70, 73
Brode, W. R. 4
Brokken-Zijp, J. 284
Bröll, D. 71, 73
Bromby, N. G. 177
Brønsted, J. N. 73, 75, 79, 237, 392
Brooker, L. G. S. 330, 335, 336, 344, 411, 414, 415, 429
Brower, K. R. 312
Brown, C. E. 213
Brown, E. C. 56
Brown, G. H. 58
Brown, H. C. 104, 431, 489
Brown, J. S. 185, 186
Brown, P. 188
Brown, R. 60, 61, 404
Brown, R. D. 113, 114
Brown, R. S. 176, 278
Brown, T. L. 369
Browning, E. 500
Brownlee, R. T. C. 103, 437
Brownstein, S. 413–415
Broxton, T. J. 156, 249, 250, 254, 257–259
Bruce, A. E. 365, 374
Bruce, M. R. M. 365, 374
Bruckenstein, S. 75, 77, 79, 499
Bruice, T. C. 254
Bruin, F. 370
Bruin, M. 370
Brunero, P. 331, 332, 340
Brüning, I. 191
Brunner, H. 200
Bruno, T. J. 351, 358
Brunschwig, B. S. 340, 345, 346
Bruvers, Z. 334
Bryce, R. A. 131
Brycki, B. 164, 166, 169, 275
Brzezinka, K.-W. 334
Buback, M. 326
Bucciarelli, M. 69
Buchachenko, A. L. 370
Buchanan, J. 310
Buchet, R. 18, 19
Buckingham, A. D. 10, 57, 368, 369, 380
Bühlmann, P. 479, 485, 487
Bulthuis, J. 386
Bunce, N. J. 211
Buncel, E. 55, 70, 77, 93, 94, 99, 101, 102, 124, 125, 152, 154, 195, 219, 242, 259, 262, 263, 315, 317, 411, 429, 432, 453, 489
Bunge, W. 478
Bunger, W. B. 50, 57, 471, 475, 477, 478
Bunnel, R. D. 259
Bunnenberg, E. 477
Bunnett, J. F. 99, 244
Bunton, C. A. 46, 165, 292, 316, 403
Burawoy, A. 117
Burchard, T. 139, 140
Burden, A. G. 364, 367, 436, 454
Burdett, J. L. 107, 112
Burfield, D. R. 478, 481
Burger, A. 113
Burger, K. 30, 147
Burgess, J. 30, 36, 40, 147, 152, 219, 237, 238, 242, 308, 314, 339
Burgos, F. S. 148, 149, 151, 217, 218
Burke, L. A. 135
Burlingame, T. G. 386, 477
Burrage, M. E. 188, 189
Burrell, H. 220, 222, 223, 225, 308, 443
Bursey, M. M. 100
Burton, N. A. 113–115
Busse, G. 40
Butin, K. P. 99, 101
Butler, R. S. 388
Butt, A. 477
Bykova, L. N. 73, 75, 79, 496, 499
Byrne, J. S. 365

c

Cafferata, L. F. R. 203
Cagniard de la Tour, C. 70
Cahen, Y. M. 36, 376
Caillot, M. 78, 79
Cainelli, G. 289
Cairns, T. L. 180, 193
Caldin, E. F. 99, 147, 238
Caldin, E. T. 229
Caldow, G. L. 369
Caldwell, G. 103, 147, 157, 158, 162, 247
Call, L. 370
Cameron, C. 306, 313, 314
Cameron, D. G. 365
Camp, R. L. 201
Campanella, C. L. 121
Campbell, D. S. 176
Campulla, M. 40
Cañizo, A. I. 203
Cao, M. 115
Cappelli, C. 10, 38, 127, 131, 329, 345, 381
Cargioli, J. D. 378
Carins, S. N. 101
Carlson, R. 84, 86, 87
Carlson, R. L. 415
Carmichael, A. J. 62, 336
Carpenter, B. K. 148, 149, 151, 217, 218
Carr, P. W. 340, 345, 346, 396, 399, 443, 452, 457–460
Carré, J. C. 281
Carrea, G. 143
Carretto, J. 84, 88, 89, 425
Carrion, F. 158
Carroll, F. I. 129, 130
Carroll, M. 376, 377
Carson, F. W. 184, 185, 449
Carter, G. E. 274, 275, 403, 405–407, 432
Caruso, J. A. 82, 95, 96, 121, 252, 475
Casadevall, E. 376, 383
Castle, R. N. 112
Castleman, A. W. 162
Catalán, J. 243, 357, 390, 393, 398, 411, 429, 439–443, 447, 448, 453, 461
Cativiela, C. 188, 401
Caude, M. 494
Cauquis, G. 500

Cavell, E. A. S. 238, 241
Cavvazzini, M. 321
Cecil, R. 28
Celebi, S. 199
Chaimovich, H. 292
Chambers, J. G. 339
Chambers, J. Q. 370
Chan, T.-H. 57, 294
Chandrasekhar, J. 158
Chang, P. 55
Chang, T.-H. 475, 500
Chang, Y.-M. 191
Channot, E. 61
Chanon, M. 84, 88, 89
Chantoni, M. K. 77, 79, 95, 96, 121, 499
Chapelet-Letourneux, G. 370
Chapman, D. 42
Chapman, N. B. 85, 390, 454
Chapman, W. H. 201
Charles, R. 69
Charlot, G. 57, 59, 95, 97, 121, 496
Charlton, M. H. 336, 345
Charton, M. 405–407
Chastrette, M. 84, 88, 89, 425
Chateauneuf, J. E. 325
Chattaraj, P. K. 80
Chattopadhyay, P. K. 499
Chauhan, M. S. 40
Chauvin, Y. 322, 323, 506
Chawla, B. 437
Chayet, L. 292
Chaykovsky, M. 260
Che, D. 417
Chen, L. 60
Chen, P. 330, 334, 340, 345
Chen, R. H. H. 345
Chen, S.-H. 29, 30
Chen, T. 25, 81, 402
Chen, X. 326
Cheng, E. S. 462
Chernyak, B. I. 365, 366
Chesta, C. 357, 358
Chevrier, B. 50
Chevrier, M. 113–115
Chibisov, A. K. 124, 125
Chimichi, S. 119
Chiosi, S. 284
Choho, K. 162
Choi, C. 354
Choux, G. 247
Christ, H. A. 378, 383
Christe, K. O. 261
Christen, H. 331
Christen, H. R. 139–141
Christensen, J. J. 268
Christian, S. D. 121
Christiansen, J. A. 237
Chronister, C. W. 453, 464
Chujo, Y. 333
Chumak, V. L. 39
Cinquini, M. 268, 360
Ciobanu, M. 308, 309, 314, 315
Ciula, J. C. 428
Claessens, M. 67, 133, 389
Claisen, L. 3, 269
Clark, J. H. 15, 251, 253, 261, 262
Clarke, J. T. 201
Clary, D. C. 159–161
Clauß, K. 182, 183
Clavijo, E. 365
Claxton, T. A. 370, 372
Clennan, E. L. 284
Clifford, A. A. 327, 328
Clifford, T. 70
Cline Love, L. J. 44
Clippinger, E. 53, 54
Coates, R. M. 197
Cocivera, M. 172
Cockerill, A. F. 170, 254
Coddens, M. E. 60
Coetzee, J. F. 30, 36, 38, 41, 54, 73, 75, 77, 79, 82, 84, 252, 253, 364, 365, 475
Cohen, H. L. 477
Coker, T. G. 60
Cole, A. G. 499
Cole, A. R. H. 367, 368
Collier, G. 436, 454, 458
Collin, G. 177
Collings, A. J. 500
Collins, C. J. 315, 317
Colnay, M.-E. 67
Colonna, S. 303, 360
Comini, S. G. 197
Compton, R. G. 304
Condon, C. U. 341, 352
Condroski, K. R. 177
Connelly, S. A. 268
Connolly, J. D. 384
Connors, K. A. 139, 147, 149, 218, 219, 426, 427
Conradi, M. S. 42, 71
Conway, B. E. 30, 38
Cook, C. D. 214
Cook, D. 51, 156
Cook, F. L. 268
Cook, M. J. 113, 114, 129, 132
Cookson, R. C. 188, 189
Coomber, D. I. 194
Cooper, K. A. 163, 165, 169–172, 215, 225, 310
Cooper, M. A. 131, 132
Cooper, R. M. 479
Coppinger, G. M. 351
Corey, E. J. 260
Corley, R. C. 304
Corni, S. 127, 131
Cornils, B. 320–322, 488, 505, 506
Corriu, R. 360
Corset, J. 438
Cortijo, M. 399
Cossio, F. P. 182, 183, 194
Costa, L. M. 292
Cöster, G. 188, 222
Coulombeau, C. 360, 498
Coulter, G. A. 370
Coulter, L. V. 499
Court, R. 479
Courtot-Coupez, J. 499
Couto, A. 398
Covington, A. K. 30, 32, 42, 57, 73, 75, 78, 79, 499
Covington, J. B. 113–115
Cowley, D. J. 335
Cox, B. 191
Cox, B. G. 30, 139, 140, 249, 250, 254, 257–259, 267
Cox, G. S. 292, 296
Cox, R. A. 99
Cox, R. H. 388
Cradock, S. 488
Craft, Q. D. 253
Craig, A. C. 335, 414, 415, 429
Craig, R. E. R. 490, 491
Cram, D. J. 139, 140, 147, 251, 253, 259–263, 265, 266
Cramb, D. T. 353
Cramer, C. J. 38, 95, 131, 329, 345, 349, 360, 381, 453, 466, 467
Cramer, R. D. 29, 84, 86, 87
Crass, G. 477
Creary, X. 52, 54, 407, 408
Cressmann, H. W. J. 336
Criegee, R. 204, 205, 207, 285, 307
Criss, C. M. 32
Crombie, L. 132

Crugeiras, J. 293
Cruz, R. 129
Csizmadia, I. G. 103
Cuevas, G. 131
Cuma, M. 17
Curran, D. P. 197, 320, 321, 488, 505
Curtiss, L. A. 19, 21

d
Dabard, O. A. G. 4
Dack, M. R. J. 10, 30, 35, 63, 65, 68, 121, 147, 216, 220, 222, 223, 225, 308, 310–312, 314, 390, 393, 411
Dahmen, A. 198
Dähne, S. 330, 331, 335, 341–345, 347, 378, 379, 429
Dalati, M. T. 432, 433, 456, 457
Dallas, A. J. 399, 460
Dalton, L. R. 369
Daltrozzo, E. 113, 115, 116, 335, 341, 345
Daniel, D. C. 18
Danneberg, J. J. 108
Darley, P. E. 36
Darvas, F. 468
Darwish, D. 62
Darwish, M. 334
Das, G. P. 38
Das, M. N. 499
Das, P. K. 42
Da Silva, L. 345
Dathe, K. 363, 369
Datta, A. 243
Daudel, R. 31
Daum, H. 477
Dau-Schmidt, J.-P. 408
Davey, R. J. 490
Davies, C. W. 237
Davies, M. 31
Davies, M. M. 73, 75, 79, 336
Davies, R. H. 113–115
Davis, A. M. 30
Davis, G. T. 376, 377
Davis, J. C. 382, 384
Davis, J. E. 477
Davis, K. M. C. 376, 417, 425
Dawber, J. G. 42, 378
Dawber, J. J. 429
Day, M. C. 397
Dayal, S. K. 437
De Boer, Th. J. 370, 372
De Gracia Nascimento, M. 428
De Jong, F. 266, 268
De Juan, A. 42
De La Mare, P. B. D. 176, 177
De Meyer, C. 113, 114
De Oliveira, G. 162
De Paz, J.-L. 429, 439, 440, 443, 447, 453, 461
De Renzo, D. J. 500
De Schryver, F. C. 43
De Silva, A. P. 355, 357
De Simone, J. M. 70, 73, 315, 505
De Souza, R. F. 322, 323
De Vijlder, M. 417, 425
De Witte, W. J. 376
De Wolf, E. 320, 321, 488
Dean, I. M. 386
Deb, K. K. 382, 384
Debenedetti, P. G. 70, 73
Debye, P. 2, 13, 237
Dechter, J. J. 41, 376, 377
Deckers, H. 336, 354
Decouzon, M. 101
Decroocq, D. 166, 227, 229, 404, 409
Deelman, B.-J. 320, 321, 488
Dega-Szafran, Z. 121
Degtereva, L. I. 329, 345, 363, 364, 369, 376, 378, 381
Deguchi, Y. 370, 372
DeHazan, Y. 333
Dehmlov, E. V. 266, 318
Dehmlow, S. S. 266, 318
Dei, A. 147
Dekant, W. 500
Del Bene, J. E. 348
Delaney, M. F. 84
Delker, D. A. 479
Della, E. W. 242
Dellinger, M. 45
Delpuech, J. J. 36, 168, 234, 249
Demange-Guérin, G. 82, 252
DeMello, N. C. 177
Dempsey, B. 99
Demus, D. 58
Demyanow, P. I. 271, 272
Deng, T. 416
Denham, H. 243
Denisov, E. T. 200, 201, 207, 208
Denisov, G. S. 17
Denny, R. W. 192, 193, 282
Dent, S. G. 336
Depaemelaere, S. 43
Depuy, C. H. 242
Derevyanko, N. A. 345
Dersch, W. 370
Dershowitz, S. 194
Dervan, P. B. 204, 351
Deshayes, H. 253
Deshpande, D. K. 337, 354
Desimoni, G. 178, 188, 197, 198
Desiraju, G. 15
Devaure, J. 365, 368
Deverell, C. 36, 40
Devlin, C. J. 128
DeVoe, R. J. 281
Dewar, M. J. S. 158, 188
Deyrup, A. J. 77, 99
Dhar, M. L. 163, 165, 169–172, 215, 225, 310
Di Fede, C. 185, 186, 292, 300
Diaz, C. 243, 429, 439, 440, 442, 443, 447, 453, 461
Dickert, F. L. 124, 303
Dickinson, H. R. 479
Dickinson, T. 30, 32, 57, 73, 75, 78, 79
Diederich, F. 121, 139, 141, 142
Diederichsen, U. 505
Diehl, P. 378, 383
Dietrich, B. 266, 268
Dietrich, P. 177
Dietz, F. 131, 331, 340, 347, 352
Dill, K. A. 29, 30
Dilling, W. L. 348–350
Dillow, A. K. 185, 186
Dimroth, K. 40, 42, 330, 332, 333, 389, 390, 411, 416, 417, 425, 426, 428, 448–451
Dimroth, O. 4, 106, 110, 135, 154
Dirksen, H.-W. 82, 252
Doan, P. E. 27
Dobi, P. 338, 339, 412
Dobis, O. 305
Docherty, R. 336, 345
Dodd, G. H. 370
Dodin, G. 116
Dodiuk, H. 355, 417, 425
Dogonadze, R. R. 27, 30

Doherty, R. M. 390, 394–396, 399, 408, 432, 443, 444, 457–461,
Dolbier, W. R. 183
Dondoni, A. 193, 303
Dong, D. C. 358, 429, 430
Donohue, J. A. 298
Donovan, W. H. 138
Doolittle, A. K. 489
Doraiswamy, L. K. 200
Doran, M. A. 50
Dore, J. C. 6, 15, 18
Doring, C. E. 177
Dörr, F. 113, 335, 341
Dorsch, D. 290, 291
Dorsey, J. G. 333, 417, 425, 426, 428, 448, 478
Došen-Mićović, L. 129, 131, 132
Dougherty, D. A. 20
Dougherty, R. C. 453
Douty, C. F. 82, 253
Dowd, W. 404
Doye, S. 264
Doyle, M. P. 68, 360, 362
Drago, R. S. 20, 22, 24, 25, 27, 80, 398, 415, 453, 467
Dreher, M. 438
Drenth, W. 333
Drexler, E. J. 107, 108
Drexler, J. 180
Dreyer, J. 334
Dreyfus, M. 116
Drickamer, H. G. 417, 425, 428
Drougard, Y. 166, 227, 229, 409
Drummond, C. J. 297, 333
D'Souza, M. J. 407, 408
Du Bose, J. C. 205
Dubini-Paglia, E. 334
Dubois, J.-E. 112–116, 348, 378, 414, 415, 429
Duce, P. P. 238, 242
Dudis, D. S. 38
Duggan, P. G. 276
Duggua, H. F. 454
Duismann, W. 203
Dulewicz, E. 121
Dulova, V. I. 75, 79
Dumbacher, B. 337
Dumont, L. 129, 131
Dunbar, B. I. 268
Dunbar, R. C. 100
Dunn, A. D. 475
Dunn, L. C. 180
Dupire, S. 334, 339
Durocher, G. 345
Durov, V. A. 18
Durst, H. D. 268
Durst, T. 70, 77, 99, 101, 102, 489
Dutkiewicz, M. 68
Duve, G. 9, 63, 500
Dvorko, G. F. 165
Dwek, R. A. 107
Dyachkov, A. I. 286
Dydykina, A. P. 201
Dye, J. L. 340, 376
Dykstra, R. R. 55
Dyson, A. J. 69, 202
Dyumaev, K. M. 117, 118, 121

e

Earl, R. A. 61
Earle, M. J. 322–324, 506
Easteal, A. J. 42, 417, 425, 426
Eastes, J. W. 335
Eaton, G. 364–366, 438
Ebel, H. F. 29, 54
Eberbach, W. 281
Eberhardt, R. 71, 424
Ebiike, H. 144
Eckell, A. 191
Eckert, C. A. 147, 185, 186 188, 219, 222, 312, 327, 328
Eckert, K. 334
Eddy, R. W. 238, 239
Edlund, U. 386, 460
Edward, J. T. 119
Edwards, A. G. 338, 339, 412
Edwards, J. O. 242, 244
Edwards, R. T. 367
Effenberger, F. 334
Efimova, T. D. 227, 230
Efremova, L. S. 39
Eggert, H. 106–109, 112, 113
Ehrenson, S. 340, 345, 346
Eiben, R. 55, 137
Eichler, S. 194
Eichler, T. 337
Eigen, M. 304
Eilers-König, N. 334
Eisenberg, D. 6
Eisenthal, K. B. 417, 425
Eistert, B. 107, 108
El Abbady, S. 113, 114
El-Ayaan, U. 330, 340, 345
El Bermani, M. F. 365
El Mouhtadi, M. 379
El Seoud, O. A. 42, 297, 333, 426
El-Zawawi, F. 336
Eldik, R. van 203, 308–310, 314, 315
Élégant, L. 24, 121, 398
Elguero, J. 15, 84, 87, 113, 115
Elias, H. 417, 425, 438
Eliasson, B. 460
Eliel, E. L. 68, 121, 126, 128–133, 288, 360, 362, 396
Eliezer, I. 6
Elkins, D. 399, 400, 460
Ellingsen, T. 264, 268
Ellis, B. 323
Ellison, G. B. 272
Ellison, J. 113–115
El-Nahas, A. M. 345
Eltsov, A. V. 168, 174, 175
Emanuel, N. M. 147, 210, 219
Emsley, J. 15, 16, 106–109, 112, 113
Emsley, J. W. 386
Emslie, P. H. 415
Enders, D. 70, 288
Endres, A. 57, 58, 320, 321, 488, 505
Engberts, J. B. F. N. 27–29, 40, 42, 108, 147, 152, 219, 237, 238, 292, 365, 370, 372, 417, 425, 426, 453, 467
Engel, G. 38
Engel, P. S. 203, 204
Engelbrecht, B. P. 478, 494
Engelhardt, G. 341–343, 378
Engelhardt, H. 478, 492
Engler, E. M. 384, 385
Engstrom, J. P. 205
Engström, M. 370, 372
Ensing, B. 162
Entelis, S. G. 147, 218, 219, 221, 222
Epperlein, J. 334
Eppert, G. J. 492
Epstein, J. 334
Epsthein, L. M. 238
Erlich, R. H. 41
Ernst, S. 339, 340, 429, 430
Ernsting, N. P. 334
Ershov, B. A. 106
Escale, R. 118

Eschner, M. 417
Eskenazi, C. 303
Esselborn, W. 479
Etemad, S. 42
Étlis, V. S. 286
Evans, A. G. 50
Evans, C. C. 48
Evans, D. F. 29, 30, 387, 388
Evans, M. G. 308
Evans, M. W. 29
Evleth, E. M. 370
Exner, O. 84, 85, 103, 115, 128, 131, 132, 136, 349, 390
Eyler, G. N. 203
Eyring, H. 5, 147, 149, 154, 179, 218, 219, 227, 230, 233, 236, 360

f

Fabian, J. 330, 331, 334, 340, 343, 352
Fabian, M. A. 131
Fabian, W. M. F. 115
Faigl, F. 468
Fainberg, A. H. 53, 54, 165, 215, 216, 226, 231, 232, 238, 239, 402–405, 408, 409
Fairclough, R. A. 188
Faita, G. 178, 188, 197, 198
Falbe, J. 488
Famini, G. R. 390, 394, 395, 399, 453, 457, 459, 460, 466, 467
Fărcasiu, D. 274, 403
Fargin, E. 24, 84, 87, 397, 398
Farhataziz 428
Farkas, J. 253
Farris, F. J. 357
Faure, J. 331, 332, 340
Faure, R. 135
Fawcett, W. R. 27, 81, 365, 398, 453, 462
Fayed, T. A. 347
Fedorov, L. A. 120
Fedorynski, M. 266, 318
Fei, Z. X. 294
Feibush, B. 69
Feichtmayr, F. 331
Feigel, M. 50, 51, 125
Feiler, L. A. 180, 181, 193
Feiner, S. 195
Felcht, U.-H. 253
Feld, M. 305
Felder, E. 108
Felger, E. 4
Fenby, D. V. 5
Fendler, E. J. 292, 297, 351
Fendler, J. H. 42, 46, 292, 297, 351, 412
Fenoglio, D. J. 126, 127
Fernandez-Mayorales, A. 144
Fernández-Prini, R. 357, 358, 500
Ferrand, E. F. 265
Ferris, D. C. 453, 467
Fessenden, R. W. 354, 357
Feuerlein, F. 253
Fialkov, Yu. Ya. 39, 57, 96
Fico, R. M. 56
Fidanza, J. 121
Fidler, J. 362
Fiedler, H. 188, 410
Fiedler, P. 128, 131, 132
Field, K. W. 107, 108
Figeys, H. P. 369
Figueras, J. 335
Filley, J. 242, 272
Fineman, M. A. 201
Finke, M. 253
Finkentey, J.-H. 334
Finston, H. L. 121, 123
Firestone, R. A. 188, 191, 309
Firman, P. 139, 140
Firth, D. W. 107
Fischer, E. 143
Fischer, H. 304, 307
Fischer, K. 333, 478
Fischer, R. D. 69
Fischer, S. 116
Fischer, T. 323
Fisera, L. 191
Fish, R. H. 58, 320, 321, 488
Fisher, N. 479
Fisher, R. D. 404
Fisichella, S. 128
Flamm, S. J. 335
Flannery, J. B. 124, 125
Flek, J. 471
Fleming, I. 178, 187, 188, 191, 195, 197
Flessner, T. 264
Fletcher, T. L. 254
Fletton, R. A. 479
Flippen, J. 179, 180, 193
Flitsch, W. 118, 119
Flohr, A. 62, 189, 268
Floriano, M. A. 306, 313, 314
Floris, F. M. 166, 168
Flowers, R. A. 52
Flunt, R. 396, 456
Fogassy, E. 468
Foglia, T. A. 251
Fomin, V. A. 286
Fompeydie, D. 124
Fong, C. W. 378
Fong, G. D. 308
Foote, C. S. 192, 193, 282
Forbes, J. W. 351
Ford, R. A. 74, 75, 79, 471, 491–493
Ford, W. T. 60, 244–246
Foreman, M. I. 376, 378, 382, 384
Forlani, L. 172
Forni, A. 69
Forsén, S. 106
Forster, H. 370
Förster, Th. 341–343
Förster, W. 131, 347
Fortier, D. 259
Foster, R. 19, 20, 334, 415
Foster, W. R. 201
Fouassier, J. P. 331, 332, 340
Fowler, F. W. 453
Fowler, P. W. 10
Fox, I. R. 376, 377, 437
Fox, M. A. 325
Fox, M. F. 340, 479
Fraenkel, G. K. 369, 370, 373
Franceschi, J. de 24, 84, 87, 397, 398
Franck, E. U. 327, 328
Franck, J. 341, 352
Franco, M. L. T. M. B. 272
Frank, H. S. 6, 29, 35, 37, 38
Frank, J. 113–115, 304
Frankel, L. S. 40
Frankland, S. J. V. 354
Franks, F. 6, 8, 27–30, 39, 73, 119
Fratiello, A. 36, 40
Fratini, G. 24, 398
Freed, J. H. 369, 370, 373
Freedman, H. H. 50
Freeman, F. 176, 177
Freeman, N. J. 106–109, 112, 113

Freemantle, M. 322, 324, 506
Freitag, W. 378
Freitas, M. P. 129, 131
Frendo, D. T. 59, 300
Frensdorff, H. K. 266
Fried, B. 494, 496
Friedl, Z. 128, 131, 132
Friege, H. 188
Frimer, A. A. 192, 193, 282, 283
Fringuelli, F. 294
Frisch, M. J. 104, 113–115, 127, 131
Fritsch, J. 17, 121, 122
Fritsch, K. D. 303
Froelich, P. 417, 425
Fromherz, P. 44
Frøyen, P. 194
Fruchier, A. 84, 87
Fruwert, J. 126, 363, 369
Fry, J. L. 166, 403
Fuchs, O. 8–10, 42, 57, 63, 471, 500
Fuchs, R. 244–247, 396, 461
Fujio, M. 408
Fujisaka, T. 308
Fujisawa, T. 23, 398
Fujiwara, K. 303
Fujiyama, T. 369
Fukuda, E. K. 276
Fukuda, Y. 330, 340, 345
Fukuto, J. M. 175
Fukuzumi, S. 175
Fulava, J. 113, 114
Fuoss, R. M. 46, 47, 52, 54, 227, 229
Furton, K. G. 60
Furukawa, N. 253
Fuson, N. 365, 367
Futrell, J. H. 99

g

Gabidulina, N. A. 499
Gabrielsen, B. 333, 428, 448, 478
Gageiro, V. 428
Gagnaire, D. 477
Gaines, G. L. 344
Gajewski, J. J. 195, 197
Gal, J.-F. 23, 24, 84, 87, 101, 103, 121, 123, 366, 397, 398, 457, 458
Galdames, J. S. 345
Gale, P. A. 243
Galin, J. C. 113
Gall, J.-S. 48, 244, 245, 263, 264
Gallagher, S. H. 331
Galletti, P. 289
Gálvez, N. 505
Galy, J. P. 135
Gan, G.-H. 478
Ganapathy, S. 299, 300, 302
Gandolfi, R. 192
Ganem, B. 195, 197
Gao, J. 113–115, 131, 162, 343
Garcia, A. E. 29, 30
Garcia, J. I. 188, 401
Garcia, M. V. 435
Garcia-Blanco, F. 243, 443, 461
Garcia-Rosas, J. 267
Garde, S. 29, 30
Gardecki, J. A. 354
Garnovskii, A. D. 126
Garrett, B. C. 149
Garst, J. F. 54
Gattermann, L. 61
Gäumann, T. 102
Gavrilov, V. V. 125, 168
Gawinecki, R. 117
Gawley, R. E. 288
Gázquez, J. L. 80
Gee, G. 64, 65
Geerlings, P. 162
Geiseler, G. 126, 369
Geiss, F. 107, 108
Geisser, P. 116
Gendell, J. 369, 370, 373
Genkina, N. K. 271
George, M. V. 191
Gerger, W. 24–26, 49, 376–378, 383, 437, 438, 465
Gerig, J. T. 378
Germer, H. A. 345
Gerson, F. 374
Gerus, I. I. 178
Gervay, J. 189, 190
Gholami, M. R. 296
Ghoneim, N. 42, 330, 340, 345, 352, 353, 357, 358
Giacomini, D. 289
Giam, C. S. 378
Gibian, M. J. 304
Gibson, H. W. 336
Gielen, M. 175, 404, 409, 410
Giera, H. 191
Giese, B. 178
Giguère, P. A. 73
Gil, J. 401
Gil-Av, E. 69
Gill, D. S. 40
Gillard, R. D. 42
Gilligan, W. H. 335
Gindin, V. A. 106
Ginsberg, H. 166
Ginsburg, O. F. 117
Giorgianni, P. 193
Girardello, M. A. 478
Gladysz, J. A. 320, 321
Glasstone, S. 147, 149, 154, 179, 218, 219, 222, 227, 230, 233, 236
Gleave, J. L. 169
Glebov, E. M. 327
Glemser, O. 61
Glew, D. N. 46
Glick, R. E. 376
Glikberg, S. 462, 468, 475
Glukhovtsev, M. N. 162
Gnamm, H. 10, 57, 471
Goddard, R. 56
Goddard, W. A. 283
Goddu, R. F. 479
Godfrey, N. B. 65, 471
Goeden, H. A. 10
Goetz, E. 348, 414, 415, 429
Goggin, P. L. 24, 25, 81, 402
Goh, E. H. 478
Goitein, R. 254
Gokel, G. W. 266, 268, 318
Gold, V. 99
Goldshtein, I. P. 19
Goldstein, J. H. 388
Golinkin, H. S. 313
Gollnick, K. 282
Golovnaya, R. V. 444, 445, 492
Golubov, N. S. 17
Gómez, J. 398
Gompper, R. 178, 179, 188, 193, 269, 273
Goncalves, R. M. C. 468
Gonikberg, M. G. 311
Gonzalez, G. 40, 365
Gonzalez, T. 268
Goodby, J. 58
Gopalan, S. 70, 73
Gorbenko, G. P. 333
Gordienko, V. G. 329, 345, 363, 364, 369, 376, 378, 381
Gordon, A. 114
Gordon, A. J. 74, 75, 79, 471, 491–493

Gordon, J. E. 30, 36, 38, 40, 42, 46, 48, 52, 54, 59, 60, 244–247, 252, 262, 292, 293, 374, 415, 443, 457
Gordon, M. H. 129
Gordon, M. S. 15, 17, 103
Gorman, A. A. 334
Görner, H. 124, 125
Gorrie, T. M. 16
Gosh, B. D. 21
Gosney, I. 489
Goto, F. 287
Gottarelli, G. 303, 362
Gotthardt, H. 334
Götz, H. 367
Gough, T. E. 329, 342, 363, 364, 369, 376, 378
Gould, E. S. 238, 402
Gowland, J. S. 412
Grabowski, E. J. J. 104, 505
Grabowski, J. J. 242
Grabowski, S. J. 104
Grabowski, Z. R. 354
Graczyk, P. P. 131
Grader, G. S. 333
Graf, E. 266
Graf, H. 180
Graf, R. 181, 193
Grahn, W. 393
Gramstad, T. 365
Grande, K. D. 109
Grant, D. J. W. 9
Grant, H. G. 282, 283, 378
Grashey, R. 192
Grate, J. W. 243
Grätzel, M. 292
Gray, G. W. 58
Grayson, M. 431
Graziano, G. 29, 30
Graziano, M. L. 284
Grdina, M. J. 193
Greco, F. A. 29
Green, C. E. 463
Green, D. P. 55, 298
Greenberg, A. 200, 207, 208
Greenberg, M. S. 41, 376, 397
Greene, F. D. 201
Greenwald, R. 260
Greiciute, D. 177
Greif, N. 199
Greim, H. 500, 503
Grellier, P. C. 166, 227, 228, 230, 257, 259
Grellier, P. L. 400, 401, 408, 449, 458
Grellmann, K. H. 354
Grenby, T. H. 19
Gressner, M. 316
Griebenow, K. 144
Grieco, P. A. 57, 62, 189, 197, 268, 294, 295, 505
Grieger, R. A. 188, 222, 312
Grieser, F. 297, 333
Griffiths, L. 129
Griffiths, T. R. 340, 390, 393, 412–414, 417
Grimm, A. G. 166
Grimm, K. 175
Grimme, S. 357
Grisdale, E. E. 244–246, 268
Gritzner, G. 33, 81, 139, 402, 500
Grob, C. A. 185, 284
Groß, V. 254
Grossi, L. 370, 374
Grotemeier, G. 290, 291
Grozema, F. C. 349, 360
Grubbs, R. H. 478
Gruda, I. 336
Grundmaier, M. 306
Grunwald, E. 93, 99, 109, 147, 149, 218, 219, 393, 402–405, 408
Grushka, E. 268
Gschwind, R. M. 56
Guérin, M. 67
Guerra, M. 370, 374
Guglielmetti, R. 337
Guibe, F. 263, 271
Guidetti, S. 198
Guillerez, J. 113–115
Guimon, C. 113–115
Guiot, S. 499
Gulbrandsen, T. 280
Gumbel, G. 417, 425
Gumby, W. L. 265
Günthard, H. H. 479
Günther, H. 136, 376, 378
Gupta, M. N. 143–145
Gupte, S. S. 188, 189
Gurney, R. W. 37
Guryanova, E. N. 19
Güsten, H. 292
Gut, R. 499
Gutbezahl, B. 99
Gutmann, V. 19–22, 24–27, 46, 48–50, 57, 58, 79–81, 123, 147, 215, 257, 263, 264, 278, 376–378, 383, 390, 393, 396–398, 411, 437, 438, 462, 465
Gvozdeva, E. A. 253, 260
Gyenes, I. 496

h

Ha, S. 131
Haak, J. R. 417, 425, 426
Habashy, M. M. 336
Haberfield, P. 103, 147, 166, 188, 189, 253, 259, 260, 270
Hadži, D. 15–17
Hädicke, E. 203
Hadizamani, K. 411
Haessner, C. 350
Hafner, K. 112
Hafter, R. 129
Hagen, R. 112
Hagii, H. 286
Hahn, K. 370, 372
Hahn, S. 23, 397
Haines, R. I. 339
Halban, H. von 166
Halbert, T. R. 303
Hale, H. K. 180
Halevi, E. A. 178, 187, 188, 191, 195, 197
Hall, J. R. 203
Hall, R. J. 131
Hallam, H. E. 364–367, 383, 488
Haller, R. 119
Hallpap, P. 468
Halpern, M. 266, 318
Ham, J. S. 345
Hamann, S. D. 310
Hames, D. A. 46
Hamlet, Z. 188, 190, 191, 280, 291, 295, 404, 410, 446
Hamlin, R. 136
Hammack, W. S. 417, 425, 428
Hammam, E. 345
Hammerich, O. 106–109, 112, 113
Hammett, L. P. 30, 77, 99, 147, 390, 392
Hammond, G. S. 306
Hammons, J. H. 77
Hampe, M. 490
Hampel, B. 478, 479
Hampel, M. 188, 410
Han, C. 428, 448
Hancock, R. D. 80
Handa, M. 11, 67
Handel, H. 268
Handy, P. R. 36, 376
Hannay, J. B. 70

Hansch, C. 30, 87, 399, 400, 460
Hansen, D. L. 479
Hansen, H.-J. 196
Hansen, P. E. 387, 388
Hanson, C. 490
Hanson, M. P. 271
Hantzsch, A. 3, 4, 76
Haque, M. S. 191
Harano, K. 197
Harbusch, E. 332, 333, 417, 425, 428
Harding, L. B. 283
Harkness, J. B. 156
Harley, J. D. 166, 230, 234
Harned, H. S. 498
Harris, E. M. 121
Harris, H. C. 274, 275
Harris, H. P. 268
Harris, J. M. 166, 169, 333, 403, 408, 444, 457, 458
Harris, P. A. 113
Hart, D. J. 60
Hart, D. M. 415
Hart, H. 112
Hartley, G. S. 44
Hartmann, H. 100, 166, 167, 227, 228, 312, 313, 330, 334
Hartmann, I.-M. 281
Hartmann, W. 281
Hartnagl, M. 175
Hartshorn, M. P. 287, 404
Hase, W. L. 162
Hasegawa, H. 290
Hashida, Y. 175
Hashimoto, H. 200, 201
Hashimoto, Y. 106–109, 112, 113
Hassan, M. A. 180
Hassel, O. 19
Hassner, A. 181, 193
Hastie, J. W. 59
Hatton, J. V. 382
Hatz, R. 58
Hauri, R. J. 60, 244–246
Hauthal, H. G. 82, 177, 188, 252, 262, 410
Havlas, Z. 15, 17
Hawkes, S. J. 46, 498
Hawkins, D. T. 6
Hawthorne, S. B. 492
Hayami, J. 147
Hayat, H. 370–372, 374
Hayes, W. P. 348, 349
Haymet, A. D. J. 29, 30
Haynes, C. 325
Hayon, E. 479
Headley, A. D. 105, 387, 453, 467
Healy, T. W. 297, 333
Heathcock, C. H. 70, 183, 184
Hecht, H. 49
Heckenkemper, G. 417
Hecker, E. 490, 492, 493
Hedaya, E. 205, 207
Hedwig, G. R. 249, 250, 254, 257–259
Heess, R. 369
Hefter, G. T. 25, 33, 34, 81, 251, 256, 313, 402, 478
Heftmann, E. 493
Heidenhain, F. 55, 120, 137
Heilbronner, E. 112, 331, 367
Heimgartner, H. 191
Heineken. F. W. 370
Heinrich, G. 292
Heiss, J. 345
Heitele, H. 356
Helbert, M. 366, 398
Helgeson, R. C. 266
Helmchen, G. 290, 291
Hempel, R. 117
Hemptinne, A. von 166
Henchmann, M. 159–161
Henderson, D. 5
Henderson, W. G. 103, 104
Hendrickson, D. N. 417, 425, 428
Hendry, D. G. 210
Henrichs, P. M. 405
Henrion, A. 87, 468
Henrion, R. 87
Hensel, F. 62
Hensel, T. 363, 369
Henty, D. W. 479
Herbrandson, H. F. 222, 443, 457
Herges, R. 281
Herk, L. 305
Herodes, K. 42
Herold, B. J. 272
Herrebout, W. A. 127, 131
Herrmann, R. 471
Herrmann, W. A. 323, 506
Herte, P. 200, 201, 222, 224, 317
Hertz, H. G. 30, 35, 36, 38
Herzfeld, R. 115, 417, 425, 426
Heseltine, D. W. 335, 336, 344, 411, 414, 415, 429
Hesse, G. 478
Hetzer, H. B. 336
Heublein, G. 50, 176, 278, 279, 468
Heydtmann, H. 166, 227
Heyes, D. M. 5
Hiatt, R. 202, 203, 286
Hiatt, R. R. 204
Hibbert, F. 15
Hickey, K. 365
Hicks, J. 417, 425
Hida, M. 334
Hidalgo, H. 370
Hidber, A. 69
Hierl, P. M. 159–161
Higuchi, T. 9
Hildebrand, J. H. 5, 9, 10, 29, 63, 65, 219–221, 443, 471
Hill, R. K. 187
Hillier, I. H. 113–115, 131
Hilmersson, G. 142, 143
Hilvert, D. 243
Hinchcliffe, A. J. 488
Hinckley, D. A. 123, 124
Hine, J. 102
Hine, M. 102
Hinman, R. L. 205, 207
Hinsberg, W. D. 204, 351
Hinshelwood, C. N. 166, 188
Hinton, J. F. 30, 35, 36, 38, 40, 84, 147, 215, 219, 226, 234, 238, 376, 377
Hiraoka, K. 103, 398, 438
Hirata, F. 103
Hirata, H. 59, 301, 302
Hiromi, K. 225
Hirose, Y. 144
Hirota, H. 240
Hirota, N. 370
Hirsch, J. A. 390, 393, 411
Hirsch, M. S. 453, 467
Hitzler, M. G. 326
Hiyama, H. 117, 118
Ho, S. S. J. 60
Ho, T.-L. 80
Hobbs, S. J. 197
Hoblack, N. 339
Hobza, P. 10, 15, 17, 19–21
Hochstrasser, R. M. 348
Hoefnagels, J. 337, 448
Hoekman, D. 399, 400, 460
Hoekstra, M. S. 477

Hofer, E. 117, 118
Hofer, O. 130, 131, 396
Hoffman, D. M. 159, 201
Hoffmann, E. de 166, 229, 238, 241, 242
Hoffmann, H. M. R. 192
Hoffmann, J. E. 253
Hoffmann, M. 6
Hoffmann, M. M. 71
Hoffmann, R. 178, 187, 188, 191, 195, 197
Hoffmann, R. W. 70, 288
Hofmann, D. 393
Hogarth, J. 70
Hogen-Esch, T. E. 54, 55
Hogg, D. R. 176
Hogiu, S. 334
Hohlneicher, G. 115, 116, 341, 343, 378
Holder, R. W. 271
Holler, M. 113
Holliday, A. K. 57
Hollmann, G. 428, 429
Hollosi, M. 360
Holmboe, E. S. 107
Holton, D. M. 370, 372
Holtz, D. 103, 104
Homer, J. 376, 378, 381, 382
Honig, B. 96
Hoogenboom, B. E. 490
Hoover, D. J. 69, 471
Hopkins, H. P. 244–246, 268
Hoq, M. F. 339
Horák, M. 364, 365, 367
Horibe, I. 385
Hormadaly, J. 428
Horne, R. A. 6, 7
Horng, M. L. 354
Horpp, D. N. 192, 203
Horrom, B. W. 264, 265
Horváth, I. T. 58, 319–321, 488
Hosomi, A. 200, 207, 208
Hoss, R. 268
Houk, K. N. 131, 177, 178, 187, 188, 191, 195, 197, 243
Houlikan, W. J. 489
Houriet, R. 102
House, H. O. 288, 489
Houssa, A. J. H. 477
Houston, P. L. 147, 215, 304, 305
Howard, J. A. 210
Howery, D. G. 84, 85
Hoz, S. 124, 125, 242
Hrovat, D. A. 201, 300, 301
Hsi, S. C. 365
Huang, P.-K. 413
Huang, X. 60
Hubbard, C. D. 308, 310
Huber, H. 69, 191, 198, 381, 383
Huber, M. 370, 372
Huber, W. 496
Hubert, A. J. 260
Hückel, E. 2, 237
Huffman, H. L. 252
Huggins, C. M. 369
Huggins, M. L. 15
Hughes, D. L. 104, 244
Hughes, E. D. 163, 165, 168–172, 215, 225, 276, 277, 310
Huisgen, R. 134, 178–181, 191–193, 198, 201, 280, 312, 477
Hulme, L. 331, 340
Hummer, G. 29, 30
Humphreys, R. R. W. 202
Hünig, S. 273, 330, 336, 337
Hunt, F. 326
Hunter, C. A. 20
Huot, J.-Y. 27, 30
Huppert, D. 36
Huque, E. M. 27, 30
Hush, N. S. 330, 334, 340, 345
Hussey, C. L. 59–62, 322
Hutchings, M. G. 334
Hutchins, R. O. 251
Hutchinson, M. H. 343
Hutchison, E. 42, 45
Hutchison, J. D. 244, 245, 263
Hutson, J. M. 10
Hütte, S. 56
Hutton, H. M. 388
Huyser, E. S. 155, 200, 201, 207, 208, 212, 213, 215, 216
Huyskens, P. 6, 10
Hvidt, A. 29, 30
Hyatt, J. A. 327, 424
Hyne, J. B. 313
Hynes, J. T. 54, 121, 147–149, 151, 217, 218, 223, 224, 352, 355, 356

i

Ibáñez, F. 428
Ichikawa, T. 39
Iesce, M. R. 284
Iffland, D. C. 269, 477
Ihrig, A. M. 288
Ikariya, T. 326
Ikegami, Y. 365, 369, 373, 374
Ikushima, Y. 327
Ikuta, S. 247
Illman, D. L. 84
Ilyasov, A. V. 370
Imai, Y. 333
Imakura, Y. 296
Imanishi, S. 348, 349, 367
Imoto, T. 312, 413, 417, 425 428
Inagaki, Y. 135
Inamoto, N. 135
Inch, T. D. 69, 471
Infante, G. A. 351
Ingold, C. K. 3, 147, 155, 163, 169–172, 215, 225, 276, 277, 310, 489
Ingold, K. U. 210–213, 304
Inman, D. 59
Inoue, T. 27, 365, 435, 436
Inuzuka, K. 348, 349, 367
Ioffe, A. 166, 168
Iqbal, M. 137
Iqbal, R. 137
Ireland, R. E. 291
Irish, D. E. 329, 342, 363, 364, 369, 376, 378
Isaacs, N. S. 183, 195, 198, 308, 314, 413, 455
Isenberg, N. 15
Ishchenko, A. A. 331, 340, 345, 352
Ishii, N. 117
Ishii, Y. 281
Ishikawa, M. 334
Ishizu, K. 370, 372
Iskhakova, G. G. 125, 168
Ismailov, A. I. 119
Isohisa, A. 188
Isozaki, T. 374
Israelachvili, J. N. 10, 11, 29, 42, 43, 46, 48
Ito, M. 348, 349, 367, 413
Ito, O. 213, 334
Ittoh, M. 168
Ives, D. J. G. 8, 39
Ivleva, L. P. 75, 79
Iweibo, I. 337, 345, 354
Izatt, R. 268
Izmailov, N. A. 47, 96
Izutsu, K. 99, 121, 496, 500

j
Jackman, L. M. 55, 269
Jacknow, B. B. 212
Jackson, J. A. 36
Jackson, N. M. 387
Jackson, P. F. 365
Jackson, W. G. 462
Jackson, W. R. 354
Jacob, S. W. 82, 252
Jacobson, P. 200
Jacobson, R. J. 351, 358
Jacobus, J. 184, 185, 449
Jacques, P. 18, 42, 336, 344, 345
Jaffé, H. H. 10
Jaggi, D. 282, 283
Jagodzinski, P. W. 365
Jähme, J. 274, 403
Jahnke, U. 177
Jakubetz, W. 30, 36, 37
Jalander, L. 69
Jallali-Heravi, M. 378, 381
James, K. C. 8
James, R. V. 378, 383
Jameson, C. J. 376, 378
Jana, D. 499
Janardhanan, S. 38
Jancsó, G. 5
Jander, G. 49, 57, 499
Jander, J. 57
Janowski, A. 351, 429, 430
Janz, G. J. 59, 60, 496, 500
Janzen, E. G. 370
Jarrouse, J. 318
Jasien, P. G. 334
Jauquet, M. 329, 363, 364, 376, 378, 380, 381, 384, 385
Jefford, C. W. 193, 282, 283
Jeffrey, G. A. 15, 17
Jehlička, V. 334
Jencks, W. P. 121, 275
Jenkins, A. D. 5
Jenkins, P. W. 335, 414, 415, 429
Jenner, G. 188, 194, 296, 308, 311, 314
Jensen, F. 162
Jensen, F. R. 175
Jensen, J. H. 15, 17, 103
Jensen, W. B. 19, 79
Jessop, P. G. 326
Jewell, D. J. 364
Jewett, J. G. 282, 283
Jhon, M. S. 5, 160
Ji, M. 326
Jinnouchi, Y. 351
Jochims, J. C. 387, 477
Joesten, M. D. 15, 16, 364
Johannesen, R. B. 431
Johansson, E. 386
John, M. 56
Johnels, D. 386, 460
Johnson, A. P. 276
Johnson, B. F. G. 4, 506
Johnson, B. P. 57, 333, 417, 425, 426, 428, 448, 478
Johnson, C. R. 339
Johnson, D. A. 333, 416
Johnson, P. G. 471
Johnson, W. C. 479
Johnston, K. P. 113, 114, 325
Johnston, M. D. 376, 378, 380, 387, 388
Jolicoeur, C. 27, 30
Jonathan, N. 365
Jones, D. C. 369
Jones, F. M. 104
Jones, G. 354
Jones, G. A. 177
Jones, H. W. 402–405, 408
Jones, M. E. 272
Jones, M. N. 42
Jones, M. T. 370
Jones, R. A. Y. 77, 147, 219, 367
Jones, W. M. 404
Jones-Hertzog, D. K. 185
Jores, L. 16
Jorgensen, W. L. 38, 53, 54, 126–128, 131, 132, 158, 185, 188, 189, 197, 201
José, S. M. 147, 158, 159
Josien, M.-L. 364, 365, 367
Jouanne, J. von 179–181, 193, 312
Jovanovic, M. V. 135
Juaristi, E. 77, 99, 101, 131
Jung, M. 71, 73
Jung, M. E. 189, 190
Jungers, J.-C. 166, 227, 229, 404, 409
Jurczak, J. 308, 314
Jurczyk, S. 406, 407
Juřina, J. 425, 448
Just, G. 177, 188, 410
Just, M. 334
Jutila, M. 386
Jutz, C. 341, 343, 378

k
Kaatze, U. 60
Kabachnik, M. I. 121, 123, 253
Kadaba, P. K. 84, 191, 192, 254
Kadish, K. M. 475
Kadobayashi, Y. 45
Kadzyauskas, P. P. 177
Kagan, H. B. 303
Kagiya, T. 27, 365, 435, 436
Kahlert, K. 188, 410
Kahn, M. 238, 240, 241
Kaifer, A. E. 139, 142
Kaifu, Y. 359
Kaim, W. 62, 339, 340
Kajimoto, O. 108, 325
Kajtár, J. 360
Kajtár, M. 360
Kakimoto, M. 369
Kálal, J. 413
Kalantar, T. H. 298
Kalas, R.-D. 116
Kalidas, C. 33, 38, 40
Kalinowski, H.-O. 477, 479, 485, 487
Kalinowski, M. K. 429
Kálmán, E. 27, 30
Kalyanam 383–385
Kalyanasundaram, K. 292, 358
Kamaev, F. G. 119
Kamil, W. A. 403
Kamimura, R. 117
Kamiya, Y. 210
Kamlet, M. J. 15, 18, 19, 24, 335, 390, 393–396, 399, 403, 429, 431, 432, 434–437, 442, 444, 447, 450–453, 456–461, 464, 465
Kamounah, F. S. 115
Kampars, V. 21, 334
Kampas, F. J. 345
Kanagasabapathy, V. M. 275
Kaneda, T. 266
Kanski, R. 330
Kapecki, J. A. 193
Kar, T. 15, 17
Karabatsos, G. J. 126, 127
Karagiannidis, P. 113, 114
Karcher, W. 49
Karelson, M. 69, 90, 91, 108, 113, 166, 168, 329, 345, 451–453
Kariya, K. 144
Karle, I. 179, 180, 193
Karplus, M. 131

Karpovich, D. S. 358
Kasha, M. 348–350
Kaspar, R. 205, 207, 285
Kass, S. R. 272
Kataoka, M. 11, 66
Kato, A. 106–109, 112, 113
Kato, M. 127, 131
Kato, S. 103
Kato, T. 296
Katritzky, A. R. 69, 72, 73, 90, 91, 108, 113–116, 164, 166, 169, 275, 367, 393, 453
Kauffman, G. B. 73
Kauffman, J. F. 13, 71, 324
Kauffmann, E. 243
Kaufmann, H. 156, 188
Kaul, B. L. 117
Kaul, C. 71, 73
Kaupp, G. 70, 73, 505
Kauranen, P. 166, 227
Kauzmann, W. 6
Kawamura, T. 370, 372
Kawata, M. 103
Kawazura, H. 135
Kawski, A. 345
Kayser, E. G. 335
Kazaryan, N. A. 496
Kazmaier, P. M. 124, 125
Kebarle, P. 99, 100, 103, 147, 157, 158, 247
Keck, A. 505
Kecki, Z. 437
Keefer, R. M. 188
Keesee, R. G. 162
Keesom, W. H. 11, 12
Keim, W. 60, 323, 324, 488, 506
Keiser, W. E. 479, 483
Keith, T. A. 127, 128, 131
Kelker, H. 58, 59
Kelm, H. 179–181, 188, 193, 203, 312, 313
Kemell, M. 42
Kemp, D. S. 243
Kende, A. S. 191
Kennard, O. 16
Kenyon, J. 477
Kerber, R. 200, 201, 210
Kerridge, D. H. 59
Kersten, B. R. 60
Kessick, M. A. 404
Kessler, H. 50, 51, 125, 292
Kessler, M. A. 357, 417
Kessler, T. E. 365
Kester, T. C. 253
Ketlinski, V. A. 168
Keutel, D. 348, 351
Kevan, L. 39
Kevill, D. N. 166, 403, 405–408
Keyes, G. H. 336, 344, 411
Khaddar, M. R. 36
Khajehpour, M. 13
Khaledi, M. G. 333, 417, 425, 426
Khalil, F. Y. 194
Khan, A. Y. 137
Khananashvili, N. L. 401
Kheifets, G. M. 106, 113
Khim, Y. H. 169
Kibler, L. M. 205, 207
Kier, L. B. 69
Kierkus, P. C. 505
Kikta, E. J. 268
Kikuchi, O. 373
Kim, C. 453, 467
Kim, H. J. 148, 149, 151, 217, 218
Kim, J. Y. 499
Kinart, C. M. 39
Kinart, W. J. 39
King, B. A. 253
King, E. J. 73, 75, 78, 79, 95, 121, 498
King, J. F. 404, 415
Kingsbury, C. A. 147, 253, 260
Kinoshita, T. 51, 52
Kintzinger, J.-P. 243
Kipkemboi, P. K. 42
Kiplinger, J. P. 103, 159, 162
Kiprianov, A. I. 330
Kirchner, B. 69
Kirk, D. N. 360
Kirkland, J. J. 445, 494, 495
Kirkwood, J. G. 225, 367
Kirmse, K. M. 325
Kirshenbaum, M. R. 107
Kisakürek, M. V. 62
Kiselev, V. D. 125, 168, 188, 311
Kishimoto, S. 117, 118
Kiss, Á. 445
Kistiakowsky, G. B. 156
Kitagawa, T. 51, 52
Kitahara, S. 117, 118
Kittila, R. S. 82, 252, 262
Kittsley, S. L. 10
Kivelyuk, R. B. 448
Kiyooka, S. 287
Kiyozumi, Y. 39
Kjaer, A. M. 334
Klaboe, P. 129
Klamt, A. 345
Klärner, F.-G. 308, 314
Klatsmányi, P. G. 500
Klawunn, P. J. 59, 300
Kleinberg, J. 57
Kleiner, H.-J. 253
Kleine-Weischede, K. 61
Kleinpeter, E. 115
Klement, I. 321
Klesper, E. 494
Klessinger, M. 336
Klibanov, A. M. 143–145
Klier, J. 298
Kliment, M. 199
Kliner, D. A. 259
Klingensmith, G. B. 437
Klippenstein, S. J. 149
Klopman, G. 273
Klose, M. 6
Kloss, A. A. 365
Klyne, W. 360
Knapman, C. E. H. 444, 445
Knauer, B. R. 370, 372, 374, 414, 415, 437
Knoche, W. 38
Knochel, P. 57, 320, 321, 488, 505
Knocks, A. 60
Knoevenagel, K. 200, 201, 222, 224, 317
Knops, P. 6
Knorr, L. 3
Knowles, J. R. 293, 294
Knox, G. R. 251, 260, 261
Knunyants, I. L. 177, 261
Knupfer, H. 192
Knyazhanskii, M. I. 337, 348, 354
Ko, E. C. F. 249, 250, 254, 257–259
Kob, N. 503
Kobayashi, H. 428
Kobayashi, M. 286
Kobilarov, N. 16
Koch, A. 115
Kochansky, J. 10
Kochi, J. K. 20, 52, 139, 175, 307
Kocsis, G. 445
Köhler, G. 357
Koehler, W. 417, 425
Könen, W. 326

Koenig, T. 205–207, 304, 307
Koenig, W. A. 69
Koepp, E. 264
Koepp, H.-M. 39
Koeppl, G. W. 316
Koh, D. S. P. 478
Kohler, F. 5
Kohlmann, S. 339, 340, 429, 430
Kohmoto, S. 193, 282, 283
Kohno, M. 351
Koizumi, M. 359
Kokars, V. 334
Kokocinska, H. 87
Kokubun, H. 121
Koll, A. 122
Kollecker, W. 84, 86
Kollie, T. O. 445
Kolling, O. W. 368, 370, 372, 374, 433, 437
Kolobkov, V. P. 429, 430
Kolodyazhnyi, O. I. 118
Kolodziejski, W. 437
Kolsaker, P. 280
Kolthoff, I. M. 75, 77, 79, 95, 96, 121, 499
Koltsov, A. I. 106, 113
Kolwyck, K. C. 403
Komatsu, K. 51, 52, 139
Komiya, K. 290
Komoto, N. 281
Kondo, Y. 166–168, 312, 313
Kong, Y. S. 160
Konishi, S. 370, 372, 373, 437
Konovalov, A. I. 125, 168, 188
Kooyman, E. C. 284
Koppel, I. A. 42, 101, 103, 231, 381, 383, 390, 393, 403, 411, 417, 425, 426, 428, 436, 453–455
Koppel, J. B. 42, 101, 103, 417, 425, 426, 428
Korewa, R. 117
Korn, J. 188, 410
Kornblum, N. 269, 270
Kornyshev, A. A. 27, 30
Korobov, M. V. 468
Korolev, B. A. 121
Kortüm, G. 18, 30, 31, 32, 46
Koskinen, A. M. P. 143–145
Kosower, E. M. 37, 53, 137, 147, 155, 162, 173, 199, 207–210, 215, 330, 334, 335, 337–339, 351, 352, 355, 356, 390, 393, 411–415, 417, 425
Kosswig, K. 42
Koutek, B. 334
Kovalenko, S. A. 334
Kováts, E. 68, 444
Kovrizhnykh, E. A. 263
Kovshev, E. I. 58
Kowalski, B. R. 84, 85
Kozankiewicz, B. 297, 333
Kozima, K. 129
Kraemer, H. P. 69, 477
Krafft, F. 43
Krämer, A. 71, 73
Krämer, R. 121
Kramers, H. A. 149
Krammer, P. 71, 73
Krasnitskaya, N. D. 429, 430
Krasnoperov, L. N. 327
Kratochvil, B. 496, 500
Kraus, C. A. 48
Krause, V. 109
Kravtsov, D. N. 120
Kredel, J. 291
Kremers, F. 269
Kresge, A. J. 316
Kreshkov, A. P. 496, 498, 499
Krestov, G. A. 30, 39
Kreysa, G. 62
Kreysig, D. 87, 334
Kriegsmann, H. 363, 369
Krienke, H. 30
Krimer, M. Z. 177
Krindel, P. 6
Krishnamurthy, S. 287
Krishtopa, L. G. 327
Kriwanek, J. 333
Kroeger, M. K. 27
Kropf, H. 177
Krygowski, T. M. 104, 398, 417, 425, 426, 433, 453, 454, 462
Kuanar, M. 345
Kubinyi, H. 399, 400, 460
Kubota, S. 369, 374
Kubota, T. 330, 345, 359
Kuboyama, A. 351
Kuc, T. 121
Kucharský, J. 496
Kucher, R. V. 205
Kuczkowski, R. L. 307, 308
Kuhn, H. 345
Kulak, H. J. 251
Kulik, N. I. 165
Kulkarni, M. G. 200
Kumar, A. 189
Kumari, N. 40
Kumoi, S. 428
Kundt, A. 4
Kundu, K. K. 499
Kunerth, D. C. 251
Kuntz, I. D. 380
Kuntz, R. R. 334
Kunz, W. 30
Kunze, O. 50
Kuokkanen, T. 139, 140
Kupfer, M. 468
Kurg, R. 101
Kuroda, T. 281, 282
Kurono, Y. 144
Kurth, Y. 339, 340
Kurts, A. L. 147, 269, 271, 272
Kusabayashi, S. 166–168, 308, 312, 313
Kuta, J. 500
Kuznetsova, O. M. 178
Kwart, H. 187

l

Laar, P. C. 3
L'abbé, G. 192
La Clair, J.-J. 357
La Manna, G. 398, 438, 466
Labes, M. M. 404
Labhart, H. 342
Labský, J. 413
Ladanyi, B. M. 54, 223, 224, 432
Lafrenz, C. 57
Lagalante, A. F. 351, 358
Lagowski, J. J. 46, 57, 59, 73, 79, 82
Lagowski, J. M. 113
Laha, A. K. 42, 357
Lai, D. T. 45
Laidlaw, W. G. 313
Laidler, K. J. 2, 147, 149, 150, 154, 179, 218, 219, 225–227, 230, 233, 236
Laila, A. A. R. 198
Laila, A. H. 183
Lamb, R. C. 201, 202
Lambert, J. B. 20
LaMer, V. K. 236
Lampe, J. 183, 184

Lancelot, C. J. 409
Landers, J. P. 211
Landini, D. 238, 241, 268
Landskroener, P. A. 225, 227, 233
Lane, E. H. 121
Lang, D. 188
Lange, B. C. 269
Langer, W. 70, 477
Langford, C. H. 40
Langhals, E. 191, 333
Langhals, H. 38, 40, 42, 357, 411, 417, 425–430, 478
Lankamp, H. 200
Lantzke, I. R. 329, 342, 363, 364, 369, 376, 378
Lapinski, L. 113, 114
Lappenberg, M. 119
Lapshova, N. I. 227, 230
Larsen, J. W. 338, 339, 412
Lascombe, J. 365
Lasperas, M. 289, 290
Lassau, C. 166, 227, 229, 404, 409
Laszlo, P. 36, 295, 329, 363, 364, 376, 378, 380–385, 388
Latimer, W. M. 15
Lattes, A. 404
Lau, E. Y. 378
Lauer, K. 331
Lauer, W. 109
Laughton, P. M. 315, 317
Laurence, C. 19, 366, 398, 417, 432–435, 437, 453, 456, 457, 459
Lawrence, G. A. 462
Lay, P. A. 331, 462
Layloff, T. 370
Laynez, J. 398
Lazana, M. C. R. L. R. 272
Lazaridis, T. 29, 30
Le Beuze, A. 336, 337, 344, 345
Le Carpentier, J. M. 50
Le Démézet, M. 499
Le Noble, W. J. 179, 180, 193, 269–272, 308–310, 314, 315
Le Rosen, A. L. 331, 335, 340
Le Roux, L. J. 168
Leach, M. R. 79
Leach, S. 45
Leahy, E. M. 505
Leary, J. A. 238, 240, 241
Lebert, K.-H. 100
Lebrilla, C. B. 437
Lecea, B. 182, 183, 194
Lee, K.-H. 478
Lee, T. V. 288
Lee, W. H. 82, 253
Lee-Ruff, E. 100
Lees, A. J. 339, 429, 430
Leffler, J. E. 93, 109, 147, 149, 201, 218, 219, 393
Legrand, M. 361
Lehman, T. A. 100
Lehn, J.-M. 139, 141, 243, 266, 268
Leibfritz, D. 51, 125
Leigh, W. J. 300, 302, 303
Leigh, W. L. 59
Lein, G. M. 266
Leis, J. R. 293
Leitner, W. 70, 73, 315, 326, 505
Leito, I. 42
Lemaire, H. 370
Lemal, D. M. 136
Lemieux, R. U. 119, 121, 126, 129
Lemire, R. J. 84
Lemons, J. F. 36
Leo, A. 30, 399, 400, 460
Leonard, J. E. 265
Leonard, P. 388
Leontev, V. B. 119
Lerf, C. 42
Leroy, G. 135
Leroi, G. E. 270
Les, A. 113, 114
Lesch, W. 365
L'Esperance, R. P. 284
Letcher, T. M. 9
Leupold, D. 330, 331, 341–343, 378
Levillain, P. 124
Levine, R. D. 147
Levy, D. H. 305
Levy, G. C. 378
Levy, J. M. 492
Levy, M. 155, 201
Lewis, D. W. 4, 506
Lewis, E. S. 187
Lewis, F. D. 281
Lewis, G. N. 2, 15, 79,
Lewis, I. C. 376, 377
Lewisky, J. A. 61, 323
Li, A. S. W. 39
Li, C. 162
Li, C.-J. 57, 294, 505
Li, H. 60
Li, J. 38, 349, 360
Li, K. P. 147, 149, 218, 219
Li, Y. 178, 187, 188, 191, 195, 197
Liang, H. 60
Liang, Y.-L. 137, 138
Lichkova, N. V. 75, 79
Lichtenthaler, R. N. 23, 397
Lichter, R. L. 378
Lichtin, N. N. 49, 50
Liddel, U. 382
Lide, D. R. 57, 60, 72, 316, 471
Lieberman, J. F. 200, 207, 208
Liebig, H. 84, 251
Lieder, C. A. 244–246
Lilley, T. H. 187
Lim, D. 188, 189, 197, 201
Limbach, H.-H. 17
Lin, G. M. L. 406, 407
Lin, S. H. 147, 149, 218, 219
Linclau, B. 505
Lincoln, L. L. 335, 414, 415, 429
Lincoln, S. F. 35, 267
Linda, P. 113
Lindeboom, G. A. 1
Lindemann, S. V. 20
Lindermeyer, R. 180
Lindgren, M. 370, 372
Lindh, R. 355
Lindman, B. 44
Lindner, B. 369
Lindner, K. 16
Lindon, J. C. 386
Lindqvist, I. 22
Linert, W. 29, 30, 36
Linkova, M. G. 177
Liotta, C. L. 185, 186, 244–246, 266, 268, 318
Lipiński, J. 334
Lippert, E. 57, 336, 345, 348, 353, 354, 359, 382, 384
Lippmaa, E. 341–343, 378
Liptay, W. 330, 342, 345–348, 354, 358
Little, L. H. 367, 368
Litvinenko, L. M. 227, 229
Liu, G. 365
Liu, G. M. L. 166
Liu, H. 162
Liu, J. 177
Liu, J. H. 300, 301
Liu, K.-T. 408
Liu, L. 376

Liu, L.-J. 412
Liu, Y. 437
Livant, P. 205, 206
Llewellyn, G. 408
Llor, J. 399
Lo, T. C. 490
Loan, J. W. 129
Loan, L. D. 307
Lobanov, V. S. 69
Löbbecke, S. 71, 424
Lobunez, W. 202
Locke, C. J. 15
Lockwood, K. L. 107
Lodzinski, A. 264
Löliger, J. 116
Lomonosov, M. V. 2
London, F. 13, 14
Longhi, P. 498
Loosen, K.-D. 336, 354
Lopata, A. 468
Lopez, A. 67, 133
Lopez Arbeola, I. 123, 124
Lopez, V. 357, 429, 439, 440, 443, 447, 453, 461
Lopez-Mardomingo, C. 128
Lorents, D. C. 8
Lorenz, G. 177
Lorenz, O. 307
Lotfi, M. 188
Lothe, J. J. 129
Lougnot, D. J. 331, 332, 340
Loupy, A. 4, 171, 237, 263, 506
Lovering, D. G. 59
Lowry, A. H. 453, 466, 467
Lowry, T. M. 73, 79
Loze, C. de 438
Lubineau, A. 57, 294–296, 505
Luc Massart, D. 85
Luchetti, J. 295
Lucius, R. 247, 248
Luck, W. A. P. 6, 10
Lucken, E. A. C. 370
Luckhurst, G. R. 370, 373, 386
Lucon, M. 366, 417, 434, 436, 437, 459
Lüder, W. 354
Ludwig, P. 370
Ludwig, R. 6
Luhmer, M. 381
Luisi, P. L. 46
Lunazzi, L. 147
Lunds, K. 129
Lundstedt, T. 84, 86, 87
Luppa, D. 369
Luque, F. J. 38
Lussier, L. S. 18, 19
Lusztyk, J. 211–213
Lutsenko, I. F. 120
Lutskii, A. E. 329, 345, 363, 364, 376, 378, 381
Lütjens, H. 321
Lüttke, W. 336
Lüttringhaus, A. 82, 252, 488
Lutz, R. E. 134
Luxon, S. G. 500
Łuczkowski, B. 437
Lyle, J. L. 378, 437
Lyon, R. K. 305

m

Ma, J. C. 20
Maas, G. 57, 58, 320, 321, 478, 488, 505
McAdoo, D. J. 305
McBain, M. E. L. 42, 45
McCabe, J. R. 312
Maccarone, E. 188, 194, 195
McCarthy, Y. 365
McClellan, A. L. 475
McConnell, H. 348
McCormac, P. B. 323
McCrindle, R. 384
McCullough, K. J. 308
McDermott, M. 268
McDonald, N. A. 127, 131
McDonald, S. R. 407, 408
McEwan, I. H. 50
McGill, R. A. 243, 333, 400, 401
Machado, C. 42, 333, 345, 428, 478
Machado, V. G. 42, 333, 428, 478
McHale, J. L. 18, 334, 356
Macias, A. 271, 272
Maciejewski, A. 57
Maciel, G. E. 376–378, 383
McIver, R. T. 18, 82, 88, 99–104, 147, 148, 157, 244, 276
Mackay, G. I. 103, 156, 158–160, 245, 246
McKelvey, D. R. 315–317
McKelvey, R. D. 416
McKenzie, E. D. 42
MacKnight, A. K. 201
Mackor, A. 370, 372
McLafferty, F. W. 272
McLauchlan, K. A. 388
McLean, A. 362
MacLean, C. 200
McMahon, T. B. 103, 162
McManus, S. P. 278, 403
McNab, H. 133
Mc Nally, M. E. P. 492
McNiven, N. L. 479
MacNulty, B. J. 163, 165, 169–172, 215, 225, 310
Macomber, R. S. 376, 378
Macquarrie, D. J. 333
McQuillin, F. J. 477
McRae, E. G. 336, 340, 342, 345, 359
McReynolds, W. O. 445
McWilliams, D. 370, 372
Madaule-Aubry, F. 49, 84, 251
Mäder, H. 134
Mäemets, V. 381, 383
Maercker, A. 136
Maeyer, L. de 499
Magaha, S. 405
Magat, M. 2, 367
Magnera, T. 147, 157, 158
Mahanty, J. H. 13
Mahendran, K. 244–247
Mahieu, V. 369
Maia, A. 238, 241, 263, 268
Maier, G. 121, 135
Mainar, A. M. 401
Maitland, G. C. 10
Maiwald, M. 71
Maizus, Z. K. 147, 210, 215, 219
Majer, Zs. 360
Maki, A. H. 370, 372
Makitra, R. G. 125, 168, 365, 366, 396, 401, 436, 444, 448, 456, 457
Makohon, O. 396, 456
Makosza, M. 266, 318
Maksimović, Z. B. 40, 42, 333, 416, 417, 425, 426, 428
Malecki, J. 120–122
Malhotra, R. 8
Malinowski, J. 505
Malinowsky, E. R. 84, 85
Malkin, V. G. 381, 383
Malkina, O. L. 381, 383
Mamantov, C. B. 59
Mamantov, G. 59
Manabe, O. 117, 118
Mancini, P. M. E. 42, 173, 174, 426

Mander, L. N. 121, 126, 128, 131–133, 288, 360, 362
Mander, S. R. 344
Mangini, A. 103
Mannucci, C. 119
Manoharan, M. 129, 130
Mantsch, H. H. 365
Manuta, D. M. 339, 429, 430
Manzur, J. 365
Mao, F. 365, 374
Maquestiau, A. 113, 114
Maran, U. 108, 166, 168
Marcandalli, B. 185, 186, 292, 300, 334
March, J. 52, 54, 106
March, N. H. 10, 57
Marchot, E. C. 138
Marcos, E. S. 38, 134
Marcus, R. A. 356
Marcus, Y. 11, 19, 22–27, 30, 32–34, 38–40, 42, 46, 53, 54, 57, 63, 64, 72, 81, 254–256, 313, 316, 390, 393, 397, 398, 401, 402, 411, 417, 425, 426, 428, 432, 433, 439, 446–448, 453, 462, 468, 471, 475, 479, 498, 500
Margolin, Z. 102
Maria, P.-C. 23, 24, 84, 87, 101, 103, 123, 397, 398, 457, 458
Marius, W. 30, 36, 37
Mark, F. 281
Mark, G. 281
Markgraf, J. H. 201
Maroldo, S. G. 272
Maroncelli, M. 334, 354
Marquez, M. 131
Marschalok, G. A. 456
Martell, A. E. 80
Martin, D. 82, 84, 86, 252, 262
Martin, G. J. 78, 79, 383
Martin, H. 93, 147
Martin, J. 379
Martin, J. C. 200, 205–208, 306, 307
Martin, J. M. L. 137, 162
Martin, M. L. 78, 79, 383
Martin, R. L. 192
Martinez, J. M. 38, 381
Martinez, R. D. 173, 174
Martinez, R. M. 401
Martini, G. 370, 372
Martinmaa, J. 82, 253
Martino, C. J. 70, 73
Martin-Villamil, R. 357, 429, 439, 440, 443, 447, 453, 461
Maryott, A. A. 50, 475
Marzin, C. 113
Maschmeier, C.-P. 62
Mashelkar, R. A. 200
Mashima, M. 18, 82, 88, 103, 244
Maskill, H. 173
Maslak, P. 201
Massa, W. 136
Masse, J. 360
Massey, A. G. 57
Masters, A. J. 131
Mataga, N. 330, 345, 359
Matern, K. 333, 478
Mathis, J. R. 148, 149, 151, 217, 218
Matschiner, H. 62
Matsuda, M. 213, 334
Matsui, K. 175
Matsui, T. 166
Matsumoto, K. 308, 309, 314, 315
Matsumoto, M. 45
Matsunami, S. 370, 372
Matsuyama, A. 121
Mattay, J. 52
Matthews, M. 137, 138
Mattox, T. F. 334
Matulenko, M. 333, 428, 448, 478
Matusch, R. 106
Matwiyoff, N. A. 36
Matyushov, D. V. 432
Maugh, T. 60, 61, 404
Maul, J. J. 505
Mauser, H. 111
Mavel, G. 376, 378
Mavrin, G. V. 188
Maw, S. A. 131
Mayer, H. 296, 297
Mayer, R. 117
Mayer, U. 23–26, 46, 49, 50, 121, 257, 264, 376–378, 383, 398, 437, 438, 453, 465, 466
Maynard, J. T. 253
Mayo, P. de 334, 429
Mayoral, J. A. 188, 401
Mayr, H. 175, 198, 247, 248 408
Mayrhofer, W. 81
Mazur, Y. 108
Mazurenko, Yu. T. 345
Mchedlov-Petrossyan, N. O. 333
Meade, M. 365
Mears, W. H. 156
Medici, A. 303
Medley, J. H. 397
Meehan, E. J. 75, 79
Meek, D. W. 21
Meerwein, H. 166
Mehlhorn, A. 331, 340, 347, 352
Mehta, A. J. 113, 114
Mei, E. 376
Meier, H. 109
Meijer, E. J. 162
Meilahn, M. K. 191
Mekelburger, H.-B. 6
Melzig, M. 113
Ménassa, P. 18, 19
Mendenhall, G. D. 93
Menger, F. M. 30, 42, 44, 46, 68, 122, 123, 129, 294, 298, 361, 400
Mennucci, B. 10, 38, 166, 168, 329, 345, 381
Menon, B. 55
Menschutkin, N. 3, 147
Mente, S. R. 334
Mentha, J. 109
Merchán, M. 355
Merin, R. 345
Mestdagh, H. 279, 280
Mészáros, S. 445
Metz, B. 266
Metz, K. R. 36, 376, 377
Metzger, J. O. 4, 506
Metzsch, F. A. von 490
Meyer, B. 488
Meyer, E. F. 14
Meyer, K. H. 4, 396
Meyer, T. J. 330, 334, 340, 345
Meyer, V. R. 492
Meyer, W. C. 181, 193
Meyers, W. W. 166, 230, 234
Mezhov, E. A. 401
Mhala, M. M. 403
Micheli, C. de 192
Michl, J. 59
Mikeš, F. 413
Mikeś, O. 493
Mikkelsen, K. V. 345
Mikolajcyk, M. 131

Mikulaschek, G. 253
Mikulecká, A. 368
Mikutta, B. 347
Milakofsky, L. 404
Milano, M. 268
Milanovic, J. 201, 202
Milart, P. 425
Milczarek, E. 433
Miles, D. W. 360
Millefiori, A. 113–115
Millefiori, S. 113–115
Miller, D. M. 115
Miller, E. G. 184, 185, 449
Miller, J. 82
Miller, J. G. 202, 205
Miller, R. 333
Miller, T. 202
Miller, W. M. 23, 397
Mills, S. G. 106–109, 111–113
Milstein, D. 137
Mimoun, H. 177
Minch, M. J. 336
Minisci, F. 370, 374
Minkin, V. I. 337, 348, 354
Minot, C. 244
Mioskowski, C. 288
Mirakyan, A. L. 468
Mirejovsky, D. 29, 30
Mirone, P. 369
Mirzoian, A. 139, 142
Mischenko, K. P. 30
Misharina, T. A. 444, 445, 492
Mishima, M. 106
Mishra, A. 331, 340, 345
Mishra, B. K. 331, 340, 345
Mislow, K. 69, 184, 185, 449, 471
Misuishi, M. 117
Mitchel, E. 16
Mitsonobu, K. 45
Mitsuhashi, T. 195, 196, 240
Mittal, K. L. 42
Mitton, C. G. 316
Miura, M. 308
Miyabo, A. 51, 52
Mizan, T. I. 70, 73
Mloston, G. 191
Modena, G. 103, 176
Moelwyn-Hughes, E. A. 93, 109, 147, 149, 156, 159, 218, 219
Moffatt, J. R. 403
Mohamed, A. A. 162
Mohammad, M. 137, 207, 209, 210, 334, 338, 339, 411–415
Mohr, G. J. 333
Möhrle, H. 119
Moldenhauer, F. 330
Molotsky, T. 36
Molter, K. E. 51, 126, 138
Montanari, F. 238, 241, 268
Monteiro, P. M. 292
Montero-Cabrera, L. A. 334
Moore, J. W. 147, 149, 218, 219, 221, 236
Morao, I. 183
More O'Ferrall, R. A. 99, 316
Moreno-Mañas, M. 505
Moretti, I. 69, 360
Morgan, K. J. 365
Morgenstern, J. 117
Moriconi, E. J. 181, 193
Morihashi, K. 373
Morill, T. C. 384
Morine, G. H. 334
Morisue, T. 45
Morita, H. 253
Morita, Y. 296
Moriyasu, M. 106–109, 112, 113
Morley, J. O. 117, 118, 336, 345
Morley, R. M. 336, 345
Mormann, W. 337
Moroi, Y. 42, 45, 46
Morokuma, K. 161
Moroni, A. F. 152, 200, 201
Morris, D. G. 366
Morrison, J. D. 69, 471, 477
Morrow, R. M. 37
Morten, D. H. 274, 275, 406
Morton, T. H. 147, 305
Mosely, R. B. 408
Mosher, H. S. 69, 471
Mosier-Boss, P. A. 267
Moskovits, M. 488
Moura-Ramos, J. J. 129, 131, 337, 354
Mousseron-Canet, M. 117
Movius, W. G. 36
Moylan, C. R. 99
Mu, L. 69
Mu, X. 475
Muanda, M. wa 42
Mueller, R. H. 291
Mueller, W. A. 188, 190, 191, 280, 291, 295, 404, 410, 446
Mukai, K. 370, 372
Mukerjee, P. 351, 429, 430
Mukhopadhyay, T. 82, 252, 253
Mukhtar, R. 179, 180, 193
Mulindabyuma, J. M. 334, 339
Müller, K. 281
Müller, K.-H. 188, 410
Müller, P. 53, 168, 259, 264, 389
Müller, R. 392–394, 428
Müller-Dethlefs, K. 10
Müller-Plathe, F. 162
Mulliken, R. S. 19–21
Mullin, A. 259
Mulski, M. J. 139
Mulzer, J. 182
Munk, M. E. 191
Munoz, A. 119
Munro, M. H. G. 287
Munson, M. S. B. 100
Murata, F. 330, 340, 345
Murcko, M. 127, 131
Murphy, D. 370, 372
Murphy, W. S. 276
Murray, C. J. 330
Murray, R. W. 307, 308
Murrell, J. N. 5, 336, 344, 378
Murthy, A. S. N. 382, 384
Murto, J. 239, 241, 242
Murto, M.-L. 254
Murty, T. S. S. R. 16
Muruyan, R. 72, 73
Mussini, P. R. 498
Mussini, T. 498
Mußmann, L. 322, 323
Mustamir 106
Mustroph, H. 117, 334, 350
Muszkat, K. A. 123, 124
Myers, A. B. 345
Myers, R. Th. 80

n

Naberuchin, J. I. 6
Nad, S. 354
Nagai, M. 51, 52
Nagai, T. 362, 410, 411
Nagakura, K. 334
Nägeli, C. W. von 43
Nagy, J. B. 42, 334, 339

Nagy, O. B. 42, 188, 334, 339
Nagy, P. 115, 417, 425, 426
Nair, P. M. 117
Nakagawa, K. 281
Nakagawa, N. J. 385, 386
Nakahara, M. 180, 312
Nakajima, A. 358
Nakamura, K. 372–375
Nakamura, M. 378
Nakamura, Y. 296
Nakano, T. 59, 301, 302
Nakaten, H. 201
Nakazaki, M. 303
Nam, J. 453, 467
Nanni, E. J. 254
Napier, J. J. 370, 372, 374, 414, 415, 437
Nascimento, M. G. 345
Nasehzadeh, A. 166, 408, 458
Nash, L. K. 63
Nasielski, J. 404, 409, 410
Natterstadt, J. J. 378, 383
Naus, P. 103
Nauta, W. T. 200
Nebenzahl, L. L. 77, 99, 101
Neddham, T. E. 27
Nedeltcheva, D. 115
Neidhart, B. 71, 424
Neilands, O. 21, 334
Neitzel, S. 417, 425, 438
Nelson, A. 266, 319
Nelson, G. L. 378
Nelson, K. L. 431
Nelson, W. M. 4
Nemethy, G. 27
Nerbonne, J. M. 301, 302
Nernst, W. 2, 47
Nesakumar, J. E. 62, 189, 268
Nesi, R. 119
Neuenschwander, M. 178
Neufeld, F. R. 222, 443, 457
Neuman, R. C. 203, 220, 309
Neumann, M. A. 45
Neumann, R. 266, 318
Nevěčná, T. 166, 417, 425
Newitt, D. M. 311
Newman, K. E. 42
Nibbering, N. M. M. 162, 172
Nicholls, C. H. 117
Nichols, D. A. 72, 73
Nickel, B. 111
Niclas, H.-J. 82, 252
Nicol, M. F. 345
Nicolet, P. 417, 432–435, 456, 457, 459
Nicoud, J. F. 303
Nielsen, A. T. 489
Nielson, J. B. 15, 17
Nigam, S. 330
Niki, E. 210
Nikki, K. 385, 386
Nikolajewski, H.-E. 331, 341–343, 378
Nilles, G. P. 471
Nilsson, M. 106
Ninham, B. W. 13
Niqam, S. 42
Nishiguchi, H. 370, 372
Nishimura, N. 124, 125, 185, 186, 292, 300, 334, 374
Nishiwaki, T. 281
Nitsch, S. 478
Nitzan, A. 36
Nodelman, N. 307
Noe, L. J. 348
Nojima, M. 308
Nolte, K.-D. 335, 341–343, 345, 347, 378, 429
Nolte, R. J. M. 333
Nome, F. 292
Norcross, B. E. 214
Nordahl, A. 84, 86, 87
Nordén, B. 360
Normant, H. 49, 82, 253, 262
Normant, J. F. 253
Norris, J. F. 166
Norris, R. K. 118
North, A. M. 304
Notario, R. 27, 101, 103, 166, 168, 230, 390, 393, 411, 432, 433, 448, 456, 457
Nöth, H. 253, 288
Novak, E. R. 477
Novak, T. J. 334
Novaki, L. P. 42, 297
Novikova, S. S. 401
Nowak, M. J. 113, 114
Nowak, W. 357
Noy, R. S. 106
Noyes, R. M. 304
Noyori, R. 70, 73, 326, 505
Nozaki, H. 70
Nozaki, K. 201
Nudelman, A. 166
Nudelman, N. S. 173, 174
Nürrenbach, A. 331
Nuti, L. 370, 372
Nuyken, O. 200, 201

o

Oae, S. 169, 253
Oakenfull, D. 293
Oakes, J. 370, 372
Oancea, A.-M. 499
Ocampo, R. 183
Oda, R. 331
O'Donnell, T. A. 77, 81
Oehler, U. M. 370
Offermann, W. 212, 213
Ogasa, M. 166
Ogata, T. 23, 398
Ogawa, M. 281
Ogawa, S. 117
Oglashennyi, Yu. I. 456
Ogo, Y. 312, 417, 425, 428
Ohcho, M. 124, 125
Ohmine, I. 6
Ohnishi, M. 166, 312
Ohno, A. 60, 61, 404
Ohta, N. 210
Ohta, S. 361
Ohtaki, H. 2, 25, 26, 30, 36, 63, 72
Oi, N. 364, 365, 367
Okada, T. 185, 186, 300
Okamoto, K. 51, 52, 54, 238, 240
Okazaki, R. 135
Okuno, Y. 276
Olah, G. A. 50, 54, 77, 81, 275
Olbrich-Deusner, B. 340, 429, 430
Olech, A. 425
Olin, G. R. 303
Oliva, A. 188
Olivier, H. 322, 323
Olivier-Bourbigou, H. 506
Ollerenshaw, J. E. 478
Olmstead, W. N. 18, 82, 88, 102, 103, 156–159, 244–246
Olofson, G. 24
Olofson, I. 24, 397, 398
Olsen, F. P. 99
Olsen, H. 192, 203
Olszewski, P. K. 425
Oncescu, T. 499
O'Neil, M. J. 475, 505
Ong, E. H. 478
Onsager, L. 342, 345, 367, 380
Ookawa, A. 287, 290

Oppermann, M. 334
Orbach, N. 355, 417, 425
Orfanopoulos, M. 193
Orgel, L. E. 370, 373
Orioli, P. 289
Orozco, M. 38
Ortega, J. 42, 331
Ortega, J. P. 180
Orth, P. 200, 201, 222, 224, 317
Orville-Thomas, W. J. 10, 21, 120–122, 126, 128
O'Shea, K. E. 325
Oshika, Y. 345
Oshima, T. 362, 410, 411
Osiecki, J. H. 370
Osipov, M. A. 362
Osipov, O. A. 126, 337, 348, 354
Osterby, B. R. 416
Osteryoung, R. A. 61
Ostheimer, M. 6
Ostrovskii, M. K. 129
Ostrowicki, A. 264
Ostrowski, P. J. 505
Ostwald, W. 2
Osugi, J. 180, 312
Oteyza de Guerrero, M. 113
Otruba, J. P. 185, 186, 300
Ottaviani, M. F. 370, 372
Otto, P. 180, 181, 193, 280
Otto, S. 294
Ottolenghi, M. 355, 417, 425
Ouellette, R. J. 222, 312
Ousset, J. B. 288
Ouvrard, C. 19
Ovchinnikova, Yu. I. 286
Overbeck, H. 9, 63, 500
Overberger, C. G. 203
Owen, B. B. 496
Owenius, R. 370, 372
Oyama, K. 428
Ozin, G. A. 488

p

Pacifici, J. G. 201, 202
Packard, M. E. 381, 382
Paczkowski, M. A. 292, 296
Padias, A. B. 180
Padova, J. I. 38, 40
Padwa, A. 188, 191
Page, A. D. 113, 114
Page, J. E. 479
Pagni, R. M. 59
Paik, C. 292
Painelli, A. 335
Paju, A. I. 436, 453
Pakkanen, T. A. 166, 168
Pal, H. 354
Paley, M. S. 333
Palit, S. K. 271, 272
Palit, S. R. 243
Palm, N. 91, 452
Palm, V. A. 91, 147, 231, 390, 393, 403, 411, 436, 452–455
Palombini, L. 67, 389
Pan, H.-L. 254
Pangborn, A. B. 478
Pannel, J. 370
Panse, P. 194
Paoloni, L. 398, 438, 466
Papadopoulos, M. 188, 194
Papazyan, A. 38
Pappalardo, R. R. 38, 134
Pappiaonnou, C. G. 201, 202
Paquette, L. A. 61, 505
Paracelsus, Th. 1
Paradisi, C. 103
Parchment, O. G. 113–115
Pardo, L. 188
Paredes, R. 183
Park, H. 46, 298
Parker, A. J. 14, 18, 82–84, 86, 147, 156, 158, 159, 217, 238, 244, 246, 249–251, 253–255, 257–259, 287, 398, 437, 438, 465
Parker, D. 471
Parks, C. R. 307
Parthiban, S. 162
Partington, J. R. 2, 37, 194
Parusel, A. B. J. 355, 357
Paschalis, P. 259
Paschke, T. 492
Patai, S. 370
Patat, F. 325
Patel, K. B. 365
Patterson, L. K. 351
Patterson, R. B. 303
Patz, M. 247
Paul, K. G. 243
Paulaitis, M. E. 29, 30, 315
Paulson, A. 139
Paulson, J. F. 159–161
Pauschmann, H. 492
Pavličkova, L. 334
Pavlish, N. V. 168
Pawelka, Z. 121, 334
Payzant, J. D. 156, 158, 245, 246
Pazos, J. F. 201
Peacock, J. 229
Pearson, B. D. 478
Pearson, J. M. 305
Pearson, R. G. 79, 80, 147, 149, 167, 215, 218, 219, 221, 236, 242, 244, 245
Peck, D. G. 113, 114
Peck, D. R. 197
Pedersen, C. J. 266
Pedersen, J. A. 370, 372
Pedersen, K. J. 392
Pedulli, G. F. 370, 374
Peguy, A. 36
Pehk, T. 341–343, 378
Pellerite, M. J. 101, 102, 147, 156–159
Pelletier, W. M. 271
Pellicciari-Di Liddo, L. 185, 186, 292, 300, 334
Pena-Nuñez, A. S. 351, 372, 437
Penski, C. A. 453, 466, 467
Pentin, Yu. A. 129
Peregudov, A. S. 120
Pereira, S. M. 505
Peretz, P. 123, 124
Pérez del, C. 42
Perez, P. 357
Pérez-Ossorio, R. 116, 128, 289, 290
Pérez-Rubalcaba, A. 116, 289, 290
Perjéssy, A. 365, 453, 467
Perraud, R. 268
Perrin, C. L. 15, 17, 131
Perrin, D. D. 99, 478
Perrini, G. 188, 194, 195
Persico, M. 38, 329, 345, 381
Person, W. B. 19, 21
Persson, I. 24, 25, 35, 81, 390, 393, 398, 402
Persson, P. 390, 393, 398, 402
Peslherbe, G. H. 54
Pessin, J. 259
Pestemer, M. 479
Peters, D. W. 478
Petersen, R. C. 201
Peterson, M. J. 194, 195
Peterson, P. E. 278, 405
Petkov, I. 113, 114
Petrosyan, V. S. 175, 263, 276, 278
Petrov, E. S. 253

Petrov, S. I. 73, 75, 79, 496, 499
Petzke, F. 336, 354
Pevzner, I. D. 496
Pews, R. G. 404, 415
Pfeffer, P. E. 251
Pfeiffer, M. 334
Pfeil, E. 188, 222
Pfister-Guillouzo, G. 113–115
Pflug, J. L. 323
Pham Van Huong 365
Pham, N. 388
Phelps, C. L. 70, 73, 492
Phuc, N. V. 297
Pickering, I. A. 458
Pickles, N. J. T. 166
Pienta, N. J. 24
Pierre, J.-L. 243, 268
Pihl, V. O. 101
Pileni, M. P. 46
Pimentel, G. C. 8, 9, 348, 369
Pincock, R. E. 185, 207, 286
Pines, A. 12
Pines, H. 260
Pinto, P. B. 107
Pirana, M. 40
Pirig, Ya. N. 125, 168, 365, 366, 396, 401, 436, 444, 448, 456, 457
Pirilä, P. 42
Pirkle, W. H. 69, 386, 471, 477
Piskala, A. 195
Pizey, S. S. 82, 252, 262
Platt, L. R. 29, 30
Platz, M. S. 199
Plesch, P. H. 229
Pleva, S. G. 334
Plieninger, H. 69, 477
Plieninger, P. 297, 333, 378, 429
Pliva, J. 364, 365, 367
Plumet, J. 116, 128
Pocker, Y. 50, 165, 428
Podsiadla, M. 429
Poh, B.-L. 122
Pohl, H. 192
Pohl, L. 58
Pohorille, A. 30
Polansky, O. E. 281
Polanyi, M. 308
Poliakoff, M. 326
Poll, T. 291
Pollack, S. K. 437
Poltoratskii, G. M. 30
Pommer, H. 331
Ponomareva, E. A. 165
Poole, C. F. 60, 445, 494, 496
Poole, S. K. 445, 494, 496
Popov, A. I. 23, 36, 40, 41, 95, 96, 121, 243, 267, 270, 376, 397, 398
Popovych, O. 30, 33, 57, 73, 147, 238, 254, 255
Popper, K. 469
Porter, G. 350
Potter, A. 212
Pourcelot, G. 290
Powell, A. L. 398, 408, 453, 462–465
Powell, J. R. 42, 426
Powell, W. R. 479
Powling, J. 111
Pracejus, H. 69, 471
Prahl, H. 260
Prajsnar, B. 276, 277
Prakash, G. K. S. 275
Pranata, J. 115
Prat, D. 290
Pratt, L. R. 27, 30
Prausnitz, J. M. 5, 9, 23, 63, 65, 220, 221, 397, 443
Pregosin, P. S. 36
Prentiss, S. W. 166
Pretsch, E. 479, 485, 487
Prévost, C. 383
Prezhdo, V. V. 329, 345, 363, 364, 369, 376, 378, 381
Price, A. 50
Price, C. C. 260
Price, E. 46, 50, 376, 377, 437
Price, J. E. 236
Priller, U. R. 376
Prince, L. M. 298
Prinzbach, H. 337
Pritzkow, W. 177, 188, 410
Proskow, S. 180, 193
Pross, A. 147, 162, 164, 166, 168, 215
Pross, E. 428
Puar, M. S. 244
Pudjaatmaka, A. H. 77
Puechberty, A. 279, 280
Pugh, D. C. 390, 393, 412–414, 447
Pullin, A. D. E. 364, 368
Pungor, E. 500
Purcell, K. F. 22, 84, 88, 89, 415
Pusey, D. F. G. 494
Pyron, R. S. 188
Pytela, O. 84, 87, 390, 453
Pyter, R. A. 351, 429, 430

q

Quadbeck-Seeger, H.-J. 205, 207, 285, 286
Quast, H. 196, 505
Quemeneur, F. 166
Queneau, Y. 294, 295, 505
Quici, S. 321
Quina, F. H. 292
Quinlan, J. E. 234, 235
Quiroga, M. L. 289, 290
Qureshi, M. L. 197
Qureshi, P. M. 96
Qureshi, R. 197

r

Rábai, J. 319–321
Raban, M. 69, 471
Raber, D. J. 166, 169, 403
Rabinowitsch, E. 304
Rabinowitz, M. 12
Rablen, P. R. 127, 128, 131
Raczyńska, E. D. 101, 117
Radeglia, R. 331, 335, 341–343, 345, 378, 379, 417, 425, 429
Radhakrishnan, T. P. 336, 354
Radnai, T. 72
Radom, L. 162
Raevsky, O. A. 121, 139, 140
Ráfols, C. 42, 331
Rahimi, P. M. 202, 203, 286
Rahman, M. A. 354
Răileanu, M. 57
Raistrick, B. 311
Raja, R. 4, 506
Rajagopal, S. 411, 429, 432, 453
Rajamohanan, P. R. 56
Rajzmann, M. 84, 88, 89
Rakshit, A. B. 103, 159, 162
Rakshit, A. K. 103
Raley, J. H. 155
Ramachandran, C. 351, 429, 430
Ramachandran, V. 308
Ramakrishna, T. V. 40

Ramamurthy, V. 292, 296, 297, 299, 302, 307, 358
Ramaswamy, S. 195
Rambeck, W. 253
Ramesh, V. 59, 301
Ramirez, F. 194
Rampoldi, A. 241
Ramsay, B. G. 337
Ramsay, N. F. 382
Raner, K. D. 212
Rannala, E. 455
Ranneva, Yu. I. 253, 260
Ransdell, R. A. 138
Rao, A. V. R. 117
Rao, B. G. 6
Rao, C. N. R. 329, 345, 358, 363, 364, 367, 376, 378, 382, 384
Rao, G. V. 378
Rao, K. S. 337, 354
Raoult, F.-M. 2
Rapp, M. W. 404
Rappoport, Z. 106, 109, 370
Rashin, A. R. 96
Rasmussen, J. K. 181, 193
Rassat, A. 360, 370
Rasteikiene, L. 177
Rastogi, P. P. 365
Ratajczak, H. 10, 21, 120–122
Ratajczak, J. F. 265
Ratcliffe, J. 479
Rathore, R. 20
Rau, H. 69
Rauf, M. A. 428
Raulins, N. R. 195
Ray, A. 29, 413
Ray, A. K. 188, 189
Rayner, L. 370
Razzaq, M. 137
Rebbert, R. E. 305
Rebek, J. 142, 143
Rebollo, H. 284
Reczek, J. 243
Reddoch, A. H. 370, 372, 373, 437
Reddy, A. C. 166, 168
Redondo, M. I. 435
Ree, T. 5
Ree, T. S. 5
Reed, A. E. 19, 21
Reed, C. A. 331
Reetz, M. T. 56, 199
Reeves, L. W. 376, 388
Rehan, A. H. 195
Reich, H. J. 55
Reich, R. 166
Reichardt, C. 40, 42, 69, 71, 85, 87, 136, 147, 155, 173, 199, 215, 330, 332–334, 337, 357, 389, 390, 392–394, 411, 416–418, 424–426, 428, 429, 446, 448–454, 475, 478, 491
Reichert, J. 333, 478
Reid, C. E. 331, 335, 340
Reimers, J. R. 330, 334, 340, 345
Reimlinger, H. 260
Reinheimer, J. D. 166, 230, 234
Reinhold, Y. 188, 410
Reinhoudt, D. N. 266, 268
Reisse, J. 67, 129, 131, 133, 337, 354, 389
Reissig, H.-U. 191
Rembaum, A. 155
Remerle, K. 40
Remizov, A. B. 127, 131
Renneboog, R. 103
Renolen, R. 177
Resch, G. 24, 25
Rettig, W. 352, 355
Reuben, J. 383
Reutov, O. A. 99, 101, 263, 269, 271, 272, 276–278
Reychler, A. 43
Reynolds, L. 354
Reynolds, M. B. 48
Reynolds, W. F. 376
Rezende, M. C. 345, 427–429
Rhoads, S. J. 195, 271
Rice, S. A. 304
Richard, J. P. 275
Richards, R. E. 382
Richardson, M. 222
Richardson, W. H. 185
Richey, H. G. 253, 505
Richter, T. 71, 73
Richter-Egger, D. L. 335
Rickborn, B. 147, 251, 253, 260
Rickert, H. 478
Rico, I. 404
Rico, M. 379
Riddick, J. A. 50, 57, 471, 475, 477, 478
Rideout, D. C. 189, 295
Ridgeway, R. W. 477
Rieger, A. L. 365, 374
Rieger, P. H. 365, 374
Riemer, M. 397
Riess, W. 107, 108
Rigby, M. 10
Righetti, P. P. 178, 188, 197, 198
Riihimäki, V. 500
Rimmelin, J. 188
Rinaldi, D. 134, 477
Rinck, G. 312, 313
Rios, A. 293
Rios, R. 108
Ritchie, C. D. 30, 36, 38, 41, 54, 73, 75, 77, 79, 84, 244, 247, 248, 437
Rittenhouse, J. R. 202
Rittner, R. 129, 131
Riva, S. 143
Rivail, J.-L. 134
Riveros, J. M. 103, 104, 147, 158, 159, 276
Rivetti, F. 176
Rizvi, H. 492
Roa, G. 182, 194
Roach, E. T. 36, 376
Roberts, G. 323
Roberts, J. D. 136, 378
Roberts, J. L. 254, 496, 500
Roberts, K. 406–408, 459
Roberts, R. M. G. 188, 276, 277
Robertson, R. E. 238, 242, 308, 314, 315
Robins, M. J. 360
Robins, R. K. 360
Robinson, G. C. 53, 54
Robinson, G. W. 6
Robinson, M. J. T. 129
Robinson, R. A. 499
Rodebush, W. H. 15
Rodehüser, L. 40
Rodewald, R. F. 244–247
Rodger, A. 360, 362
Rodger, P. M. 362
Rodriguez, I. 435
Rodriguez, J. G. 357, 429, 439, 440, 443, 447, 453, 461
Rodriguez, M. 386
Roe, D. K. 60, 61, 404
Rogasch, P. E. 365, 367
Rogers, B. D. 197
Rogers, M. T. 107, 112
Rohatgi-Mukherjee, K. K. 123, 124
Rohrschneider, L. 444, 445
Rolla, F. 238, 241
Röllig, M. 177, 188, 410

Rolston, J. H. 278, 279
Roman, M. N. 261
Romanowski, S. J. 39
Romm, I. P. 19
Romm, R. 166
Romsted, L. S. 292
Ronayne, J. 376, 378, 381, 382, 384
Rondinini, S. 498
Rooney, D. W. 60–62, 500, 506
Roos, B. O. 355
Roper, G. C. 499
Rosalky, J. M. 266
Rosen, R. K. 478
Rosenbaum, E. E. 82, 252
Rosenfarb, J. 82, 252, 475
Rosenfeld, S. M. 109
Rosenthal, I. 123, 124
Rosenthal, O. 336
Rosés, M. 42, 331, 426, 427
Rosotti, H. 123
Rospenk, M. 17, 122
Rosquete, G. A. 334, 343
Ross, M. D. 42
Ross, P. 162
Ross, S. D. 201, 404
Rossky, P. J. 334, 340
Rößler, G. 4
Roth, F. 269
Roth, L. 502, 503
Roth, T. 429, 430
Rotkiewicz, K. 354
Rottman, C. 333
Roussel, R. 113
Rowley, A. G. 489
Rowlinson, J. S. 5
Rozas, I. 15
Rozeboom, M. D. 103, 159, 162
Rozhkov, I. M. 261
Ruasse, M.-F. 176, 278
Rubio, E. 144
Rüchardt, C. 194, 202–205, 207, 285, 286, 306, 403, 477
Rücker, C. 188
Rüdiger, V. 121, 139, 140
Rudkevich, D. M. 142, 143
Rudorf, W.-D. 475
Ruf, H. 166
Ruiz-López, M. F. 134
Rule, H. G. 362
Rummens, F. H. A. 380
Ruoff, R. S. 8
Rupp, G. 503
Rurländer, R. 288
Rush, D. J. 128, 131
Russel, A. J. 143
Russel, G. A. 210, 211, 370
Russel, P. B. 109
Russell, J. C. 46
Russo, T. J. 265
Rust, F. F. 155
Rutan, S. C. 42, 330, 494
Rutherford, R. J. D. 453
Rüttinger, H. H. 62
Ryabokobylko, Yu. S. 276, 277
Rychtman, A. C. 121, 123
Rylander, P. N. 489
Rzepa, H. S. 124
Rzeszotarska, J. 429

s

Sabashita, K. 129
Sabatino, A. 398, 438, 466
Sackmann, E. 58
Sadek, H. 54
Sadek, P. C. 492
Sadykov, A. S. 119
Saenger, W. 15, 16
Saeva, F. D. 303
Šafařik, L. 73, 496
Safarzadeh-Amiri, A. 334, 429
Saffar, S. G. 188
Safi, B. 162
Sahyun, M. R. V. 260, 261
Saidov, G. V. 337, 348, 354
Saint-Gilles, L. P. de 2
Saito, S. 6
Saiz, J. L. 398
Sajus 404, 409
Sakai, M. 45
Sakano, T. K. 50, 57, 471, 475, 477, 478
Sakurai, H. 200, 207, 208, 281, 282
Saleh, N. 252
Salman, S. R. 388
Salomon, M. 32, 496
Saluja, P. P. S. 306, 313, 314
Salvatella, L. 188, 401
Sam, D. J. 268
Samanta, A. 62, 336, 354, 357
Samat, A. 337
Samii, A. A. 404
Samoilov, O. Ya. 30, 36, 37
Samori, B. 303
Samoshin, V. V. 126, 128, 131, 132
Sana, M. 135
Sanakararaman, S. 189, 268
Sanchez, A. M. 68
Sandell, E. B. 75, 79
Sandorfy, C. 18, 19, 345, 479
Sandstede, G. 62
Sandström, J. 134
Sandström, M. 24, 25, 81, 390, 393, 398, 402
Sangwan, N. K. 295, 401
Sankar, G. 4, 506
Sankararaman, S. 62
Santamaria, J. 142, 143
Santesmases, M. J. 116
Sapiro, R. H. 311
Sapunov, V. N. 24, 147, 218, 428, 438, 439, 447, 466
Sargent, F. P. 370
Sargeson, A. M. 462
Sarpola, S. 139, 140
Sarthou, P. 271
Sasaki, I. 144
Sasaki, M. 180
Sasaki, Y. 11, 66
Sasson, Y. 266, 318
Sauer, E. 188, 334, 410
Sauer, J. 180, 188, 189, 191, 260, 268, 281, 291, 296, 297, 404
Saunders, W. H. 9, 170, 254
Saupe, A. 58
Sauvage, J.-P. 266, 506
Sauvageau, P. 479
Savage, G. P. 505
Savage, P. E. 70–73, 328, 506
Savéant, J.-M. 164
Savedoff, L. G. 48, 244, 245, 263, 264
Savelli, G. 292
Savelova, V. A. 227, 229
Savignac, A. de 404
Savvina, L. P. 333
Sawada, H. 286
Sawyer, D. T. 254, 496, 500
Scaiano, J. C. 211
Scanlan, M. J. 113–115
Scarpati, R. 284
Scatchard, G. 220, 221, 236
Schaad, L. J. 15, 16, 364
Schaal, R. 499
Schaap, W. B. 498, 499
Schädler, H. D. 60, 62, 397
Schadt, F. L. 405–407, 409
Schaefer, A. D. 265

Schaefer, H. F. 202
Schaefer, T. 378, 380, 388
Schäfer, G. 417
Schäfer, L. 115
Schäfer, V. 337
Schaffernicht, W. 82
Schank, K. 107, 108
Schanze, K. S. 334
Schaub, B. 195
Schaumann, E. 193
Scheer, W. 134
Scheffer, J. 299
Scheffler, K. 119, 370
Scheffold, R. 116
Scheibe, G. 4, 29, 30, 115, 116, 341, 343, 345, 378
Scheibe, P. 335, 341
Scheiner, S. 10, 15, 17
Schenck, G. O. 281, 282
Scheraga, H. A. 27
Scherhaufer, A. 22, 24, 27, 49, 252, 396–398, 462, 465
Scheurle, B. 59
Schiemenz, G. P. 262
Schiess, P. 195
Schiess, P. W. 185, 284
Schiewe, J. 62
Schiffrin, D. J. 52
Schildknecht, H. 478
Schilf, W. 117
Schilling, G. 505
Schilling, S. L. 272
Schläfer, H. L. 82
Schlag, J. 331
Schlegel, H. B. 158, 164
Schleitzer, G. 254
Schleyer, P. von R. 16, 50, 54, 166, 187, 274, 275, 365, 369, 403, 405–407, 409, 414, 415, 435
Schlosser, H.-J. 337
Schlosser, M. 54, 195, 286
Schmehl, R. H. 46
Schmeltz, I. 251
Schmid, G. H. 412
Schmid, H. 196
Schmid, R. 10, 23–25, 29, 30, 147, 219, 257, 309, 397, 398, 428, 430, 432, 437–439, 447, 456, 466
Schmidt, A. P. 166, 167, 227, 228, 313
Schmidt, F. C. 498, 499
Schmidt, P. P. 18
Schmidt, R. 192
Schmidt, R. E. 136
Schmidt, R. L. 388
Schmidt, W. 352, 353
Schmidtchen, F. P. 243
Schmierer, R. 290, 291
Schmittel, M. 203
Schmuttenmaer, C. A. 333
Schneider, G. M. 71
Schneider, H. 38–40, 139, 140, 267
Schneider, H.-J. 121, 139, 140, 295, 378, 401
Schneider, R. L. 471
Schneider, S. 113, 335, 341
Schneider, W. G. 376, 378, 380
Schnurpfeil, D. 188, 410
Schob, F. 335, 345, 429
Schoening, R. C. 376
Schöffski, K. 478
Scholz, E. 478
Scholz, R. 334
Schomäcker, R. 298
Schowen, R. L. 315–317
Schrader, B. 364, 367
Schrader, W. 60
Schreiber, M. 334
Schreiner, P. R. 187
Schriesheim, A. 253
Schriver, G. W. 29, 30, 272
Schröder, A. 334
Schroth, W. 60, 62, 397
Schuetz, R. D. 471
Schug, R. 179, 180, 193
Schulman, J. H. 298
Schulman, S. G. 352, 353
Schultz, E. 46
Schultz, P. G. 204, 351
Schultz, P. W. 270
Schultze, O. W. 3
Schulz, A. 203, 340, 429, 430
Schulze, U. 69
Schurhammer, R. 33
Schurz, J. 479
Schuster, F. 191, 296, 297
Schuster, G. B. 52
Schuster, P. 15, 17, 30, 36, 37
Schwager, I. 208, 209
Schwarz, H. 478, 500
Schwarzenbach, G. 108
Schwarzer, H. 205, 207
Schweig, A. 337, 343, 428
Schwendener, S. 43
Schwetlick, K. 147, 317, 402, 490
Sclove, D. B. 201
Scorrano, G. 40, 78, 79, 99, 103
Scott, C. B. 244
Scott, J. A. 102, 103
Scott, J. M. W. 308, 314
Scott, J. W. M. 238, 242
Scott, R. L. 5, 9, 63, 65, 219–221, 443, 471
Scotton, M. 119
Sears, P. G. 84, 252
Sebastián, R. M. 505
Seddon, K. R. 4, 15, 17, 60–62, 322–324, 336, 500, 506
Šedivec, V. 471
Seebach, D. 55, 69, 70, 82, 252, 253, 289, 477, 505
Seebacher, M. 335
Seefelder, M. 196
Seehra, J. K. 365
Seeliger, A. 477
Seib, A. 200, 201, 222, 224, 317
Seidel, M. 192
Seidl, H. 191
Seiffert, W. 341, 343, 378
Selim, A. 290, 291
Selmi, M. 127, 131
Seltzer, R. 270
Seltzer, S. 188
Sen, J. N. 306
Sen, P. 31
Sendhoff, N. 6
Senthilnathan, V. P. 329, 345, 358, 363, 364, 376, 378
Senyushkina, T. A. 334
Seppelt, K. 56, 251, 254
Sera, A. 308, 309, 314, 315
Serjeant, E. P. 99
Serrano-Andrés, L. 355
Seta, P. 45
Sethi, A. 323
Severance, D. L. 188, 189, 197, 201
Seybold, P. G. 123, 124
Seyden-Penne, J. 171
Shabana, R. 87
Shah, D. O. 42, 298
Shah, S. S. 336
Shaik, S. S. 158, 164, 166, 168
Shamsipur, M. 267
Shao, L. 113–115
Sharaf, M. A. 84
Sharagin, I. A. 188
Sharma, A. K. 61, 352, 353

Sharp, J. H. 255, 370, 374, 375
Sharpe, P. E. 303
Sharts, C. M. 251
Shashidhar, M. A. 337, 354
Shatenshtein, A. I. 253, 260, 263
Shaw, J. E. 251
Shaw, R. W. 327, 328, 333, 416
Sheban, G. V. 117, 118
Sheikhet, I. I. 219
Sheinker, V. N. 126
Shen, H.-C. 408
Shenderovich, I. G. 17
Shepherd, R. E. 339
Sheppard, S. E. 4
Sheppard, W. A. 251
Sherif, A. K. 336
Sherma, J. 494, 496
Sherry, J. J. 251
Sheu, W.-S. 340
Shevelev, S. A. 269, 272
Shibuya, T. 345
Shigematsu, Y. 290
Shih, P.-S. 351
Shilov, I. Yu. 18
Shin, D.-M. 185, 186, 300
Shine, H. J. 159, 201
Shiner, V. J. 316, 404
Shinohara, K. 117
Shipps, G. 388
Shirai, M. 243
Shiraishi, S. 117
Shmidt, V. S. 401
Shorter, J. 84, 85, 390, 392, 394, 436, 453–455, 458
Shpinel', Ya. I. 118
Shtyrlin, Yu. G. 125, 168
Shum, Y.-Y. 345
Sicher, J. 286
Sicinska, W. 378
Sicking, W. 188
Siefker, J. R. 499
Siegfried, B. 168, 259, 264
Siepmann, T. 332, 333, 416, 417, 449–451
Sievers, S. 69
Sigemoto, H. 385
Signal, J. 341–344, 347, 378
Silbert, L. S. 251
Sild, S. 90, 451, 453
Silver, B. L. 370–372, 374
Silver, P. A. 187
Silversmith, E. F. 333, 416
Silverstein, T. P. 27, 30
Simánek, V. 378
Simchen, G. 114
Simkin, B. Ya. 219
Simmons, C. J. 348
Simmons, H. E. 180, 193, 268
Simon, J. D. 147–149, 151, 217, 218
Simon, L. C. 322, 323
Simonetta, M. 46
Simpson, G. W. 505
Sims, J. 191
Sinanoğlu, O. 345–347, 358
Sinclair, J. R. 499
Singh, R. 250
Singh, S. 6, 329, 345, 358, 363, 364, 376, 378
Singh, T. P. 122, 123
Singh, U. C. 6
Sinou, D. 506
Siow, H.-L. 122
Siskin, M. 72, 73
Sisler, H. H. 57
Siverns, T. M. 128
Sjöström, M. 85, 390, 394, 395, 434, 459, 460
Skvortsov, S. A. 174, 175
Skwierczynski, R. D. 42, 426, 427
Slates, R. V. 55
Sloan, J. P. 202
Smart, N. G. 70, 73, 492
Smets, G. 192, 337, 448
Smid, J. 46, 47, 52, 54, 55, 243
Smiley, R. A. 269
Smirnov, S. N. 17
Smit, W. A. 177
Smith, A. L. 468
Smith, D. W. 32
Smith, E. B. 10
Smith, E. R. 50, 475
Smith, G. P. 59
Smith, K. 268
Smith, L. W. 297
Smith, M. B. 52, 54, 106
Smith, P. 138
Smith, R. D. 253
Smith, R. G. 251
Smith, R. S. 185
Smith, S. 48, 62, 244, 245, 263, 264, 490
Smith, S. F. 158
Smith, S. G. 113–115, 197, 238, 239, 244–246, 271, 402, 404, 408, 409
Smith, S. L. 376, 378, 387, 388
Smithers, R. H. 478
Smolova, N. T. 499
Snatzke, G. 360
Snieckus, V. 55
Snyder, J. P. 128, 192, 203
Snyder, L. R. 65, 445, 471, 490, 494, 495
Snyder, W. H. 260
Soai, K. 287, 290
Sobczyk, L. 122, 334
Sobel, H. 244, 245
Sokolov, V. I. 276, 277
Solá, M. 166, 168, 230
Solladié, G. 288
Solov'ev, V. P. 121, 139, 140
Sommer, J. 77, 81
Somolinos, C. 435
Sone, K. 330, 340, 345
Songstad, J. 244, 245
Soong, L. T. 287
Soper, F. G. 222
Souček, M. 334
Soujana, T. 357
Soukop, R. W. 22, 25, 397, 430
Soumillion, J. P. 210
Southall, N. T. 29, 30
Sowada, R. 82, 252
Sowinski, A. F. 252
Spada, G. P. 362
Spandau, H. 49, 57
Spange, S. 333, 348, 351
Spanget-Larsen, J. 370, 372
Speed, J. A. 238, 241
Spegt, P. 113
Speiser, B. 164
Spencer, J. N. 107, 121
Sperker, S. 81
Speziale, A. J. 194
Spichiger-Keller, U. E. 333, 478
Spiess, H.-W. 58
Spirić, A. 40, 42, 333, 416, 417, 425, 426, 428
Spirin, Yu. L. 200, 207, 208
Spiteller, G. 471
Spiteller, M. 471
Spitzner, R. 60, 62
Spotswood, T. M. 267
Sprague, R. H. 335
Springer, H. J. 341, 343, 378
Squiller, E. P. 253
Srp, N. E. 479
Staab, H. A. 106, 200, 330

Stace, A. J. 40
Stadnicka, K. 425
Stahl, N. 121
Stahlbush, J. R. 27
Stalder, H. 195
Stangl, H. 191
Starks, C. M. 266, 318
Starnes, S. D. 453, 467
Staudigl, B. 120
Stefaniak, L. 117
Stegmann, H. B. 119, 370
Steigel, A. 108
Steigman, J. 95, 96, 121
Steinberg, M. 155, 201
Steinebrunner, G. 381, 383
Steiner, G. 179, 180, 193
Steiner, T. 15
Steinhausen, R. 200, 201
Steinmetz, R. 281
Steinsträßer, R. 58
Stengle, T. R. 40
Stensland, P. 259
Stepanov, V. 327
Stephenson, L. M. 193
Stepien, B. 104
Stepukhovich, A. D. 227, 230
Sterk, H. 113, 118
Sterner, O. 4
Sternhell, S. 118
Stevens, B. 405
Stevens, I. D. R. 48, 188, 189, 244, 245, 263, 264
Stevenson, C. D. 56
Stevenson, D. P. 133, 351
Stevenson, G. R. 370
Stewart, G. W. 37
Stewart, R. 121
Steybe, F. 334
Stien, M.-L. 67, 129, 131, 133, 337, 354, 389
Stirling, C. J. M. 177
Stiz, S. 40
Stobbe, H. 3
Stöckigt, J. 262
Stockis, A. 36, 376, 383
Stockmayer, W. H. 201
Stoeckenius, W. 298
Stolařová, M. 433, 434, 447
Stolov, A. A. 127, 131
Stone, A. J. 10
Stone, E. W. 370, 372
Storch, D. M. 158
Storesund, H.-J. 280
Storey, B. T. 205, 207, 285
Stork, K. 370, 372
Story, P. R. 307
Stotz, D. S. 265
Stout, G. 370, 372
Stoyanov, S. 113, 114
Stoyanova, T. 113, 114
Stoye, D. 57, 471, 500, 505
Strack, T. 326
Strakhova, N. N. 121, 139, 140
Strandberg, R. 69
Stránský, Z. 73, 496
Straub, H. 336, 344, 345
Straub, P. A. 112, 331
Streck, R. 378
Street, K. W. 358, 430
Strehlow, H. 38–40, 500
Streitwieser, A. 77, 99, 101, 238, 245, 408
Streuli, C. A. 496
Strohbusch, F. 121, 123
Strohmeier, W. 106, 107, 111
Štrop, P. 413
Stubbs, R. J. 42
Stübchen, H. 479
Stukalin, E. B. 468
Sturm, H. J. 191
Su, T. 162
Suarez, C. 388
Sucharda-Sobczyk, A. 122
Sucrow, W. 61
Sudholt, W. 355
Sueishi, Y. 124, 125, 185, 186, 292, 300, 334, 374
Sugden, S. 48, 61
Suginome, H. 360
Sugiura, M. 381, 384, 385
Sugiyama, Y. 124
Suhara, M. 370
Suhr, H. 249
Suleman, N. K. 200, 207, 208, 212
Sumida, Y. 27, 365, 435, 436
Sunamoto, J. 45
Sundermeyer, W. 59, 61
Sundheim, B. R. 59
Suner, G. A. 124
Sunko, D. E. 404
Sunner, J. 147, 157, 158, 247
Sunol, A. K. 70, 73, 492, 494, 505
Sunol, S. G. 70, 73, 492, 494, 505
Suppan, P. 42, 330, 337, 340, 345, 350, 352–354, 357, 358
Surawski, H. 499
Surya Prakash, G. K. 77, 81
Sustmann, R. 188, 189, 191
Sutin, N. 340, 345, 346
Sutton, L. E. 343
Suzuki, K. 287, 336
Svartman, E. L. 203
Svedberg, T. 298
Svoboda, P. 84, 87
Swain, C. G. 60, 61, 201, 238, 239, 244, 316, 398, 404, 408, 453, 462–465
Swain, J. 345
Swain, M. S. 398, 408, 453, 462–465
Swart, E. R. 168
Swieton, G. 180, 181, 188, 193, 312
Swinton, F. L. 5
Swiss, K. A. 309
Symons, E. A. 259
Symons, M. C. R. 6, 36, 40, 84, 238, 242, 351, 363–366, 369, 370, 372, 374–378, 437, 438
Szabadváry, F. 1
Szele, I. 175, 285, 404
Szmant, H. H. 260, 261
Szombathely, M. 363, 369
Szuleiko, J. E. 162
Szwarc, M. 46, 52–55, 100, 155, 201, 262, 263, 305
Szymanski, H. A. 479, 483

t

Taagepera, M. 103, 104
Tabacik, V. 117
Tabner, B. J. 369
Tacconi, G. 197
Tada, E. B. 42, 297
Tadjer, A. 331, 340, 352
Taffer, I. M. 251
Taft, R. W. 15, 18, 19, 24, 82, 88, 99, 101–104, 106, 244, 335, 376, 377, 390, 393–396, 399, 403, 429, 431, 432, 434–437, 442, 443, 447, 450, 453, 456, 457, 459–461, 464, 465
Taguchi, T. 197
Taigel, G. 477
Takács, J. M. 445
Takahashi, F. 253

Takaki, H. 370, 372
Takamuku, S. 281, 282
Takao, N. 381, 384, 385
Takase, H. 373
Takashima, K. 147, 158, 159, 276
Takemura, T. 112
Takenaka, N. E. 136
Takenaka, S. 362
Takeuchi, K. 51, 52, 408
Takka, R. M. 242
Tamamushi, R. 25, 26, 30
Tamm, T. 90, 91, 451–453
Tamura, K. 312, 413, 417, 425, 428
Tan, A. S. C. 478
Tanabe, F. 175
Tanaka, K. 4, 156, 158, 245, 246, 506
Tanaka, N. 23, 25, 26, 30, 398
Tanaka, S. 110
Tanaka, T. 185, 186, 292, 300
Tanaseichuk, B. S. 201
Tanford, C. 27, 42, 225
Tang, R. 184, 185, 449
Tanganov, B. B. 499
Taniguchi, Y. 127, 131
Tanizawa, K. 355, 417, 425
Tanko, J. M. 200, 207, 208, 212, 326, 327
Tapia, O. 38, 219
Tarbell, T. S. 477
Tarján, G. 445
Tarnopolskii, Yu. I. 118
Taticchi, A. 294
Taube, H. 36
Taufer, I. 290, 291
Tawaki, S. 144
Taylor, J. W. 306
Taylor, M. D. 104
Taylor, R. 16
Tchoubar, B. 147, 237, 238, 242, 263
Teague, S. J. 30
Tedder, J. M. 212
Teixeira, J. 6, 15, 18
Te-kang, H. 462
Temnova, I. S. 253
Tenno, S. 103
Teppen, B. J. 115
Terenziani, F. 335
Tero-Kubota, S. 373
Terrier, F. 259
Tesfai, A. 335
Tester, J. W. 327
Tézé, A. 499
Theilacker, W. 477
Thelen, H. 3
Theodoropulos, S. 205, 207
Thiébaut, D. 494
Thiel, W. 55, 137
Thielemans, A. 85
Thieme, R. 131, 136
Thomas, B. E. 131
Thomas, J. H. 50
Thomas, J. K. 46
Thomas, J. M. 4, 506
Thomas, V. K. 84
Thomasson, L. M. 268
Thompson, A. R. 117
Thompson, H. W. 364, 369
Thompson, R. S. 287
Thomson, J. J. 47
Thulstrup, E. W. 59
Tidor, B. 131
Tidwell, T. T. 275
Tietze, E. 269
Tiffon, B. 378
Tiger, R. P. 147, 218, 219, 221, 222
Tikhonova, L. G. 201
Tikunova, I. V. 115
Tilley, J. 360
Timasheff, S. N. 30
Timmers, F. J. 478
Timmons, C. J. 349
Timoneda, J. J. 121
Tipper, C. F. H. 304
Tipson, R. S. 488
Tišler, E. M. 112
Titov, V. V. 58
Toda, F. 4, 506
Töke, L. 468
Tokunaga, Y. 142, 143
Tokura, N. 166, 312, 360, 362
Tolmachev, A. P. 345
Toma, L. 178, 188
Tomaselli, G. A. 188
Tomasi, J. 10, 38, 127, 131, 329, 345, 381
Tomič, M. 404
Tomkins, R. P. T. 30, 33, 57, 73, 147, 238, 254, 255, 496, 500
Tommila, E. 166, 227, 254
Tonnellato, U. 176
Topol, I. A. 101
Topsom, R. D. 103, 393
Tori, K. 385
Tormena, C. F. 129, 131
Torre, G. 69, 360
Torre, M. 188
Tosi, M. P. 10, 57
Tóth, K. 500
Touchstone, J. C. 494–496
Toullec, J. 112
Trachtman, M. 205
Trappe, W. 495
Traynham, J. G. 393
Treanor, R. L. 300, 301
Trémillon, B. 57, 59, 95, 97, 121, 496, 498
Tribble, M. T. 393
Trösken, J. 370
Trotman-Dickenson, A. F. 105
Troughton. E. B. 51, 126, 272
True, N. S. 388
Truhlar, D. G. 38, 95, 131, 149, 329, 345, 349, 360, 381
Tsao, M.-L. 199
Tse, D. S. 8
Tsikanchuk, Ya. 365
Tsuda, Y. 180
Tsvetkov, E. N. 253
Tucker, C. J. 298
Tucker, S. A. 42, 335, 426
Tuji, H. 271, 272
Tuleen, D. L. 205, 206
Tundo, P. 268, 317, 325
Tunnicliff, D. D. 479
Tunuli, M. S. 428
Turner, A. B. 134
Turner, D. W. 21
Turner, G. M. 333
Turovskii, A. A. 205
Turowska-Tyrk, I. 351, 429, 430
Turro, N. J. 292, 296, 300, 301, 307, 349, 350
Tvaroška, I. 128, 132
Tyler, D. R. 365, 374
Tyryshkin, N. I. 125, 168
Tyutyulkov, N. 331, 340, 352

u

Ublacker, G. A. 505
Uchida, M. 312
Uchida, T. 308, 309, 314, 315
Uchytilová, V. 128, 132
Ueji, S. 378, 381, 384, 385
Ueno, K. 428

Uffmann, H. 118
Ugalde, J. M. 182, 194
Ugelstad, J. 177, 264, 268
Ugi, I. 281
Ulbrich, R. 13, 365, 368
Ulfvarson, U. 500
Ullmann, E. F. 370
Ulstrup, J. 27, 30, 334
Ulyanova, O. D. 129
Umemoto, K. 485
Umemura, J. 365
Urbanska, H. 117
Urieta, J. S. 401
Ustinova, A. M. 205

v
Vajs, V. 276
Valeur, B. 352, 353, 357, 358
Vallombroso, T. M. 201
Vallribera, A. 505
Valters, R. E. 118, 119
Vamvakas, S. 500
Van Beijnen, A. J. M. 333
Van der Heijde, H. B. 496
Van der Sluys, W. G. 9
Van der Veeken, B. J. 127, 131
Van Dormael, A. 330
Van Duijnen, P. Th. 349, 360
Van Emster, K. 166
Van Gunsteren, W. F. 162
Van Haverbeke, Y. 113, 114
Van Helmont, J. B. 1
Van Houten, J. 2
Van Koten, G. 320, 321, 488
Van Phuc, N. 333
Van Scoy, R. M. 155, 201, 215, 216
Van Stam, J. 43
Van Zwet, H. 284
Van't Hoff, J. H. 2, 154
Vandeginste, B. G. M. 84
Vandensavel, J.-M. 192
Vandersall, M. 417, 425
Vandewyer, P. H. 337, 448
VanDyke, R. H. 336
VanLare, E. 336
Vanterpool, A. 251
VanZandt, G. 336
Varadarajan, T. K. 40
Varma, R. S. 4, 386, 506
Varshney, R. K. 96
Varughese, P. 244–246
Vary, S. 479
Vasseur, A. 67
Vasyutin, Ya. M. 396, 401, 436, 456
Vaughan, J. 287
Vaughan, W. E. 155
Vaughn, J. W. 82
Vavruch, I. 65
Vdovenko, S. I. 178
Večeřa, M. 84, 87
Vedamuthu, M. 6
Vedejs, E. 194, 195
Veerawat, R. 453, 467
Veith, R. W. 108
Vellaccio, F. 243
Venkataram, U. V. 30
Venkataraman, K. 117, 400
Venkatasubramanian, N. 378
Verbit, L. 303
Verducci, J. 118
Vereshchagin, L. F. 311
Verhoek, F. H. 499
Verhoeven, J. W. 356
Vernon, F. 444, 445
Vichutinskii, A. A. 210
Vidrine, D. W. 405
Vidulich, G. A. 36
Viggiano, A. A. 159–161
Vill, B. 58
Villiger, V. 428
Vincent, E. J. 135
Vincent, M. A. 113–115
Vinogradov, S. N. 121
Vinter, J. G. 129
Viola, H. 117
Viout, P. 263, 265, 268
Virtanen, P. O. I. 244, 247
Visapää, A. 479
Vishnyakov, V. M. 178
Vitha, M. F. 42, 333, 416, 428, 478
Vlasov, V. M. 101, 103
Vodolazkaya, N. A. 333
Vogel, E. 136
Vogel, G. C. 27
Vogel, H. 71, 73
Vogt, M. 320
Vögtle, F. 6, 139–141, 212, 213, 264, 266, 268, 428, 429
Volz, H. 417, 425, 438
Vondenhof, M. 52
Voss, S. 191
Vostrowsky, O. 194, 195
Vottero, L. R. 42, 173, 174
Vymětalová, J. 166
Vyprachticky, D. 413

w
Waack, R. 50
Waddington, T. C. 57, 59
Wade, L. G. 61
Wagenhofer, H. 191
Waghorne, W. E. 30, 255, 256, 365
Wagner, H.-U. 269, 273
Wagner, P. 213
Wagner, P. J. 415
Wagner, R. E. 14
Wähnert, M. 341–344, 378
Wai, C. M. 70, 73, 326, 492
Waits, H. P. 201, 202
Wajer, Th. A. J. W. 370, 372
Wakaumi, W. 11, 67
Wakeham, W. A. 10
Wakisaka, A. 39
Walbrick, J. M. 404
Walden, P. 1, 49, 61, 322
Waldmann, H. 62, 189, 268
Walker, B. J. 128
Walker, R. A. C. 408, 458
Walker, S. M. 373
Wallace, T. J. 253
Wallbillich, G. 192
Wallenfels, K. 393
Waller, F. J. 405
Walling, C. 200–202, 207, 208, 212, 213, 215, 216, 415
Wallis, S. R. 360
Walter, A. 51, 125
Walter, W. 429, 430
Walther, D. 429, 430
Walton, J. C. 212
Walz, G. 345–347
Wamser, C. C. 138
Wanczek, K.-P. 100
Wang, H. 162
Wang, M.-L. 266, 318, 319
Wang, Y. 90, 91, 437, 451–453
Wang, Z. 60
Warshel, A. 38
Wasey, A. 124, 125
Wassermann, A. 156, 188
Wasserscheid, P. 60, 62, 323, 324, 488, 506
Watanabe, M. 227, 229
Watanabe, S. 388
Waterbeemd, H. van de 27
Waters, J. A. 159, 201

Watson, B. A. 201
Watts, D. W. 249, 250, 254, 257–259
Watts, V. S. 388
Wawrzyniewicz, W. 266, 318
Wayland, B. B. 25, 27, 398
Weast, R. C. 475
Weathersby, P. K. 461
Weaver, W. M. 244, 245, 263
Webb, G. A. 117, 378, 381, 378
Webb, H. M. 104
Weber, B. 289
Weber, E. 264, 266, 268
Weber, G. 357
Weber, K. 200, 201, 222, 224, 317
Weber, L. L. 334
Weber, W. P. 266, 318
Wehner, G. 273
Wehrli, R. 196
Wehrly, K. 192
Weigang, O. E. 340, 345, 362
Weingärtner, H. 60
Weinhold, F. 19, 21
Weis, A. L. 117
Weise, A. 82, 252
Weiss, C. 131, 136, 347
Weiß, P. B. 444
Weiss, R. 50, 266
Weiss, R. G. 59, 185, 186, 299–302
Weiss, W. 370, 372
Weissberger, A. 9, 50, 68
Weissermel, K. 253
Weissman, S. I. 55
Weitl, F. L. 403
Weller, U. 502
Wells, P. R. 390
Wells, S. L. 70, 73, 315, 505
Welton, T. 60, 62, 323, 324, 488, 506
Wen, W. Y. 6, 35, 37, 38
Wenclawiak, B. 492
Wendisch, D. 119
Wentworth, P. 243
Werder, R. D. 479
Werncke, W. 334
Werner, H. 80
Wescott, C. R. 143
Wesdemiotes, C. 272
West, W. 367
Westaway, K. C. 172, 173
Westbrook, C. K. 369
Westheimer, F. 225
Wetzler, D. E. 357, 358
Whalley, E. 306, 313, 314
Whiffen, D. H. 5
Whim, B. P. 57, 471
White, F. L. 336
White, J. M. 113–115
White, M. S. 15
White, W. N. 195–197, 415
Whitesides, G. M. 250, 252
Whiting, J. 119
Whitten, D. G. 46, 185, 186, 300, 334
Whittingham, D. J. 168
Wiberg, K. B. 104, 113–115, 127, 128, 131, 147, 149, 218, 219, 230
Wicke, E. 6
Wideqvist, S. 238, 241, 242
Wiedenhoft, A. 177
Wieland, H. 61
Wieland, Th. 61
Wigfield, D. C. 195
Wijayanayake, R. H. 340
Wilcox, C. F. 150
Wild, D. D. 340, 345
Wilen, S. H. 68, 121, 126, 128, 131–133, 288, 360, 362
Wilka, E.-M. 477
Wilkes, J. S. 61, 322
Wilkins, H. 61, 323
Willard, A. K. 291
Willi, A. V. 315, 317
Williams, D. H. 376, 378, 381, 382, 384
Williams, D. J. 406, 407
Williams, G. H. 201
Williams, R. 137, 138
Williams, R. A. 378, 429
Williams, R. F. 298
Williams, R. L. 364, 365, 367, 383
Williams, S. H. 222, 312
Wills-Johnson, G. 349
Wilson, G. J. 64, 65
Wilson, H. 93, 94, 152, 154, 195, 219, 259
Wilson, L. 18, 19
Wilson, L. Y. 453, 466, 467
Wilson, R. A. 61
Wilson, W. W. 261
Wingler, A. 500
Winguth, L. 177
Winkler, H. G. 477
Winkley, M. W. 360
Winnik, M. A. 358, 429, 430
Winstein, S. 48, 53, 54, 62, 165, 172, 215, 226, 231, 232, 238, 239, 244, 245, 263, 264, 402–405, 408, 409
Winter, R. 62
Winther, C. 362
Wipff, G. 33
Wislicenus, W. 3
Witanowski, M. 117, 378
Witt, O. N. 329
Wittig, G. 194
Witzel, M. 505
Wøien, G. 280
Wohar, M. M. 365
Wohlleben, G. 478, 494
Woischnik, E. 337
Wójcik, J. 178
Wojtyk, T. C. 124, 125
Wolbach, W. S. 121
Wold, S. 84, 85, 390, 394, 395, 434, 459, 460
Wolf, F. J. 490
Wolf, K. L. 31
Wolfarth, E. F. 195–197, 415, 479
Wolfbeis, O. S. 352, 357, 358, 417
Wolfe, S. 158, 164
Wolfenden, R. 137, 138
Wolff, H. 166
Wolff, R. 343, 378, 379
Wolfgarth, P. 288
Wondraczek, R. 468
Wong, K. F. 188, 222
Wong, M. W. 104, 113–115
Wong, S. K. 334
Wood, D. C. 82, 252
Wood, P. D. 334
Wood, R. E. 133
Wood, R. H. 253
Wood, W. C. 304
Woodward, A. J. 365
Woodward, R. B. 178, 187, 188, 191, 195, 197
Woolf, J. 323
Woolf, L. I. 163, 165, 169–172, 215, 225, 310
Worley, S. D. 278
Wörz, O. 29, 30, 345
Wozniak, K. 101
Wright, G. F. 477
Wrona, P. K. 351, 417, 425, 426, 429, 430, 433, 454
Wu, F. 60
Würthner, F. 334

Würthwein, E.-U. 136
Wyatt, P. A. H. 498
Wychera, E. 396–398, 462, 465
Wypych, G. 4, 9, 38, 39, 57, 60, 61, 63, 65, 66, 68, 70, 73, 95, 121, 147, 215, 238, 251, 317, 443, 471, 488, 507
Wyss, H. R. 479

x
Xodo, L. E. 129, 132

y
Yalkowski, S. H. 9, 45
Yamada, K. 135
Yamaguchi, I. 376, 378
Yamaguchi, Y. 72
Yamamoto, G. 195, 240
Yamamoto, K. 303
Yamamoto, O. 200, 201
Yamamoto, S. 124, 334, 374
Yamamoto, T. 82, 253
Yamamoto, Y. 39
Yamasaki, K. 108
Yamashita, J. 200, 201
Yamataka, H. 106
Yamazaki, M. 334
Yamdagni, R. 103
Yangjeh, A. H. 296
Yano, T. 428
Yarwood, J. 19
Yasuda, M. 357
Yatchishin, I. I. 456
Yates, K. 99, 278, 279
Yavari, I. 378
Yazdanbachsch, M. R. 177
Yeager, H. L. 500
Yerger, E. A. 121
Yoder, C. H. 14
Yogev, A. 108
Yokoi, H. 23
Yonezawa, T. 370, 372
Yoshida, H. 39
Yoshida, M. 286
Yoshimura, T. 253, 365
Yoshimura, Y. 312
Young, L. B. 100, 158
Young, R. H. 192
Yukawa, Y. 188
Yun, K.-Y. 136
Yutronic, N. 40
Yvernault, G. 22, 397
Yvernault, T. 22, 397

z
Zachariasse, K. A. 297, 333
Zackrisson, M. 22
Zahradnik, R. 19–21
Zaikov, G. E. 147, 210, 215, 219
Zaitsev, B. E. 117, 118
Zalewski, R. I. 84, 87
Zamchuk, Z. S. 178
Zana, R. 44
Zandstra, P. J. 370, 372
Zanka, A. 167, 312, 313
Zaschke, H. 58
Zaugg, H. E. 264, 265
Závada, J. 286
Zaworotko, M. J. 61
Zeegers-Huyskens, T. 10, 122
Zefirov, N. S. 126, 128, 131, 132, 177
Zehner, P. 71, 73
Zeidler, M. D. 5, 6
Zeigler, J. M. 113–115
Zelinskii, V. V. 429, 430
Zewail, A. H. 150
Zewdu, M. 360
Zhang, B.-L. 176
Zhang, S. M. 20
Zhang, X. 254
Zhao, Y. 20
Zhmyreva, I. A. 429, 430
Zhong, M. 272
Zhong-yuan, Z. 462
Zhu, T. 38, 320, 321
Zhukovskii, V. Ya. 365, 366, 436
Ziegler, K. 6, 176, 200, 201, 222, 224, 317
Zielkowska, U. 417, 425, 426, 454
Ziessow, D. 376, 377
Žigman, V. 129, 131, 132
Zilian, U. 68
Zimmermann, G. 188, 410
Zimmermann, H. 17, 18, 30
Zimmermann, H. W. 334
Zimmermann, R. G. 299, 300, 302
Zincke, Th. 3
Zingaro, R. A. 57
Zink, J. I. 376, 377
Zippel, M. 182
Zipse, H. 243
Zollinger, H. 173, 175, 285
Zook, H. D. 265
Zosel, K. 492
Zubkov, V. A. 106
Zubrick, J. W. 268
Zundel, G. 17, 30, 121, 122
Zutaut, S. E. 403
Zwan, G. van der 352, 355, 356