



Blast Injuries: Essential Facts



Key Concepts

- Bombs and explosions can cause unique patterns of injury seldom seen outside combat
- Expect half of all initial casualties to seek medical care over a one-hour period
- Most severely injured arrive after the less injured, who bypass EMS triage and go directly to the closest hospitals
- Predominant injuries involve multiple penetrating injuries and blunt trauma
- Explosions in confined spaces (buildings, large vehicles, mines) and/or structural collapse associated with greater morbidity and mortality
- Primary blast injuries in survivors are predominantly seen in confined space explosions
- Follow standard protocols for triage, trauma resuscitation, treatment, and transfer
- Repeatedly examine and assess patients exposed to a blast

Blast Injuries

- Primary: Injury from over-pressurization force (blast wave) impacting the body surface
 - TM rupture, pulmonary damage and air embolization, hollow viscus rupture
- Secondary: Injury from projectiles (bomb fragments, flying debris)
 - Penetrating trauma, fragmentation injuries, blunt trauma
- Tertiary: Injuries from displacement of victim by the blast wind, and structural collapse
 - Crush injuries, blunt/penetrating trauma, fractures and traumatic amputations
- Quaternary: All other injuries from the blast
 - Burns, asphyxia, toxic exposures, etc.

Primary Blast Injury

- Lung Injury
 - Signs usually present at time of initial evaluation, but may be delayed up to 48 hrs
 - Reported to be more common in patients with skull fractures, >10% BSA burns, and penetrating injury to the head or torso
 - Varies from scattered petechiae to confluent hemorrhages
 - Suspect in anyone with dyspnea, cough, hemoptysis, or chest pain following blast
 - CXR: “butterfly” pattern
 - High flow O₂ sufficient to prevent hypoxemia via NRB mask, CPAP, or ET tube
 - Fluid management similar to pulmonary contusion; ensure tissue perfusion but avoid volume overload
 - Endotracheal intubation for massive hemoptysis, impending airway compromise or respiratory failure
 - Consider selective bronchial intubation for significant air leaks or massive hemoptysis
 - Positive pressure may risk alveolar rupture or air embolism

- Prompt decompression for clinical evidence of pneumothorax or hemothorax
- Consider prophylactic chest tube before general anesthesia or air transport
- Air embolism can present as stroke, MI, acute abdomen, blindness, deafness, spinal cord injury, claudication
 - High flow O₂; prone, semi-left lateral, or left lateral position
 - Consider transfer for hyperbaric O₂ therapy
- **Abdominal Injury**
 - Gas-filled structures most vulnerable (esp. colon)
 - Bowel perforation, hemorrhage (small petechiae to large hematomas), mesenteric shear injuries, solid organ lacerations, and testicular rupture
 - Suspect in anyone with abdominal pain, nausea, vomiting, hematemesis, rectal pain, tenesmus, testicular pain, unexplained hypovolemia
 - Clinical signs can be initially silent until acute abdomen or sepsis is advanced
- **Ear Injury**
 - Tympanic membrane most common primary blast injury
 - Signs of ear injury usually evident on presentation (hearing loss, tinnitus, otalgia, vertigo, bleeding from external canal, otorrhea)
 - Isolated TM rupture not a marker for morbidity

Other Injury

- Traumatic amputation of any limb is a marker for multi-system injuries
- Concussions are common and easily overlooked
- Consider delayed primary closure for grossly contaminated wounds, and assess tetanus immunization status
- Compartment syndrome, rhabdomyolysis, and acute renal failure are associated with structural collapse, prolonged extrication, severe burns, and some poisonings
- Consider possibility of exposure to inhaled toxins (CO, CN, MetHgb) in both industrial and terrorist explosions
- Significant percentage of survivors will have serious eye injuries

Disposition

- No definitive guidelines for observation, admission, or discharge
- Discharge decisions will also depend upon associated injuries
- Patients with no complaints suggestive of primary blast injury, normal CXR, and not hypoxic may be considered for discharge after 6-8 hours observation
- Admit 2nd and 3rd trimester pregnancies for monitoring
- Close follow-up of wounds, head injury, eye, ear, and stress-related complaints
- Patients with ear injury may have tinnitus or deafness; communications and instructions may need to be written

Additional information: www.bt.cdc.gov/masstrauma/index.asp