

**MATH 1401 SPRING 2000 CHEAT SHEET
FINAL**

JAN MANDEL

1. Important formulas from algebra. $\sin(a+b) = \sin(a)\cos(b) + \cos(a)\sin(b)$, $\sin^2 x + \cos^2 x = 1$, $a^{b+c} = a^b a^c$, $a^{m/n} = \sqrt[n]{a^m}$, $a^b = e^{(\log a)b}$. Solution of $ax^2 + bx + c = 0$ is $x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

2. Limits and continuity. $\lim_{x \rightarrow c} f(x) = f(c) \iff f$ is continuous at c
 $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$, $\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = 0$, $\lim_{x \rightarrow 0} (1+x)^{1/x} = e$
 $\lim_{x \rightarrow c} f(x) = L \iff \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = L$
 Intermediate value theorem: If f is continuous on $[a, b]$ and k is between $f(a)$ and $f(b)$, then there exists $c \in [a, b]$ such that $f(c) = k$.
 Infinite limits: The formulas for the limit of sum, product, and quotient apply unless they lead to undefined expressions of the form $\infty - \infty$, $\infty \cdot 0$, $L/0$, ∞/∞ .
 If $\lim_{x \rightarrow c} f(x) \neq 0$ and $\lim_{x \rightarrow c} g(x) = 0$, with $g(x) \neq 0$ on a neighborhood of c , then the graph of f/g has vertical asymptote $x = c$.

3. Differentiation. The equation of the line passing through (x_0, y_0) with slope s is $y - y_0 = s(x - x_0)$. The equation of the tangent to the graph of f at (x_0, y_0) , $y_0 = f(x_0)$, is $y - y_0 = f'(x_0)(x - x_0)$.
 $f'(c) = \lim_{x \rightarrow c} (f(x) - f(c))/(x - c)$. If $f'(c)$ exists, f is continuous at c .
 $(x^n)' = nx^{n-1}$, $(\sin x)' = \cos x$, $(\cos x)' = -\sin x$, $(\ln x)' = 1/x$, $(e^x)' = e^x$
 $\sin' x = \cos x$, $\cos' x = -\sin x$, $(\arctan x)' = 1/(1+x^2)$, $\arcsin' x = \frac{1}{\sqrt{1-x^2}}$, $\operatorname{arcsec}' x = \frac{1}{|x|\sqrt{1-x^2}}$, $(uv)' = u'v + uv'$, $(u/v)' = (u'v - uv')/v^2$, $f(g(x))' = f'(g(x))g'(x)$
 If $g = f^{-1}$ and $y = g(x)$, $f'(y) \neq 0$, then $g'(x) = 1/f'(y)$.

4. Applications and extrema. If f is continuous on $[a, b]$, then f attains maximum and minimum on $[a, b]$. f can attain extremum on $[a, b]$ only at endpoints or critical numbers (where f' does not exist or $f' = 0$). f can attain relative extremum in (a, b) only at a critical number.
 Mean value theorem: If f is continuous on $[a, b]$ and differentiable on (a, b) then there exists $c \in (a, b)$ such that $f'(c) = (f(b) - f(a))/(b - a)$. (The case when $f(a) = f(b)$ is Rolle's theorem.)
 If $f' > 0$ in (a, b) and f is continuous on $[a, b]$, then f is increasing on $[a, b]$.
 If f is continuous at c , $f'(x) < 0$ for $x < c$ and $f'(x) > 0$ for $x > c$, then f has relative minimum $(c, f(c))$. (Or, relative minimum $f(c)$ at $x = c$.)
 If f' is increasing in interval I , then f is concave upward in I .
 If $f'' > 0$ in (a, b) , then f is concave upward in (a, b) .
 If $f'(c) = 0$ and $f''(c) > 0$, then f has relative minimum at c .

5. Hyperbolic functions. $\sinh x = \frac{e^x - e^{-x}}{2}$, $\cosh x = \frac{e^x + e^{-x}}{2}$, $\cosh^2 x - \sinh^2 x = 1$, $\cosh' x = \sinh x$, $(\tanh^{-1})' = 1/(1-x^2)$

6. Integration. $\int f(x) dx = F(x) + C$, $F' = f$.
 $\int x^n dx = x^{n+1}/(n+1) + C$, $n \neq -1$, $\int f(g(x))g'(x) dx = \int f(u) du$, $u = g(x)$
 $\int \frac{1}{\sqrt{a^2-x^2}} = \arcsin \frac{x}{a} + C$, $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \arctan \frac{x}{a} + C$
 $\int \frac{1}{\sqrt{a^2+x^2}} = \sinh^{-1} \frac{x}{a} + C$, $\int \frac{1}{a^2-x^2} dx = \frac{1}{a} \tanh^{-1} \frac{x}{a} + C$
 $\int_a^b f(x) dx = F(b) - F(a)$, $F' = f$.
 $(d/dx) \int_a^x f(t) dt = f(x)$