TABLE 5.—SUMMARY OF DICHLORVOS RISKS AND BENEFITS—Continued

Uses	Non-Dietary Margin of Exposure: Cholinesterase Inhibition	Dietary Upper Bound Cancer Risk	Benefits	Proposed Action
Reentry				
After 24 hours	21 (no respirator)			Reentry Restrictions
After 48 hours	289 (no respirator)			Limited reentry during first 48 hours following treatment. No entry within first 4 hours; limited reentry (one hour per 24 hours) for handling activities only.
Greenhouse				
Applicator	Majority of MOEs less than 100 and 30% less than 50	1.6×10^{-7} (lettuce) 2.6×10^{-8} (cucumbers) 1.4×10^{-8} (tomatoes) 8.8×10^{-10} (radishes)	Not expected to be significant	**Allowable Application Methods** -Automatic foggers -Thermal foggers through a port **Application Methods Not Allowed (Cancel)** -Hand application or any method in which the applicator remains inside the greenhouse during application.
Reentry				
After 24 hours	21 (no respirator)			Reentry Restrictions
After 48 hours	289 (no respirator)			Limited reentry during first 48 hours following treatment. No entry within first 4 hours; limited reentry (one hour per 24 hours) for handling activities only.
Domestic food/nonfood animals (non-poultry)				
Applicator	6.1	6.2×10^{-7} (milk)	Probable regional impacts	Cancel all hand application methods to both food and nonfood animals